

Synemin and Vimentin are Components of Intermediate Filaments in Avian Erythrocytes

BRUCE L. GRANGER, ELIZABETH A. REPASKY, and ELIAS LAZARIDES
Division of Biology, California Institute of Technology, Pasadena, California 91125

ABSTRACT Synemin, a high-molecular-weight protein associated with intermediate filaments in muscle, and vimentin, an intermediate-filament subunit found in many different cell types, have been identified by immunologic and electrophoretic criteria as components of intermediate filaments in mature avian erythrocytes. Desmin, the predominant subunit of intermediate filaments in muscle, has not been detected in these cells. Two-dimensional immunofluorescence of proteolytic fragments of synemin and vimentin demonstrates that the erythrocyte proteins are highly homologous, if not identical, to their muscle counterparts. Double immunofluorescence reveals that erythrocyte synemin and vimentin co-localize in a cytoplasmic network of sinuous filaments that extends from the nucleus to the plasma membrane and resists aggregation by Colcemid. Erythrocytes that are attached to glass cover slips can be sonicated to remove nuclei and nonadherent regions of the plasma membrane; this leaves elliptical patches of adherent membrane that retain mats of vimentin- and synemin-containing intermediate filaments, as seen by immunofluorescence and rotary shadowing. Similarly, mechanical enucleation of erythrocyte ghosts in suspension allows isolation of plasma membranes that retain a significant fraction of the synemin and vimentin, as assayed by electrophoresis, and intermediate filaments, as seen in thin sections. Both synemin and vimentin remain insoluble, along with spectrin and actin, in solutions containing nonionic detergent and high salt. However, brief exposure of isolated membranes to distilled water releases the synemin and vimentin together in nearly pure form, before the release of significant amounts of spectrin and actin. These data suggest that avian erythrocyte intermediate filaments are somehow anchored to the plasma membrane; erythrocytes may thus provide a simple system for the study of intermediate filaments and their mode of interaction with membranes. In addition, these data, in conjunction with previous data from muscle, indicate that synemin is capable of associating with either desmin or vimentin and may thus perform a special role in the structure or function of intermediate filaments in erythrocytes as well as in muscle.

Mature avian erythrocytes are nucleated, biconvex, elliptical discs that contain relatively few cytoplasmic organelles. Removal of hemoglobin from these cells by hypotonic lysis (15) reveals an equatorial bundle of microtubules known as the marginal band (5), a submembranous spectrin-actin shell (9), as well as a residual network of cytoplasmic filaments that surrounds the mitochondria and extends from the nucleus to the plasma membrane (27). This latter network of filaments is probably a component of the "trans marginal band material" noted in many nonmammalian vertebrate erythrocytes (13). These filaments appear to be of the type known as intermediate filaments (33, 39), due to their characteristic ultrastructural morphology and insolubility in nonionic detergents (59, 63).

The close association of these filaments with the plasma membrane and nucleus, as shown by electron microscopy, suggests that they might function to maintain the shape of the cell or position the nucleus within the cell (13, 27, 59, 63).

We have examined these filaments biochemically, immunologically, and ultrastructurally and have determined that they are composed predominantly of vimentin, an intermediate filament subunit common to many different cell types (18, 39). The other major component of these filaments is synemin, a high molecular weight protein originally isolated from avian smooth muscle in association with desmin and subsequently shown to co-localize with desmin and vimentin in skeletal muscle (25). Double immunofluorescence shows that erythro-

cyte synemin and vimentin also coexist in a network of cytoplasmic filaments. The electrophoretic and immunologic criteria used to identify vimentin and synemin in these cells fail to detect desmin, the major intermediate filament subunit of muscle (40, 53) that is also found in some nonmuscle cells (19, 55, 58).

Various cell fractionation procedures based on differential centrifugation have indicated that synemin and vimentin sediment with both nuclear and membrane fractions. In this study we have concentrated on those intermediate filaments that remain associated with the erythrocyte plasma membrane after mechanical enucleation of the cells. These filaments resist dissociation from the membranes by sonication and treatment with high salt and nonionic detergent, suggesting that they are in some way anchored to the membrane cytoskeleton, perhaps to the spectrin-actin network. However, we have found that the filaments can be selectively removed from the membranes by treatment with low ionic strength solutions and that the predominant proteins thus released are vimentin and synemin.

The evidence presented here that synemin associates with vimentin in erythrocytes, in conjunction with the evidence presented previously that synemin associates with desmin in smooth muscle and desmin and vimentin in skeletal muscle (25), suggests that synemin may be capable of associating with different intermediate filament subunits in different cell types. Because synemin is not detectable in all cell types, it may play a special role in the structure or function of certain types of intermediate filaments or filaments in certain cell types.

MATERIALS AND METHODS

Preparation of Erythrocyte Membranes

White leghorn chickens were given intravenous injections of 3–4 mg (500–700 USP units) of heparin (17) and then ~50 mg of sodium pentobarbital. Blood was collected from the neck vein into a solution of 0.01% heparin, 5 mM HEPES (N-2-hydroxyethylpiperazine-N'-2-ethanesulfonic acid), 155 mM choline chloride (pH 7.1 at room temperature). Phosphate-buffered saline was simultaneously injected into a wing vein, and the perfusate was also collected until it became relatively clear. Alternatively, blood for some experiments was drawn from wing veins or neck veins of uninjected chickens and collected in 1–2 vol of the above heparin-containing solution. Blood cells were pelleted by centrifugation for 5 min at 1,000 g, and the top white layer of cells (buffy coat) and supernatant were removed by aspiration. The erythrocyte pellet, exclusive of a dark-red layer adhering to the bottom of the centrifuge tube, was resuspended in 155 mM choline chloride, 5 mM HEPES (pH 7.1 at room temperature) and recentrifuged. Again, the supernatant, buffy coat, and dark-red layer were discarded. This cycle was repeated for a total of 4–8 washes, and was performed either at room temperature or at 4°C.

The final pellet of erythrocytes was rapidly resuspended in at least 10 vol of ice-cold hypotonic lysis buffer (Buffer H) [5 mM Tris-Cl (pH 7.5), 5 mM Na₂S₂O₈, 5 mM MgCl₂, 1 mM EGTA (ethyleneglycol-bis(β-amino-ethyl ether) N,N'-tetraacetic acid), 1 mM dithioerythritol (DTE) or dithiothreitol (DTT), 0.5 mM phenylmethyl sulfonyl fluoride (PMSF)]. MgCl₂ was included to keep the nuclei intact, EGTA and PMSF were included as protease inhibitors, and DTE (or DTT) was found to increase both the yield and size of the plasma membrane fragments (see below). The resulting nucleated erythrocyte ghosts were pelleted by centrifugation for 5 min at approximately 10,000 g, then resuspended in at least 10 vol of the same solution. This cycle was repeated for a total of three or four washes in Buffer H.

For the preparation of plasma membranes, the final pellet was resuspended in 2–4 vol of Buffer H, loaded into a syringe, and forced rapidly through a 23 gauge hypodermic needle bent into the shape of a Z (with two 30° angles). Centrifugation for 10–20 min at 1,000 g in a swinging bucket rotor resulted in three layers: a firm, white pellet of free nuclei on the bottom, a loose, pink layer of undisrupted cells in the middle, and a supernatant containing soluble proteins and membrane fragments. The middle layer was resuspended in Buffer H and again forced through the needle and centrifuged to give the three layers. This was usually repeated three times; the yield of membrane fragments increased with each cycle. The supernatants from each centrifugation were combined and recentrifuged for 30 min at 100–200 g to remove any remaining nuclei. The supernatant of this low

speed centrifugation was respun for 10 min at 20,000 g to pellet the membranes which were subsequently resuspended in 10–20 vol of Buffer H and stored on ice. Contaminating nuclei visible by phase contrast microscopy could be removed by another low speed centrifugation. After storage for a month on ice, little change was seen in the electrophoretic protein profile as judged by one dimensional SDS-polyacrylamide gel electrophoresis (SDS-PAGE) (see below).

Selective Extraction of Synemin and Vimentin From Erythrocyte Membranes

The suspension of erythrocyte membranes was mixed with 10–30 vol of 2 mM EDTA 10 mM Tris (pH adjusted with HCl to 7.4 at 0°C); after 10 min on ice the membranes were pelleted by centrifugation at 20,000 g for 10 min. The pellet was resuspended (with a Pasteur pipette) in distilled water at 0°C and centrifuged as above. The supernatant was then lyophilized directly for electrophoretic analysis, or first recentrifuged for 1–5 h in a Beckman SW 50.1 swinging bucket rotor at 50,000 RPM (Beckman Instruments, Inc., Fullerton, Calif.).

Polyacrylamide Gel Electrophoresis

One-dimensional SDS-PAGE was based on the discontinuous, Tris-glycine system of Laemmli (37), as modified and described previously (29). Separating gels were 11 × 14 × 0.16 cm and contained 12.5% acrylamide and 0.11% N,N'-methylene-bisacrylamide. Samples were solubilized with 1% SDS, 125 mM Tris-Cl (pH 6.8), 10% glycerol, 1% 2-mercaptoethanol, 1 mM EDTA, 0.004% bromophenol blue ("1% SDS sample buffer"), and immediately placed in a boiling water bath for ~1 min.

Two-dimensional isoelectric focusing (IEF)/SDS-PAGE was performed according to the method of O'Farrell (48), as modified and described previously (29), except that nonidet P-40 (NP-40) was omitted from all gels and samples (this enhanced the resolution of the isoelectric variants of several proteins of interest in this system). Second-dimension SDS slab gels were as described above.

Phosphorylation

Incorporation of [³²P]-phosphate into erythrocyte proteins was performed essentially as described by Beam et al. (4) and Alper et al. (2). Blood was collected from the wing vein of an adult hen turkey in cold heparinized choline chloride buffer and washed as described above. Erythrocytes were then washed once at room temperature with 40 vol of 157.5 mM NaCl, 2.5 mM KCl, 11.1 mM D-glucose, 10 mM HEPES (brought to pH 7.65 at room temperature with NaOH). 1 ml of packed cells was resuspended in 9 ml of this solution; 2 mCi of ³²P-phosphoric acid (New England Nuclear, Boston, MA; 100 μl of carrier-free in 20 mM HCl) were then added and the suspension was incubated in an orbital shaker bath at 39°C. After 3.5 h the suspension was divided in half; to one-half was added 50 μl of 0.1 mM DL-isoproterenol-HCl (Sigma Chemical Co., St. Louis, MO) in the above solution (final concentration 1 μM). Both aliquots were incubated for another 20 min at 39°C and then processed in parallel for gel electrophoresis. The erythrocytes were spun down and washed once with 40 vol of the above solution at 39°C, then lysed with 80 vol of 10 mM Tris-Cl (pH 8.0), 5 mM MgCl₂, 1 mM EGTA, 1 mM o-phenanthroline, 0.5 mM PMSF, 1 mM NaF at 0°C. The nucleated ghosts were spun down and washed once with this solution, then disrupted by one passage through a bent hypodermic needle as described above. Intact cells and nuclei were removed by low-speed centrifugation, and free membrane fragments were collected by high-speed centrifugation. The membrane pellet was boiled in 1% SDS sample buffer and analyzed by SDS-PAGE. After staining and destaining, the gel was dried on filter paper and exposed to Kodak X-Omat R XR5 film at room temperature for 10 d without an intensifying screen.

In a separate experiment, erythrocytes were collected, washed and labeled as described above, except that the labeling period was 20 h and no isoproterenol was added. Cells were lysed with 5 mM sodium phosphate, 5 mM MgCl₂, 1 mM EGTA, 1 mM PMSF (pH 7.4 at 4°C) and disrupted as described above. The membrane fraction was dissolved in 10 M urea containing 1% 2-mercaptoethanol, analyzed by IEF/SDS-PAGE, and autoradiographed.

Immunoautoradiography

Immunoautoradiography was performed as described previously (25). Gels were incubated with antisera diluted 1,000-fold, followed by radioiodinated protein A (4–7 μCi ¹²⁵I/μg protein A; each gel was incubated with 20–30 μCi in 100 ml of solution). The dried gels were exposed to x-ray film for the following times with (+) or without (–) an intensifying screen: Fig. 3 b, 27 h (–); Fig. 3 c, 15 h (+); Fig. 4 b, 36 h (–); Fig. 4 c, 55 h (+); Fig. 5 b, 405 h (–); Fig. 5 d, 42 h (+).

Samples for electrophoresis were prepared as follows. Fig. 3: The white cell layer (buffy coat) was collected from the first centrifugation of heparinized chicken blood and recentrifuged twice in HEPES-buffered saline containing 1 mM EDTA to give an erythrocyte-free preparation. The cells were mixed with 40 vol of ethanol at 0°C, pelleted, and boiled in 1% SDS sample buffer. Whole adult chicken-gizzard smooth-muscle tissue was frozen and pulverized in liquid nitrogen, thawed in ethanol, pelleted, and boiled in 1% SDS sample buffer. Turkey erythrocyte membranes were extracted at 0°C for 20 min with 10 mM Tris, 1 mM EDTA, 1 M NaCl, 1% 2-mercaptoethanol, 1% Triton X-100, and the residue was pelleted and boiled in 1% SDS sample buffer. Chicken erythrocyte membranes were extracted at 0°C for 70 min with 10 mM Tris, 2 mM EDTA, 0.5 M KCl, 1% Triton X-100 (pH 7.4), and the residue was pelleted and boiled in 1% SDS sample buffer. Another aliquot of chicken erythrocyte membranes was washed at 0°C with 10 mM Tris-2 mM EDTA (pH 7.4) and then treated for 4 h at 0°C with 10 mM Tris-1 mM EGTA (pH 7.4); the extract was dialyzed against water, lyophilized, and boiled in 1% SDS sample buffer.

Fig. 4: Chicken erythrocyte membranes were treated with 60 vol of ethanol to permeabilize the vesicles, pelleted, and dissolved at room temperature in a saturated urea solution containing 1% 2-mercaptoethanol.

Fig. 5: Chicken erythrocytes were extracted twice at 0°C with 1% Triton X-100, 150 mM NaCl, 2 mM EGTA, 5 mM MgCl₂, 1 mM sodium tetrathionate, 1 mM *ε*-amino-*n*-caproic acid, 1 mM *o*-phenanthroline, 1 mM PMSF, 10 mM Tris-Cl (pH 7.2) and rinsed twice with this solution without the Triton X-100. The resulting cytoskeletons were then extracted with this latter solution containing 6 M urea for 1.5 h at 0°C. This extract was dialyzed against water, and the resulting precipitate was collected and dissolved in a saturated urea solution containing 0.5% 2-mercaptoethanol. Whole adult chicken-gizzard smooth-muscle tissue was frozen and pulverized in liquid nitrogen, thawed in ethanol, pelleted, boiled for 30 s in 10 μl of 1% SDS, and then dissolved in 90 μl of saturated urea containing 1% 2-mercaptoethanol.

Immunofluorescence

Glass cover slips were pretreated with Alcian Blue to promote erythrocyte adhesion (54). Cover slips were cleaned, simmered for 5 min in 0.1% Alcian Blue 8GX (Sigma Chemical Co., St. Louis, Mo.), rinsed with distilled water, and air dried. Washed erythrocytes in the choline chloride/HEPES buffer were allowed to settle on the cover slips for 5–10 min at room temperature; nonadherent cells were removed by rinsing with the same solution.

For the sonication experiments, cover slips with attached erythrocytes were hypotonically lysed at room temperature in Buffer H and then placed in Buffer F [130 mM KCl, 5 mM NaCl, 1 mM NaN₃, 5 mM MgCl₂, 1 mM EGTA, 20 mM potassium phosphate (pH 7.5)]. Cover slips were laid face-up in a beaker of buffer F and sonicated for 20 s at 20 watts using a Braunsonic 1510 sonicator (B. Braun Instruments, San Francisco, Calif.) with a 4-mm titanium probe tip positioned 3–4 cm above the cells. A glass rod was used to hold the cover slips in position on the bottom of the beaker during sonication. Incubations with antisera and subsequent washes were all performed at room temperature in buffer TM (0.5% Triton X-100, 130 mM NaCl, 5 mM KCl, 5 mM NaN₃, 5 mM MgCl₂, 1 mM EGTA, 10 mM Tris-Cl (pH 7.5)).

Alternatively, sonication was performed in Buffer H and antibody incubations and washes in Buffer F without the MgCl₂ and EGTA (Fig. 8*a,b,c,h*).

To investigate the Colcemid sensitivity of the filaments, turkey erythrocytes attached to cover slips were incubated for 16 h at 37°C in growth medium (Eagle's minimal essential medium, nonessential amino acids, 15% horse serum, 5% chick embryo extract, 0.01% streptomycin and 100 U penicillin/ml) containing 5 μM Colcemid (demecolcine; Calbiochem-Behring Corp., La Jolla, Calif.). Control cells were treated identically, except that they were not exposed to Colcemid. Cover slips were placed in Buffer TM containing 0.5 mM PMSF for 1 min at room temperature to make cytoskeletons, then fixed for 10 min at room temperature in Buffer F containing 2% formaldehyde. Subsequent incubations and washes were done in Buffer TM containing a 0.5 mM PMSF. Similar results were obtained with unfixed cells, with cells incubated in suspension rather than attached to cover slips, and with cells incubated with 100 μM Colcemid.

Rabbit anti-vimentin was prepared using antigen from embryonic chicken skeletal muscle cytoskeletons (24). Rabbit anti-desmin was prepared using desmin present in a low-salt extract of chicken-gizzard smooth muscle (25). Rabbit anti-synemin was prepared using antigen obtained from chicken-gizzard intermediate-filament proteins that had been solubilized and precipitated three times in acetic acid (25). All antigens were ultimately purified by preparative SDS-PAGE before injection.

Conjugation of anti-vimentin with rhodamine B was performed as described (25). Double immunofluorescence was performed by the indirect/direct method (32). Fluorescein-conjugated goat anti-rabbit IgG was purchased from Miles-Yeda, Ltd. (Rehovot, Israel) and diluted 150-fold for use. Primary antisera were partially purified by precipitation with ammonium sulfate at 50% saturation and used at ~1/30 serum concentration.

Cover slips were mounted in 90% glycerol in Tris-buffered saline and photographed with Kodak Tri-X film using a Leitz phase/epifluorescence microscope and filter modules K and N2. The patches of plasma membrane in Fig. 8*s* are visible by phase-contrast microscopy because an air bubble was present under that portion of the cover slip.

Electron Microscopy

For thin sectioning, chicken erythrocyte membranes were pelleted and fixed with 1% glutaraldehyde, 5 mM MgCl₂, 10 mM sodium phosphate (pH 7.5), postfixed in 1% OsO₄, 0.1 M sodium cacodylate (pH 7.4), stained at 60°C with 1% uranyl acetate, 0.1 M sodium maleate (pH 5.15), dehydrated in ethanol and propylene oxide, and embedded in a 9/16/1 mixture of 1,2,7,8-diepoxyoctane (Aldrich; Milwaukee, Wis.)/nonenyl succinic anhydride (ICN K/K Laboratories, Inc., Plainview, N. Y.)/DMP-30. The resin was cured for 2 d at 60°C and sectioned with glass knives on a Reichert OmU2 ultramicrotome. Sections were stained for 3 min with 0.2% lead citrate, examined with a Philips EM201 at 80 kV, and photographed on 35-mm film.

Shadowed replicas were made as follows: Chicken erythrocytes adhering to Alcian Blue-coated cover slips were lysed and sonicated as described under Immunofluorescence. The cover slips were immersed in buffered 1% glutaraldehyde followed by 1% OsO₄, dehydrated with ethanol, dried in a carbon dioxide critical-point drier, rotary shadowed with platinum/palladium (80/20) at an angle of 6°, and then carbon coated. Replicas were separated from cover slips with 5% HF, washed with water, mounted on 300-mesh copper grids, and viewed by transmission electron microscopy.

RESULTS

Membrane Fractionation and Ultrastructure

The study of minor protein components of avian erythrocytes is hampered by the relative abundance of hemoglobin and chromatin in these cells. Removal of these two components makes biochemical, immunological and ultrastructural characterization of the remaining structures easier. Hypotonic lysis (to remove hemoglobin) and subsequent mechanical enucleation (to remove chromatin) gives a preparation of membranes that can be studied in a manner analogous to the study of the simpler mammalian erythrocyte ghosts. To this end, chicken and turkey erythrocytes were isolated from fresh blood by differential centrifugation and lysed in a low osmolarity buffer containing magnesium ions to keep the nuclei intact (27). These nucleated ghosts were then disrupted by passage through a bent hypodermic needle, and a membrane fraction was separated from the nuclei and unbroken cells by differential centrifugation. This membrane fraction is the main object of this study and will hereafter be referred to as "erythrocyte membranes."

Representative thin sections of the erythrocyte membranes are shown in Fig. 1. The preparation is composed primarily of plasma membranes, both complete and in pieces (compare with sections of whole cells in references 4, 5, 63, 66). Close examination reveals that in a given thin section many of the membranes have filaments associated with them. These filaments are ~9 nm in diameter and are therefore classified as intermediate filaments. They are present on the cytoplasmic side of the plasma membrane fragments and often appear to be in close apposition to the protein network (analogous to the spectrin network in mammalian erythrocytes) just inside the lipid bilayer. This is especially apparent in grazing sections of the membrane in these relatively thick sections. The filaments are usually curved and randomly distributed and do not exhibit any obvious association with specific cell structures.

Thin sections of the free nuclei (not shown) also reveal intermediate filaments associated with these structures, as previously shown by Woodcock (63). It has not been determined what proportion of the filaments remains with the membranes and what proportion with the nuclei, due to the difficulty of




FIGURE 1 Thin sections of chicken erythrocyte membranes. (a) Medium-power view of pelleted membranes showing that filaments can be seen associated with many, but not all, membranes in a given section ($\times 11,000$). (b, c) High-power views. Note cytoplasmic filaments and frequent close associations between filaments and membranes (b: $\times 38,000$; c: $\times 47,000$). Bars, 500 nm.

enucleating the ghosts quantitatively and completely separating the resulting fractions. Such a quantitation is also complicated by the fact that the proportion of the filaments associated with the nucleus before cell disruption that remains associated with the nuclei after fractionation is probably a function of the severity of the disruption procedure and the extent of loss of the outer nuclear membrane. However, based on various biochemical and ultrastructural data (see below), it appears that on the average less than half of the cell's intermediate filaments end up in the membrane fraction.

It is apparent from these thin sections that the erythrocyte membrane fraction contains low levels of contamination by fragments of structures other than the plasma membrane. Even though this enucleation procedure results in free nuclei that appear to be intact by phase-contrast microscopy, the relatively fragile outer nuclear membrane may become partially fragmented and fractionate with the plasma membranes (27, 65). Fragments of mitochondrial membranes may also be present in this fraction. However, because our studies were concerned primarily with intermediate filaments rather than specific membrane proteins, further purification of the membrane fraction was not deemed necessary for subsequent biochemical studies. The purpose of the fractionation was to remove chromatin that would have physically interfered with the membrane extraction experiments, and this was accomplished. Negligible amounts of histone could be seen when the membrane fraction was analyzed by SDS-PAGE, and nuclear membrane lamins (21, 52) could not be detected by IEF/SDS-PAGE, showing that the level of contaminating material was low.

Electrophoretic Analysis of Membrane Fraction

Analysis of the protein composition of avian erythrocyte membranes was performed with regard to the voluminous work on mammalian erythrocyte ghosts. Similarities between the two systems include two major high-molecular-weight proteins in avian membranes that correspond to the mammalian erythrocyte spectrins (see Figs. 2 and 6). Avian α -spectrin comigrates by SDS-PAGE with mammalian α -spectrin, but the β variant has a higher mobility and can be resolved into a closely spaced doublet on underloaded gels (not shown). Both systems contain

actin at 42,000 daltons as well as a broad band of membrane proteins around 100,000 daltons (Band 3; reference 16). Among the characteristic differences are the presences in avian membranes of goblin, a hormonally-regulated phosphoprotein (4), and of the intermediate filament proteins, vimentin and synemin. The presence of these two intermediate filament components in association with avian erythrocyte membranes was demonstrated by two-dimensional gel electrophoresis (IEF/SDS-PAGE; see Fig. 4a). Erythrocyte vimentin coelectrophoreses in this gel system with vimentin identified in other avian cell types (19, 24); the identification of synemin was tentative at this stage and required immunological and biochemical confirmation, as described below. Desmin was not detected on these electrophoretograms.

Initial biochemical studies of erythrocyte membranes began with attempts to remove peripherally bound proteins from the membrane lipid bilayer. We found that the solubilization or release of any protein components from the erythrocyte membranes, without the use of detergents or strongly chaotropic agents, required the removal of divalent cations. Therefore, before most biochemical experiments, the magnesium ions present in the membrane suspension (in the hypotonic lysis buffer) were removed from the membranes by washing with a low-salt buffer containing EDTA.

Treatment of membranes with solutions of very low ionic strength was expected to release spectrin, by analogy to the mammalian erythrocyte system (42, 43). However, it was observed that if such an extraction was performed briefly at 0°C , then the primary protein released was vimentin. Fig. 2a shows a two-dimensional gel of the extract obtained by treatment of chicken erythrocyte membranes with distilled water for 30 min on ice. In addition to the four or five isoelectric variants of vimentin at 52,000 daltons, are synemin at 230,000 daltons and actin at 42,000 daltons. Identification of the 230,000 dalton polypeptide as synemin is based on its immunological cross-reactivity with smooth muscle synemin and its immunoradiographic peptide map, both as detailed below, as well as its copurification with vimentin. No desmin can be detected on this gel. Distilled water was found to be the optimal solvent for extraction of relatively pure vimentin and synemin, but other low ionic strength solutions (eg., 1–2 mM EDTA or EGTA




FIGURE 2 Water extracts of chicken erythrocyte membranes. (a) Two-dimensional gel of a distilled water extract of erythrocyte membranes showing predominance of vimentin (V) and synemin (S). The β and γ variants of actin (A) are present. (b) One-dimensional gel; extract as in (a) was centrifuged for 1 h at 240,000 *g* (av.); lane 1 contains the supernatant and lane 2 the pellet. For comparison, lane 3 shows a preparation of gizzard desmin (D) and synemin that was cycled twice by solubilization and precipitation using acetic acid. (c) Erythrocyte membrane residue after two sequential extractions with distilled water over 2-h period. G, goblin (see Fig. 6 for documentation of this identification); 1 and 2, α and β spectrin, A, actin. (d, e, f) Sequential water extracts of a single aliquot of erythrocyte membranes; extraction periods were for 1 min, 0.5 h and 9 h, respectively, and each gel represents all of the protein extracted at each step. All membrane extractions in this figure were performed at 0°C. Panel a represents the protein extracted from 35–40 μ l of packed erythrocyte membranes, and lane 1 from 20–25 μ l of membranes.

brought to pH 7.4 with Tris) will produce extracts of comparable purity but lower yield (approximately one-half). If the pH of the extracting solution is raised, release of several components in addition to vimentin is favored, resulting in a less pure preparation of vimentin and synemin. At pH 11, the membranes are nearly quantitatively stripped of non-integral proteins (not shown; see reference 56). Similarly, many proteins can be released from the membranes by treatment with acetic acid (cf. reference 41), but this results in a low-yield, very impure preparation of vimentin and synemin that exhibits extensive proteolytic degradation. Acetic acid thus appears to be undesirable for use in the extraction and purification of intermediate filaments from erythrocytes as it has been used previously for smooth muscle (29, 53).

This technique of obtaining highly enriched preparations of vimentin and synemin takes advantage of the fact that spectrin and actin are released very slowly from the membranes in distilled water or low salt buffers at 0°C. Fig. 2d-f show sequential extracts of a membrane aliquot made with distilled water at 0°C for 1 min, 30 min and 9 h. Most of the vimentin and synemin are released from the membranes within 1 min. The ratio of actin to vimentin increases with each extraction, and spectrin becomes a major component of the extract after a few hours. After prolonged extraction, though, even in the presence of EDTA and reducing agents, most of the spectrin (and much of the actin) is still associated with the membrane (not shown). If the extractions are performed at 37°C instead of 0°C, all of these proteins are released more rapidly; spectrin is solubilized rapidly enough to become the major component of short-interval extracts, making vimentin a minor component (not shown).

After repeated and prolonged extractions with distilled water at 0°C, the membranes tend to break up into smaller fragments and vesicles. Partial loss of the spectrin network may account for this.

Lane 1 of Fig. 2b is the one-dimensional gel profile of a distilled water extract identical to that in Fig. 2a, except that it was centrifuged for 1 h at 170,00–310,000 *g* (240,000 *g* av.); the supernatant was lyophilized and run in Lane 1, and the pellet was run in Lane 2. It is apparent that under these conditions little of the protein is sedimented. A similar result was obtained with a 5-h centrifugation. Minor polypeptides that do not focus discretely in the two-dimensional gel system can be visualized here. Electrophoresis of this material on less porous polyacrylamide gels shows that most of the material migrating with the dye front in Lane 1 is residual hemoglobin. Lane 3 is a preparation of chicken-gizzard smooth-muscle intermediate filaments that shows synemin, vimentin, desmin, and actin for molecular weight comparison. Synemin and α spectrin are difficult to resolve from one another on normally loaded one-dimensional gels but are clearly resolved on underloaded gels and two-dimensional gels.

Not all of the vimentin is extracted from erythrocyte membranes with a single distilled water treatment. Fig. 2c shows a sample of membranes that was extracted twice with water over a 2-h period; the amount of vimentin is less, but some still remains. Residual vimentin is evident even after four extractions over a 5-d period (not shown). However, in thin sections of membranes treated with distilled water for 5 min, cytoplasmic intermediate filaments cannot be found. This suggests that the residual vimentin may not be in the form of free cytoplasmic filaments (see Discussion).

Immunological Characterization of Erythrocyte Intermediate Filaments

The technique of immunautoradiography (10), which uses antibodies to detect protein antigens in polyacrylamide gels, was used in this study for three purposes: (a) to determine whether the erythrocyte intermediate-filament subunits were antigenically crossreactive with their muscle counterparts; (b) as a form of peptide map analysis to determine whether the subunits in erythrocytes were homologous or identical to their muscle counterparts; and (c) to detect these antigens in gels with a sensitivity much greater than that afforded by Coomassie Blue staining. Antisera used in this study were all elicited against chicken muscle proteins, purified by SDS-PAGE, and each appears to be specific for its respective antigen as assayed by two-dimensional immunautoradiography (24, 25).

Fig. 3 shows the presence of immunoreactive forms of both vimentin and synemin in various fractions of chicken and turkey erythrocyte membranes. Fig. 3a shows a Coomassie Blue-stained SDS-polyacrylamide gel of a variety of samples; this gel was labeled with anti-synemin followed by radioiodinated protein A, and the corresponding autoradiogram is shown in Fig. 3c. A duplicate gel was processed with anti-vimentin, and its autoradiogram is shown in Fig. 3b.

Lane 1 represents whole white cells from chicken blood, examined to ensure that the vimentin and synemin being studied in the erythrocyte preparation were not originating from the extremely low level of contamination by white cells. Little if any synemin is detectable, and the quantity of vimentin is low relative to the amount of actin present in these cells.

Lane 2 is whole chicken-gizzard smooth-muscle tissue. It is the tissue from which synemin was originally purified and was the source of the synemin used for immunization (25). Vimentin is also present in this tissue (25; Fig. 3b). The two most

prominent bands near the top of the lane are filamin and myosin; the autoradiogram of Fig. 3c shows that synemin migrates between these two proteins (see also reference 25). α -Actinin is visible at 100,000 daltons, and the two major bands in the middle are desmin and actin (50,000 and 42,000 daltons).

The remaining lanes demonstrate the presence of vimentin and synemin in erythrocyte membranes, membrane cytoskeletons, and low salt extracts. Lanes 3 and 8 display chicken erythrocyte membranes, and lane 4 displays turkey erythrocyte membranes. Both samples contain polypeptides that have molecular weights and antigenicities similar to those of muscle vimentin and synemin. Lanes 5 and 6 represent turkey and chicken erythrocyte membranes that have been extracted with high salt and Triton X-100. Both vimentin and synemin remain insoluble in the cytoskeletal residue, as do the spectrin and actin. Goblin, the 100,000 dalton cluster, a 44,000 dalton polypeptide and many minor components are partially or completely solubilized (compare lanes 5 and 6 with lanes 3 and 4). Lane 7 contains a low-salt extract of chicken erythrocyte membranes that is highly enriched in vimentin and synemin; the greater amount of handling and processing of these samples relative to the other samples on the gel probably accounts for the increased quantity of proteolytic fragments of vimentin and synemin evident in these lanes of the autoradiogram.

Anti-desmin does not label any of the proteins in these samples, except for the desmin present in the gizzard tissue (not shown).

Fig. 4a depicts chicken erythrocyte membrane proteins resolved by IEF/SDS-PAGE. This gel was processed in immunautoradiography with anti-vimentin, and the resulting autoradiogram is shown in Fig. 4b. An identical gel was processed with anti-synemin, and its autoradiogram is in Fig. 4c. Proteins readily identified by Coomassie Blue staining include goblin, α and β spectrin, synemin, the multiple isoelectric variants of




FIGURE 3 Immunautoradiography using antivimentin and antisynemin. (a) Polyacrylamide gel stained with Coomassie Blue after labeling with antisynemin and radioiodinated protein A; (b) Autoradiogram of duplicate gel labeled with anti-vimentin; (c) Autoradiogram of gel in (a). Samples are from chickens unless otherwise noted. Lane 7: Buffy coat. The identity of the prominent labeled polypeptide just above the dye front is not known. It labels comparably with both antisera and may therefore be an IgG receptor or protein A receptor that survives SDS denaturation and acetic acid/ethanol fixation. Lane 2: Whole gizzard muscle. Lane 3: Erythrocyte membranes. Lane 4: Turkey erythrocyte membranes. Lane 5: High salt plus detergent residue of turkey erythrocyte membranes. Lane 6: High salt plus detergent residue of erythrocyte membranes. Lane 7: Low salt extract of erythrocyte membranes. Lane 8: Erythrocyte membranes. A, actin; V, vimentin. Synemin is the major high molecular weight protein in lane 7; it migrates just beneath α spectrin.




FIGURE 4 Two-dimensional immunofluorescence of chicken erythrocyte membranes using antivimentin and antisynemin. (a) Two-dimensional gel of chicken erythrocyte membranes stained with Coomassie Blue after labeling with antivimentin and radioiodinated protein A. (b) Autoradiogram of same gel. (c) Autoradiogram of duplicate gel labeled with antisynemin. A, actin; V, vimentin; S, synemin; 1, α spectrin 2, β spectrin; G, goblin.

vimentin, and nearly equal amounts of β and γ actin. Antivimentin and anti-synemin label only their respective proteins on the gel and do not crossreact with other proteins in this system. Desmin is not detectable by Coomassie Blue staining, nor by immunofluorescence with anti-desmin (not shown). The diagonal string of polypeptides smaller and more acidic than vimentin, visible in Fig. 4b, represents breakdown products of vimentin (19, 24); the same probably holds true for the numerous polypeptides under synemin that label with anti-synemin. Because the vimentin and synemin used for immunization were excised from an SDS-polyacrylamide gel, it seems unlikely that all of these minor polypeptides could be unrelated contaminants. The relative quantity of these other polypeptides increases with increased processing of the samples. They can be reduced in amount or completely eliminated if special precautions are taken to inhibit proteolytic enzymes (19, 25). Also, the same patterns are seen in samples from different tissues (references 19, 24, 25, and below).

To determine the antigenic homology of erythrocyte synemin and gizzard synemin, a fortuitous form of peptide mapping was used. Fragments of synemin generated by endogenous proteases during processing of the tissues were detected with antibodies to synemin as visualized by two-dimensional immunofluorescence. An antiserum specific for a given protein thus allows visualization of the protein's peptide map without the necessity of prior purification of that protein. The degradation pattern of erythrocyte vimentin as seen in Fig. 4b is similar to that already published for muscle vimentin (24).

Fig. 5 compares the synemin present in chicken erythrocyte cytoskeletons and that in chicken gizzard smooth muscle tissue. Fig. 5a shows the proteins that remain after extraction of erythrocytes with 1% Triton X-100 in a physiological salt buffer; a major difference in the protein pattern, compared with the membranes in Fig. 4a, is the presence of the nuclear lamins. There is also more vimentin relative to the amount of actin present, as this preparation also includes the nucleus-associated intermediate filaments. The autoradiogram of this gel after processing with anti-synemin is shown in Fig. 5b. Similarly, a two-dimensional gel of whole gizzard tissue processed with anti-synemin, and its autoradiogram, are shown in Fig. 5c and d. The similarities in the two synemin patterns are striking (see also Fig. 4c); in each case, the parent molecule is most heavily labeled, and the arcs of daughter products terminate in what appears to be a particularly stable fragment at $\sim 34,000$ daltons (pI ~ 4.9).

We have noted no consistent differences between erythrocyte

synemin and smooth muscle synemin. Minor differences in the fragment patterns may be attributable to different endogenous proteases, different processing schemes, or slight differences in electrophoresis; the latter two would explain the very minor differences between the erythrocyte synemins in Figs. 4c and 5d. There is a slight variation in the observed isoelectric points of erythrocyte and smooth-muscle synemin, but there is variation even among different samples of erythrocyte synemin (compare Figs. 2a and 4a). The focusing of synemin seems to be influenced by the amount of protein loaded on the isoelectric focusing gel; the apparent isoelectric point of synemin is often the same as that of desmin or vimentin if either of the latter is present in large quantities on the gel (Fig. 2a and reference 25). We conclude from these immunofluorescent data that erythrocyte synemin and muscle synemin are similar if not identical; similarities in solubility properties and cellular distribution (below) strengthen the conclusion that the erythrocyte polypeptide may be regarded as synemin as defined previously in smooth muscle.

Phosphorylation

Goblin is a high molecular weight protein of the turkey erythrocyte plasma membrane characterized by hormone dependent phosphorylation (4). Both goblin and synemin have reported molecular weights of $\sim 230,000$ daltons; although their solubility properties and distributions appeared to differ, we thought it was necessary to conclusively determine whether goblin and synemin were indeed different proteins. We identified goblin by its characteristic properties of being a large membrane-associated protein and the most hyperphosphorylated polypeptide in turkey erythrocytes treated briefly with the β -adrenergic agonist, isoproterenol (4). Fig. 6a shows a Coomassie Blue-stained gel of membranes of turkey erythrocytes labeled with [32 P] inorganic phosphate; those on the left were also treated with isoproterenol, whereas those on the right were not. Fig. 6b is the corresponding autoradiogram. By the above criteria, we conclude that the band designated in the figure is goblin. In this gel system, goblin migrates more slowly than the two spectrin variants, rather than migrating between them as in the system of Beam et al. (4). Using their gel system, we found that the electrophoretic pattern of our samples was indeed different: the relative positions of some bands was different, and goblin and the spectrins were not resolved as well.

A 44,000 dalton polypeptide is also noticeably hyperphos-




FIGURE 5 Comparison of erythrocyte and gizzard synemins by two-dimensional immunautoradiography. (a, c) Two-dimensional gels stained with Coomassie Blue after labeling with antisynemin and radioiodinated protein A. (b, d) Corresponding autoradiograms. Samples are (a) a Triton X-100 insoluble cytoskeleton of chicken erythrocytes, and (c) whole gizzard smooth muscle tissue. A, actin; D, desmin; V, vimentin; L, lamins A, B, and C (references 21 and 52); M, myosin; S, synemin; F, filamin.

phorylated in the presence of isoproterenol (4), and this hormone-dependent phosphorylation seems to apply to a lesser degree to several other proteins as well. Vimentin appears to be one of these, because a slight increase in labeling of this protein with isoproterenol can be detected on the autoradiogram (Fig. 6*b*). This is consistent with the hormone-dependent phosphorylation of vimentin that has been observed in other cell types (D. L. Gard and E. Lazarides, manuscript submitted for publication). Synemin migrates too close to α spectrin on one-dimensional gels to be able to determine whether it is phosphorylated, but two-dimensional gels of erythrocytes phosphorylated to a steady state (1) suggest that synemin is indeed phosphorylated (not shown). These gels also show that all but the most basic isoelectric variant of vimentin are phosphorylated, as is the case with vimentin in other cell types (46, 47).

Localization of Synemin and Vimentin by Immunofluorescence

With antibodies specific for synemin and vimentin, it was possible to determine the spatial distributions of these antigens within the avian erythrocyte by immunofluorescence. The abundance of hemoglobin in these cells was not a problem because synemin and vimentin remained insoluble after removal of the hemoglobin by hypotonic lysis or detergent lysis. Typically, erythrocytes were allowed to attach to Alcian Blue-coated glass cover slips (54), then lysed with a physiological salt solution containing Triton X-100 and magnesium ions, and processed for examination by immunofluorescence microscopy without any fixation. The result with both anti-vimentin and

anti-synemin was a cytoplasmic network of sinuous filaments extending from the nucleus to the plasma membrane. This network gave the impression of being composed of a small number of long filaments, since relatively few free ends could be seen. A high concentration of filaments was often noted near the poles of the nuclei (Fig. 7*d*). However, the small size and relatively great depth of field of the cells, coupled with the dense packing of the filaments, prevented adequate resolution of the individual filaments after photographic reproduction of the network. Nevertheless, this technique of immunofluorescence allowed us to determine the effects of Colcemid on the filaments in these cells.

Colcemid has been found to cause an aggregation and bundling of intermediate filaments in a variety of cultured cell types (8, 18, 23, 31, 33, 38). When turkey erythrocytes were incubated with Colcemid under conditions that are normally used for cultured cells, their intermediate filaments did not aggregate. Chicken embryo fibroblasts incubated in the same petri plate showed normal filament aggregation. Fig. 7 shows the distribution of the erythrocyte intermediate filaments after this incubation, as revealed by immunofluorescence using anti-vimentin. The top row depicts cells that received Colcemid, and the bottom row shows control cells that were treated identically, except that they received no Colcemid. After incubation with or without Colcemid, the cells were briefly lysed with Triton X-100 to remove hemoglobin, then fixed with formaldehyde to ensure preservation of filament distribution. There was no obvious Colcemid-induced aggregation of the filaments, and what appeared to be individual filaments could clearly be seen extending to the plasma membrane. No consistent difference between the Colcemid-treated cells and con-




FIGURE 6 Phosphorylation of turkey erythrocyte plasma membrane proteins: identification of goblin. Erythrocytes were incubated for 4 h with ^{32}P -phosphate; half were treated with isoproterenol for the final 20 min. Plasma membranes were isolated and the proteins were resolved by SDS-PAGE. The Coomassie Blue-stained gel was dried (a) and autoradiographed (b). Goblin (G), the high-molecular-weight protein that is most noticeably hyperphosphorylated in the presence (+) of isoproterenol, is distinct from synemin, which runs just beneath α spectrin (1). Abbreviations: G, goblin; 1 and 2, α and β spectrin; V, vimentin; A, actin; + and - refer to the presence and absence of isoproterenol in the incubation mixture.

trols could be detected. (This filament distribution is the same as in erythrocytes that have not been subjected to an *in vitro* incubation but examined soon after removal from the animal.) Similar results were obtained with anti-synemin. Unfixed cells exhibit a similar filament distribution, but the nuclei shrink somewhat and become more dense during processing. High concentrations of Colcemid (100 μM rather than 5 μM) also had no effect on the intermediate filaments, and cells treated in suspension rather than after attachment to cover slips were similarly unaffected.

To examine the association of the filaments with the membrane and allow visualization of the filament network more clearly, a technique was developed that removed the erythrocyte nuclei and most of the filaments to leave a residual mat of membrane-associated filaments in which the individual strands could be resolved and photographed. This technique is based on ideas that stemmed from a number of sources (3, 26, 35, 44, 54). It involves attaching erythrocytes to Alcian Blue-coated glass cover slips, then disrupting the cells by cavitation (with a sonicator) forcefully enough to remove nuclei and other cellular structures not firmly anchored to the cover slip. This results in a cover slip covered with residual elliptical patches of membrane, each with its most firmly associated structures. Immu-

nofluorescence shows that these patches often have vimentin- and synemin-containing filaments attached to them. This was the first good indication that intermediate filaments might in some way be physically anchored to the erythrocyte plasma membrane. Fig. 8 is a montage of fluorescence micrographs showing the presence and distribution of vimentin and synemin on these residual patches of membrane. Tangled and wavy networks of filaments and fragments of filaments can be visualized. The uniform-diameter filaments probably represent individual intermediate filaments rather than bundles (see electron microscopic correlates in Fig. 9). Their measured diameter of 200–300 nm is close to the resolution limit of the light microscope and consistent with the immunofluorescence image of individual microtubules (49). Anti-desmin gives negligible fluorescence, which is consistent with our inability to detect it electrophoretically. Synemin and vimentin preimmune sera also give negligible fluorescence, and preadsorption of the antisera with the corresponding purified proteins has been shown previously to block fluorescence (25).

The quantity of filaments remaining on the membrane patches is probably a function of the degree of sonication. Sonication was monitored by phase-contrast microscopy and performed at a level that removed most of the nuclei. This produced a wide range of anucleate membrane patches; most patches retained no intermediate filaments (see Fig. 8s), whereas the rest displayed patterns ranging from short fragments (Fig. 8l–p) to complex networks of filaments (Fig. 8a–d). Fig. 8s is a combination phase-contrast/fluorescence micrograph showing bare membrane patches as well as filament-containing patches. It is noteworthy that the remaining filaments are not always distributed uniformly over the membrane patch.

Double immunofluorescence was performed to directly compare the distributions of vimentin and synemin on these patches. Fig. 8q and r show that the distributions are the same. Both antigens appear along the same filaments, and nowhere at this level of resolution is one antigen present and the other absent. However, the synemin fluorescence sometimes gives the impression of being slightly punctate along filaments that show uniform vimentin fluorescence.

We have obtained comparable immunofluorescence results with cells sonicated in low ionic strength and physiologic salt buffers, by following sonication with a detergent/high-salt extraction, and with subsequent incubations and rinses in the presence or absence of Triton X-100. In none of the preparations shown in Fig. 8 were the samples fixed with protein crosslinkers or denaturants.

Ultrastructure of Sonicated Membranes

Sonicated erythrocyte membrane patches on glass cover slips, similar to those used for immunofluorescence, were fixed and rotary-shadowed for examination by transmission electron microscopy. Fig. 9 shows portions of three such membrane patches with their adherent filaments. The pattern of the filaments is similar to the pattern seen in immunofluorescence; the filaments tend to be relatively long, and can be straight, wavy or curved into loops. Occasionally, the filaments extend beyond the edge of the membrane patch (Fig. 9b and c), presumably as a result of the sonication. Similar patterns are obtained if the erythrocytes are sonicated in a physiologic salt buffer without previous hypotonic lysis, showing that the filaments are not precipitated on the membranes as a result of the low-salt treatment. Treatment with Triton X-100 also does not




FIGURE 7 Distribution of vimentin in Colcemid-treated erythrocytes. Phase (a, c) and corresponding fluorescence micrographs (b, d) of turkey erythrocytes incubated in culture medium with (a, b) or without (c, d) Colcemid. Intermediate filaments were visualized in fixed cytoskeletons by indirect immunofluorescence using anti-vimentin. Similar patterns were obtained with antisynemin. Bar, 10 μm . $\times 1350$.

affect this pattern. Strands of chromatin originating from the erythrocyte nuclei have a distinctive morphology and are evident in the replicas only when magnesium ions are omitted from the lysis or sonication buffers. The relatively flat, coarse, upper face of the membrane patch probably represents the spectrin network that lines the cytoplasmic surface of the plasma membrane. The intermediate filaments are of fairly uniform diameter; it is not known whether the frequent, slight bulges, constrictions, and discontinuities have a molecular basis or are just an artifact of the shadowing procedure (cf. reference 28). Detail cannot be resolved sufficiently to determine whether the filaments branch or merely associated laterally in various places. We have not been able to positively identify structures that might anchor the filaments to the membrane patch, and it is not evident from these micrographs how abundant such linkers might be. From the distribution of the filaments on the sonicated membranes, it appears likely that they are not randomly distributed over the plasma membrane.

DISCUSSION

Intermediate Filaments in Avian Erythrocytes

Avian erythrocytes provide a simple system for the study of

intermediate filaments. These cells are terminally differentiated, nondividing, nonadherent, and nonmotile. They are easily obtained and purified. Their cytoplasm is relatively simple and nondynamic. Interaction of intermediate filaments with both the nucleus and plasma membrane can thus be examined without the variability and complexity inherent in most other cell types. Comparison of the avian erythrocyte with the well characterized mammalian erythrocyte provides insight into the functions and relationships between components that are not common to both, such as nuclei, microtubules and intermediate filaments.

In this study we demonstrate that vimentin and synemin are the major components of avian erythrocyte intermediate filaments. These filaments had previously been postulated to be composed of vimentin (63), and vimentin indeed appears to be their major subunit. Vimentin and synemin appear to coexist uniformly in the filaments, which form a looping, intertwined network in the cytoplasm. A portion of this filament network is associated with the plasma membrane firmly enough to resist detachment by physical disruption forces that are sufficient to remove the nucleus and fragment the plasma membrane. However, synemin and vimentin can be selectively released from the plasma membrane by treatment with divalent cation-free, low ionic strength solutions.

We have used chicken and turkey erythrocytes as representatives of avian erythrocytes in general for this work. Immunofluorescence was performed on turkey erythrocytes, because they are slightly larger than chicken erythrocytes. The protein goblin had been defined in turkey erythrocytes (4), so we used these cells for the phosphorylation experiments. Biochemical studies were based primarily on chicken erythrocytes, which were more readily available than turkey erythrocytes; electron microscopy of thin sections was performed on chicken erythrocytes to correlate with the biochemistry. However, it is evident from electrophoretograms such as in Fig. 3a that chicken and turkey erythrocytes are not identical. For example, turkey erythrocyte membranes have a smaller β -spectrin and less protein in the 100,000 dalton cluster and 44,000 dalton band than chicken erythrocyte membranes. Vimentin and synemin, though, appear to be very similar, if not identical, as judged by IEF/SDS-PAGE and immunautoradiography (not shown). We believe, therefore, that the generalizations made about vimentin and synemin in this paper are likely to be applicable to all avian erythrocytes.

Studies of Intermediate Filaments Associated with the Plasma Membrane

Our studies of the properties of erythrocyte intermediate filaments have focused on those filaments that fractionate with the plasma membrane. They have the same antigenicity and electrophoretic mobility as the filaments that fractionate with the nuclei, so we regard them as equivalent in terms of basic composition and properties. Whether a given segment of filament will fractionate with the nucleus or plasma membrane may be variable and depend as much on the homogenization conditions as on the spatial arrangement of the filaments.

We have developed an enucleation technique that gives fair yields of whole plasma membranes and large pieces of membrane (referred to here simply as "erythrocyte membranes"). Many of these membranes retain large networks of intermediate filaments; in a given thin section, though, many do not appear to possess any associated filaments. This may be a result of loss during enucleation, initial lack of attachment of the filaments to certain regions of the plasma membrane, absence




FIGURE 8 Immunofluorescence of turkey erythrocyte intermediate filaments. Erythrocytes adhering to cover slips were hypotonically lysed and sonicated to remove overlying membranes and nuclei. Intermediate filaments remaining attached to the resulting patches of plasma membrane were visualized by immunofluorescence using antibodies to synemin (*a, r*) or vimentin (*b-g, s*). Specimens were not fixed, and all but *a, b, c,* and *h* were treated with Triton X-100. Micrographs *a-p* are indirect immunofluorescence images; *q* and *r* demonstrate colocalization of vimentin (*q*) and synemin (*r*) by double immunofluorescence. Micrograph *s* is a combination phase/fluorescence image showing the distribution of vimentin on the elliptical patches of plasma membrane; note that many patches are devoid of filaments. Bars, 5 μm . *a-r*, $\times 3040$; *s*, $\times 1330$.




FIGURE 9 Platinum replicas of sonicated chicken erythrocyte ghosts. Samples were prepared as in Fig. 8, then fixed, dried, and rotary shadowed with platinum for examination by transmission electron microscopy. Intermediate filaments can be seen on patches of plasma membrane that remained attached to the cover slip during sonication. In *b* and *c*, a portion of the filaments have fallen beyond the edge of the membrane patch. Magnification: Bar, 1 μm . $\times 16,000$.

from certain regions of the erythrocyte cytoplasm, or a close apposition to the plasma membrane that renders the filaments unresolvable.

Previous studies involving isolation of the avian erythrocyte plasma membrane by differential centrifugation after mechanical disruption of the cells have relied on pressure-release homogenization (7, 14, 61), sonication (4, 27), rotating blades (11, 66), or a tight-fitting Dounce (12, 22) or Potter-Elvehjem homogenizer (6). However, the presence of filaments associated with the isolated membrane fragments was noted only rarely

(27), and, in comparisons to mammalian erythrocyte membranes, the presence of an extra polypeptide, similar in molecular weight to vimentin, was rarely mentioned (11). Some of these disruption techniques produce very small membrane fragments that may be largely stripped of filaments; alternatively, the filaments may assume a distribution or configuration in which they are not readily identifiable by electron microscopy. The gentler disruption techniques appear to produce membrane fragments similar to those in this study, but associated filaments have tended to escape detection. Intermediate

filaments have been most apparent in detergent-insoluble cytoskeletons of whole erythrocytes examined by thin sectioning or negative staining (59, 63).

Treatment of avian erythrocyte membranes with certain low ionic strength solutions removes the associated intermediate filaments. Filaments can no longer be found with the membranes in thin sections, and the low-salt extract contains nearly pure vimentin and synemin. This release seems to depend on low ionic strength and absence of divalent cations and be independent of reducing agents or nonionic detergents. Our highest yields have been obtained using distilled water. Roughly 60–90% of the vimentin is released after 1 min of extraction with distilled water. Selective release of vimentin and synemin, as compared to spectrin and actin, is enhanced by low temperature and brevity of treatment. Because the released vimentin and synemin cannot be sedimented by centrifugation for 5 h at 240,000 g, yet appear to comigrate in a gel filtration column with an exclusion limit of 15 million daltons (unpublished observations), they must exist in solution as some sort of multimeric complex or oligomer. This implies that the filaments break down or partially depolymerize during or after release from the membranes. Solubility in low salt has similarly been described for other preparations of native intermediate filaments (29, 30, 50, 51, 55, 57). These extraction conditions may thus be resulting in a dissolution of the filaments rather than a dissociation of the filaments from the membranes. It is conceivable that these extraction conditions have no disruptive effect on the anchorage points of the filaments to the membranes, which would explain why some of the vimentin remains with the membranes after extensive extraction with water. This vimentin may be a distinct population associated with anchorage points in the form of tightly bound monomers or oligomers or short segments of filament not resolvable in thin sections. These extraction data thus do not permit a conclusion about the nature of attachment of the filaments to the membranes. It can only be stated, based on the physical data of enucleation and sonication, that at least some of the intermediate filaments in avian erythrocytes are somehow anchored to the plasma membrane, and that this attachment is stable in the presence of physiologic salt, high salt, and nonionic detergent.

Comparison of Erythrocyte Proteins

An aspect of comparative biochemistry exemplified by this study is the difficulty of comparing protein profiles of a given preparation by different SDS-PAGE systems. Although useful for general comparisons, different gel systems may not be directly comparable with regard to specific polypeptides. There has classically been disagreement between different investigators about calculated molecular weights; even the relative positions of different polypeptides may not be consistent in different gel systems (for example, the high molecular weight proteins shown in this paper—see Results). This stresses caution in identifying a polypeptide solely by its mobility on an SDS-polyacrylamide gel. Our electrophoretic profiles of avian erythrocyte membrane proteins differ from those of other laboratories, which also differ among themselves (2, 4, 7, 11, 12, 34, 60, 61); some of these differences have been noted and attributed to endogenous proteases or proteases present in contaminating leukocytes (11, 34, 61). Extrapolation from one class to another (for example, mammalian (16) to avian erythrocyte membranes) may not be justified either and may lead to erroneous identification of polypeptides. Two-dimensional gel

electrophoresis makes polypeptide identification less ambiguous, because another parameter (isoelectric point) is taken into account and has proved useful for several proteins in this study. Nevertheless, other (nonelectrophoretic) evidence for the identity of a protein band on a gel is essential. We have used immunologic and solubility properties, in addition to electrophoretic characteristics, to identify synemin and vimentin in avian erythrocytes, and phosphorylation characteristics to identify goblin (4). Determination of why similarly prepared samples show not only different relative mobilities but also different relative amounts using different gel systems awaits further study.

The Effects of Colcemid

One indication of a functional or interactive difference between the intermediate filaments of avian erythrocytes and most other cell types grown *in vitro* is the insensitivity of the former to Colcemid. Treatments with Colcemid that will cause aggregation and perinuclear bundling of intermediate filaments in most cultured cells (8, 18, 23, 31, 33, 38) appear to have no effect on the filaments of erythrocytes. Colcemid sensitivity might thus be a function of how dynamic a cell is, or perhaps its state of differentiation, as appears to be likely for skeletal muscle cells (20, 25) but not be an intrinsic property of the filaments themselves. Related to this may be the observation that chick erythrocyte marginal band microtubules are resistant to depolymerization by Colcemid (5).

Intermediate Filament Proteins

Synemin was originally found to be associated with intermediate filaments in smooth and skeletal muscle (25). Here we show that synemin is not a muscle-specific protein but is present as well in at least one nonmuscle cell, the mature avian erythrocyte. The original study also raised the possibility that synemin was a desmin-associated polypeptide; here we show that synemin can also exist and associate with vimentin. In both muscle cells and erythrocytes, synemin appears to be a component of the same filaments that contain desmin and vimentin, as determined by double immunofluorescence. Densitometric scans of Coomassie Blue-stained polyacrylamide gels of preparations of vimentin and synemin from chicken erythrocytes give a vimentin-to-synemin ratio of ~50:1. This is similar to the ratio obtained for desmin and synemin in smooth muscle and suggests a constant stoichiometry between synemin and intermediate filaments of different subunit composition. This ratio is a very rough estimate, not taking into account differential proteolysis of the proteins during processing and possible nonlinearity in dye binding and densitometry, and should therefore not be regarded as the true ratio. It is useful, however, for rough comparisons of different systems.

We have taken advantage of a novel form of two-dimensional peptide mapping to compare synemins from different tissues. This combination of partial hydrolysis of tissue proteins by endogenous proteases and two-dimensional immunoradiography has demonstrated a high degree of homology between synemins from avian smooth muscle and erythrocytes. Both molecules exhibit an S-shaped string of fragments that terminates in a proteolytically stable peptide of 34,000 daltons. This technique is extremely sensitive, detecting peptides much too scarce to be seen by Coomassie Blue staining, but does not resolve the high molecular weight peptides sufficiently to allow detailed comparisons. Also, slight variations from gel to gel do

not allow us to conclude that the synemins we are examining are identical. Minor differences in the maps may be artifactual or may reflect functional differences in the molecule, perhaps related to whether synemin is found in association with desmin or with vimentin.

These data do not address the question of whether synemin is an integral or an associated filament protein, or what its properties are independent of desmin and vimentin. The large size and paucity of synemin relative to desmin and vimentin tend to favor a role for synemin as an associated polypeptide. Perhaps it is analogous to the high molecular weight polypeptide of neurofilaments, which appears to be wrapped helically around the core filament (62), where it may function to stabilize the filament, promote assembly (45), or mediate interactions with other molecules or organelles.

The presence of nonmicrotubular filaments in nucleated erythrocytes has been known for some time (27, 36), but only recently were they identified as intermediate filaments (59, 63). These filaments were usually noted and studied in relation to the nucleus or nuclear membrane. In this paper we show that they also exhibit a close association with and apparent anchorage to the plasma membrane, and that they contain the intermediate-filament subunits vimentin and synemin. Nucleated erythrocytes may thus be an ideal model system for the study of filament-membrane interactions and for examining intermediate filament nucleation, assembly, and deployment during differentiation.

We thank David L. Gard for his helpful comments on the manuscript, and Dr. Jean-Paul Revel and Mr. Patrick F. Koen for their help with the electron microscopy.

This work was supported by grants from the National Institutes of Health (NIH) (GM 06965), National Science Foundation, Muscular Dystrophy Association of America, and a Biomedical Research Support Grant to the Division of Biology, California Institute of Technology. B. L. Granger was also supported by a Predoctoral Training Grant from the NIH (GM 07616), and E. A. Repasky by a Postdoctoral Fellowship from the NIH (GM 07401). E. Lazarides is the recipient of a Research Career Development Award from the NIH.

Received for publication 13 July 1981, and in revised form 5 October 1981.

REFERENCES

- Alper, S. L., K. G. Beam, and P. Greengard. 1980. Hormonal control of Na⁺-K⁺ co-transport in turkey erythrocytes. Multiple site phosphorylation of goblin, a high molecular weight protein of the plasma membrane. *J. Biol. Chem.* 255:4864-4871.
- Alper, S. L., H. C. Palfrey, S. A. DeRiemer, and P. Greengard. 1980. Hormonal control of protein phosphorylation in turkey erythrocytes. Phosphorylation by cAMP-dependent and Ca²⁺-dependent protein kinases of distinct sites in goblin, a high molecular weight protein of the plasma membrane. *J. Biol. Chem.* 255:11029-11039.
- Batten, B. E., J. J. Aalberg, and E. Anderson. 1980. The cytoplasmic filamentous network in cultured ovarian granulosa cells. *Cell.* 21:885-895.
- Beam, K. G., S. L. Alper, G. E. Palade, and P. Greengard. 1979. Hormonally regulated phosphoprotein of turkey erythrocytes. Localization to plasma membrane. *J. Cell Biol.* 83: 1-15.
- Behnke, O. 1970. A comparative study of microtubules of disk-shaped blood cells. *J. Ultrastruct. Res.* 31:61-75.
- Bilezikian, J. P., and G. D. Aurbach. 1973. A β -adrenergic receptor of the turkey erythrocyte. I. Binding of catecholamine and relationship to adenylate cyclase activity. *J. Biol. Chem.* 248:5577-5583.
- Blanchet, J. P. 1974. Chicken erythrocyte membranes: Comparison of nuclear and plasma membranes from adults and embryos. *Exp. Cell Res.* 84:159-166.
- Blose, S. H., and S. Chacko. 1976. Rings of intermediate (100 Å) filament bundles in the perinuclear region of vascular endothelial cells. Their mobilization by colcemid and mitosis. *J. Cell Biol.* 70:459-466.
- Branton, D., C. M. Cohen, and J. Tyler. 1981. Interaction of cytoskeletal proteins on the human erythrocyte membrane. *Cell.* 24:24-32.
- Burridge, K. 1978. Direct identification of specific glycoproteins and antigens in sodium dodecyl sulfate gels. *Methods Enzymol.* 50:54-64.
- Caldwell, A. B. 1976. Proteins of the turkey erythrocyte membrane. *Biochemistry.* 15:2711-2718.
- Chan, L.-N. L. 1977. Changes in the composition of plasma membrane proteins during differentiation of embryonic chick erythroid cells. *Proc. Natl. Acad. Sci. U. S. A.* 74:1062-1066.
- Cohen, W. D. 1978. Observations on the marginal band system of nucleated erythrocytes. *J. Cell Biol.* 78:260-273.
- Davoren, P. R., and E. W. Sutherland. 1963. The cellular location of adenyl cyclase in the pigeon erythrocyte. *J. Biol. Chem.* 238:3016-3023.
- Dodge, J. T., C. Mitchell, and D. J. Hanahan. 1963. The preparation and chemical characteristics of hemoglobin-free ghosts of human erythrocytes. *Arch. Biochem. Biophys.* 100:119-130.
- Fairbanks, G., T. L. Steck, and D. F. H. Wallach. 1971. Electrophoretic analysis of the major polypeptides of the human erythrocyte membrane. *Biochemistry.* 10:2606-2617.
- Fourie, F. leR. 1977. Effects of anticoagulants on the hematocrit, osmolarity and pH of avian blood. *Poult. Sci.* 56:1842-1846.
- Franke, W. W., E. Schmid, M. Osborn, and K. Weber. 1978. Different intermediate-sized filaments distinguished by immunofluorescence microscopy. *Proc. Natl. Acad. Sci. U. S. A.* 75:5034-5038.
- Gard, D. L., P. B. Bell, and E. Lazarides. 1979. Coexistence of desmin and the fibroblastic intermediate filament subunit in muscle and nonmuscle cells: identification and comparative peptide analysis. *Proc. Natl. Acad. Sci. U. S. A.* 76:3894-3898.
- Gard, D. L., and E. Lazarides. 1980. The synthesis and distribution of desmin and vimentin during myogenesis in vitro. *Cell.* 19:263-275.
- Gerace, L., and G. Blobel. 1980. The nuclear envelope lamina is reversibly depolymerized during mitosis. *Cell.* 19:277-287.
- Ginsberg, B. H., C. R. Kahn, and J. Roth. 1976. The insulin receptor of the turkey erythrocyte. Characterization of the membrane-bound receptor. *Biochim. Biophys. Acta.* 443:227-242.
- Goldman, R. D. 1971. The role of three cytoplasmic fibers in BHK-21 cell motility. I. Microtubules and the effects of Colchicine. *J. Cell Biol.* 51:752-762.
- Granger, B. L., and E. Lazarides. 1979. Desmin and vimentin coexist at the periphery of the myofibril Z disc. *Cell.* 18:1053-1063.
- Granger, B. L., and E. Lazarides. 1980. Synemin: a new high molecular weight protein associated with desmin and vimentin filaments in muscle. *Cell.* 22:727-738.
- Hall, C. E. 1956. Visualization of individual macromolecules with the electron microscope. *Proc. Natl. Acad. Sci. U. S. A.* 42:801-806.
- Harris, J. R., and J. N. Brown. 1971. Fractionation of the avian erythrocyte: an ultrastructural study. *J. Ultrastruct. Res.* 36:8-23.
- Heuser, J. E., and M. W. Kirschner. 1980. Filament organization revealed in platinum replicas of freeze-dried cytoskeletons. *J. Cell Biol.* 86:212-234.
- Hubbard, B. D., and E. Lazarides. 1979. Copurification of actin and desmin from chicken smooth muscle and their copolymerization in vitro to intermediate filaments. *J. Cell Biol.* 80:166-182.
- Huiatt, T. W., R. M. Robson, N. Arakawa, and M. H. Stromer. 1980. Desmin from avian smooth muscle. Purification and partial characterization. *J. Biol. Chem.* 255:6981-6989.
- Hynes, R. O., and A. T. Destree. 1978a. 10 nm filaments in normal and transformed cells. *Cell.* 13:151-163.
- Hynes, R. O., and A. Destree. 1978b. Relationships between fibronectin (LETS protein) and actin. *Cell.* 15:875-886.
- Ishikawa, H., R. Bischoff, and H. Holtzer. 1968. Mitosis and intermediate-sized filaments in developing skeletal muscle. *J. Cell Biol.* 38:538-555.
- Jackson, R. C. 1975. The exterior surface of the chicken erythrocyte. *J. Biol. Chem.* 250: 617-622.
- Jacobson, B. S., and D. Branton. 1977. Plasma membrane: rapid isolation and exposure of the cytoplasmic surface by use of positively charged beads. *Science (Wash. D. C.).* 195: 302-304.
- Koehler, J. K. 1968. Freeze-etching observations on nucleated erythrocytes with special reference to the nuclear and plasma membranes. *Z. Zellforsch. Mikrosk. Anat.* 85:1-17.
- Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature (Lond.).* 227:680-685.
- Lazarides, E. 1978. The distribution of desmin (100 Å) filaments in primary cultures of embryonic chick cardiac cells. *Exp. Cell Res.* 112:265-273.
- Lazarides, E. 1980. Intermediate filaments as mechanical integrators of cellular space. *Nature (Lond.).* 283:249-256.
- Lazarides, E., and B. D. Hubbard. 1976. Immunological characterization of the subunit of the 100 Å filaments from muscle cells. *Proc. Natl. Acad. Sci. U. S. A.* 73:4344-4348.
- Maddy, A. H., and P. G. Kelly. 1971. Acetic acid as a solvent for erythrocyte membrane proteins. *Biochim. Biophys. Acta.* 241:290-301.
- Marchesi, S. L., E. Steers, V. T. Marchesi, and T. W. Tillack. 1970. Physical and chemical properties of a protein isolated from red cell membranes. *Biochemistry.* 9:50-57.
- Marchesi, V. T., and E. Steers. 1968. Selective solubilization of a protein component of the red cell membrane. *Science (Wash. D. C.).* 159:203-204.
- Marek, L. F., R. O. Kelley, B. D. Perdue, and P. O. Zamora. 1979. Role of the cytoskeleton in fibroblast attachment to poly-L-lysine treated substrates. *J. Cell Biol.* 83 (2, Pt. 2): 324a (Abstr.).
- Moon, H. M., T. Wisniewski, P. Merz, J. DeMartini, and H. M. Wisniewski. Partial purification of neurofilament subunits from bovine brains and studies on neurofilament assembly. *J. Cell Biol.* 89:560-567.
- O'Connor, C. M., D. R. Balzer, and E. Lazarides. 1979. Phosphorylation of subunit proteins of intermediate filaments from chicken muscle and nonmuscle cells. *Proc. Natl. Acad. Sci. U. S. A.* 76:819-823.
- O'Connor, C. M., D. L. Gard, and E. Lazarides. 1981. Phosphorylation of intermediate filament proteins by cAMP-dependent protein kinases. *Cell.* 23:135-143.
- O'Farrell, P. H. 1975. High resolution two-dimensional electrophoresis of proteins. *J. Biol. Chem.* 250:4007-4021.
- Osborn, M., R. E. Webster, and K. Weber. 1978. Individual microtubules viewed by immunofluorescence and electron microscopy in the same PK2 cell. *J. Cell Biol.* 77:R27-R34.
- Rueger, D. C., J. S. Huston, D. Dahl, and A. Bignami. 1979. Formation of 100 Å filaments from purified glial fibrillary acidic protein in vitro. *J. Mol. Biol.* 135:53-68.
- Schlaepfer, W. W. 1978. Observations on the disassembly of isolated mammalian neurofilaments. *J. Cell Biol.* 76:50-56.
- Shelton, K. R., L. L. Higgins, D. L. Cochran, J. J. Ruffolo, and P. M. Egle. 1980. Nuclear lamins of erythrocyte and liver. *J. Biol. Chem.* 255:10978-10983.
- Small, J. V., and A. Sobieszek. 1977. Studies on the function and composition of the 10-nm (100-Å) filaments of vertebrate smooth muscle. *J. Cell Sci.* 23:243-268.
- Sommer, J. R. 1977. To cationize glass. *J. Cell Biol.* 75 (2, Pt. 2): 245a (Abstr.).
- Starger, J. M., W. E. Brown, A. E. Goldman, and R. D. Goldman. 1978. Biochemical and immunological analysis of rapidly purified 10-nm filaments from baby hamster kidney (BHK-21) cells. *J. Cell Biol.* 78:93-109.
- Steck, T. L., and J. Yu. 1973. Selective solubilization of proteins from red blood cell membranes by protein perturbants. *J. Supramol. Struct.* 1:220-232.
- Steinert, P. M., W. W. Idler, F. Cabral, M. M. Gottesman, and R. D. Goldman. 1981. In

- vitro assembly of homopolymer and copolymer filaments from intermediate filament subunits of muscle and fibroblastic cells. *Proc. Natl. Acad. Sci. U. S. A.* 78:3692-3696.
58. Tuszynski, G. P., E. D. Frank, C. H. Damsky, C. A. Buck, and L. Warren. 1979. The detection of smooth muscle desmin-like protein in BHK₂₁/C₁₃ fibroblasts. *J. Biol. Chem.* 254:6138-6143.
59. Virtanen, I., M. Kurkinen, and V.-P. Lehto. 1979. Nucleus-anchoring cytoskeleton in chicken red blood cells. *Cell Biol. Int. Rep.* 3:157-162.
60. Weise, M. J., and L.-N. L. Chan. 1978. Membrane protein synthesis in embryonic chick erythroid cells. *J. Biol. Chem.* 253:1892-1897.
61. Weise, M. J., and V. M. Ingram. 1976. Proteins and glycoproteins of membranes from developing chick red cells. *J. Biol. Chem.* 251:6667-6673.
62. Willard, M., and C. Simon. 1981. Antibody decoration of neurofilaments. *J. Cell Biol.* 89: 198-205.
63. Woodcock, C. L. F. 1980. Nucleus-associated intermediate filaments from chicken erythrocytes. *J. Cell Biol.* 85:881-889.
64. Yu, J., D. A. Fischman, and T. L. Steck. 1973. Selective solubilization of proteins and phospholipids from red blood cell membranes by nonionic detergents. *J. Supramol. Struct.* 1:233-248.
65. Zentgraf, H., B. Deumling, and W. W. Franke. 1969. Isolation and characterization of nuclei from bird erythrocytes. *Exp. Cell Res.* 56:333-337.
66. Zentgraf, H., B. Deumling, E.-D. Jarasch, and W. W. Franke. 1971. Nuclear membranes and plasma membranes from hen erythrocytes. I. Isolation, characterization, and comparison. *J. Biol. Chem.* 246:2986-2995.