

RESEARCH ARTICLE

Reproductive characteristics modify the association between global DNA methylation and breast cancer risk in a population-based sample of women

Lindsay J. Collin^{1*}, Lauren E. McCullough¹, Kathleen Conway², Alexandra J. White³, Xinran Xu⁴, Yoon Hee Cho⁵, Sumitra Shantakumar⁶, Susan L. Teitelbaum⁷, Alfred I. Neugut^{8,9}, Regina M. Santella¹⁰, Jia Chen^{7,11,12}, Marilie D. Gammon²

1 Department of Epidemiology, Emory University, Atlanta, GA, United States of America, **2** Department of Epidemiology, University of North Carolina, Chapel Hill, NC, United States of America, **3** Epidemiology Branch, National Institute of Environmental Health Science, Research Triangle Park, NC, United States of America, **4** Roche Product Development in Asia-Pacific, Shanghai, China, **5** Department of Biomedical and Pharmaceutical Sciences, University of Montana, Missoula, MT, United States of America, **6** Glaxo-Smith-Kline, Inc., Singapore, **7** Department of Preventive Medicine, Icahn School of Medicine at Mount Sinai, New York, NY, United States of America, **8** Department of Epidemiology, Columbia University, New York, NY, United States of America, **9** Department of Medicine, Columbia University, New York, NY, United States of America, **10** Department of Environmental Health, Columbia University, New York, NY, United States of America, **11** Department of Pediatrics, Icahn School of Medicine at Mount Sinai, New York, NY, United States of America, **12** Department of Oncological Science, Icahn School of Medicine at Mount Sinai, New York, NY, United States of America

* lindsay.jane.collin@emory.edu



OPEN ACCESS

Citation: Collin LJ, McCullough LE, Conway K, White AJ, Xu X, Cho YH, et al. (2019) Reproductive characteristics modify the association between global DNA methylation and breast cancer risk in a population-based sample of women. PLoS ONE 14 (2): e0210884. <https://doi.org/10.1371/journal.pone.0210884>

Editor: Aamir Ahmad, University of South Alabama Mitchell Cancer Institute, UNITED STATES

Received: August 28, 2018

Accepted: January 3, 2019

Published: February 14, 2019

Copyright: © 2019 Collin et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: Data cannot be made publicly available due to ethical and legal restrictions. Ethically, public availability would compromise participants confidentiality as the minimal data-set would include information on the age, sex, and reproductive characteristics of women in a specific geographic region within a limited time-frame. Legally, the investigation began in 1993 under Public Law 103-43, with funding and coordination provided by the National Cancer Institute (NCI), in collaboration with the National

Abstract

DNA methylation has been implicated in breast cancer aetiology, but little is known about whether reproductive history and DNA methylation interact to influence carcinogenesis. This study examined modification of the association between global DNA methylation and breast cancer risk by reproductive characteristics. A population-based case-control study assessed reproductive history in an interviewer-administered questionnaire. Global DNA methylation was measured from white blood cell DNA using luminometric methylation assay (LUMA) and pyrosequencing assay (long interspersed elements-1 (LINE-1)). We estimated adjusted odds ratios (ORs) and 95% confidence intervals (CIs) among 1 070 breast cancer cases and 1 110 population-based controls. Effect modification was assessed on additive and multiplicative scales. LUMA methylation was associated with elevated breast cancer risk across all strata (comparing the highest to the lowest quartile), but estimates were higher among women with age at menarche ≤ 12 years (OR = 2.87, 95%CI = 1.96–4.21) compared to >12 years (OR = 1.66, 95%CI = 1.20–2.29). We observed a 2-fold increase in the LUMA methylation-breast cancer association among women with age at first birth >23 years (OR = 2.62, 95%CI = 1.90–3.62) versus ≤ 23 years (OR = 1.32, 95% CI = 0.84–2.05). No modification was evident for parity or lactation. Age at menarche and age at first birth may be modifiers of the association between global DNA methylation and breast cancer risk.

Institute of Environmental Health Sciences (NIEHS). Given that our data are sensitive, federally mandated, and were collected prior to NIH public access policies, we are restricted not to disclose data to a third party. While data sharing agreements prohibit us from making the data set publicly available, access may be granted by request to the Study PI (Dr. Marilie Gammon, gammon@unc.edu) who is ultimately responsible for maintaining the confidentiality and integrity of the data. Per Institutional Review Board and data agreements that occurred at the beginning of the study, Dr. Marilie Gammon is required to be the point of contact for data requests. Approvals should be obtained by the Institutional Review Boards of the University of North Carolina at Chapel Hill (irb_questions@unc.edu), Columbia University (irboffice@columbia.edu), and Mt. Sinai (IRB@mssm.edu).

Funding: This work was supported, in part, by grants from the National Cancer Institute at the National Institutes of Health [Grant numbers U01CA/ES66572 (PI: MG), R01CA109753 (PI: JC) and 3R01CA109753-04S1 (PI JC)] and the intramural program of the NIH Z99 ES99999. The funders had no role in the study design, data collection, data analysis, decision to publish, or preparation of the manuscript. X. Xu is employed by Roche Product Development. Roche Product Development provided support in the form of salary for author XX, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. S. Shantakumar is employed by Glaxo-Smith-Kline, Inc. Glaxo-Smith-Kline, Inc. provided support in the form of salary for author SS, but did not have any additional role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript. The specific roles of these authors are articulated in the 'author contributions' section.

Competing interests: I have read the journal's policy and the authors of this manuscript have the following competing interests: S. Shantakumar is employed at GlaxoSmithKline and owns shares. X. Xu is employed by Roche Product Development. A. Neugut serves as a consultant for Otsuka Pharmaceuticals, Pfizer, Eisai, Hospira, Teva, and United Biosource Corp, and is also a member of the medical advisory board of EHE Intl. There are no patents, products in development or marketed products to declare. This does not alter our adherence to all the PLOS ONE policies on sharing data and materials.

Introduction

Breast cancer remains the most commonly diagnosed invasive cancer among women in the United States [1]. Breast carcinogenesis is a multi-stage process involving both genetic and epigenetic changes [2], the latter defined as changes in gene expression, independent of modifications to the gene sequence [3]. Epigenetic aberrations have been implicated in breast carcinogenesis [4–6] and, unlike genetic alterations, these modifications may be altered across the lifespan by both exogenous and endogenous factors [7]. One commonly studied epigenetic modification is DNA methylation, characterized as the addition of a methyl group (-CH₃) to the 5-carbon position of cytosine CpG dinucleotides [8]. DNA methylation is a regulatory mechanism for gene expression and may influence cancer development through activation or silencing of genes involved in tumorigenesis [8]. In addition to gene-specific DNA methylation, global DNA hypomethylation in regions that are normally methylated (i.e., tandem repeats or transposable elements) can lead to genomic instability and oncogene expression in breast tissue [8]. We previously reported that breast cancer risk was increased among women in the highest level of luminometric methylation assay (LUMA), but no association was observed when we considered long interspersed elements-1 (LINE-1) methylation in white blood cell (WBC) DNA [9].

Reproductive history, which comprises a group of well-established risk factors for breast cancer, contributes modestly to overall risk [10]; affecting the lifetime cumulative exposure of the breast epithelium to endogenous ovarian steroid hormones [11, 12]. Prior studies have shown that age at first birth and parity alter the methylation profile of CpGs related to ER α expression in non-malignant breast tissue [13]. Another previous investigation in 376 healthy women showed reproductive characteristics were inversely associated with markers of global methylation as measured by LUMA [14]. However, this study was limited to menarcheal age and did not explore the potential links to breast cancer incidence. While both reproductive characteristics and aberrant DNA methylation are known to be relevant to breast cancer, no previous research has investigated their potential interaction with respect to breast cancer risk. Cyclical DNA methylation/demethylation may be activated by steroid hormones [15], modulating the expression of steroid receptors. It is therefore biologically plausible that reproductive characteristics and DNA methylation have a joint effect on breast carcinogenesis.

In this current study, we sought to expand our previous work on the association between global DNA methylation and breast cancer risk by examining the potential modification of the association between global DNA methylation and breast cancer risk by age at menarche, age at first birth, parity, and lactation. We used two independent methods to assess global DNA methylation in WBC DNA. The first is a pyrosequencing assay, which measures methylation levels in repetitive, LINE-1; this method is commonly used as a surrogate marker of genome-wide DNA methylation [16, 17]. The second method, LUMA, measures the levels of 5-^mC in the C^mCGG motif which allows approximation of methylation levels at gene promoters rather than the total genome [9]. We also investigated whether reproductive characteristics were associated with global methylation in controls. We hypothesized that reproductive factors would modify the association between global DNA methylation and breast cancer risk.

Materials and methods

Existing resources from the population-based case-control Long Island Breast Cancer Study Project (LIBCSP) were used to conduct our ancillary study. Details of participant recruitment, study design, and cohort characteristics have been described elsewhere [18]. All participating institutions obtained Institutional Review Board approval.

Study population and study interview

Study participants in the LIBCSP were English-speaking female residents of Nassau and Suffolk counties of Long Island, New York [18]. Cases were defined as women with a newly diagnosed first primary breast cancer between August 1, 1996 and July 31, 1997, and were identified through daily or weekly contact with the local hospital pathology departments. Population-based controls were randomly selected using random digit dialling for women under age 65 and Health Care Finance Administration rosters for women ages 65 and older. Controls were frequency matched to the expected age distribution of case women by 5-year age groups.

Interviews were completed for 82.1% (n = 1 508) of eligible cases and 62.8% (n = 1 556) of controls [18] (Fig 1). As abstracted from the medical/pathology records of the case women, 84.4% (n = 1 273) were diagnosed with invasive breast cancer, and 73.3% (of the 990 for whom this information was available) presented with estrogen receptor positive (ER+) breast cancer. Also, among all case-control participants, age at diagnosis (at identification for controls) ranged from 20 to 98 years, 67.1% were determined to be postmenopausal, and 92.8% were white (which is consistent with the underlying race distribution of the study counties at the time of




Fig 1. Selection and response of study population based on inclusion criteria of subjects in the population-based case control study of women age 20–98, in the Long Island Breast Cancer Study Project.

<https://doi.org/10.1371/journal.pone.0210884.g001>

data collection) [18] (Table 1). We previously reported that, among the LIBCSP population, increased breast cancer risk was associated with: early age at menarche, parity or nulliparity, late age at first birth, and little or no lactation [19]; and higher average lifetime alcohol intake [20]. Age- and menopausal-specific associations were also found for: use of oral contraceptives and hormone replacement therapy [21], adult weight gain [22]; increased body size and little or no physical activity [23, 24].

At the time of the case-control interview (which for cases occurred approximately two to three months after diagnosis), trained and licensed phlebotomists/nurses obtained a non-fasting blood sample from 73.1% of cases (N = 1 102) and 73.3% (N = 1 140) of control participants, which for most women occurred prior to adjuvant therapy. Study participants who agreed to donate a blood sample were generally younger than those who declined [18]. DNA was isolated from the donated blood samples using methods previously described [25].

Written informed consent was obtained from all participants prior to the study questionnaire, and prior to phlebotomy.

Table 1. Distribution of demographic and clinical characteristics by case-control status among the 1070 cases and 1110 controls with available global methylation data, Long Island Breast Cancer Study Project.

	Case (1070)	Control (1110)
	N (%)	N (%)
Age at diagnosis		
<50 years	294 (27.5)	371 (33.4)
≥50 years	776 (72.5)	739 (66.6)
Menopausal status		
Premenopausal	376 (35.1)	376 (33.9)
Postmenopausal	703 (65.7)	688 (66.1)
Missing	25	46
Cigarette Smoking		
Never	491 (45.9)	502 (45.3)
Current	205 (19.1)	209 (18.9)
Former	374 (35.0)	397 (35.8)
Missing	0	2
Age at menarche		
Less than 12 years	457 (43.1)	484 (43.9)
12 years or older	604 (56.9)	619 (56.1)
Missing	9	7
Parity		
Nulliparous	140 (13.1)	124 (11.2)
Parous	930 (86.9)	986 (88.8)
Age at first birth (among parous women only)		
Less than 23 years	314 (33.8)	355 (36.0)
23–27 years	310 (33.4)	345 (35.0)
Over 27 years	305 (32.8)	286 (29.0)
Missing	1	0
Lactation (among parous women only)		
Never	566 (60.9)	580 (58.8)
Any	364 (39.1)	406 (41.2)
ER Status		
Positive	422 (61.1)	
Negative	269 (38.9)	
Missing	379	

<https://doi.org/10.1371/journal.pone.0210884.t001>

Global methylation

Details of global methylation assessment within the LIBCSP study population have been described in detail [9]. Briefly, LUMA methylation levels were expressed as a percentage based on the following equation: $\text{methylation (\%)} = [1 - (\text{HpaII } \Sigma G/\Sigma T)/(\text{MspI } \Sigma G/\Sigma T)] * 100$. [26] LINE-1 methylation levels were assessed using a previously validated pyrosequencing assay to assess 4 CpG sites in its promoter region [27, 28]. Methylation status at each of the four loci were analysed individual as a T/C single nucleotide polymorphism using QCpG software (Qiagen) and then averaged to provide an overall percentage 5^mC status. As a quality control measure, to examine possible batch effects, samples were randomly selected for replication. Cases and controls were assayed simultaneously with the laboratory personnel blinded to both case-control status and quality control status. Of the 1 102 cases and 1 140 controls with available blood, 1 070 cases (97.1%) and 1 110 controls (97.4%) had valid global methylation data and were included in the study reported here.

Reproductive characteristics

The reproductive characteristics of interest for our study were age at menarche, parity, age at first birth, and lactation. These variables were collected as part of the main 100-minute interviewer-administered questionnaire, which included a reproductive calendar to enhance recall [29]. For the purposes of this ancillary study, age at menarche was dichotomized at 12 years of age, and parity was categorized as nulliparous versus parous. Among parous women, age at first birth was dichotomized at 23 years of age, and women with any lactation were compared to women without lactation. These categorizations were based on previously published data in our study population [19], optimization of cell counts, and incorporation of biologic plausibility. We explored a three-level category for age at first birth (≤ 23 , 23–27, ≥ 27 years of age), but observed similar effect estimates in the 23–27 and ≥ 27 categories. We therefore collapsed these two groups in our final analysis.

Statistical methods

Using unconditional logistic regression we first estimated odds ratios (ORs) and 95% confidence intervals (CI) for the associations between reproductive characteristics and global methylation levels, among control women only ($n = 1\ 110$, 88.8% parous), given previous evidence that women with breast cancer (compared to women without breast cancer) are likely to have lower global methylation levels [30]. Then, using information from both cases and controls (2 180 women, 87.9% parous), we evaluated whether the association between global methylation and breast cancer risk was modified by reproductive history. Finally, as a sensitivity analysis, we also explored whether reproductive characteristics modified the association between global DNA methylation and ER+ breast cancer, given that sex-steroid hormones are a potential unifying mechanism for DNA methylation, reproductive characteristics and breast cancer risk [15].

We estimated the odds of breast cancer within quartiles of LUMA and LINE-1 global methylation markers and within strata of each of the four reproductive characteristics. For associations stratified by age at menarche and parity, models included all women ($n = 2\ 180$), but for age at first birth or lactation, models included parous women only ($n = 1\ 916$). Quartiles (Q) of LUMA and LINE-1 methylation were categorized based on distribution of the percent methylation among controls. Given that LUMA, a global measurement of promoter methylation, was positively associated with overall breast cancer risk in our study population (9), we used the lowest quartile as the referent category. In contrast, LINE-1 hypomethylation is hypothesized

to be a marker of decreased genomic integrity, we therefore used the highest quartile of LINE-1 as the referent category.

Effect modification between global methylation and reproductive characteristics in association with breast cancer incidence was assessed on multiplicative and additive scales. Multiplicative interactions were evaluated using stratification and by comparing nested models with and without the interaction term using a Likelihood Ratio Test (*a priori* p -value < 0.05). Additive interactions were evaluated using a common referent and the relative excess risk due to interaction (RERI) [31, 32] and corresponding 95% CIs calculated using the delta method [32].

For all models, potential confounders were first selected using directed acyclic graphs (DAGs). Potential covariates included: family history of breast cancer (no/yes), history of benign breast disease (no/yes), race (white/black/other), use of oral contraceptives (never/ever), body mass index (BMI, < 25 kg/m², 25–29.9 kg/m², ≥ 30 kg/m²), recreational physical activity (RPA, high/moderate/inactive), alcohol history (never/ever), and smoking history (never/former/current). We further considered each possible confounder for inclusion in the final models based on a $> 10\%$ change in estimate approach [33], but none met this criteria. Thus, final models presented here only include the frequency matching factor age (5-year age groups).

Results

Among control women, we observed no associations between any reproductive characteristic and LUMA or LINE-1 methylation levels. Observed estimates were at or near the null, with all confidence intervals including the null (S1 Table).

Age at menarche

As shown in Table 2, among women with age of menarche ≤ 12 years of age, breast cancer risk increased within each increasing quartile of LUMA methylation, compared to the lowest quartile (OR_{Q2} = 1.47, 95%CI = 0.98–2.20; OR_{Q3} = 2.23, 95%CI = 1.51–3.29; OR_{Q4} = 2.87, 95%CI = 1.96–4.21). A non-monotonic increase in breast cancer risk with increasing LUMA methylation was also observed among those with age at menarche > 12 years of age, although the effect estimate was less pronounced (OR = 1.66, 95%CI = 1.20–2.29, comparing Q4 vs. Q1). This effect modification of the LUMA methylation-breast cancer association by age at menarche was evident on the multiplicative ($p_{\text{interaction}} = 0.05$), but not the additive scale [the RERI for additive interaction comparing those reporting an age at menarche ≤ 12 years and the highest level of global methylation assessed by LUMA (doubly exposed) to those with age at menarche > 12 years and the lowest LUMA methylation levels (referent) was 0.19 (95%CI = -0.38–0.76)], [S2 Table].

As shown in Table 3, the association between LINE-1 methylation and breast cancer risk (comparing Q1 vs. Q4) did not appear to differ between those with age at menarche ≤ 12 years of age (OR = 1.04, 95%CI = 0.72–1.49) and those who reported an age at menarche > 12 years (OR = 1.08, 95%CI = 0.78–1.45).

Parity

The association between LUMA methylation and breast cancer did not appear to differ substantially between strata of parity. When comparing the highest to the lowest quartile, we observed similar ORs between LUMA methylation and breast cancer risk among nulliparous (OR = 2.13, 95%CI = 1.04–4.37) and parous (OR = 2.08, 95%CI = 1.60–2.69) women

Table 2. Age-adjusted odds ratios (ORs) and 95% confidence intervals (CIs) for the multiplicative interaction between the luminometric methylation assay (LUMA) and reproductive characteristics (age at menarche, age at first birth, parity, lactation) with breast cancer risk in a population-based sample of 2180 women with available global methylation data, Long Island Breast Cancer Study Project.

LUMA (Quartiles)	Cases/ Controls	OR	95% CI	Cases/ Controls	OR	95% CI
Age at Menarche						
>12 years			≤12 years			
Q1 (<0.43)	115/142	1.00	reference	66/131	1.00	reference
Q2 (0.43<0.56)	92/156	0.72	0.50–1.02	90/118	1.47	0.98–2.20
Q3 (0.56<0.66)	170/154	1.36	0.97–1.89	133/119	2.23	1.51–3.29
Q4 (≥0.66)	215/161	1.66	1.20–2.29	165/113	2.87	1.96–4.21
<i>multiplicative p = 0.05</i>						
Parity						
Parous			Nulliparous			
Q1 (<0.43)	160/250	1.00	reference	23/26	1.00	reference
Q2 (0.43<0.56)	161/240	1.02	0.77–1.36	22/35	0.74	0.34–1.63
Q3 (0.56<0.66)	274/244	1.74	1.34–2.27	32/30	1.32	0.61–2.83
Q4 (≥0.66)	324/243	2.08	1.60–2.69	59/33	2.13	1.04–4.37
<i>multiplicative p = 0.70</i>						
Age at First Birth						
≤23 years			>23 years			
Q1 (<0.43)	58/82	1.00	reference	102/168	1.00	reference
Q2 (0.43<0.56)	61/88	0.93	0.58–1.49	99/152	1.06	0.74–1.51
Q3 (0.56<0.66)	101/84	1.65	1.05–2.58	173/160	1.78	1.28–2.47
Q4 (≥0.66)	91/97	1.32	0.84–2.05	233/146	2.62	1.90–3.62
<i>multiplicative p = 0.02</i>						
Lactation						
Any			Never			
Q1 (<0.43)	98/148	1.00	reference	62/102	1.00	reference
Q2 (0.43<0.56)	98/136	1.06	0.74–1.53	63/104	0.97	0.62–1.51
Q3 (0.56<0.66)	172/142	1.83	1.30–2.57	102/102	1.62	1.07–2.47
Q4 (≥0.66)	191/150	1.91	1.37–2.67	122/93	2.35	1.55–3.55
<i>multiplicative p = 0.50</i>						

<https://doi.org/10.1371/journal.pone.0210884.t002>

(Table 2). Similarly, parity did not appear to modify the association between LINE-1 methylation and breast cancer risk (Table 3).

Age at first birth

As shown in Table 2, among women with an age at first birth >23 years the risk of breast cancer increased with increasing quartiles of LUMA methylation (OR_{Q2 vs Q1} = 1.06, 95% CI = 0.74–1.51; OR_{Q3 vs Q1} = 1.78, 95%CI = 1.28–2.47; OR_{Q4 vs Q1} = 2.62, 95%CI = 1.90–3.62). By comparison, among women reporting early age at first birth (≤23 years) the association with breast cancer among women in the highest LUMA methylation quartile was less pronounced and included the null value (OR = 1.32, 95%CI = 0.84–2.05). This effect modification by age at first birth on the LUMA methylation–breast cancer association was evident on the multiplicative ($p_{\text{interaction}} = 0.02$) and additive scales [the observed RERI was 1.07 (95% CI = 0.43–1.71) comparing women with a first birth >23 years of age and the highest quartile

Table 3. Age-adjusted odds ratios (ORs) and 95% confidence intervals (CIs) for the multiplicative interaction between the long-interspersed elements-1 (LINE-1) assay and reproductive characteristics (age at menarche, age at first birth, parity, lactation) with breast cancer risk in a population-based sample of 2180 women with available global methylation data, Long Island Breast Cancer Study Project.

LINE-1 (Quartiles)	Cases/Controls	OR	95% CI	Cases/Controls	OR	95% CI
Age at Menarche						
	>12 years			≤12 years		
Q4 (≥80.4)	154/160	1.00	reference	123/116	1.00	reference
Q3 (78.7<80.4)	135/141	1.02	0.73–1.41	101/133	0.72	0.50–1.02
Q2 (77.0<78.7)	149/148	1.06	0.77–1.46	110/123	0.83	0.58–1.19
Q1 (<77.0)	162/163	1.06	0.78–1.45	122/110	1.04	0.72–1.49
	<i>multiplicative p = 0.43</i>					
Parity						
	Parous			Nulliparous		
Q4 (≥80.4)	239/249	1.00	reference	40/27	1.00	reference
Q3 (78.7<80.4)	203/240	0.89	0.69–1.15	35/34	0.77	0.39–1.54
Q2 (77.0<78.7)	225/246	0.95	0.73–1.22	36/29	0.91	0.45–1.85
Q1 (<77.0)	257/243	1.12	0.87–1.44	29/32	0.62	0.30–1.26
	<i>multiplicative p = 0.41</i>					
Age at First Birth						
	≤23 years			>23 years		
Q4 (≥80.4)	82/90	1.00	reference	156/159	1.00	reference
Q3 (78.7<80.4)	62/84	0.84	0.54–1.32	141/156	0.92	0.67–1.26
Q2 (77.0<78.7)	72/91	0.88	0.57–1.35	153/155	0.99	0.73–1.36
Q1 (<77.0)	96/88	1.25	0.82–1.90	161/155	1.06	0.67–1.26
	<i>multiplicative p = 0.75</i>					
Lactation						
	Any			Never		
Q4 (≥80.4)	80/90	1.00	reference	159/159	1.00	reference
Q3 (78.7<80.4)	93/103	0.80	0.57–1.12	110/137	1.02	0.68–1.55
Q2 (77.0<78.7)	83/105	1.01	0.73–1.39	142/141	0.87	0.57–1.32
Q1 (<77.0)	107/105	1.11	0.80–1.52	150/138	1.15	0.77–1.73
	<i>multiplicative p = 0.53</i>					

<https://doi.org/10.1371/journal.pone.0210884.t003>

of LUMA methylation (doubly exposed) to those with a first birth ≤23 years and the lowest quartile of LUMA methylation (referent).

Age at first birth did not appear to modify the association between LINE-1 methylation and breast cancer risk (Table 3). For women in the lowest quartile of LINE-1 methylation, there was a 25% increased risk of breast cancer within the strata of age at first birth ≤23 years, (OR = 1.25, 95%CI = 0.82–1.90) as compared with a 6% increased risk within the strata of age at first birth >23 (OR = 1.06, 95%CI = 0.67–1.26). We also considered a three-category variable (<23, 23–27, >27) for age at first birth, but found similar effect estimates among those with a first birth between 23–27 years of age and those >27 years of age (S3 Table).

Lactation

The association between LUMA methylation and breast cancer risk was relatively homogenous across strata of lactation (Table 2). Comparing women in the highest quartile of LUMA methylation to the lowest, we observed around a 2-fold increase in the risk for breast cancer within the never lactating strata (OR = 2.35, 95%CI 1.55–3.55), as well as the ever lactating strata

(OR = 1.91, 95%CI = 1.37–2.67) women. We additionally found no evidence for heterogeneity by lactation for the association between LINE-1 methylation and breast cancer (Table 3).

Sensitivity analysis

In models restricted to ER+ breast cancer only, our results were similar to those shown, although the effect estimates were less precise (S4 Table).

Discussion

In this population-based case-control study, we found that the association between global methylation, measured by LUMA in peripheral blood DNA, and breast cancer risk may depend on select reproductive characteristics, namely age at menarche and age at first birth (but not parity or lactation). Specifically, among women with age at menarche ≤ 12 years, there was nearly a 200% increase in breast cancer risk in association with high LUMA methylation levels; in contrast, the LUMA methylation-breast cancer association was only modestly increased by 66% among women with age at menarche > 12 years (multiplicative interaction $p = 0.05$). Similarly, among women with a first birth > 23 years, the association between high LUMA methylation levels and breast cancer risk was increased by greater than 150%; whereas among women with a first birth ≤ 23 years, the LUMA methylation-breast cancer association was increased by only 32% (multiplicative interaction p value = 0.02) No modification of the LINE-1 methylation-breast cancer association was observed by any of the four reproductive characteristics considered.

Numerous studies have been published on DNA methylation changes in breast tissue [34],[35]. Less is known about WBC DNA methylation and breast cancer risk, although several studies have emerged. Given that blood biomarkers can be measured repeatedly over time in large populations, they represent a useful tool for uncovering aetiology and ultimately early detection [36]. To our knowledge, this is the first population-based study to examine modification of the association between global methylation and breast cancer risk by multiple reproductive characteristics using two independent global methylation assays measured in WBC. DNA hypomethylation (estimated via LINE-1) increases genomic instability leading to the activation of oncogenes, is likely to operate early in carcinogenesis, and is known to be affected by multiple factors (including age) [37]. In contrast, LUMA provides an overall quantitation of methylation levels at gene promoters [9] which may more directly reflect gene expression, cell proliferation, and other pathways important in cancer promotion or progression. The specificity of this marker may make it a better surrogate for understanding the role of well-defined exposures (*i.e.*, reproductive characteristics) in cancer risk.

Contrary to our findings among control women, data from the EPIC-Italy sub-cohort showed that for each yearly increase in age at menarche, the odds of having genome-wide methylation (assessed by LUMA) below median level was increased by 32% (OR = 1.32, 95% CI = 1.14–1.53) [14]. The association between global methylation (assessed via LUMA) and breast cancer risk are mixed. While we previously estimated a 2.41-fold increased risk of breast cancer (OR_{quintile 5 vs. 1} = 95%CI = 1.83–3.16) comparing quintile 5 vs. 1,[9] others report greater breast cancer risk among women in the lowest tertile of LUMA (OR = 2.86, 95% CI = 1.85–4.44, comparing tertile 1 vs. 3),[38] and still others found no association between measures by LUMA and breast cancer risk [39]. While the distribution of LUMA methylation levels in these studies may be an important consideration, the observed inconsistencies highlight the importance of considering reproductive characteristics in combination with peripheral blood methylation and breast cancer [40]. Our data provide preliminary evidence that the

presence of early menarche or late age at first birth, and elevated LUMA methylation may be particularly deleterious for breast cancer risk.

The divergence of the epigenome as a function of age due to stochastic changes in methylation (a phenomena termed “epigenetic drift” [41, 42]) may contribute to tumorigenic transformation [42] including chromosomal instability, mutations, genetic recombination, large deletions, or translocations [43]. This divergence may be influenced by different environmental and lifestyle factors throughout an individual’s life course [41]. Exposure of the genome to factors such as smoking and alcohol consumption have been shown to modify the biologic aging process [7]. Chronic stress has also been found to lead to epigenetic alterations and accelerated aging of tissue samples [44]. Furthermore, environmental and other factors, including hormonal changes during an individual’s lifespan, may contribute to the epigenetic alterations [41]. Our hypotheses that reproductive characteristics modify the association of DNA methylation and breast cancer risk relate to the conceptual framework that hormonal changes contribute to epigenetic drift. Our findings, particularly around age-related reproductive characteristics, support this hypothesis.

Our study has many strengths, namely the availability of global methylation assessments in a well-characterized population-based case-control study with a relatively large sample size. The study included detailed exposure assessment, particularly for reproductive history. Questionnaires were administered by trained interviewers using validated methods to mitigate recall error, [45] however, exposure was assessed retrospectively and may suffer from differential misclassification by case-control status. Nonetheless, it is unlikely that women knew their methylation status and thus any differential recall of reproductive characteristics would not likely bias the interaction parameters. Because global methylation was assessed in cases shortly after diagnosis, there is concern for reverse causality. However, in a nested case-control study, investigators of the Sister Study Cohort showed that aberrant DNA methylation (assessed in blood) was present in women some 5 years prior to diagnosis and was predictive of breast cancer risk [46]. Another consideration of the current study is the racial and ethnic homogeneity of our study population. This limits the generalizability of our findings to primarily white women, who are at the highest risk of developing breast cancer in the United States [47]. Further research in prospective studies with diverse study populations is needed to confirm our findings.

Conclusions

Using resources from a large population-based case-control study, we observed that high methylation levels, assessed using the LUMA platform, were differentially associated with breast cancer risk among women with an earlier age at menarche and a later age at first birth. This study provides etiologic insight into how age-related reproductive factors may influence breast cancer risk through its interaction with the DNA methylome. Our findings may also help in the identification of an early biomarker among women who are at an increased risk of breast cancer based on their reproductive history.

Supporting information

S1 Table. Evaluation of the association between reproductive characteristics and DNA methylation among controls.

(DOCX)

S2 Table. Evaluation of the modification of the effect of DNA methylation on breast cancer risk by reproductive characteristics on the additive scale.

(DOCX)

S3 Table. Use of three category age at first birth to evaluate effect modification of the association between DNA methylation and breast cancer risk.

(DOCX)

S4 Table. Sensitivity analysis of the modification of the effect of DNA methylation on breast cancer risk among ER+ breast cancer cases.

(DOCX)

Author Contributions

Conceptualization: Lauren E. McCullough.

Data curation: Xinran Xu.

Formal analysis: Lindsay J. Collin, Alexandra J. White.

Funding acquisition: Jia Chen, Marilie D. Gammon.

Investigation: Yoon Hee Cho, Sumitra Shantakumar, Susan L. Teitelbaum, Alfred I. Neugut, Regina M. Santella, Jia Chen, Marilie D. Gammon.

Methodology: Lauren E. McCullough, Kathleen Conway, Xinran Xu, Yoon Hee Cho, Sumitra Shantakumar, Susan L. Teitelbaum, Alfred I. Neugut, Regina M. Santella, Jia Chen.

Project administration: Jia Chen.

Supervision: Susan L. Teitelbaum, Marilie D. Gammon.

Validation: Xinran Xu.

Writing – original draft: Lindsay J. Collin, Lauren E. McCullough.

Writing – review & editing: Lindsay J. Collin, Lauren E. McCullough, Kathleen Conway, Alexandra J. White, Xinran Xu, Yoon Hee Cho, Sumitra Shantakumar, Susan L. Teitelbaum, Alfred I. Neugut, Regina M. Santella, Jia Chen, Marilie D. Gammon.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. *CA: Cancer J Clin*. 2016; 66(1):7–30.
2. Lustberg MB, Ramaswamy B. Epigenetic Therapy in Breast Cancer. *Curr Breast Cancer Rep*. 2011; 3(1):34–43. <https://doi.org/10.1007/s12609-010-0034-0> PMID: 23097683
3. Widschwendter M, Jones PA. DNA methylation and breast carcinogenesis. *Oncogene*. 2002; 21(35):5462–82. <https://doi.org/10.1038/sj.onc.1205606> PMID: 12154408
4. Jovanovic J, Rønneberg JA, Tost J, Kristensen V. The epigenetics of breast cancer. *Molecular Oncol*. 2010; 4(3):242–54.
5. Baylin SB, Jones PA. A decade of exploring the cancer epigenome—biological and translational implications. *Nat Rev Cancer*. 2011; 11(10):726. <https://doi.org/10.1038/nrc3130> PMID: 21941284
6. Huang Y, Nayak S, Jankowitz R, Davidson NE, Oesterreich S. Epigenetics in breast cancer: what's new? *Breast Cancer Res*. 2011; 13(6):225. <https://doi.org/10.1186/bcr2925> PMID: 22078060
7. Beach SR, Dogan MV, Lei MK, Cutrona CE, Gerrard M, Gibbons FX, et al. Methylomic Aging as a Window onto the Influence of Lifestyle: Tobacco and Alcohol Use Alter the Rate of Biological Aging. *J Am Geriatr Soc*. 2015; 63(12):2519–25. <https://doi.org/10.1111/jgs.13830> PMID: 26566992
8. Basse C, Arock M. The increasing roles of epigenetics in breast cancer: Implications for pathogenicity, biomarkers, prevention and treatment. *Int J Cancer*. 2015; 137(12):2785–94. <https://doi.org/10.1002/ijc.29347> PMID: 25410431

9. Xu X, Gammon MD, Hernandez-Vargas H, Herceg Z, Wetmur JG, Teitelbaum SL, et al. DNA methylation in peripheral blood measured by LUMA is associated with breast cancer in a population-based study. *The FASEB Journal*. 2012; 26(6):2657–66. <https://doi.org/10.1096/fj.11-197251> PMID: [22371529](https://pubmed.ncbi.nlm.nih.gov/22371529/)
10. Madigan MP, Ziegler RG, Benichou J, Byrne C, Hoover RN. Proportion of breast cancer cases in the United States explained by well-established risk factors. *JNCI: J Natl Cancer Inst*. 1995; 87(22):1681–5. PMID: [7473816](https://pubmed.ncbi.nlm.nih.gov/7473816/)
11. Cancer CGoHFIB. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50 302 women with breast cancer and 96 973 women without the disease. *The Lancet*. 2002; 360(9328):187–95.
12. Ban KA, Godellas CV. Epidemiology of breast cancer. *Surg Oncol Clin N Amer*. 2014; 23(3):409–22.
13. Izadi P, Noruzinia M, Fereidooni F, Mostakhdemine Hosseini Z, Kamali F. Epigenetic marks in estrogen receptor alpha CpG island correlate with some reproductive risk factors in breast cancer. *Mol Biol Rep*. 2014; 41(11):7607–12. <https://doi.org/10.1007/s11033-014-3650-3> PMID: [25135164](https://pubmed.ncbi.nlm.nih.gov/25135164/)
14. Demetriou CA, Chen J, Polidoro S, Van Veldhoven K, Cuenin C, Campanella G, et al. Methylome analysis and epigenetic changes associated with menarcheal age. *PLoS One*. 2013; 8(11):e79391. <https://doi.org/10.1371/journal.pone.0079391> PMID: [24278132](https://pubmed.ncbi.nlm.nih.gov/24278132/)
15. Metivier R, Gallais R, Tiffocche C, Le Peron C, Jurkowska RZ, Carmouche RP, et al. Cyclical DNA methylation of a transcriptionally active promoter. *Nature*. 2008; 452(7183):45–50. <https://doi.org/10.1038/nature06544> PMID: [18322525](https://pubmed.ncbi.nlm.nih.gov/18322525/)
16. Weisenberger DJ, Campan M, Long TI, Kim M, Woods C, Fiala E, et al. Analysis of repetitive element DNA methylation by MethyLight. *Nucleic Acids Res*. 2005; 33(21):6823–36. <https://doi.org/10.1093/nar/gki987> PMID: [16326863](https://pubmed.ncbi.nlm.nih.gov/16326863/)
17. Yang AS, Estéicio MR, Doshi K, Kondo Y, Tajara EH, Issa JPJ. A simple method for estimating global DNA methylation using bisulfite PCR of repetitive DNA elements. *Nucleic Acids Res*. 2004; 32(3):e38–e. <https://doi.org/10.1093/nar/gnh032> PMID: [14973332](https://pubmed.ncbi.nlm.nih.gov/14973332/)
18. Gammon MD, Neugut AI, Santella RM, Teitelbaum SL, Britton JA, Terry MB, et al. The Long Island Breast Cancer Study Project: description of a multi-institutional collaboration to identify environmental risk factors for breast cancer. *Breast Cancer Res Treat*. 2002; 74(3):235–54. PMID: [12206514](https://pubmed.ncbi.nlm.nih.gov/12206514/)
19. Shantakumar S, Terry MB, Teitelbaum SL, Britton JA, Millikan RC, Moorman PG, et al. Reproductive factors and breast cancer risk among older women. *Breast Cancer Res Treat*. 2007; 102(3):365–74. <https://doi.org/10.1007/s10549-006-9343-4> PMID: [17033925](https://pubmed.ncbi.nlm.nih.gov/17033925/)
20. Terry MB, Zhang FF, Kabat G, Britton JA, Teitelbaum SL, Neugut AI, et al. Lifetime alcohol intake and breast cancer risk. *Ann Epidemiol*. 2006; 16(3):230–40.
21. Shantakumar S, Terry MB, Paykin A, Teitelbaum SL, Britton JA, Moorman PG, et al. Age and menopausal effects of hormonal birth control and hormone replacement therapy in relation to breast cancer risk. *Am J Epidemiol*. 2007; 165(10):1187–98. <https://doi.org/10.1093/aje/kwm006> PMID: [17337757](https://pubmed.ncbi.nlm.nih.gov/17337757/)
22. Eng SM, Gammon MD, Terry MB, Kushi LH, Teitelbaum SL, Britton JA, et al. Body size changes in relation to postmenopausal breast cancer among women on Long Island, New York. *Am J Epidemiol*. 2005; 162(3):229–37. <https://doi.org/10.1093/aje/kwi195> PMID: [15987723](https://pubmed.ncbi.nlm.nih.gov/15987723/)
23. McCullough LE, Eng SM, Bradshaw PT, Cleveland RJ, Teitelbaum SL, Neugut AI, et al. Fat or fit: the joint effects of physical activity, weight gain, and body size on breast cancer risk. *Cancer*. 2012; 118(19):4860–8. <https://doi.org/10.1002/cncr.27433> PMID: [22733561](https://pubmed.ncbi.nlm.nih.gov/22733561/)
24. McClain KM, McCullough LE, Bradshaw PT, Shantakumar S, Terry MB, Neugut AI, et al. Age-Specific Indicators of a Healthy Lifestyle and Postmenopausal Breast Cancer. *J Women's Health*. 2017.
25. Gammon MD, Santella RM, Neugut AI, Eng SM, Teitelbaum SL, Paykin A, et al. Environmental toxins and breast cancer on Long Island. I. Polycyclic aromatic hydrocarbon DNA adducts. *Cancer Epidemiol Prev Biomarkers*. 2002; 11(8):677–85.
26. Bjornsson HT, Sigurdsson MI, Fallin MD, Irizarry RA, Aspelund T, Cui H, et al. Intra-individual change over time in DNA methylation with familial clustering. *JAMA*. 2008; 299(24):2877–83. <https://doi.org/10.1001/jama.299.24.2877> PMID: [18577732](https://pubmed.ncbi.nlm.nih.gov/18577732/)
27. McShane LM, Altman DG, Sauerbrei W, Taube SE, Gion M, Clark GM. Reporting recommendations for tumor marker prognostic studies. *J Clin Oncol*. 2005; 23(36):9067–72.
28. Eads CA, Danenberg KD, Kawakami K, Saltz LB, Danenberg PV, Laird PW. CpG island hypermethylation in human colorectal tumors is not associated with DNA methyltransferase overexpression. *Cancer Res*. 1999; 59(10):2302–6. PMID: [10344733](https://pubmed.ncbi.nlm.nih.gov/10344733/)
29. Belli RF. The structure of autobiographical memory and the event history calendar: Potential improvements in the quality of retrospective reports in surveys. *Memory*. 1998; 6(4):383–406. <https://doi.org/10.1080/741942610> PMID: [9829098](https://pubmed.ncbi.nlm.nih.gov/9829098/)

30. Brennan K, Flanagan JM. Is there a link between genome-wide hypomethylation in blood and cancer risk? *Cancer Prev Res*. 2012; 5(12):1345–57.
31. Knol MJ, VanderWeele TJ. Recommendations for presenting analyses of effect modification and interaction. *Int J Epidemiol*. 2012; 41(2):514–20. <https://doi.org/10.1093/ije/dyr218> PMID: 22253321
32. Assmann SF, Hosmer DW, Lemeshow S, Mundt KA. Confidence intervals for measures of interaction. *Epidemiology*. 1996; 7(3):286–90. PMID: 8728443
33. David K, Mitchel K. Logistic regression: A self learning text. New York: Springer–Verlag Inc; 1994.
34. Taby R, Issa JP. Cancer epigenetics. *CA: Cancer J Clin*. 2010; 60(6):376–92.
35. Choi JY, James SR, Link PA, McCann SE, Hong CC, Davis W, et al. Association between global DNA hypomethylation in leukocytes and risk of breast cancer. *Carcinogenesis*. 2009; 30(11):1889–97. <https://doi.org/10.1093/carcin/bgp143> PMID: 19584139
36. Terry MB, Delgado-Cruzata L, Vin-Raviv N, Wu HC, Santella RM. DNA methylation in white blood cells: association with risk factors in epidemiologic studies. *Epigenetics*. 2011; 6(7):828–37. <https://doi.org/10.4161/epi.6.7.16500> PMID: 21636973
37. Costello JF, Plass C. Methylation matters. *J Med Genet*. 2001; 38(5):285–303. <https://doi.org/10.1136/jmg.38.5.285> PMID: 11333864
38. Kuchiba A, Iwasaki M, Ono H, Kasuga Y, Yokoyama S, Onuma H, et al. Global methylation levels in peripheral blood leukocyte DNA by LUMA and breast cancer: a case-control study in Japanese women. *Brit J Cancer*. 2014; 110(11):2765–71. <https://doi.org/10.1038/bjc.2014.223> PMID: 24786600
39. Delgado-Cruzata L, Wu HC, Perrin M, Liao Y, Kappil MA, Ferris JS, et al. Global DNA methylation levels in white blood cell DNA from sisters discordant for breast cancer from the New York site of the Breast Cancer Family Registry. *Epigenetics*. 2012; 7(8):868–74. <https://doi.org/10.4161/epi.20830> PMID: 22705975
40. Zhu ZZ, Hou L, Bollati V, Tarantini L, Marinelli B, Cantone L, et al. Predictors of global methylation levels in blood DNA of healthy subjects: a combined analysis. *Int J Epidemiol*. 2012; 41(1):126–39. <https://doi.org/10.1093/ije/dyq154> PMID: 20846947
41. Langevin SM, Pinney SM, Leung Y-K, Ho S-M. Does epigenetic drift contribute to age-related increases in breast cancer risk? *Epigenomics*. 2014; 6(4):367–9. <https://doi.org/10.2217/epi.14.28> PMID: 25333845
42. Fraga MF, Esteller M. Epigenetics and aging: the targets and the marks. *Trends in Genetics*. 2007; 23(8):413–8. <https://doi.org/10.1016/j.tig.2007.05.008> PMID: 17559965
43. Brennan K, Flanagan JM. Is there a link between genome-wide hypomethylation in blood and cancer risk? *Cancer Prev Res*. 2012; 5(12):1345–57.
44. Simons RL, Lei MK, Beach SR, Philibert RA, Cutrona CE, Gibbons FX, et al. Economic hardship and biological weathering: the epigenetics of aging in a US sample of black women. *Soc Sci Med*. 2016; 150:192–200. <https://doi.org/10.1016/j.socscimed.2015.12.001> PMID: 26765221
45. Caspi A, Moffitt TE, Thornton A, Freedman D, Amell JW, Harrington H, et al. The life history calendar: a research and clinical assessment method for collecting retrospective event-history data. *Int J Methods Psychiatr Res*. 1996.
46. Xu Z, Bolick SC, DeRoo LA, Weinberg CR, Sandler DP, Taylor JA. Epigenome-wide association study of breast cancer using prospectively collected sister study samples. *JNCI: J Natl Cancer Inst*. 2013; 105(10):694–700. <https://doi.org/10.1093/jnci/djt045> PMID: 23578854
47. DeSantis C, Ma J, Bryan L, Jemal A. Breast cancer statistics, 2013. *CA: Cancer J Clin*. 2014; 64(1):52–62.