

A microtubule dynamics reconstitutive convention

Kevin C. Slep

Department of Biology, University of North Carolina, Chapel Hill, NC 27599

In vitro reconstitution is the fundamental test for identification of the core components of a biological process. In this issue, Moriwaki and Goshima (2016). *J. Cell Biol.* <https://doi.org/10.1083/jcb.201604118> reconstitute all phases of microtubule dynamics through the inclusion of five key regulators and demonstrate that Polo kinase activity shifts the system from an interphase mode into an enhanced mitotic mode.

Microtubules are cytoskeletal structures that serve as tracks for motor-based intracellular transport and underlie the organization of biological apparatuses, including the mitotic spindle, cilia, and the phragmoplast. In vivo, microtubules are highly dynamic and interconvert between phases of polymerization, pause, and depolymerization. In vitro, microtubules are dynamic, but their behavior poorly mimics that observed in living cells. Identifying the regulators responsible for tuning in vitro microtubule dynamics to match the in vivo behavior has been a quest in the reconstitution field. In a new study published in this issue, Moriwaki and Goshima have now identified the five key components that are necessary for the recapitulation of all three phases of microtubule dynamics in vitro.

Nearly a century ago, mitotic spindle dynamics were observed. Determining the mechanisms that underlie these movements has remained a topic of active research. A key advance occurred in 1967 when Inoué and Sato (1967) used polarization microscopy to observe dynamic linear elements in the spindle. What were these linear elements made out of and how did they change their length? The discovery that colchicine disrupted the spindles positioned the field to use tritiated colchicine as a marker to identify tubulin, the protein formerly known as the colchicine binding protein (Borisy and Taylor, 1967). Tubulin is an obligate heterodimer consisting of α -tubulin and β -tubulin. The tubulin heterodimer polymerizes to form the microtubule lattice, mediated by both lateral and longitudinal interactions. The polarized nature of the tubulin heterodimer is propagated along the microtubule lattice, conferring the polymer with polarity: at one end, β -tubulin is exposed (the plus end), and at the other end, α -tubulin is exposed (the minus end). Each end of the polymer exhibits distinct dynamic behavior.

Given that tubulin was identified as the microtubule building block, scientists should then be able to use purified tubulin to generate dynamic polymers that recapitulate the dynamic behavior observed in cells. Key steps toward achieving this goal required identifying the requisite nucleotide (GTP), divalent ion (magnesium), and optimal buffer conditions. In the mid-1980s, Mitchison and Kirschner (1984) polymerized tubulin in vitro,

imaged fixed time points during the polymerization phase, and then diluted the system and imaged fixed time points during the depolymerization phase. They termed the nonequilibrium polymerization dynamics they observed “dynamic instability.” Under this process, microtubules do not reach a steady-state equilibrium length, but instead transition between phases of polymerization, pause, and depolymerization (Fig. 1 A). Underlying dynamic instability is the polymerization-dependent GTPase cycle of the tubulin heterodimer. Real-time visualization of dynamic instability came a few years later when dark field and superresolution video-enhanced differential interference contrast microscopy were used to observe microtubules transitioning between phases of polymerization and depolymerization (Horio and Hotani, 1986; Walker et al., 1988). Microtubule dynamics had been reconstituted, but there was one catch: the rates observed in vitro and the percentage of time microtubules spent in each phase (polymerization, depolymerization, and pause) did not correlate well with in vivo observations. Were there other factors required to regulate and tune dynamic instability? Enter the microtubule-associated proteins (MAPs).

Two key MAP families were identified that promoted microtubule polymerization and depolymerization. In 1987, Gard and Kirschner (1987) purified XMAP215 and characterized its ability to potentiate microtubule polymerization. In 1999, Desai et al. (1999) identified Kinesin-13 family members as microtubule depolymerization factors. With these factors now in hand, the Hyman laboratory set out to reconstitute microtubule dynamics using purified tubulin, XMAP215, and Kinesin-13 (Kinoshita et al., 2001). With the inclusion of these MAPs, microtubule dynamics started to approach in vivo rates, but the limited sampling of the microtubule pause state in vitro suggested that yet another factor was required to stabilize microtubules in the pause state.

Key steps forward included the identification of the CLIP-associating protein (CLASP) family that promotes the microtubule pause state (Akhmanova et al., 2001; Sousa et al., 2007), as well as a microtubule plus end tracking complex involving EB1 and SLAIN2/Sentin that recruits XMAP215 and CLASP to growing microtubule tips (van der Vaart et al., 2011; Li et al., 2012). With these molecular machines identified, the field was now positioned to test whether XMAP215, Kinesin-13, CLASP, Sentin, and EB1 could collectively reproduce microtubule dynamics in vitro. Moriwaki and Goshima (2016) have now addressed this challenge. Given the extent to which these regulators had been characterized in *Drosophila melanogaster* cell culture, the authors decided to use purified

Correspondence to Kevin C. Slep: kslep@bio.unc.edu

© 2016 Slep This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms>). After six months it is available under a Creative Commons license [Attribution–Noncommercial–Share Alike 3.0 Unported license, as described at <http://creativecommons.org/licenses/by-nc-sa/3.0/>].






Figure 1. **Five MAPs reconstitute all phases of microtubule dynamics in vitro.** (A) Diagram depicting the phases of microtubule dynamic instability, including polymerization (growth), pause, and depolymerization (shrinkage). Three proteins, XMAP215^{MSPS}, CLASP^{Mast/Orbit}, and Kinesin-13^{Klp10A}, promote polymerization, pause, and depolymerization, respectively. Although these regulators engage the microtubule directly, EB1 and Sentin recruit CLASP^{Mast/Orbit} and XMAP215^{MSPS} to polymerizing microtubule plus ends and affect their activity. (B) Microtubule dynamics diamond graphs showing in vitro microtubule behavior without (left) and with (right) Plk1^{Polo} treatment. The black diamond graph at left is overlaid on the purple graph at right and shown in dark gray for comparison.

Drosophila homologues for their in vitro analyses. This would enable them to compare and contrast cellular microtubule dynamics in wild-type and factor-depleted cultures with the effects of all or a subset of these MAPs on in vitro microtubule dynamics. They purified *Drosophila* XMAP215^{MSPS}, Kinesin-13^{Klp10A}, CLASP^{Mast/Orbit}, EB1, and Sentin, as well as fly tubulin.

To determine whether CLASP would promote microtubule pause, Moriwaki and Goshima (2016) first analyzed the behavior of CLASP and tubulin alone and found that it decreased the microtubule growth rate and catastrophe frequency as compared with a tubulin control. CLASP bound along microtubules and limited their overall dynamics. They next investigated whether inclusion of the CLASP binding protein EB1 would recruit CLASP to microtubule plus ends. EB1 did recruit CLASP to the microtubule plus end, but the relocalization of CLASP did not dramatically alter microtubule dynamics. The authors then examined CLASP in concert with all four MAPs: Sentin, EB1, XMAP215^{MSPS}, and Kinesin-13^{Klp10A}. With this ensemble of MAPs, Moriwaki and Goshima (2016) witnessed the full regimen of microtubule dynamics: microtubules interconverting between phases of polymerization, pause, and depolymerization with rates and frequencies that were on the order of those observed in cell culture (Fig. 1 B). Interestingly, many of the pause and rescue events correlated with sites on the microtubule lattice where CLASP was bound, highlighting the role of CLASP as a pause and rescue factor. Systematically

removing a single component from the milieu showed parallels to microtubule dynamics parameters observed when the corresponding factor was depleted from *Drosophila* S2 cells. These results indicate that these five MAPs play a core role in regulating microtubule dynamics and strongly suggest that their concerted action is not caused by the sum of their individual activities, but rather that interactions and synergy between the components is critical.

As cells transition from interphase into mitosis, the parameters of microtubule dynamic instability change. The polymerization rate of astral microtubules increases, as does the catastrophe frequency, whereas the rescue frequency decreases (Belmont et al., 1990; Rusan et al., 2001). If Moriwaki and Goshima (2016) had identified the core components required for dynamic instability, would there be other MAPs required to establish the mitotic parameters, or could there be a factor that modified the activity of the core five regulators? Although mitotic MAPs play key roles in this process, Moriwaki and Goshima (2016) inquired whether a mitotic kinase could modulate the activity of their reconstituted system. The prime candidate was the kinase Cdk1 that had been shown in cells to trigger the transition from interphase microtubule dynamics to mitotic microtubule dynamics (Verde et al., 1990). When the authors incubated Cdk1 and ATP with their five MAPs, Kinesin-13^{Klp10A} and XMAP215^{MSPS} were phosphorylated, but these posttranslational modifications had little effect on microtubule dynamics in

their reconstituted system. Perhaps Cdk1 was acting upstream of another mitotic kinase that directly drove the mitotic transition. Their next candidate was Polo kinase (Plk1). In vitro, Plk1 phosphorylated Kinesin-13^{Klp10A}, Sentin, and CLASP^{Mast/Orbit}. In this Polo-dependent phosphorylated state, the five-component system showed dramatically different microtubule dynamics. The polymerization rate increased, as did the catastrophe frequency, whereas the rescue frequency decreased (Fig. 1 B). The authors also noted that CLASP lattice association decreased and there was a concomitant decrease in the time microtubules spent in the pause state. The collective changes in microtubule dynamics the authors observed in their reconstituted system paralleled the way the parameters also changed in cells transitioning from interphase to mitosis. Collectively, this strongly suggests that these five components are indeed the core regulators of microtubule dynamic instability and that Polo kinase tunes the system for mitosis.

Moriwaki and Goshima (2016) have established a foundation upon which additional regulators of microtubule dynamics can be investigated. Of note, whereas the seven parameters of dynamic instability have been reconstituted, they do not perfectly align with in vivo rates and frequencies. Differences are caused, in part, by experimental constraints: not all components could be analyzed at physiological concentrations, crowding agents were not present, and there were likely limits imposed by protein activity as well as in vitro environmental constraints. Clearly limiting is the fact that only five MAPs were present. Adding higher order complexity through the addition of other regulators will be key for future studies. As the complexity of the reconstituted system grows and synergistic effects of components are observed, it will be important to approach the system with quantitative rigor and to systematically titrate components. What is the future for microtubule dynamics reconstitution? Key areas to explore include the identification and mechanistic analysis of phosphoregulatory sites on MAPs, analysis of tubulin posttranslational modifications on the systems' behavior, and the addition of other motors and MAPs (augmin, katanin, tetrameric kinesins, etc.). Incorporating other biological systems into the fray will also be a key trajectory: how do dynamic microtubules engage targets such as membrane anchor points or kinetochores, and how do cytoskeletal cross-linkers affect the concerted dynamics of the microtubule and actin network? Many researchers are actively tackling these questions, but the groundwork laid to date by numerous laboratories, including the work by Moriwaki and Goshima (2016), will help direct these efforts.

Acknowledgments

The author wishes to apologize to colleagues whose studies have not been cited due to space restrictions.

The Slep laboratory is funded by March of Dimes grant 1FY14247 and National Institutes of Health grant R03HD084980.

The author declares no competing financial interests.

Submitted: 19 October 2016

Accepted: 19 October 2016

References

- Akhmanova, A., C.C. Hoogenraad, K. Drabek, T. Stepanova, B. Dortland, T. Verkerk, W. Vermeulen, B.M. Burgering, C.I. De Zeeuw, F. Grosveld, and N. Galjart. 2001. Clasps are CLIP-115 and -170 associating proteins involved in the regional regulation of microtubule dynamics in motile fibroblasts. *Cell*. 104:923–935. [http://dx.doi.org/10.1016/S0092-8674\(01\)00288-4](http://dx.doi.org/10.1016/S0092-8674(01)00288-4)
- Belmont, L.D., A.A. Hyman, K.E. Sawin, and T.J. Mitchison. 1990. Real-time visualization of cell cycle-dependent changes in microtubule dynamics in cytoplasmic extracts. *Cell*. 62:579–589. [http://dx.doi.org/10.1016/0092-8674\(90\)90022-7](http://dx.doi.org/10.1016/0092-8674(90)90022-7)
- Borisy, G.G., and E.W. Taylor. 1967. The mechanism of action of colchicine. Colchicine binding to sea urchin eggs and the mitotic apparatus. *J. Cell Biol.* 34:535–548. <http://dx.doi.org/10.1083/jcb.34.2.535>
- Desai, A., S. Verma, T.J. Mitchison, and C.E. Walczak. 1999. Kin I kinesins are microtubule-destabilizing enzymes. *Cell*. 96:69–78. [http://dx.doi.org/10.1016/S0092-8674\(00\)80960-5](http://dx.doi.org/10.1016/S0092-8674(00)80960-5)
- Gard, D.L., and M.W. Kirschner. 1987. A microtubule-associated protein from *Xenopus* eggs that specifically promotes assembly at the plus-end. *J. Cell Biol.* 105:2203–2215. <http://dx.doi.org/10.1083/jcb.105.5.2203>
- Horio, T., and H. Hotani. 1986. Visualization of the dynamic instability of individual microtubules by dark-field microscopy. *Nature*. 321:605–607. <http://dx.doi.org/10.1038/321605a0>
- Inoué, S., and H. Sato. 1967. Cell motility by labile association of molecules. The nature of mitotic spindle fibers and their role in chromosome movement. *J. Gen. Physiol.* 50(Suppl):259–292. <http://dx.doi.org/10.1085/jgp.50.6.259>
- Kinoshita, K., I. Arnal, A. Desai, D.N. Drechsel, and A.A. Hyman. 2001. Reconstitution of physiological microtubule dynamics using purified components. *Science*. 294:1340–1343. <http://dx.doi.org/10.1126/science.1064629>
- Li, W., T. Moriwaki, T. Tani, T. Watanabe, K. Kaibuchi, and G. Goshima. 2012. Reconstitution of dynamic microtubules with *Drosophila* XMAP215, EB1, and Sentin. *J. Cell Biol.* 199:849–862. <http://dx.doi.org/10.1083/jcb.201206101>
- Mitchison, T., and M. Kirschner. 1984. Dynamic instability of microtubule growth. *Nature*. 312:237–242. <http://dx.doi.org/10.1038/312237a0>
- Moriwaki, T., and G. Goshima. 2016. Five factors can reconstitute all three phases of microtubule polymerization dynamics. *J. Cell Biol.* <https://doi.org/10.1083/jcb.201604118>
- Rusan, N.M., C.J. Fagerstrom, A.M. Yvon, and P. Wadsworth. 2001. Cell cycle-dependent changes in microtubule dynamics in living cells expressing green fluorescent protein- α tubulin. *Mol. Biol. Cell*. 12:971–980. <http://dx.doi.org/10.1091/mbc.12.4.971>
- Sousa, A., R. Reis, P. Sampaio, and C.E. Sunkel. 2007. The *Drosophila* CLASP homologue, Mast/Orbit regulates the dynamic behaviour of interphase microtubules by promoting the pause state. *Cell Motil. Cytoskeleton*. 64:605–620. <http://dx.doi.org/10.1002/cm.20208>
- van der Vaart, B., C. Manatschal, I. Grigoriev, V. Olieric, S.M. Gouveia, S. Bjelic, J. Demmers, I. Vorobjev, C.C. Hoogenraad, M.O. Steinmetz, and A. Akhmanova. 2011. SLAIN2 links microtubule plus end-tracking proteins and controls microtubule growth in interphase. *J. Cell Biol.* 193:1083–1099. <http://dx.doi.org/10.1083/jcb.201012179>
- Verde, F., J.C. Labbé, M. Dorée, and E. Karsenti. 1990. Regulation of microtubule dynamics by cdc2 protein kinase in cell-free extracts of *Xenopus* eggs. *Nature*. 343:233–238. <http://dx.doi.org/10.1038/343233a0>
- Walker, R.A., E.T. O'Brien, N.K. Pryer, M.F. Soboeiro, W.A. Voter, H.P. Erickson, and E.D. Salmon. 1988. Dynamic instability of individual microtubules analyzed by video light microscopy: rate constants and transition frequencies. *J. Cell Biol.* 107:1437–1448. <http://dx.doi.org/10.1083/jcb.107.4.1437>