

Research article

Open Access

## Does practicing hatha yoga satisfy recommendations for intensity of physical activity which improves and maintains health and cardiovascular fitness?

Marshall Hagins\*<sup>†1</sup>, Wendy Moore<sup>†2</sup> and Andrew Rundle<sup>†2</sup>

Address: <sup>1</sup>Division of Physical Therapy, Long Island University, One University Plaza, Brooklyn, NY 11201, USA and <sup>2</sup>Mailman School of Public Health, Columbia University, 622 W. 168th St. New York, NY 10032, USA

Email: Marshall Hagins\* - mhagins@liu.edu; Wendy Moore - mcdowell\_wendy@hotmail.com; Andrew Rundle - ag3r@columbia.edu

\* Corresponding author †Equal contributors

Published: 30 November 2007

Received: 13 June 2007

Accepted: 30 November 2007

*BMC Complementary and Alternative Medicine* 2007, **7**:40 doi:10.1186/1472-6882-7-40

This article is available from: <http://www.biomedcentral.com/1472-6882/7/40>

© 2007 Hagins et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

**Background:** Little is known about the metabolic and heart rate responses to a typical hatha yoga session. The purposes of this study were 1) to determine whether a typical yoga practice using various postures meets the current recommendations for levels of physical activity required to improve and maintain health and cardiovascular fitness; 2) to determine the reliability of metabolic costs of yoga across sessions; 3) to compare the metabolic costs of yoga practice to those of treadmill walking.

**Methods:** In this observational study, 20 intermediate-to-advanced level yoga practitioners, age  $31.4 \pm 8.3$  years, performed an exercise routine inside a human respiratory chamber (indirect calorimeter) while wearing heart rate monitors. The exercise routine consisted of 30 minutes of sitting, 56 minutes of beginner-level hatha yoga administered by video, and 10 minutes of treadmill walking at 3.2 and 4.8 kph each. Measures were mean oxygen consumption ( $VO_2$ ), heart rate (HR), percentage predicted maximal heart rate (%MHR), metabolic equivalents (METs), and energy expenditure (kcal). Seven subjects repeated the protocol so that measurement reliability could be established.

**Results:** Mean values across the entire yoga session for  $VO_2$ , HR, %MHR, METs, and energy/min were 0.6 L/kg/min; 93.2 beats/min; 49.4%; 2.5; and 3.2 kcal/min; respectively. Results of the ICCs (2,1) for mean values across the entire yoga session for kcal, METs, and %MHR were 0.979 and 0.973, and 0.865, respectively.

**Conclusion:** Metabolic costs of yoga averaged across the entire session represent low levels of physical activity, are similar to walking on a treadmill at 3.2 kph, and do not meet recommendations for levels of physical activity for improving or maintaining health or cardiovascular fitness. Yoga practice incorporating sun salutation postures exceeding the minimum bout of 10 minutes may contribute some portion of sufficiently intense physical activity to improve cardio-respiratory fitness in unfit or sedentary individuals. The measurement of energy expenditure across yoga sessions is highly reliable.

## Background

Physical activity conveys multiple well established health benefits including decreased rates of coronary artery disease [1-3], hypertension [4,5], non-insulin dependent diabetes mellitus [6], osteoporosis [7], colon cancer [8], anxiety and depression [9,10], as well as decreased risk of overall mortality [1]. Despite these clear benefits less than half of U.S. adults meet the current recommendations for physical activity [11].

The most recent recommendations from the American College of Sports Medicine (ACSM) and the American Heart Association (AHA) [12] suggest that to promote and maintain health all healthy adults aged 18–65 years old need moderate-intensity aerobic (endurance) activity for a minimum of 30 minutes on five days each week or vigorous-intensity activity for a minimum of 20 minutes on three days each week. These recommendations further suggest that combinations of moderate and vigorous exercise can be used to meet the requirements and that bouts of moderate-intensity exercise lasting a minimum of 10 minutes can be accumulated to achieve the 30 minute minimum. Recommendations for developing and maintaining cardiovascular fitness from the ACSM Position Stand recommend that exercise achieve 55–90% of the maximal heart rate, be performed 3–5 days per week, and last 20–60 minutes (continuous or intermittent) [13].

Yoga is an alternative form of physical activity which may assist in achieving recommended levels of physical activity for some individuals. Yoga is increasing in popularity [14-16] with a recent report suggesting that 15 million Americans have practiced yoga at least once in their lifetime [17]. Yoga classes are now offered at 75% of all U.S. health clubs [18], as well as at yoga studios and in private homes. Yoga may be attractive as an alternative to traditional aerobic and strength training programs because it requires little space, virtually no equipment, has limited harmful side effects [14,16,19,20] and, with its focus on relaxation, body awareness, and meditation provides a qualitatively different exercise experience which may be perceived as less strenuous and more pleasurable. Given these characteristics, yoga satisfies many of the conditions which have been shown to be strongly related to participation in physical activity, such as low perceived barrier to participation [21], being enjoyable [21], and having a low-to-moderate intensity [22].

Although yoga has received a considerable amount of study to date demonstrating significant cardio-respiratory [23-42], musculoskeletal [43,44] and metabolic health benefits [42], the degree to which the physical activity component of yoga may have contributed to these benefits has received virtually no attention. Surveying the English speaking literature available via Pubmed and

CINAHL revealed eight studies examining the metabolic costs of yoga. Of these studies [45-52], only one [51] examined the variety of postures seen in a typical yoga class (standing, sitting, and lying). Additionally, all previous studies used measurement techniques that physically encumber the subject by requiring masks or mouthpieces to measure oxygen consumption. Such techniques may alter the performance of the yoga activities and therefore provide invalid estimates of metabolic costs.

Therefore, the purposes of this study were: 1) to determine whether an approximately one hour yoga session incorporating a variety of postures meets the current recommendations for physical activity sufficient to improve and maintain health [12] and cardiovascular fitness [13] as measured within a respiratory chamber (indirect calorimeter); 2) to determine the reliability with which the metabolic costs of yoga can be measured; 3) to compare the metabolic costs of yoga practice to the metabolic costs of steady state treadmill walking at two speeds (3.2 and 4.8 kph).

## Methods

### Subjects

Subjects were recruited via fliers posted at various yoga studios around New York, NY, USA, and on the Columbia University Health Sciences campus. Each participant was administered an eligibility questionnaire by telephone prior to enrolment. Exclusion criteria included: a) age < 18 years; b) < 1 year experience doing yoga; b) negative response to the question "are you comfortable doing Sun Salutation and basic yoga standing poses such as Triangle, Warriors I and II, and Tree?"; c) negative response to the question "are you able to do inversions (headstand, shoulder stand) with or without a wall?"; d) presence of any physical conditions or limitations that would prevent the individual from performing the tasks involved in the study. These exclusion criteria ensured that the participants would be able to engage in the yoga routine safely and that the measurements taken would not be influenced by inexperience.

### Measurement Instruments

The study took place at the Obesity Research Center at St. Luke's-Roosevelt Hospital, New York, N.Y., USA. After providing informed consent, each participant was outfitted with one Actiheart™ activity and heart rate logger (Mini-Mitter, OR, USA). The Actiheart™ is a small, light-weight device that was affixed to each participant's chest using standard EKG pads. It collected continuous heart rate (HR) data and produced data points at 1-minute intervals, reflecting the average heart rate for each minute.

Each participant performed all activities inside the respiratory chamber. The performance characteristics of the res-

piratory chamber have been described elsewhere [53,54]. Briefly, the system is an air-tight, temperature-controlled room in which rates of oxygen consumption ( $VO_2$ ) and carbon dioxide production ( $VCO_2$ ) are calculated as the difference between the composition of entering and exiting air. Analysis uses magnetopneumatic oxygen (Magnos 4G) and carbon dioxide (Magnos 3G) analyzers (Hartmann & Braun, Frankfurt, Germany), and the data is displayed and stored by the on-line computer system. The respiratory chamber provided the average values for  $VO_2$ ,  $VCO_2$ , and energy expenditure (kcal/min) every two minutes. Energy expenditure was calculated via the respiratory chamber software using a modified Weir equation:  $EE = (3.94 \times VO_2) + (1.1 \times (VCO_2))$  [55].

Resting metabolic rate (RMR) for each participant was calculated by taking the mean energy expenditure during the last 20 minutes of the initial resting period. Each subsequent value for energy was divided by the RMR to determine metabolic equivalents (METs). To obtain the predicted maximum heart rate for each individual, the following formula was used:  $208 - (0.7 \times \text{age})$  [56]. Heart rate data was multiplied by 100 and divided by the predicted maximum heart rate to determine the percentage of maximum predicted heart rate (%MHR) achieved by each individual for each 1-minute interval.

**Procedures**

Subjects initially performed a 30 minute resting period during which they were seated and motionless except for the movements associated with reading (e.g., turning pages). Subjects then began a 56-minute beginner Ashtanga yoga DVD [57]. The yoga session began with 28 minutes of sun salutation poses (a standard series of moving poses found in many styles of yoga), followed by approximately 20 minutes of standing poses, and lastly, approximately 8 minutes of sitting and lying poses (relaxation). Table 1 describes the postures and time for each portion of the yoga session. For additional details see the commercially available "Ashtanga Yoga Beginner's Practice DVD [57]."

After completing the yoga practice, subjects walked on a treadmill at two different speeds in the following order: 3.2 kph (2 mph) and 4.8 kph (3 mph) for 10 minutes each. To determine across-session reliability, a convenience sample of seven participants returned between one week and one month later to perform an identical test protocol.

The first four minutes of HR and respiratory chamber data for each activity was discarded to allow each participant time to achieve a steady state of energy demand. Therefore, the data used for analysis included the final 20 min-

**Table 1: Description of yoga practice**

<b>Surya Namaskar (Sun Salutation)</b>	
28 minutes	
Pose	Name
Stand at Attention	<i>Samasthiti</i>
Hands above head	<i>Ekam</i>
Forward Bend	<i>Uttanasana</i>
Forward Bend/Head up	<i>Trini</i>
Down-dog	<i>Adhomukha Svanasana</i>
Four-legged staff pose	<i>Chaturanga Dandasana</i>
Up-dog	<i>Urdhvamukha Svanasana</i>
Chair	<i>Utkatasana</i>
Warrior I	<i>Virabhadrasana</i>
<b>Standing Poses</b>	
20 minutes	
Hands-to-feet	<i>Padahastasana</i>
Triangle	<i>Trikonasana</i>
Revolved triangle	<i>Parivrtta Trikonasana</i>
Side angle stretch	<i>Parivrtta Parsvakonasana</i>
Chest stretch	<i>Parsvottanasana</i>
<b>Sitting/Lying Poses</b>	
8 minutes	
Lotus	<i>Padmasana</i>
Corpse	<i>Savasana</i>

utes of the resting period, the final 52 minutes of the yoga session, and the final 6 minutes of each treadmill period.

**Statistical Analyses**

SAS version 9.1 (SAS Institute Inc., Cary, NC, USA.) was used for all statistical analysis. Description of the means and standard deviations for all variables across the entire time periods of the resting, yoga, and treadmill activities were determined. Data from the seven subjects who repeated the measurement session were tested for reliability using Intra Class Correlation Coefficients (ICC) (2,1). Two separate one-sample t-tests were used to determine whether the mean METs across the entire yoga session and across the sun salutation portion of the yoga practice differed from the minimum values for METs (3.0) recommended by ACSM and AHA [12]. Additionally, two separate one-sample t-tests were used to determine whether the mean %MHR across the entire yoga session and across the sun salutation portion of the yoga practice differed from the minimum values of 55% MHR recommended by the ACSM Position Stand [13].

Three separate ANOVAs were used to compare the mean rate of energy expenditure (kcal), METs, and %MHR across the following activities: 1) rest; 2) entire yoga session; 3) treadmill walking at 3.2 kph; 4) treadmill walking at 4.8 kph. The sun salutation portion of the yoga session was not included in the first set of ANOVAs as the sun salutation portion and the entire yoga session are correlated and as such an analysis would have violated the assumptions of the ANOVA. Consequently, in order to compare the identical variables (kcal, METs, and %MHR) during the sun salutation portion of the yoga session a second set of three separate ANOVAs were performed on 1) rest; 2) sun salutation portion of the yoga session; 3) treadmill walking at 3.2 kph; 4) treadmill walking at 4.8 kph.

Because the order of the activities was not randomized and the increase in metabolism from the yoga activity may have influenced the metabolic rate during the ensuing treadmill activity, post hoc testing was performed to determine if the METs and %MHR during the final three minutes of the yoga program (during supine relaxation) had returned to values equal to those of the resting state.

**Results**

Twenty participants were recruited (18 female and 2 male). Results of the t-tests and ANOVAs were not different whether the two men were excluded from analysis or not, and therefore we report values from all 20 participants. The mean demographic values for the participants were: age 31.4, (± 8.3) years, height 165.2 (± 7.9) cm, weight 64.3 (± 9.0) kg, and BMI 23.58 (± 3.03). Means and standard deviations of HR, calories, VO<sub>2</sub>, METs, and %MHR relative to the entire yoga session, sun salutation postures, non-sun salutation postures, sitting/lying postures, and treadmill walking are described in Table 2. In a single subject, two values of energy expenditure during the final two minutes of the yoga session (during supine resting) were identified as outliers due to technical error (METs increased by a factor of ~5 between minutes 50 and 51 during supine resting (1.31 to 5.06 METs)). The outliers were replaced using the subject's value for the minute immediately prior to their occurrence (e.g., value at

minute 50:1.31 METs). Figures 1 and 2 graphically describe the mean and SD of METs and %MHR, respectively, for each minute during the entire yoga session.

Results of the ICCs (2,1) (n = 7) for mean values across the entire yoga session for metabolic cost (kcal), METs, and %MHR were 0.979, 0.973, and 0.865, respectively. The ICC value for %MHR only was based on 5 subjects, rather than 7, due to technical errors arising during data collection.

Results of the one-sample t-tests comparing mean METs and %MHR to recommended values are presented in Table 3. The mean METs across the entire yoga session were significantly less than the moderate level of physical activity reference value of 3.0 METs recommended for improving and maintaining health by the ACSM and AHA [12]. However, the mean METs during the sun salutation portion of the yoga session was not significantly different from the reference value of 3.0 METs [12]. The mean %MHR during the entire yoga session was significantly less than the 55% of MHR described as a minimum threshold by the ACSM Position Stand [13] to achieve cardio-respiratory fitness. However, the mean %MHR during the sun salutation portion of the yoga session (24 minutes) was not significantly different from the 55% MHR recommended by the ACSM Position Stand [13].


Results of the three separate repeated measures ANOVAs examining mean rate of energy (kcal), METs, and %MHR across resting, the entire yoga session and two treadmill speeds found that all three variables differed significantly across the activities (p < 0.0001). All pair wise comparisons of the three variables were significantly different across all activities, with the exception that all three variables were not significantly different when comparing the entire yoga session to walking on a treadmill at 3.2 kph.

Results of the second set of three separate repeated measures ANOVAs (kcal, METs, and %MHR) in which the entire yoga session activity was replaced with the sun salutation activity found that all three variables differed sig-

**Table 2: Means and standard deviations for energy and heart rate parameters across all activities**


Task	Task	Heart Rate (bpm)*	Energy (kcal/min)	VO <sub>2</sub> (l/kg/min)	METs	%MHR*
Resting	(20 mins)	71.0 (13.0)	1.2 (0.5)	0.3 (0.1)	1.0 (0.3)	38.2 (7.0)
Yoga	Across entire Session (52 mins)	93.2 (25.9)	3.2 (1.1)	0.6 (0.2)	2.5 (0.8)	49.4 (12.2)
	Sun Salutation (24 mins)	103.5 (25.2)	3.73 (1.01)	0.76 (0.21)	2.9 (0.77)	54.8 (11.8)
	Non-Sun Salutation Standing Poses (20 mins)	89.7 (23.9)	3.01 (0.810)	0.61 (0.16)	2.34 (0.60)	47.3 (10.4)
	Sitting/Lying poses (8 mins)	72.5 (16.1)	1.93 (0.78)	0.40 (0.18)	1.5 (0.58)	38.7 (8.7)
	Treadmill	3.2 kph/Level (6 mins)	97.8 (21.2)	3.1 (0.6)	0.7(0.1)	2.5(0.4)
	4.8 kph/Level (6 mins)	110.3 (23.1)	4.2 (0.4)	0.9 (0.1)	3.3 (0.4)	58.1 (10.7)

\*All subjects (n = 20) performed all activities. However, due to missing values, secondary to technical issues, data from only 18 subjects were included in %MHR calculations. MHR = Maximum predicted heart rate; METs = Metabolic equivalents;



**Figure 1**  
**Mean METs and standard deviation per minute during the yoga session (n = 20).** Data from the first four minutes of the 56 minute yoga practice are not displayed. The postures used within each section of the yoga practice (see Table 1) varied in sequence and timing. Minutes 1–24 were sun salutation postures with postures changing every 3–8 seconds; minutes 25–44 were standing poses with postures changing every 10–48 seconds, and minutes 45–52 were sitting and lying postures lasting 3 minutes and 4.5 minutes respectively. Readers are referred to the yoga DVD for a complete description of the yoga session [57].

nificantly across the activities ( $p < 0.0001$ ). All pair wise comparisons of the three variables were significantly different across all activities; with the exception that %MHR was not significantly different when comparing the sun salutations postures to walking on a treadmill at 4.8 kph.



**Figure 2**  
**Mean %MHR and standard deviation per minute during the yoga session (n = 18).** Data from the first four minutes of the 56 minute yoga practice are not displayed. The postures used within each section of the yoga practice (see Table 1) varied in sequence and timing. Minutes 1–24 were sun salutation postures with postures changing every 3–8 seconds; minutes 25–44 were standing poses with postures changing every 10–48 seconds, and minutes 45–52 were sitting and lying postures lasting 3 minutes and 4.5 minutes respectively. Readers are referred to the yoga DVD for a complete description of the yoga session [57].

By the end of the yoga session (during *Savasana*, resting supine posture) the mean METs had returned to 1, suggesting that the subjects had returned to the metabolic equivalent of a resting state. Post hoc statistical testing found that METs and %MHR during the final three minutes of the yoga session were not significantly different from mean values found during the initial rest period.

**Discussion**

The current study found that the metabolic costs of yoga averaged across the entire session represent low levels of physical activity [58]. Mean values for kcal/min ( $3.2, \pm 1.1$ ), METs ( $2.5, \pm 0.8$ ), and %MHR ( $49.4, \pm 12.2$ ) were not significantly different from mean values during walking at 3.2 kph on a level treadmill. This level of physical activity was significantly lower than the recommendations for "moderate" levels of physical activity recommended by ACSM and AHA (3.0 METs) [12] and as recommended by the ACSM Position stand (55%MHR) [13].

Only four studies exist that may provide some comparison to the current study as they examined a variety of yoga postures which best represent a typical yoga class, similar to the current study [49-52]. Of these, one [52] provided insufficient description of the postures for valid comparison stating only that there were "six yoga positions." The remaining studies provided data on a series of well-described standing postures, however, the only study for comparison which included standing, sitting, and lying postures as in the current study is that of Clay *et al.* [51]. Clay *et al.* [51] studied a much shorter session (30 mins), a different selection of postures, younger subjects (difference of means: approximately 8 years), and used an indirect calorimeter requiring collection of gases from a mask or mouthpiece. These differences in methodology may explain why Clay *et al.* [51] found the mean caloric expenditure across the entire yoga session to be approximately 30% less than that found in the current study (2.23 vs 3.2 kcal/min).

Only two studies have provided %MHR during yoga for comparison to the current study [51,52]. The mean %MHR of 49.4% seen in this study appeared to be similar to that Clay *et al.* [51] which yielded a mean %MHR of 56.9%, and considerably less than that of Carroll *et al.* at 77%. However, the study by Carroll *et al.* represents standing postures only in a Vinyasa style (flowing movement) measured for 15 minutes, and therefore is unlikely to represent the mean %MHR across a typical yoga session.

It is important to recognize that the mean values across the entire yoga session described above represent a combination of standing, sitting, and lying postures and that

**Table 3: Results of one sample t-tests of mean METS and %MHR compared to recommended constant minimums**

Variable	Comparison value	Difference between mean and comparison value	Standard Deviation	t statistic	P value
Mean %MHR* across entire yoga session (n = 18)	55% MHR [13]	-5.6	12.24	-14.38	< 0.0001
Mean %MHR* during sun salutation portion of yoga session (n = 18)	55% MHR [13]	-0.17	11.77	-0.30	0.7664
Mean METs across entire yoga session (n = 20)	3.0 METs [12]	-0.54	0.83	-20.81	< 0.0001
Mean METs during sun salutation portion of yoga session (n = 20)	3.0 METs [12]	-0.09	0.78	-2.80	< 0.0053

Note: \*All values for MHR based on n = 18. Values for METs based on n = 20. MHR = Maximum predicted heart rate; METs = Metabolic equivalents;

substantial variation in metabolic costs occurs during the yoga session as can be seen in Figures 1 and 2. A substantial time period of the yoga practice (54%) was engaged in sun salutation postures which had metabolic values which were significantly higher than values for treadmill walking at 3.2 kph, were not significantly different than the recommended minimum values for moderate levels of exercise intensity (3.0 METs) [12], and were not significantly different from the minimum recommendations of the ACSM Position Stand for %MHR (55%, [13]) for cardiovascular fitness. Therefore, yoga practice with a portion of sun salutation postures exceeding the recommended minimum bout of 10 minutes [13] may contribute some portion of sufficiently intense physical activity to improve cardio-respiratory fitness. This being said, there are two important caveats regarding this suggestion. First, the criterion used for comparison in this study is the minimum value suggested by the ACSM Position Stand [13] (55%MHR). This value applies only to sedentary or unfit individuals, and therefore may not be a sufficiently high threshold for individuals who are currently fit (for more fit individuals the recommendations suggest a minimum of 65% MHR [13]). Second, ACSM Position Stand recommendations use %MHR as an indirect measure of cardio-respiratory demand based on the assumption of a relatively linear relationship between %MHR and oxygen consumption. However, there is evidence that during yoga, %MHR overestimates the amount of oxygen consumption used [49,51] and consequently %MHR may not be a sufficiently accurate measure of oxygen consumption during a yoga practice. Therefore, %MHR used in this study may be overestimating the intensity of the cardio-respiratory demand, and caution should be taken in consideration of this suggestion.

This study also demonstrated that individuals performing an identical yoga session (n = 7) on two occasions expend very similar amounts of energy (ICC = .979). This finding suggests that the use of mean metabolic values for quantifying energy cost of yoga across sessions is sufficiently consistent to provide a basis for valid estimates.

As is typical of most yoga classes, the yoga practice in this study ended with several minutes of supine relaxation (e.g., shavasana). The post hoc findings that METs and %MHR in the final three minutes of this relaxation were not significantly different from the mean value found in the initial resting period suggest that the increased metabolic demand of yoga was unlikely to have contributed to metabolic measures during the treadmill activity. In further support of this, the mean value (3.3) for the treadmill METs found in this study during the 4.8 kph was identical to the normative value suggested by Ainsworth for level treadmill walking at this same speed [58].

Placing the findings of the current study into a larger discussion of the potential for the physical activity component of yoga to provide health benefits requires consideration of the continuum of benefits that can be gained from increases in physical activity. The ACSM Position Stand [13] recommendations acknowledge that lower levels of physical activity than those recommended may provide health benefits for those who are quite unfit, particularly in the area of metabolic fitness [13]. Metabolic fitness [59,60] describes the ability of metabolic systems predictive of risk of cardiovascular disease (e.g., diabetes) to improve through intensities of physical activity which do not produce change in aspects of performance such as VO<sub>2</sub> max. For example, in relation to obesity and Type II diabetes, overweight/obese individuals who increase physical activity can reduce insulin resistance even in the absence of concurrent weight loss [61-64]. Similarly among the non-obese, physical activity performed at levels insufficient to influence body mass or maximal oxygen uptake, can still improve insulin action [65]. The relatively low level of physical activity found in this study may serve to increase metabolic fitness despite being insufficient to create gains in cardio-respiratory fitness. This suggestion is in agreement with studies which have found yoga to improve indices associated with metabolic fitness such as insulin resistance [66,67] and lipid profiles [68,69].

There is a great deal of evidence that yoga or yoga based interventions reduce blood pressure and heart rate [23-42], and a recent systematic study suggests that over 85% of relevant studies published since 1970 demonstrate evidence that yoga reduces sympathetic tone, increases parasympathetic tone and improves cardiovagal function [42].

The mechanisms by which yoga conveys health benefits have received little study. However, given the findings of this study, as well as the study by Clay et al., it is unlikely that the well documented health effects across multiple systems are due solely to the low levels of physical activity within yoga practice. Yoga is a mind-body practice involving multiple integrated elements (e.g., conscious control of the breath, postures, meditation) [70]. The relative influence and interactions of these elements on health outcomes are unknown. In short, yoga may convey health benefits which are unrelated, or only partially related, to the physical activity component of yoga practice.

This study was limited in that it was a sample of convenience in which 90% of the subjects were women, and all were recruited from a private university and various urban yoga studios. Although, as mentioned previously, removing the two male participants from the data did not alter any of the results, it remains unclear whether a sample including more male participants would have altered the findings. All subjects were relatively young, healthy, and of normal weight and therefore the results may not be generalizable to persons suffering from adverse health conditions, those with very high or very low BMI, the elderly, or children under eighteen. The yoga DVD was chosen to represent the typical length, intensity, and variety of poses in a beginner yoga class, however, we have no information to determine the degree to which it is genuinely representative of yoga classes performed in the United States. Certainly, the results cannot be used to generalize to moderate or advanced level yoga classes. Also, as all subjects were moderate-to-advanced yoga practitioners, it is unclear how yoga experience may have influenced the findings in comparison to those who may be yoga novices. The lack of randomization of activities is a limitation in this study, suggesting that even though the study found that metabolic rate and %MHR returned to baseline after the yoga practice, possible influence upon the treadmill activities cannot be totally eliminated. Finally, in addition to the limitations described earlier relative to %MHR, we acknowledge that the true relative intensity of physical activity for each participant was not known as the  $VO_2$  max was not measured in this study, and consequently, more accurate data regarding each participant's percentage of  $VO_2$  max was not available.

## Conclusion

Metabolic costs of yoga averaged across the entire session represent low levels of physical activity, similar to walking on a treadmill at 3.2 kph, and do not meet recommendations for levels of physical activity for improving or maintaining health [12] or cardiovascular fitness [13]. Yoga practice with a portion of sun salutation postures exceeding the minimum bout of 10 minutes may contribute some portion of sufficiently intense physical activity to improve cardio-respiratory fitness in unfit or sedentary individuals. The measurement of energy expenditure across yoga sessions is highly reliable.

## Competing interests

The author(s) declare that they have no competing interests.

## Authors' contributions

MH and AR conceived of the study, participated in its design and coordination, and assisted with data analysis and drafting of the manuscript. WM performed data collection, assisted with organization and analysis of the data and drafting of the manuscript. All authors read and approved the final manuscript.

## Acknowledgements

The authors would like to thank the following for their contributions: the staff of the Energy Expenditure Sub-Core at St. Luke's Roosevelt Hospital for their patience and flexibility; all of the Manhattan yoga studios which provided participants: Bikram Yoga NYC, Levitate Yoga, Naga Yoga, and World Yoga Center; and lastly, the study participants themselves, without whom this work would not have been possible.

This research was supported by a grant from the National Cancer Institute:U54; CA101598-01

## References

1. Paffenbarger RS Jr., Hyde RT, Wing AL, Hsieh CC: **Physical activity, all-cause mortality, and longevity of college alumni.** *N Engl J Med* 1986, **314(10)**:605-613.
2. Morris JN, Clayton DG, Everitt MG, Semmence AM, Burgess EH: **Exercise in leisure time: coronary attack and death rates.** *Br Heart J* 1990, **63(6)**:325-334.
3. Powell KE, Thompson PD, Caspersen CJ, Kendrick JS: **Physical activity and the incidence of coronary heart disease.** *Annu Rev Public Health* 1987, **8**:253-287.
4. Hagberg JM: **Exercise, fitness, and hypertension.** In *Exercise, Fitness and Health* Edited by: Bouchard C, Shephard RJ, Stephens T, Sutton JR, McPherson BD. Champaign, IL, Human Kinetics; 1990:455-566.
5. Blair SN, Goodyear NN, Gibbons LW, Cooper KH: **Physical fitness and incidence of hypertension in healthy normotensive men and women.** *Jama* 1984, **252(4)**:487-490.
6. Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr.: **Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus.** *N Engl J Med* 1991, **325(3)**:147-152.
7. Cummings SR, Kelsey JL, Nevitt MC, O'Dowd KJ: **Epidemiology of osteoporosis and osteoporotic fractures.** *Epidemiol Rev* 1985, **7**:178-208.
8. Lee IM, Paffenbarger RS Jr., Hsieh C: **Physical activity and risk of developing colorectal cancer among college alumni.** *J Natl Cancer Inst* 1991, **83(18)**:1324-1329.
9. King AC, Taylor CB, Haskell WL, DeBusk RF: **Influence of regular aerobic exercise on psychological health: a randomized, con-**

- trolled trial of healthy middle-aged adults. *Health Psychol* 1989, **8(3)**:305-324.
10. Taylor CB, Sallis JF, Needle R: **The relation of physical activity and exercise to mental health.** *Public Health Rep* 1985, **100(2)**:195-202.
  11. Macera CA, Ham SA, Yore MM, Jones DA, Ainsworth BE, Kimsey CD, Kohl HW 3rd: **Prevalence of physical activity in the United States: Behavioral Risk Factor Surveillance System, 2001.** *Prev Chronic Dis* 2005, **2(2)**:A17.
  12. Haskell WL, Lee IM, Pate RR, Powell KE, Blair SN, Franklin BA, Macera CA, Heath GW, Thompson PD, Bauman A, American College of Sports Medicine, American Heart Association: **Physical activity and public health: updated recommendation for adults from the American College of Sports Medicine and the American Heart Association.** *Circulation* 2007, **116(9)**:1081-1093.
  13. Pollock ML Gaesser, Glenn A., Butcher, Janus D., Despres, Jean-Pierre, Dishman, Rod K., Franklin, Barry A., Garber, Carol Ewing: **ACSM Position Stand: The Recommended Quantity and Quality of Exercise for Developing and Maintaining Cardiorespiratory and Muscular Fitness, and Flexibility in Healthy Adults.** *Med Sci Sports Exerc* 1998, **30(6)**:975-991.
  14. Garfinkel M, Schumacher HR Jr.: **Yoga.** *Rheum Dis Clin North Am* 2000, **26(1)**:125-32, x.
  15. Chandler K: **The emerging field of yoga therapy.** *Hawaii Med J* 2001, **60(11)**:286-287.
  16. Raub JA: **Psychophysiological effects of Hatha Yoga on musculoskeletal and cardiopulmonary function: a literature review.** *J Altern Complement Med* 2002, **8(6)**:797-812.
  17. Saper RB, Eisenberg DM, Davis RB, Culpepper L, Phillips RS: **Prevalence and patterns of adult yoga use in the United States: results of a national survey.** *Altern Ther Health Med* 2004, **10(2)**:44-49.
  18. Corliss R Funderburg, Lise: **The Power of Yoga.** *Time* 2001, **157(16)**:54-62.
  19. Labarthe D, Ayala C: **Nondrug interventions in hypertension prevention and control.** *Cardiol Clin* 2002, **20(2)**:249-263.
  20. Gimbel MA: **Yoga, meditation, and imagery: clinical applications.** *Nurse Pract Forum* 1998, **9(4)**:243-255.
  21. Sallis JF, Hovell MF, Hofstetter CR, Faucher P, Elder JP, Blanchard J, Caspersen CJ, Powell KE, Christenson GM: **A multivariate study of determinants of vigorous exercise in a community sample.** *Prev Med* 1989, **18(1)**:20-34.
  22. Pollock ML: **Prescribing exercise for fitness and adherence.** In *Exercise Adherence: Its Impact on Public Health* Edited by: Dishman RK. Champaign, IL, Human Kinetic Books; 1988:259-277.
  23. Joseph S, Sridharan K, Patil SK, Kumaria ML, Selvamurthy W, Joseph NT, Nayar HS: **Study of some physiological and biochemical parameters in subjects undergoing yogic training.** *Indian J Med Res* 1981, **74**:120-124.
  24. Udupa KN, Singh RH, Settiwar RM: **Physiological and biochemical studies on the effect of yogic and certain other exercises.** *Indian J Med Res* 1975, **63(4)**:620-624.
  25. Schmidt T, Wijga A, Von Zur Muhlen A, Brabant G, Wagner TO: **Changes in cardiovascular risk factors and hormones during a comprehensive residential three month kriya yoga training and vegetarian nutrition.** *Acta Physiol Scand Suppl* 1997, **640**:158-162.
  26. Telles S, Nagarathna R, Nagendra HR, Desiraju T: **Physiological changes in sports teachers following 3 months of training in Yoga.** *Indian J Med Sci* 1993, **47(10)**:235-238.
  27. Latha KK: **Yoga, pranayama, thermal biofeedback techniques in the management of stress and high blood pressure.** *J Indian Psychol* 1991, **9**:36-46.
  28. Anantharaman R Kabir, R: **A study of Yoga.** *J Psychol Res* 1984, **28**:97-101.
  29. Mogra A, Singh G: **Effect of biofeedback and yogic relaxation exercise on the blood pressure levels of hypertensives: a preliminary study.** *Aviation Med* 1986, **30**:68-75.
  30. Selvamurthy W, Sridharan K, Ray US, Tiwary RS, Hegde KS, Radhakrishnan U, Sinha KC: **A new physiological approach to control essential hypertension.** *Indian J Physiol Pharmacol* 1998, **42(2)**:205-213.
  31. Patel C: **12-month follow-up of yoga and bio-feedback in the management of hypertension.** *Lancet* 1975, **1(7898)**:62-64.
  32. Bagga OP, Gandhi A: **A comparative study of the effect of Transcendental Meditation (T.M.) and Shavasana practice on cardiovascular system.** *Indian Heart J* 1983, **35(1)**:39-45.
  33. Cusumano JA, Robinson SE: **The short-term psychophysiological effects of hatha yoga and progressive relaxation on female Japanese students.** *Appl Psychol* 1992, **42**:77-90.
  34. Ray US, Mukhopadhyaya S, Purkayastha SS, Asnani V, Tomer OS, Prashad R, Thakur L, Selvamurthy W: **Effect of yogic exercises on physical and mental health of young fellowship course trainees.** *Indian J Physiol Pharmacol* 2001, **45(1)**:37-53.
  35. Harinath K, Malhotra AS, Pal K, Prasad R, Kumar R, Kain TC, Rai L, Sawhney RC: **Effects of Hatha Yoga and Omkar Meditation on Cardiorespiratory Performance, Psychologic Profile, and Melatonin Secretion.** *The Journal of Alternative and Complementary Medicine* 2004, **10(2)**:261-268.
  36. Damodaran A, Malathi A, Patil N, Shah N, Suryavanshi, Marathe S: **Therapeutic potential of yoga practices in modifying cardiovascular risk profile in middle aged men and women.** *J Assoc Physicians India* 2002, **50(5)**:633-640.
  37. Lakshmikanthan C, Alagesan R, Thanikachalam S, Ramamurthy B, Elangovan D, Viswanathan TR, Kumar S: **Long term effects of yoga on hypertension and/or coronary artery disease.** *J Assoc Physicians India* 1979, **27(12)**:1055-1058.
  38. Patel C, Datey KK: **Relaxation and biofeedback techniques in the management of hypertension.** *Angiology* 1976, **27(2)**:106-113.
  39. Aaronson N: **The european organization for reserach and treatment of cancer QLQ-C30: A quality-of-life instrument for use in international clinical trials in oncology.** *J Natl Cancer Inst* 1993, **85**:365-376.
  40. Patel C: **Reduction of serum cholesterol and blood pressure in hypertensive patients by behaviour modification.** *J R Coll Gen Pract* 1976, **26(164)**:211-215.
  41. Yogendra J, Yogendra HJ, Ambardekar S, Lele RD, Shetty S, Dave M, Husein N: **Beneficial effects of yoga lifestyle on reversibility of ischaemic heart disease: caring heart project of International Board of Yoga.** *J Assoc Physicians India* 2004, **52**:283-289.
  42. Innes KE, Bourguignon C, Taylor AG: **Risk indices associated with the insulin resistance syndrome, cardiovascular disease, and possible protection with yoga: a systematic review.** *J Am Board Fam Pract* 2005, **18(6)**:491-519.
  43. Ray US Hegde, KS, Selvamurthy, W: **Improvement in muscular efficiency as related to a standard task after yogic exercises in middle aged men.** *Indian J Med Res* 1986, **83**:343-348.
  44. Tran MD, Holly RG, Lashbrook J, Amsterdam EA: **Effects of Hatha Yoga Practice on the Health-Related Aspects of Physical Fitness.** *Prev Cardiol* 2001, **4(4)**:165-170.
  45. Prasad KVV, Venkata Ramana Y, Raju PS, Venkata MR, Murthy KJR: **Energy Cost and Physiological Efficiency in Male Yoga Practitioners.** *Journal of Exercise Physiology* 2001, **4(3)**:38-44.
  46. Rai L, Ram K: **Energy expenditure and ventilatory responses during Virasana--a yogic standing posture.** *Indian J Physiol Pharmacol* 1993, **37(1)**:45-50.
  47. Rai L, Ram K, Kant U, Madan SK, Sharma SK: **Energy expenditure and ventilatory responses during Siddhasana--a yogic seated posture.** *Indian J Physiol Pharmacol* 1994, **38(1)**:29-33.
  48. Rao S: **The Metabolic Cost of Head-Stand Posture.** *J Appl Physiol* 1962, **17(1)**:117-118.
  49. DiCarlo LJ, Sparling PB, Hinson BT, Snow TK, Roskopf LB: **Cardiovascular, Metabolic, and Perceptual Responses to Hatha Yoga Standing Poses.** *Med Exerc Nutr Health* 1995, **4**:107-112.
  50. Sinha B, Ray US, Pathak A, Selvamurthy W: **Energy cost and cardiorespiratory changes during the practice of Surya Namaskar.** *Indian J Physiol Pharmacol* 2004, **48(2)**:184-190.
  51. Clay CC, Lloyd LK, Walker JL, Sharp KR, Pankey RB: **The metabolic cost of hatha yoga.** *J Strength Cond Res* 2005, **19(3)**:604-610.
  52. Carroll J, Blansit A, Otto RM, Wygand JW, FACSM: **The Metabolic Requirements of Vinyasa Yoga.** *Med Sci Sports Exerc* 2003, **35(5)**:S155.
  53. Zhang K, Sun M, Werner P, Kovera AJ, Albu J, Pi-Sunyer FX, Boozer CN: **Sleeping metabolic rate in relation to body mass index and body composition.** *Int J Obes Relat Metab Disord* 2002, **26(3)**:376-383.
  54. Heymsfield SB, Allison DB, Pi-Sunyer FX, Sun Y: **Columbia respiratory-chamber indirect calorimeter: a new approach to airflow modelling.** *Med Biol Eng Comput* 1994, **32(4)**:406-410.



55. Fujii TK, Phillips BJ: **Quick Review: The Metabolic Cart.** *The Internet Journal of Internal Medicine* 2003, **3(2)**).
56. Tanaka H, Monahan KD, Seals DR: **Age-predicted maximal heart rate revisited.** *J Am Coll Cardiol* 2001, **37(1)**:153-156.
57. Landon T: **Ashtanga Yoga Beginners Practice.** Gaiam, Inc..
58. Ainsworth BE, Haskell WL, Whitt MC, Irwin ML, Swartz AM, Strath SJ, O'Brien WL, Bassett DR Jr, Schmitz KH, Emplainscourt PO, Jacobs DR Jr, Leon AS: **Compendium of physical activities: an update of activity codes and MET intensities.** *Med Sci Sports Exerc* 2000, **32(9 Suppl)**:S498-S504.
59. Despres JP, Pouliot MC, Moorjani S, Nadeau A, Tremblay A, Lupien PJ, Theriault G, Bouchard C: **Loss of abdominal fat and metabolic response to exercise training in obese women.** *Am J Physiol* 1991, **261(2 Pt 1)**:E159-67.
60. Despres JP, Tremblay A, Moorjani S, Lupien PJ, Theriault G, Nadeau A, Bouchard C: **Long-term exercise training with constant energy intake. 3: Effects on plasma lipoprotein levels.** *Int J Obes* 1990, **14(1)**:85-94.
61. Brown MD, Moore GE, Korytkowski MT, McCole SD, Hagberg JM: **Improvement of insulin sensitivity by short-term exercise training in hypertensive African American women.** *Hypertension* 1997, **30(6)**:1549-1553.
62. Dengel DR, Pratley RE, Hagberg JM, Rogus EM, Goldberg AP: **Distinct effects of aerobic exercise training and weight loss on glucose homeostasis in obese sedentary men.** *J Appl Physiol* 1996, **81(1)**:318-325.
63. Reitman JS, Vasquez B, Klimes I, Nagulesparan M: **Improvement of glucose homeostasis after exercise training in non-insulin-dependent diabetes.** *Diabetes Care* 1984, **7(5)**:434-441.
64. Angelopoulos TJ, Schultz RM, Denton JC, Jamurtas AZ: **Significant enhancements in glucose tolerance and insulin action in centrally obese subjects following ten days of training.** *Clin J Sport Med* 2002, **12(2)**:113-118.
65. Oshida Y, Yamanouchi K, Hayamizu S, Sato Y: **Long-term mild jogging increases insulin action despite no influence on body mass index or VO2 max.** *J Appl Physiol* 1989, **66(5)**:2206-2210.
66. Malhotra V, Singh S, Singh KP, Gupta P, Sharma SB, Madhu SV, Tandon OP: **Study of yoga asanas in assessment of pulmonary function in NIDDM patients.** *Indian J Physiol Pharmacol* 2002, **46(3)**:313-320.
67. Monro R, Power J, Coumar A, Dandona P: **Yoga therapy for NIDDM: a controlled trial.** *Complementary Medicine Research* 1992, **6**:66-68.
68. Vyas R, Dikshit N: **Effect of meditation on respiratory system, cardiovascular system and lipid profile.** *Indian J Physiol Pharmacol* 2002, **46(4)**:487-491.
69. Naruka JS, Mathur R, Mathur A: **Effect of pranayama practices on fasting blood glucose and serum cholesterol.** *Indian J Med Sci* 1986, **40(6)**:149-152.
70. Iyengar BKS: **Light on Yoga.** Paperback revised edition. New York, Schocken Books; 1979.

### Pre-publication history

The pre-publication history for this paper can be accessed here:

<http://www.biomedcentral.com/1472-6882/7/40/prepub>

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:

[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

