



Prognostic assessment of ^{18}F -boronophenylalanine positron emission tomography (BPA-PET) in salvage boron neutron capture therapy for malignant brain tumors

Ko-Han Lin¹, Yi-Wei Chen², Ling-Wei Wang², Yuh-Feng Wang¹, Lien-Hsin Hu^{1,3}, Chien Hsin Ting¹, Tse-Hao Lee¹, Jia-Cheng Lee², Nan-Jing Peng^{1,3}

¹Department of Nuclear Medicine, Taipei Veterans General Hospital, Taipei; ²Department of Radiation Oncology, Taipei Veterans General Hospital, Taipei; ³School of Medicine, National Yang-Ming Chiao Tung University, Taipei

Contributions: (I) Conception and design: KH Lin, YW Chen, NJ Peng; (II) Administrative support: YW Chen, LW Wang, YF Wang, LH Hu, CH Ting, TH Lee, JC Lee; (III) Provision of study materials or patients: KH Lin, YW Chen, LW Wang, JC Lee; (IV) Collection and assembly of data: KH Lin, YW Chen, LW Wang, JC Lee, NJ Peng; (V) Data analysis and interpretation: KH Lin, YW Chen, JC Lee, NJ Peng; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Nan-Jing Peng, MD. Department of Nuclear Medicine, Taipei Veterans General Hospital, 201 Shih-Pai Road, Sec. 2, Taipei 11217; School of Medicine, National Yang-Ming Chiao Tung University, Taipei. Email: njpeng@vghtpe.gov.tw.

Background: Boron neutron capture therapy (BNCT) stands out as a propitious anti-cancer modality. ^{18}F -boronophenylalanine positron emission tomography (BPA-PET) holds the potential to ascertain the concentration of BPA within the tumor, enabling meticulous treatment planning and outcome evaluation. However, no studies have been conducted on comparing the outcomes of those treated with BNCT to those who did not undergo this therapy. This study endeavors to analyze the correlation between BPA-PET and BNCT in the context of malignant brain tumors, and assess the survival outcomes following BNCT.

Methods: A cohort study was performed on patients who underwent BPA-PET between February 2017 and April 2022 in our hospital. Patients were stratified into two groups: those subjected to BNCT (Group 1) and those not (Group 2). The tumor to normal tissue (T/N) ratio derived from BPA-PET was set at 2.5. The findings were scrutinized based on clinical follow-up. Student's *t*-test and Chi-squared test were employed to discern differences between the groups. A cumulative survival curve was constructed employing the Kaplan-Meier method. Differences were considered statistically significant at $P < 0.05$.

Results: In total, 116 patients with T/N ratios obtained from BPA-PET were enrolled. BNCT was administered to 58 patients, while mortality was observed in 100 patients. The median overall survival (OS) for the two groups was 8.5 and 6.0 months, respectively. The cumulative OS exhibited no significant discrepancy between the two groups, nor in their T/N ratios. Within Group 1, 44 out of 58 (75.9%) patients exhibited T/N ratios exceeding 2.5. Excluding 3 patients who expired within 3 months, 55 out of 58 patients were evaluated for response after BNCT. The objective response rate (ORR) was 30.9%. Patients achieving ORR displayed substantially higher survival rates compared to those without (median OS 13.5 *vs.* 8.3 months, $P = 0.0021$), particularly when T/N ratio exceeded 2.5 (median OS 14.8 *vs.* 9.0 months, $P = 0.0199$).

Conclusions: BNCT does not appear indispensable for prolonging the survival of patients afflicted with malignant brain tumors. Nevertheless, it proves advantageous when ORR is attained, a condition closely linked to the values of T/N ratio derived from BPA-PET.

Keywords: Boron neutron capture therapy (BNCT); ^{18}F -boronophenylalanine (^{18}F -BPA); objective response rate (ORR); positron emission tomography (PET); tumor to normal tissue ratio (T/N ratio)

Submitted Dec 13, 2023. Accepted for publication Apr 22, 2024. Published online May 20, 2024.

doi: 10.21037/qims-23-1769

View this article at: <https://dx.doi.org/10.21037/qims-23-1769>

Introduction

Boron neutron capture therapy (BNCT) emerges as a promising approach in the domain of anti-cancer methods. Its basis lies in the nuclear capture reaction, wherein non-radioactive boron (^{10}B) strongly absorbs neutrons of thermal energy ($E < 1$ eV), leading to the generation of high-energy alpha particles ($^4\text{He}^{2+}$) and lithium (^7Li) nuclei. The short range of these high linear-transfer alpha and ^7Li particles (5–9 μm) is confined to one cell diameter (< 10 μm), facilitating selective tumor eradication while sparing the adjacent normal tissue, thus heralding targeted radiation therapy (1-3). The efficacy of BNCT hinges upon the preferential accumulation of sufficient quantities of ^{10}B in cancerous cells compared to their normal counterparts. This selective uptake of boron is achieved through the utilization of boron carriers, notably boronophenylalanine (BPA) derivatives (3).

BPA represents a tumor-specific probe, attributed to its interaction with L-type amino acid transporter 1 (LAT1). Notably, the LAT1 system facilitates the import of BPA primarily into proliferating cells. Consequently, the expression of LAT1 plays a pivotal role in tumor progression, as BPA actively traverses the tumor cell membrane at an elevated rate in these proliferating cells (4).

BPA, being a boron-containing drug with established clinical use, warrants a thorough examination due to its incorporation into clinical dosimetry and treatment planning through ^{18}F -boronophenylalanine positron emission tomography (BPA-PET) data. This imaging modality enables the estimation of BPA concentration within the tumor, thereby enabling the assessment of treatment feasibility and efficacy. A usual suggestion for targeted radiation therapy is achieving a tumor to normal tissue (T/N) ratio of boron-contained drug concentration greater than a specified threshold, with a patient exhibiting a ratio exceeding 2.5 being deemed an optimal candidate for BNCT (5,6). Nevertheless, quantitative assessment techniques in BPA-PET remain to be established. The T/N ratio and the tumor to blood (T/B) ratio were computed based on the maximum standardized uptake value (SUV) of the brain, tumor, and heart.

Prior investigations have demonstrated favorable treatment outcomes with BNCT, particularly in patients

afflicted with brain tumors (6-10). Notably, ^{18}F -BPA injection holds the potential to achieve this goal due to the elevated T/N ratios of BPA uptake observed in cases of newly-diagnosed glioblastoma, recurrent malignant glioma, and brainstem tumors (6-10). Moreover, there have been reports of similar promising results in other tumor types such as head and neck cancers, melanoma, sarcoma, and hepatocellular carcinoma (11-15).

However, no studies have been conducted on patients who underwent BPA-PET evaluation but did not receive BNCT treatment. The present study endeavors to evaluate the relationship between BPA-PET and BNCT treatment in patients and compare the outcomes of those treated with BNCT to those who did not undergo this therapy. We present this article in accordance with the STROBE reporting checklist (available at <https://qims.amegroups.com/article/view/10.21037/qims-23-1769/rc>).

Methods

Patients

This study was centered around a cohort of patients who underwent BPA-PET for the purpose of proceeding with BNCT for malignant brain tumors at Taipei Veterans General Hospital. A retrospective analysis was conducted (Figure 1). The inclusion criteria were those who underwent BPA-PET between February 2017 and April 2022 ($n=460$), and who were diagnosed with brain tumors ($n=149$). Patients were followed up for at least 1 year. The exclusion criteria were those who were loss of follow-up ($n=20$), or who were found the benign nature of tumors or metastases originating from elsewhere ($n=13$). Consequently, a total of 116 patients were included in this study, and they were categorized into two cohorts: those who underwent BNCT (Group 1, $n=58$) and those who did not (Group 2, $n=58$). The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study was approved by the institutional review board of Taipei Veterans General Hospital (IRB No. 2022-07-0029CC). Due to the retrospective nature of the study, written or verbal informed consent was waived.

The BNCT protocol used in this study was modified based on the regimen developed by Kyoto University in




Figure 1 A flowchart of patients who underwent ^{18}F -BPA PET and their classifications. ^{18}F -BPA PET, ^{18}F -boronophenylalanine positron emission tomography; BNCT, boron neutron capture therapy.

Japan. The total infused dose was 450 mg/kg body weight. The infusion formula was full infusion for the first two hours (180 mg/kg) and then infusion at half of the initial infusion rate (90 mg/kg) during neutron irradiation to maintain a constant boron concentration in the blood. The epithermal neutron source for BNCT was the Tsing-Hua Open-Pool Reactor (THOR) at National Tsing-Hua University. The blood boron concentration and minimum, maximum, and mean tumor doses required for the gross tumor volume (GTV) were estimated and recorded. In this study, the mean tumor dose required for the GTV was estimated to be 20–40 Gray-Equivalents (Gy-E) (9). Bevacizumab was administered at a dosage of 10 mg/kg promptly following BNCT and at biweekly intervals until disease progression, severe treatment-related toxicity (grade 3), or patient mortality.

BPA-PET imaging and image analysis

To assess the T/N ratio and facilitate treatment planning, BPA-PET imaging was conducted following the Standard Operating Procedure (SOP) at Taipei Veterans General Hospital (6,9-12). ^{18}F -BPA was utilized as the radiotracer with an intravenous injection of $185\% \pm 10\%$ MBq dose. The BPA-PET imaging was acquired using integrated

PET/MRI or PET/CT scanners 60 minutes post-injection of the radiotracer. PET imaging data was acquired from the head with a 3-minute acquisition period per bed position. The raw PET imaging data was reconstructed into a 128×128 matrix using an iterative reconstruction algorithm. Evaluation and quantification of the SUV were performed using the Xeleris Workstation (GE Healthcare). Regions of interest (ROIs) were drawn on the highest uptake area within the tumor and the normal tissue of the buccal mucosa near the tumor to measure the T/N ratio. The T/N ratio was evaluated from static emission scans acquired 60 minutes after tracer injection, with a 20-minute emission time.

The T/N and T/B ratios of BPA-PET were computed and documented. The contralateral brain lobe was designated as normal tissue for T/N ratio computation, while the heart ventricle served as a blood pool for T/B ratio determination. The mean standard uptake value (SUV_{mean}) was used for the tumor lesion. Herein, Gy-E was used to indicate the biologically equivalent dose administered to patients during BNCT. Gy-E values were calculated by multiplying the physical dose given by the relative biological effectiveness (RBE). The compound biological effectiveness (CBE) was used to assess the boron dose derived from effects of the $^{10}\text{B}(n, \alpha)^7\text{Li}$ reaction employing ^{10}B -L-BPA as the boron carrier. The following values were used in this study: RBE-thermal neutron = 3.2, RBE = 1.0, CBE = 3.8 for the tumor, CBE = 2.5 for normal skin, CBE = 1.35 for the nerves, brain max dose <10 Gy-E, and brain mean dose <2 Gy-E.

Response evaluation

Magnetic resonance imaging (MRI) scans were performed before and every 3 months after the BNCT procedure to assess radiologic response according to the Response Assessment in Neuro-Oncology criteria (RANO) (16). Complete response (CR) was defined as the complete disappearance of the tumor within the brain after treatment. Partial response (PR) was defined as a $\geq 50\%$ reduction in tumor volume after treatment (the largest cross-sectional tumor area) sustained for at least 4 weeks. Progressive disease (PD) was defined as a $\geq 25\%$ increase in tumor volume after treatment, while stable disease (SD) was defined when the tumor volume after BNCT treatment did not meet the criteria for CR, PR, or PD. Objective response rate (ORR) was defined as the sum of CR and PR, while disease control rate (DCR) was defined as the sum of CR, PR, and SD. All patients were followed up every 3 months

Table 1 Patient characteristics

Total patient (n=116)	BNCT (n=58)	No BNCT (n=58)	P value
Age (year-old)			0.8455
Mean \pm SD	42.1 \pm 20.2	42.9 \pm 23.4	
Median (range)	45 (4–88)	50 (3–79)	
Gender, n (%)			0.7115
Male	28 (48.3)	30 (51.7)	
Female	30 (51.7)	28 (48.3)	
Survival, n (%)			0.2835
Survival	6 (10.3)	10 (17.2)	
Expired	52 (89.7)	48 (82.8)	
Tumor to normal tissue ratio			
Mean \pm SD	2.8 \pm 0.9	2.1 \pm 0.7	<0.0001
Median (range)	2.6 (1.7–5.6)	2.0 (1.0–4.0)	
\geq 2.5, n (%)	44 (75.9)	26 (44.8)	0.0007
<2.5, n (%)	14 (24.1)	32 (55.2)	
Tumor type, n (%)			0.8458
Glioblastoma	38 (65.5)	39 (67.2)	
Others	20 (34.5)	19 (32.8)	
Astrocytoma	13 (22.4)	8 (13.8)	
Oligoastrocytoma	3 (5.2)	4 (6.9)	
Brainstem glioma	3 (5.2)	4 (6.9)	
Anaplastic ependymoma	1 (1.7)	1 (1.7)	
Pineal gland tumor	0 (0.0)	1 (1.7)	
Medulloblastoma	0 (0.0)	1 (1.7)	

BNCT, boron neutron capture therapy; SD, standard deviation.

with MRI imaging and neurological examination.

Statistical analysis

The clinical outcomes were ascertained from medical records, while potential prognostic factors for survival were assessed. Overall survival (OS) was defined as the time elapsed from the day of intervention to the date of death or the last follow-up, with no restriction on the cause of death. Consequently, the inception of survival was designated as the day of BPA-PET. The T/N ratio derived from BPA-PET was set at 2.5. A cumulative survival curve

was generated using the Kaplan-Meier method. Continuous variables were expressed as mean \pm standard deviation. Categorical variables were expressed as percentages n (%). Differences between the groups were analyzed using two-sided Student's *t*-test, log-rank and the chi-squared test, with statistical significance defined at $P < 0.05$. All calculations were performed using SPSS software, version 20.0 (SPSS, Chicago, IL, USA).

Results

A total of 116 patients with T/N ratios derived from BPA-PET were included in this study. Their demographic information, concerning those who underwent BNCT for malignant brain tumors, was presented in *Table 1*. The patients were divided into two groups: Group 1 (n=58) comprising those who underwent BNCT and Group 2 (n=58) consisting of those who did not. All of them were recurrent cases. No significant differences were observed in terms of age, gender, or survival rate between the two groups. However, a significant difference was found in T/N ratios derived from BPA-PET ($P < 0.0001$). Specifically, 44 of 58 (75.9%) patients in Group 1 and 26 of 58 (44.8%) patients in Group 2 exhibited T/N ratios greater than 2.5, and this difference was statistically significant ($P = 0.0007$). The tumor types included glioblastoma (n=77), astrocytoma (n=21), oligoastrocytoma (n=7), brainstem glioma (n=7), anaplastic ependymoma (n=2), pineal gland tumor (n=1), and medulloblastoma (n=1). Glioblastoma constituted the major tumor type in both groups, and there was no significant difference in tumor type distribution between the two groups (glioblastoma *vs.* others, $P = 0.8458$).

Mortality was observed in 100 patients from the entire cohort. The median OS for the entire cohort was 7.0 months (range, 0.7–38.7 months, *Table 2*). Upon comparing the two groups, Group 1 had a median OS of 8.5 months (range, 2.9–36.1 months), while Group 2 had a median OS of 6.0 months (range, 0.7–38.7 months; $P = 0.3623$, *Figure 2A*). Among the 116 patients, 70 had T/N ratios greater than 2.5, while 46 had T/N ratios lower than 2.5. However, there was no significant difference in patient survival based on T/N ratios greater than 2.5 or not (median OS 7.1 *vs.* 7.0 months, $P = 0.8955$, *Figure 2B*), neither in Group 1 (median OS 8.5 *vs.* 8.5 months, $P = 0.2630$, *Figure 2C*), nor in Group 2 (median OS 4.3 *vs.* 6.1 months, $P = 0.1602$, *Figure 2D*). The cumulative OS of patients with T/N ratios exceeding 2.5 in Group 1 was compared to those in Group 2 (median OS 8.5 *vs.* 4.3 months, $P = 0.0783$, *Figure 2E*), albeit no statistically

Table 2 Survival analysis in patients with different treatment response (n=116)

Variables	Patients, n (%)	Mean OS, months	Median OS (range), months	Survival, n (%)	P value
Total patients	116 (100)	9.4	7.0 (0.7–38.7)	16 (13.8)	
BNCT					0.3623
Yes	58 (50.0)	10.4	8.5 (2.9–36.1)	6 (10.3)	
No	58 (50.0)	8.6	6.0 (0.7–38.7)	10 (17.2)	
T/N ratio ≥ 2.5					0.8955
No	46 (39.7)	9.5	7.0 (0.8–34.5)	7 (15.2)	
Yes	70 (60.3)	9.3	7.1 (0.7–38.7)	9 (12.9)	
BNCT	44 (62.9)	10.4	8.5 (2.9–36.1)	6 (13.6)	0.0783
No BNCT	26 (37.1)	7.4	4.3 (0.7–38.7)	3 (11.5)	
BNCT					0.2630
T/N ratio ≥ 2.5	44 (75.9)	10.4	8.5 (2.9–36.1)	6 (13.6)	
T/N ratio < 2.5	14 (24.1)	9.2	8.5 (4.3–17.3)	0 (0.0)	
No BNCT					0.1602
T/N ratio ≥ 2.5	26 (44.83)	7.4	4.3 (0.7–38.7)	3 (11.5)	
T/N ratio < 2.5	32 (55.17)	9.6	6.1 (0.8–34.5)	7 (21.9)	

BNCT, boron neutron capture therapy; OS, overall survival; T/N ratio, tumor to normal tissue ratio.




Figure 2 Patient survival analysis, and the association between patient survival and tumor to normal tissue ratios derived from ^{18}F -boronophenylalanine positron emission tomography in patients with and without boron neutron capture therapy. (A) The correlation between patient survival and boron neutron capture therapy, and (B) tumor to normal tissue ratios derived from ^{18}F -boronophenylalanine positron emission tomography, as well as (C-E) the comparison of these ratios in patients undergoing boron neutron capture therapy versus those without boron neutron capture therapy. No statistically significant difference was observed. BNCT, boron neutron capture therapy; T/N ratio, tumor to normal tissue ratio.




Figure 3 Representative images of partial response of brain tumor after boron neutron capture therapy. (A) Before boron neutron capture therapy, ^{18}F -boronophenylalanine positron emission tomography revealed the location of tumor activity (tumor to normal tissue ratio of 2.6) in the right fronto-parietal lobe. (B) Magnetic resonance imaging of the patient before boron neutron capture therapy. A recurrent glioblastoma with peritumoral edema was observed in the right fronto-parietal lobe (orange irregular border). (C) Three months after boron neutron capture therapy, a follow-up magnetic resonance imaging of the patient showed interval regression of the tumor volume.

Table 3 Tumor response after boron neutron capture therapy, and their association with T/N ratio, T/B ratio and absorbed doses

Response	BNCT treatment response (n=55)			
	N (%)	T/N ratio	T/B ratio	Doses (Gy-E)
CR	7 (12.73)	2.71±0.56	3.16±0.56*	24.95±13.48
PR	10 (18.18)	3.18±0.88	2.72±1.03	27.56±09.55
SD	9 (16.36)	3.16±1.22	2.82±0.87	22.49±11.64
PD	29 (52.73)	2.66±0.61	2.52±0.91	28.06±14.06
ORR (CR + PR)	30.91%			
DCR (CR + PR + SD)	47.27%			

Data are N (%) or mean ± standard deviation. *, $P=0.0323$, CR versus PD. T/N ratio, tumor to normal tissue ratio; T/B ratio, tumor to blood ratio; BNCT, boron neutron capture therapy; Gy-E, Gray-Equivalents; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; ORR, objective response rate (ORR = CR + PR); DCR, disease control rate (DCR = CR + PR + SD).

significant distinction was observed.

Tumor response in patients was assessed by cranial MRI examination before and 3 months after BNCT (Figure 3). Among Group 1 patients, three patients could not undergo treatment response evaluation due to expiration within 3 months after BNCT. Consequently, 55 patients were included in the response evaluation. The results showed CR in 7 patients, PR in 10 patients, SD in 9 patients, and PD in 29 patients. The proportions of patients who achieved CR, PR, SD, and PD were 12.7%, 18.2%, 16.4%, and 52.7%, respectively. The ORR was 30.9%, and the DCR was 47.3% (Table 3). The T/N ratio, T/B ratio, and absorbed doses were delineated in Table 3. No significant differences were observed among them in terms of T/N ratio (CR vs. PR, $P=0.2060$; CR vs. SD, $P=0.3497$; CR vs. PD, $P=0.8359$;

PR vs. SD, $P=0.9743$; PR vs. PD, $P=0.1127$; SD vs. PD, $P=0.2676$) and absorbed doses (CR vs. PR, $P=0.6689$; CR vs. SD, $P=0.7074$; CR vs. PD, $P=0.5994$; PR vs. SD, $P=0.3182$; PR vs. PD, $P=0.9009$; SD vs. PD, $P=0.2508$; respectively). Regarding the T/B ratio, there was no significant difference among them (CR vs. PR, $P=0.2778$; CR vs. SD, $P=0.3666$; PR vs. SD, $P=0.8159$; PR vs. PD, $P=0.5933$; SD vs. PD, $P=0.3805$; respectively), except for CR vs. PD in the T/B ratio ($P=0.0323$).

The survival analyses were based on various treatment responses. Mortality was observed in 49 of 55 patients in Group 1. Patient survival analysis, and the association between patient survival and T/N ratios derived from BPA-PET in patients with CR, PR, SD, ORR, and DCR after BNCT were shown in Table 4. The median OS for these

Table 4 Survival analysis in patients with different treatment response after BNCT (n=55)

Variables	Patients, n (%)	Mean OS, months	Median OS (range), months	Survival, n (%)	P value
Total patients	55 (100)	10.8	8.5 (2.9–36.1)	6 (10.9)	
T/N ratio ≥ 2.5	42 (76.4)	10.8	6.5 (2.9–36.1)	6 (14.3)	0.2705
T/N ratio < 2.5	13 (23.6)	9.6	8.5 (4.3–17.3)	0 (0.0)	
CR					
CR	7 (12.7)	17.5	15.7 (5.8–36.1)	2 (28.6)	0.0181
PR + SD + PD	48 (87.3)	9.5	8.5 (2.9–30.6)	4 (8.3)	
T/N ratio ≥ 2.5	6 (85.7)	19.0	16.9 (5.8–36.1)	2 (33.3)	0.0253
T/N ratio < 2.5	1 (14.3)	8.5	8.5 (8.5–8.5)	0 (0.0)	
PR					
PR	10 (18.2)	12.0	11.6 (4.9–20.1)	2 (20.0)	0.2385
CR + SD + PD	45 (81.8)	10.2	8.5 (2.9–36.1)	4 (8.9)	
T/N ratio ≥ 2.5	8 (80.0)	12.6	11.8 (7.9–20.1)	2 (25.0)	0.1527
T/N ratio < 2.5	2 (20.0)	9.2	9.2 (4.9–13.5)	0 (0.0)	
SD					
SD	9 (16.4)	8.8	7.9 (2.9–16.8)	1 (11.1)	0.3724
CR + PR + PD	46 (83.6)	10.8	8.9 (3.8–36.1)	5 (10.9)	
T/N ratio ≥ 2.5	7 (77.8)	7.5	5.3 (2.9–16.8)	1 (14.3)	0.5329
T/N ratio < 2.5	2 (22.2)	13.3	13.3 (9.7–16.8)	0 (0.0)	
PD					
PD	29 (52.7)	8.8	8.4 (3.8–30.6)	1 (3.5)	0.0133
CR + PR + SD	26 (47.3)	12.3	10.1 (2.9–36.1)	5 (19.2)	
T/N ratio ≥ 2.5	21 (72.4)	8.8	8.4 (3.8–30.7)	1 (4.8)	0.9335
T/N ratio < 2.5	8 (27.6)	8.9	8.1 (4.3–17.3)	0 (0.0)	
ORR					
CR + PR	17 (30.9)	14.2	13.5 (4.9–36.1)	4 (23.5)	0.0021
SD + PD	38 (69.1)	8.8	8.3 (2.9–30.7)	2 (5.3)	
T/N ratio ≥ 2.5	14 (82.4)	15.4	14.8 (5.8–36.1)	4 (28.6)	0.0199
T/N ratio < 2.5	3 (17.6)	7.3	9.0 (4.9–13.5)	0 (0.0)	
DCR					
CR + PR + SD	26 (47.3)	12.3	10.1 (2.9–36.1)	5 (19.2)	0.0133
PD	29 (52.7)	8.8	8.4 (3.8–30.6)	1 (3.5)	
T/N ratio ≥ 2.5	21 (80.8)	12.7	10.5 (2.9–36.1)	5 (23.8)	0.2072
T/N ratio < 2.5	5 (19.2)	10.7	9.8 (4.9–16.8)	0 (0.0)	

BNCT, boron neutron capture therapy; OS, overall survival; T/N ratio, tumor to normal tissue ratio; CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; ORR, objective response rate; DCR, disease control rate.




Figure 4 Patient survival analysis, and the association between patient survival and tumor to normal tissue ratios derived from ^{18}F -boronophenylalanine positron emission tomography in patients after boron neutron capture therapy. (A-C) Patients who achieved complete response, objective response rate, and disease control rate demonstrated significantly higher median overall survival than those who did not. (D,E) For patients with complete response and objective response rate, tumor to normal tissue ratios greater than 2.5 were associated with significantly higher overall survival than those with tumor to normal tissue ratios lower than 2.5. CR, complete response; PR, partial response; SD, stable disease; PD, progressive disease; T/N ratio, tumor to normal tissue ratio; ORR, objective response rate.

patients was 8.5 months (range, 2.9–36.1 months). Among patients with CR, PR, SD, PD, ORR and DCR, the median OS times were 15.7, 11.6, 7.9, 8.4, 13.5 and 10.1 months, respectively. Patients who achieved CR, ORR, and DCR demonstrated significantly higher median OS than those who did not (median OS 15.7 *vs.* 8.5 months, 13.5 *vs.* 8.3 months, and 10.1 *vs.* 8.4 months, $P=0.0181$, 0.0021 , and 0.0133 , respectively, *Figure 4A-4C*). For patients with CR and ORR, T/N ratios greater than 2.5 were associated with significantly higher OS than those with T/N ratios lower than 2.5 (median OS 16.9 *vs.* 8.5 months and 14.8 *vs.* 9.0 months, $P=0.0253$ and 0.0199 , respectively, *Figure 4D,4E*).

Discussion

BNCT represents a contemporary approach to biochemically targeted radiation treatment, allowing precise delivery of high-dose particle irradiation exclusively to neoplastic cells, thereby sparing adjacent normal tissues. The use of BPA-PET as a reliable tool for patient screening enables noninvasive prediction of tumor uptake and the T/N ratio,

aiding physicians in selecting patients most likely to benefit from BNCT. Treatment planning and dosimetry are tailored based on the tumor uptake of the radioisotope adjusted to BPA concentration in normal tissue. A high T/N ratio is preferred to ensure maximum radiation dose to the tumor and minimal background radiation. BNCT is commonly applied if the target lesion demonstrates a T/N ratio on BPA-PET greater than 2.5 (5,6,17). Prior research by Chen *et al.* evaluated 34 malignant brain tumor patients and revealed improved survival outcomes in patients with a T/N ratio greater than or equal to 4 compared to those with a ratio less than 4 (9). In our current study, there was a significant difference in T/N ratios derived from BPA-PET between patients who received BNCT and those who did not ($P<0.0001$). When considering a T/N ratio of 2.5, 44 of 58 (75.9%) patients in the BNCT group and 26 of 58 (44.8%) patients in the non-BNCT group exhibited T/N ratios greater than 2.5. This difference reached statistical significance ($P=0.0007$). Therefore, a T/N ratio greater than 2.5 is commonly regarded as the standard for BNCT among most physicians in our hospital. Within our database,

merely 3 out of 55 instances in Group 1 exhibited T/N ratios surpassing 4.0 (4.48, 4.70, and 5.56), corresponding to the PR, PR, and SD categories, respectively. The potential significance might emerge with a larger cohort exhibiting T/N ratios exceeding 4.

The expression of LAT1 may play a significant role in tumor progression, as it acts as a tumor-specific probe of LAT1. Wittig *et al.* investigated the correlation between the proportion of tumor cells expressing LAT1 and/or Ki67 and BPA uptake in three tumor entities. Their results showed that neither LAT1 nor Ki67 expression predicted BPA uptake (18). Watanabe *et al.* discovered that the strength of LAT1 expression strongly influenced the BPA uptake ability of tumor cells, with squamous cell carcinomas expressing high levels of LAT1 regardless of the primary tumor site. Higher LAT1 expression in tumors correlated with elevated expression of cell proliferation markers and poorer patient prognosis. Patients with low LAT1 expression in the tumor tissue were more likely to have a better prognosis than those with high LAT1 expression. BPA exhibited higher concentration in tumors with high LAT1 expression, suggesting potential effectiveness of BNCT in tumors with poor prognosis and higher proliferative potential and nutritional requirements (4). However, there was no evident difference in the survival curves of patients with malignant brain tumors based on LAT1 expression (4). In our study, there was no significant difference in patient survival based on T/N ratios greater than 2.5 or not (median OS 7.1 *vs.* 7.0 months, $P=0.8955$). In patients who did not receive BNCT, a higher T/N ratio seemed to be associated with a poorer prognosis (median OS 4.3 *vs.* 6.1 months, $P=0.1602$). However, this observation did not apply to patients who received BNCT (median OS 8.5 *vs.* 8.5 months, $P=0.2630$). The cumulative OS of patients with T/N ratios greater than 2.5 who received BNCT did not show a significant difference compared to those who did not receive BNCT (median OS 8.5 *vs.* 4.3 months, $P=0.0783$). Therefore, our findings suggest that the T/N ratio may not strongly correlate with the survival of patients with malignant brain tumors. However, it remains uncertain whether high LAT1 expression correlates with poor prognosis in solid tumor patients (18,19).

Numerous prior studies have presented comprehensive outcomes for patients treated with BNCT (20-24). However, no comparative data have been explored for patients who had tumors evaluated by BPA-PET but did not receive BNCT treatment. Miyatake *et al.* conducted BNCT in cases of recurrent high-grade gliomas (HGG)

(5). Given the absence of an established standard treatment for recurrent HGG, evaluating the survival benefit after BNCT can be challenging. They compared their results with other published series of recurrent malignant gliomas in a combined cohort ($n=129$) proposed by Carson *et al.*, which revealed a median survival of 4.4 months [95% confidence interval (CI): 3.6–5.4 months] (25). In contrast, in the patients of Miyatake *et al.* with similar prognostic factors ($n=11$), the median survival was 9.1 months (95% CI: 4.4–11.0 months), indicating the potential effectiveness of BNCT in managing recurrent HGG even in cases with a poor prognosis (26). In their initial series, more than 50% reduction of the contrast-enhancing lesion volume was demonstrated during follow-up in 8 of 12 patients (27). Upon juxtaposing the two cohorts, the BNCT group exhibited a median OS of 8.5 months (range, 2.9–36.1 months), whereas the non-BNCT group displayed a median OS of 6.0 months (range, 0.7–38.7 months). The cumulative OS of patients with T/N ratios exceeding 2.5 in Group 1 surpassed that of Group 2 (median OS 8.5 *vs.* 4.3 months, $P=0.0783$). Regrettably, this disparity failed to attain statistical significance. While we did not observe a difference in the entire cohort of malignant brain tumors, patients with an ORR of 30.9% in our study showed significantly higher median OS than those who did not (median OS 13.5 *vs.* 8.3 months, $P=0.0021$). Among the 17 patients with ORR, 14 (82.4%) had a T/N ratio greater than 2.5, indicating significantly higher survival than those without (median OS 14.8 *vs.* 9.0 months, $P=0.0199$, respectively). Therefore, ORR appears to be closely associated with the values of T/N ratio derived from BPA-PET. Interestingly, in patients who did not receive BNCT, a higher T/N ratio was associated with a poorer prognosis. However, after effective BNCT, these patients experienced improved survival. Consequently, BNCT with ORR seems necessary to prolong the survival of patients with malignant brain tumors.

Using accelerator-based BNCT combined with BPA as a boron-carrying drug to treat locally recurrent head and neck cancer was approved in Japan in 2020 based on clinic trial results (28-30). In BPA-based BNCT, BPA-PET is crucial to estimate Boron-concentration quantitatively in the target tumors and surrounding normal tissue before BNCT (31-33). Our results could emphasize the use of BPA-PET before BNCT. Moreover, over-expression of LAT-1 in cancer outside the brain, head, and neck now can be quantitatively imaged by BPA-PET (34,35), thereby extending the role of BPA-PET. Using the theranostic combination of BPA and BPA-PET, we not only can assess

the aggressiveness of tumor using BPA-PET but also can treat them with BNCT.

Nevertheless, our study possesses certain limitations. Firstly, the dataset was derived from a specific population of critically ill patients with end-stage malignant brain tumors who met the criteria for Emergency and Compassionate Use (Expanded Access Program). Because we are the only medical institute performing BPA-PET and BNCT treatment, the pathologic diagnoses in our patients group became mixed and indetermined. Unfortunately, for patients referred from other institutes, KPS data of some patients were lacking. The histology, especially the tumor type “astrocytoma”, “oligoastrocytoma”, and “brainstem glioma” and the grading, cannot be confirmed. Also, some patients suffered from recurrent brain tumors did not receive re-biopsy for the recurrent lesions, despite high-grade transformation were highly suspected on the surveillant images. BNCT was only utilized as a salvage treatment for severe and life-threatening brain tumors. The exact role of management options in multimodality treatment strategies was not discussed in this study. Secondly, the sample size was relatively small and originated from a single institution, introducing potential biases. Thirdly, Bevacizumab was administered promptly following BNCT and at biweekly intervals until disease progression, severe treatment-related toxicity (grade 3), or patient mortality. The administration of Bevacizumab may influence the MRI and RANO assessments. Additionally, comparative data for treatment response were obtained from critically ill patients who voluntarily bore the high costs of fees or not. Further validation is warranted to establish the predictive efficacy of prognostic factors in treatment response and OS.

Conclusions

In conclusion, patients with higher T/N ratios derived from BPA-PET may initially exhibit a poor prognosis, but this outcome can potentially be reversed after BNCT. While BNCT may not be necessary to prolong the survival of all patients with malignant brain tumors, it becomes crucial when achieving an ORR, which appears to be closely associated with the values of T/N ratio. The use of BPA-PET as a screening tool can aid in patient selection for BNCT, ensuring that those most likely to benefit from the treatment receive it, thereby assisting clinicians in identifying patients with the highest likelihood of benefiting from BNCT. Further research is necessary to establish the quantitative assessment techniques in BPA-PET, explore

the relationship between LAT1 expression and patient prognosis in solid tumor cases. Additionally, comparative data between patients who received BNCT and those who did not should be examined to gain further insights into treatment outcomes.

Acknowledgments

We extend our gratitude to Miss Wan-Tien Hsieh for her invaluable statistical assistance and her contributions to patient arrangement and data management throughout the study.

Funding: This work was made possible with the generous support of grant V112C-192 from the esteemed Taipei Veterans General Hospital. This study received partial support from grants provided by the Ministry of Science and Technology of Taiwan (MOST110-2314-B-075-030-MY3) and the Melissa Lee Cancer Foundation.

Footnote

Reporting Checklist: The authors have completed the STROBE reporting checklist. Available at <https://qims.amegroups.com/article/view/10.21037/qims-23-1769/rc>

Conflicts of Interest: All authors have completed the ICMJE uniform disclosure form (available at <https://qims.amegroups.com/article/view/10.21037/qims-23-1769/coif>). All authors report that this work was made possible with the generous support of grant V112C-192 from the esteemed Taipei Veterans General Hospital. This study received partial support from grants provided by the Ministry of Science and Technology of Taiwan (MOST110-2314-B-075-030-MY3) and the Melissa Lee Cancer Foundation. The authors have no other conflicts of interest to declare.

Ethical Statement: The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. The study was conducted in accordance with the Declaration of Helsinki (as revised in 2013). The study was approved by the institutional review board of Taipei Veterans General Hospital (IRB No. 2022-07-0029CC). Due to the retrospective nature of the study, written or verbal informed consent was waived.

Open Access Statement: This is an Open Access article

distributed in accordance with the Creative Commons Attribution-NonCommercial-NoDerivs 4.0 International License (CC BY-NC-ND 4.0), which permits the non-commercial replication and distribution of the article with the strict proviso that no changes or edits are made and the original work is properly cited (including links to both the formal publication through the relevant DOI and the license). See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>.

References

1. Slatkin DN. A history of boron neutron capture therapy of brain tumours. Postulation of a brain radiation dose tolerance limit. *Brain* 1991;114:1609-29.
2. He H, Li J, Jiang P, Tian S, Wang H, Fan R, Liu J, Yang Y, Liu Z, Wang J. The basis and advances in clinical application of boron neutron capture therapy. *Radiat Oncol* 2021;16:216.
3. Malouff TD, Seneviratne DS, Ebner DK, Stross WC, Waddle MR, Trifiletti DM, Krishnan S. Boron Neutron Capture Therapy: A Review of Clinical Applications. *Front Oncol* 2021;11:601820.
4. Watanabe T, Sanada Y, Hattori Y, Suzuki M. Correlation between the expression of LAT1 in cancer cells and the potential efficacy of boron neutron capture therapy. *J Radiat Res* 2023;64:91-8.
5. Miyatake SI, Kawabata S, Hiramatsu R, Kuroiwa T, Suzuki M, Ono K. Boron Neutron Capture Therapy of Malignant Gliomas. *Prog Neurol Surg* 2018;32:48-56.
6. Lan TL, Chou FI, Lin KH, Pan PS, Lee JC, Huang WS, Liu YM, Chao Y, Chen YW. Using salvage Boron Neutron Capture Therapy (BNCT) for recurrent malignant brain tumors in Taiwan. *Appl Radiat Isot* 2020;160:109105.
7. Kageji T, Nagahiro S, Mizobuchi Y, Matsuzaki K, Nakagawa Y, Kumada H. Boron neutron capture therapy (BNCT) for newly-diagnosed glioblastoma: comparison of clinical results obtained with BNCT and conventional treatment. *J Med Invest* 2014;61:254-63.
8. Furuse M, Kawabata S, Wanibuchi M, Shiba H, Takeuchi K, Kondo N, Tanaka H, Sakurai Y, Suzuki M, Ono K, Miyatake SI. Boron neutron capture therapy and add-on bevacizumab in patients with recurrent malignant glioma. *Jpn J Clin Oncol* 2022;52:433-40.
9. Chen YW, Lee YY, Lin CF, Pan PS, Chen JK, Wang CW, et al. Salvage Boron Neutron Capture Therapy for Malignant Brain Tumor Patients in Compliance with Emergency and Compassionate Use: Evaluation of 34 Cases in Taiwan. *Biology (Basel)* 2021;10:334.
10. Chen YW, Lee YY, Lin CF, Huang TY, Ke SH, Mu PF, et al. Compassionate Treatment of Brainstem Tumors with Boron Neutron Capture Therapy: A Case Series. *Life (Basel)* 2022;12:566.
11. Wang LW, Chen YW, Ho CY, Hsueh Liu YW, Chou FI, Liu YH, Liu HM, Peir JJ, Jiang SH, Chang CW, Liu CS, Lin KH, Wang SJ, Chu PY, Lo WL, Kao SY, Yen SH. Fractionated Boron Neutron Capture Therapy in Locally Recurrent Head and Neck Cancer: A Prospective Phase I/II Trial. *Int J Radiat Oncol Biol Phys* 2016;95:396-403.
12. Wang LW, Wang SJ, Chu PY, Ho CY, Jiang SH, Liu YW, Liu YH, Liu HM, Peir JJ, Chou FI, Yen SH, Lee YL, Chang CW, Liu CS, Chen YW, Ono K. BNCT for locally recurrent head and neck cancer: preliminary clinical experience from a phase I/II trial at Tsing Hua Open-Pool Reactor. *Appl Radiat Isot* 2011;69:1803-6.
13. Hiratsuka J, Kamitani N, Tanaka R, Tokiya R, Yoden E, Sakurai Y, Suzuki M. Long-term outcome of cutaneous melanoma patients treated with boron neutron capture therapy (BNCT). *J Radiat Res* 2020;61:945-51.
14. Fujimoto T, Suzuki M, Sudo T, Fujita I, Sakuma T, Sakurai Y, et al. Boron neutron capture therapy for clear cell sarcoma. *Appl Radiat Isot* 2020;166:109324.
15. Lin SY, Lin CJ, Liao JW, Peir JJ, Chen WL, Chi CW, Lin YC, Liu YM, Chou FI. Therapeutic efficacy for hepatocellular carcinoma by boric acid-mediated boron neutron capture therapy in a rat model. *Anticancer Res* 2013;33:4799-809.
16. Wen PY, Macdonald DR, Reardon DA, Cloughesy TF, Sorensen AG, Galanis E, Degroot J, Wick W, Gilbert MR, Lassman AB, Tsien C, Mikkelsen T, Wong ET, Chamberlain MC, Stupp R, Lamborn KR, Vogelbaum MA, van den Bent MJ, Chang SM. Updated response assessment criteria for high-grade gliomas: response assessment in neuro-oncology working group. *J Clin Oncol* 2010;28:1963-72.
17. Chang TY, Chang WY, Chen YW, Ho MC, Chang CW, Lau SO, Peng NJ, Wu CY. Comparison of the synthesis and biological properties of no-carrier-added and carrier-added 4-borono-2-[¹⁸F]fluorophenylalanine ([¹⁸F]FBPA). *Nucl Med Biol* 2023;116-117:108313.
18. Wittig A, Sheu-Grabellus SY, Collette L, Moss R, Brualla L, Sauerwein W. BPA uptake does not correlate with LAT1 and Ki67 expressions in tumor samples (results of EORTC trial 11001). *Appl Radiat Isot* 2011;69:1807-12.
19. Lu JJ, Li P, Yang Y, Wang L, Zhang Y, Zhu JY, Zhu XR, Chen MB. Prognostic value of LAT-1 status in solid cancer: A systematic review and meta-analysis. *PLoS One*

- 2020;15:e0233629.
20. Chanana AD, Capala J, Chadha M, Coderre JA, Diaz AZ, Elowitz EH, Iwai J, Joel DD, Liu HB, Ma R, Pendzick N, Peress NS, Shady MS, Slatkin DN, Tyson GW, Wielopolski L. Boron neutron capture therapy for glioblastoma multiforme: interim results from the phase I/II dose-escalation studies. *Neurosurgery* 1999;44:1182-92; discussion 1192-3.
 21. Henriksson R, Capala J, Michanek A, Lindahl SA, Salford LG, Franzén L, Blomquist E, Westlin JE, Bergenheim AT; Swedish Brain Tumour Study Group. Boron neutron capture therapy (BNCT) for glioblastoma multiforme: a phase II study evaluating a prolonged high-dose of boronophenylalanine (BPA). *Radiother Oncol* 2008;88:183-91.
 22. Kawabata S, Miyatake S, Nonoguchi N, Hiramatsu R, Iida K, Miyata S, Yokoyama K, Doi A, Kuroda Y, Kuroiwa T, Michiue H, Kumada H, Kirihata M, Imahori Y, Maruhashi A, Sakurai Y, Suzuki M, Masunaga S, Ono K. Survival benefit from boron neutron capture therapy for the newly diagnosed glioblastoma patients. *Appl Radiat Isot* 2009;67:S15-8.
 23. Kageji T, Mizobuchi Y, Nagahiro S, Nakagawa Y, Kumada H. Clinical results of boron neutron capture therapy (BNCT) for glioblastoma. *Appl Radiat Isot* 2011;69:1823-5.
 24. Miyatake S, Kawabata S, Hiramatsu R, Kuroiwa T, Suzuki M, Kondo N, Ono K. Boron Neutron Capture Therapy for Malignant Brain Tumors. *Neurol Med Chir (Tokyo)* 2016;56:361-71.
 25. Carson KA, Grossman SA, Fisher JD, Shaw EG. Prognostic factors for survival in adult patients with recurrent glioma enrolled onto the new approaches to brain tumor therapy CNS consortium phase I and II clinical trials. *J Clin Oncol* 2007;25:2601-6.
 26. Miyatake S, Kawabata S, Yokoyama K, Kuroiwa T, Michiue H, Sakurai Y, Kumada H, Suzuki M, Maruhashi A, Kirihata M, Ono K. Survival benefit of Boron neutron capture therapy for recurrent malignant gliomas. *J Neurooncol* 2009;91:199-206.
 27. Miyatake S, Kawabata S, Kajimoto Y, Aoki A, Yokoyama K, Yamada M, Kuroiwa T, Tsuji M, Imahori Y, Kirihata M, Sakurai Y, Masunaga S, Nagata K, Maruhashi A, Ono K. Modified boron neutron capture therapy for malignant gliomas performed using epithermal neutron and two boron compounds with different accumulation mechanisms: an efficacy study based on findings on neuroimages. *J Neurosurg* 2005;103:1000-9.
 28. Advances in boron neutron capture therapy. Non-serial publications, IAEA, Vienna, 2023.
 29. Hirose K, Konno A, Hiratsuka J, Yoshimoto S, Kato T, Ono K, et al. Boron neutron capture therapy using cyclotron-based epithermal neutron source and borofalan (^{10}B) for recurrent or locally advanced head and neck cancer (JHN002): An open-label phase II trial. *Radiother Oncol* 2021;155:182-7.
 30. Matsumura A, Asano T, Hirose K, Igaki H, Kawabata S, Kumada H. Initiatives Toward Clinical Boron Neutron Capture Therapy in Japan. *Cancer Biother Radiopharm* 2023;38:201-7.
 31. Monti Hughes A, Hu N. Optimizing Boron Neutron Capture Therapy (BNCT) to Treat Cancer: An Updated Review on the Latest Developments on Boron Compounds and Strategies. *Cancers (Basel)* 2023;15:4091.
 32. Ishiwata K. 4-Borono-2-(^{18}F -fluoro-L-phenylalanine PET for boron neutron capture therapy-oriented diagnosis: overview of a quarter century of research. *Ann Nucl Med* 2019;33:223-36.
 33. Kashiwara T, Mori T, Nakaichi T, Nakamura S, Ito K, Kurihara H, Kusumoto M, Itami J, Yoshimoto S, Igaki H. Correlation between L-amino acid transporter 1 expression and 4-borono-2-(^{18}F -fluoro-phenylalanine accumulation in humans. *Cancer Med* 2023;12:20564-72.
 34. Minoji T, Takahashi H, Watabe T, Ishikawa S, Kato H, Ohgaki R, Kanai Y, Sekido Y, Hata T, Hamabe A, Ogino T, Miyoshi N, Uemura M, Yamamoto H, Doki Y, Eguchi H. The Clinical Potential of FBPA-PET/CT Imaging for Colorectal Cancer and its Liver Metastases Expressing LAT1. *Anticancer Res* 2023;43:3913-21.
 35. Watabe T, Ose N, Naka S, Fukui E, Kimura T, Kanou T, Funaki S, Sasaki H, Kamiya T, Kurimoto K, Isohashi K, Tatsumi M, Shimosegawa E, Kato H, Ohgaki R, Kanai Y, Shintani Y. Evaluation of LAT1 Expression in Patients With Lung Cancer and Mediastinal Tumors: ^{18}F -FBPA PET Study With Immunohistological Comparison. *Clin Nucl Med* 2023;48:853-60.

Cite this article as: Lin KH, Chen YW, Wang LW, Wang YF, Hu LH, Ting CH, Lee TH, Lee JC, Peng NJ. Prognostic assessment of ^{18}F -boronophenylalanine positron emission tomography (BPA-PET) in salvage boron neutron capture therapy for malignant brain tumors. *Quant Imaging Med Surg* 2024;14(6):4177-4188. doi: 10.21037/qims-23-1769