



The role of astrocytes in prion-like mechanisms of neurodegeneration

Phillip Smethurst,^{1,2} Hannah Franklin,^{1,2,†} Benjamin E. Clarke,^{1,2,†} Katie Sidle¹ and Rickie Patani^{1,2}

[†]These authors contributed equally to this work.

Accumulating evidence suggests that neurodegenerative diseases are not merely neuronal in nature but comprise multicellular involvement, with astrocytes emerging as key players. The pathomechanisms of several neurodegenerative diseases involve the deposition of misfolded protein aggregates in neurons that have characteristic prion-like behaviours such as template-directed seeding, intercellular propagation, distinct conformational strains and protein-mediated toxicity. The role of astrocytes in dealing with these pathological prion-like protein aggregates and whether their responses either protect from or conspire with the disease process is currently unclear. Here we review the existing literature implicating astrocytes in multiple neurodegenerative proteinopathies with a focus on prion-like behaviour in this context.

1 Department of Neuromuscular Disease, UCL Queen Square Institute of Neurology, Queen Square, London WC1N 3BG, UK

2 The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK

Correspondence to: Rickie Patani

The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK

E-mail: rickie.patani@ucl.ac.uk

Correspondence may also be addressed to: Katie Sidle

E-mail: ksidle@doctors.org.uk

Keywords: astrocytes; prion-like; Alzheimer's disease; Parkinson's disease; amyotrophic lateral sclerosis

Abbreviations: ALS = amyotrophic lateral sclerosis; CBD = corticobasal degeneration; FTLN = frontotemporal lobar degeneration; PSP = progressive supranuclear palsy

Introduction

The main component tying together many neurodegenerative diseases is the accumulation and deposition of misfolded proteins. The pathogenic factors underlying the cytotoxic role of aggregated and misfolded proteins are predicted by the prion hypothesis. There are four main behavioural characteristics that make a protein 'prion-like': (i) template-directed seeding and amplification of misfolded proteins to form aggregates; (ii) intercellular spreading

of aggregates and propagation of the templated seeding reaction to nearby cells; (iii) conformational strains that produce distinct clinical and biochemical disease entities; and (iv) protein-mediated cellular toxicity.¹ Indeed, on this last point there is mounting evidence to show that oligomeric intermediate species of these proteins may be responsible for the resultant neurotoxicity driving the disease.^{2,3} Importantly, although many neurodegenerative disease-related proteins possess 'prion-like' behavioural characteristics,¹ they do not, however, typically possess the same high levels

Received March 19, 2021. Revised August 20, 2021. Accepted September 03, 2021

© The Author(s) (2022). Published by Oxford University Press on behalf of the Guarantors of Brain.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

of infectivity and aggressive phenotype as seen with actual prion diseases, and are thus termed ‘prion-like’.

Currently, the differential response of neurons and astrocytes to aggregated proteins with prion-like behaviour remains relatively understudied. Specifically, there is conflicting evidence about the presence of aggregated proteins in astrocytes at post-mortem, how these aggregates affect astrocytes and what role they play in neurodegeneration.^{4–9} Importantly, previous landmark studies have characterized the neurotoxic and neuroprotective roles of astrocytes.^{10–12} These have been followed up by innumerable reports further demonstrating these findings and their possible mechanistic basis across different neurodegenerative diseases.^{13–18} Recent work also suggests that astrocyte reactive transformation can occur cell autonomously in the context of neurodegeneration-causing gene mutations.¹⁹ The purpose of this article is to review the literature that directly explores the role of astrocytes in proteinopathies with prion-like behaviour in the following neurodegenerative diseases: prion disease, Alzheimer’s disease, tauopathies, synucleinopathies, Huntington’s disease and amyotrophic lateral sclerosis (ALS).

Prion disease

Prion diseases, also known as transmissible spongiform encephalopathies, are a group of fatal neurodegenerative diseases characterized by the deposition of misfolded and aggregated prion protein (PrP^{Sc}) in the CNS, often causing severe cerebellar ataxia and rapidly progressive dementia. This is pathologically characterized at post-mortem by spongiosis (focal or diffuse clusters of small vacuoles in the neuropil of the deep cortical layers, subcortical grey matter or cerebellar cortex), reactive transformation of astrocytes, microglial activation and neuronal death. These pathologies result in a clinically heterogeneous set of conditions, whereby certain strains of the protein result in different misfolded conformations of aggregated PrP^{Sc} in distinct CNS regions.²⁰ These different strains of misfolded prion protein result in distinct diseases characterized by unique biochemical and clinical features that include: Creutzfeldt–Jakob disease, kuru, fatal familial insomnia and Gerstmann–Sträussler–Scheinker syndrome. In addition to these human diseases, prions also cause disease in other species including cattle (bovine spongiform encephalopathy), deer (chronic wasting disease), sheep and goats (scrapie). There is now increasing evidence to suggest a dissociation between the aggregation of PrP^{Sc} and toxicity. This disconnect may be explained by the formation of intermediate toxic oligomeric species that have been described both *in vitro* and *in vivo* for the prion protein.²¹ Indeed, aggregates are unlikely to be the active pathological species, but rather a marker that a cell has been exposed to a toxic, misfolded protein. These pathological protein states may consist of a pool of fibrils or oligomers and are most appropriately termed ‘assemblies’ to distinguish these toxic entities from the inert end state aggregates.¹ Importantly, it has also been suggested that PrP toxicity can be triggered by the monomeric form, independently of its oligomeric state.²² Furthermore, the N-terminus of PrP has been shown to function as a toxic effector mediating disease-related spontaneous ionic currents, regulated by the C-terminus.²³

PrP^{Sc} aggregates were first detected in astrocytes in scrapie infected sheep and were noted to form prior to neuronal aggregates and before the development of the cardinal scrapie pathological features (reactive transformation of astrocytes, vacuolation, neuronal loss and amyloid deposition).²⁴ This alluded to the potential role of astrocytes in attempting to alleviate pathology or in promoting the replication and spread of the pathological protein. A later study in rodents identified high levels of PrP^C expression in astrocytes and oligodendrocytes and provided further

evidence to support the role of glia in PrP^{Sc} propagation.²⁵ In mice expressing a hamster PrP under an astrocyte-specific promoter, infection with hamster scrapie led to high levels of PrP^{Sc} infection with the mice exhibiting a severe disease phenotype.²⁶ Indeed, detailed microscopy studies of prion pathology in the cerebellum of scrapie infected sheep have also highlighted the ability of astrocytes to sustain active prion infection with glial PrP^{Sc} pathology being detected in all cases.^{27,28} A further technical advance using flow cytometry allowed the identification of PrP^{Sc} in neurons, astrocytes and microglia early on during prion infection in mice.²⁹ Collectively, these findings highlight a clear involvement of astrocytes in the propagation of infectivity and induction of prion disease. Another study using hamster scrapie as a model showed that the 263K strain induced PrP^{Sc} in ~90% of reactive astrocytes suggesting that astrocytes may be able to produce PrP^{Sc} themselves, initiating a mechanism that may further trigger reactive transformation of astrocytes and drive the spread of both pathology and toxicity.³⁰ Interestingly, however, a follow-up study from the same group using the same model also reported PrP^{Sc} aggregation in neurons (in the absence of PrP expression) and in the extracellular neuropil space. While they detected reactive changes in the astrocytes, no further astrocytic phenotypes were reported.⁴ Further work has subsequently demonstrated cell type specific differential toxicity when comparing neuron and astrocyte-specific expression of PrP and scrapie infection. Notably, animals with neuron specific expression succumbed to disease in 100 days compared to 360 days for astrocyte-specific expression.⁵ These data suggest that while neurons remain the predominant pathologically affected cell types in prion disease, astrocytes are also involved in driving this pathology, at least to some degree.

Astrocyte PrP^{Sc} infectivity has also been replicated in rodent *in vitro* models that induce neuronal death,³¹ and PrP^{Sc} pathology is actively taken up in human astrocyte cell lines from CJD patients where degradation of both proteinase K (PK) sensitive (PrP^{Sen}) and PK resistant (PrP^{Res}) forms of PrP^{Sc} was noted, highlighting a putative protective role for astrocytes in clearing prion disease pathology.³² Indeed, these data were corroborated from studies of hamster PrP^{Res} protein, which demonstrated that astrocytes and meningeal fibroblasts uptake PrP^{Res} more efficiently than neurons, potentially as a protective mechanism for protease digestion and clearance of infection.³³ As a potential secondary effect of this, astrocytes could be propagating the resistant prion proteins to other cell types and aiding further infection. A more recent study showed that, in primary cultures of cerebellar neurons and astrocytes, the astrocytes were more prone to prion replication and accumulation of aggregated PrP^{Sc} and that prion pathology can be transferred between astrocytes and neurons through cell–cell contact occurring via tunnelling nanotubes.³⁴ This tunnelling nanotube mechanism of spread and propagation may serve as both neuroprotective on one hand and also aid in sustaining prion infection on the other.

In the human prion diseases, the presence of a polymorphic variant (methionine/methionine, methionine/valine or valine/valine) at codon 129 of the prion gene (*PRNP*) can determine disease susceptibility and disease phenotype in patients with prion disease³⁵ through the production of distinct conformers of misfolded PrP (‘strains’). One study demonstrated that multiple CJD prion strains can be replicated in human induced pluripotent stem cell (iPSC)-derived astrocytes in a *PRNP* codon 129 genotype dependent manner,³⁶ providing further support to the role of astrocytes in potentially influencing disease susceptibility or phenotype. Together, these findings demonstrate that not only do astrocytes accumulate PrP^{Sc} in prion diseases, but they may also be driving early neuroprotective responses by clearing PrP^{Sc} infection. These neuroprotective responses may at the same time serve as a double-edged

sword that can assist in sustaining the prion infection for long enough until the astrocytes become neurotoxic and exacerbate the pathological process by propagating infection at the later stages of disease (Fig. 1).

Alzheimer's disease: amyloid- β

Alzheimer's disease is a progressive neurodegenerative disease, causing irreversible loss of neurons leading to a progressive cognitive decline. One of the hallmarks of Alzheimer's disease is the deposition of aggregated misfolded amyloid- β (A β) protein as extracellular plaques in the neuropil. Amyloid- β is processed via amyloid precursor protein (APP) at the cell membrane where it is cleaved by gamma secretase into various fragments including A β 40 and A β 42. It is now well established that aggregated amyloid- β has prion-like behaviour due to its capacity for seeding, ability to spread and to propagate, form distinct strains and exert toxicity via its aggregated oligomeric intermediates both *in vitro* and *in vivo* (reviewed in Watts and Prusiner³⁷). However, an important distinction in the context of amyloid- β , when compared to other proteins described in this review, is that it is accumulating extracellularly. Indeed, the most toxic species consists of soluble oligomeric assemblies, which are generated intracellularly and then secreted extracellularly, propagating via cell to cell spread.^{38–41}

As amyloid- β plaques are extracellular in the human Alzheimer's disease brain, they do not form in astrocytes. However, astrocytes have well known roles in the secretion, uptake and clearance of aggregated neuronal amyloid- β peptides.^{41–44} Furthermore, the soluble oligomeric aggregated species of amyloid- β , rather than insoluble fibrillar amyloid- β , are thought to be responsible for triggering neurotoxicity.^{40,46–50} While the impact of these oligomers is well understood in neurons, their effect on astrocytes is still not well characterized.

Reactive transformation of astrocytes is one of the earliest detectable features in Alzheimer's disease. Astrocyte reactivity is accompanied by the release of pro-inflammatory cytokines such as TNF- α , IL-6 and IL1 β , which are thought to mediate neurotoxic effects.^{11,51} Evidence suggests that amyloid- β oligomers can trigger the release of these cytokines more acutely than fibrillar forms, producing higher levels of inducible nitric oxide synthase (iNOS), nitric oxide (NO) and TNF- α in primary astrocyte cultures.⁵² Apolipoprotein E4 (ApoE4) is the most prevalent genetic risk factor for Alzheimer's disease and is expressed in more than half of patients. Studies suggest that low levels of total apoE, as exhibited by ϵ 4 carriers, may directly contribute to the disease progression.⁵³ Furthermore, it has also been shown that toxicity of A β 42 oligomers is exacerbated when neurons are co-cultured with ApoE4 knockout astrocytes.⁵⁴ Interestingly, another study demonstrated that an injection of A β 42 oligomers caused neuronal toxicity via mechanisms initiated by astrogliosis and reactive transformation rather than direct effects on neurons.⁵⁵ Furthermore, A β 42 oligomers were shown to induce astrocyte reactivity and also reduce astrocyte viability in primary cultures in a dose dependent manner.^{55,56} A more recent study from the same group identified specific nonamer and dodecamer soluble amyloid- β oligomers in the brains of mutant PSEN1 mice that exhibit cognitive decline. In this case, these oligomers were found to specifically amplify in astrocytes, where they induced and augmented neuronal damage non-cell autonomously through reactive astrocyte transformation.⁵⁷

Mechanistically, this increase in astrocyte reactivity may be driven by A β 42 oligomer disruption of ER calcium homeostasis leading to ER stress and reactive astrocyte transformation.⁵⁸ Indeed, oligomeric A β 40 and A β 42 were found to specifically increase calcium signalling in astrocytes and not neurons. These

oligomers were also found to increase levels of reactive oxygen species (ROS), induce mitochondrial depolarization, reduce levels of the antioxidant GSH and cause metabolic dysfunction in astrocytes.^{59,60} This increase in ROS secretion from oligomer treated astrocyte cultures was also found to increase neuronal calcium signalling that in turn reduced protective STAT3 signalling.⁶¹ Some of this toxicity may be driven via mGluR5 receptor dependent ATP release from astrocytes,^{61,62} and activated scavenger receptors may be responsible for the phagocytic uptake of amyloid- β oligomers that drives glial toxicity via the hexapeptide XD4.⁶³

Amyloid- β oligomers have been found to induce reactivity and reduce neuroprotective capacity in primary astrocyte cultures and *in vivo* via alteration in the secretion of the synaptogenic factor TGF- β 1.⁶ Further neuroprotective benefits of astrocytes have been shown to be mediated by IGF1, which can release neuron bound amyloid- β oligomers and astrocyte oligomer treatment can reduce IGF1 signalling to neurons.⁶⁴ In contrast, another study demonstrated that oligomeric amyloid- β was specifically toxic to neurons and not astrocytes even at the highest concentrations and that this toxicity was JUN-kinase dependent.⁷

The neuroprotective capacity of astrocytes against amyloid- β pathology may lie in some partial uptake of the aggregates for clearance during the early phases of amyloid- β aggregation, but once levels of aggregation are high enough, a neurotoxic role seems to supersede this protection and they become instrumental in the degenerative process. However, the role of astrocytes in the processing, seeding and transmission of amyloid- β pathology remains incompletely resolved and requires further investigation.

Tauopathies

The term tauopathy refers to a collection of progressive age-related neurodegenerative diseases that are clinically and biochemically heterogeneous, but commonly involve the deposition of misfolded, aggregated and phosphorylated tau inclusions in the cells of the CNS. The tau protein aggregates in the form of neurofibrillary tangles (NFTs), pre-tangle deposits and neuropil threads within neuronal and astrocytic processes. The primary tauopathies include frontotemporal lobar degeneration (FTLD) with tau, corticobasal degeneration (CBD), Pick's disease (PiD) and progressive supranuclear palsy (PSP), each of which can express various isoform ratios (3R and 4R), conformations and subcellular localization of tau pathology. While tau is predominantly expressed in neurons^{65,66} it is expressed at lower levels in glia (oligodendrocytes > astrocytes).⁶⁷ Multiple lines of evidence now demonstrate that tau exhibits prion-like behaviour,⁶⁸ and that astrocytes can actively participate in the spread and/or clearance of aggregated tau species.⁶⁵

Alzheimer's disease-related tau pathology may be distinct from other tauopathies given that tau aggregation in this context is considered a secondary event to the deposition of extracellular amyloid- β plaques. Furthermore, wild-type tau aggregation is present in Alzheimer's disease-related tau pathology rather than tau aggregation driven by MAPT gene mutations that can cause other tauopathies.⁶⁹ Tau aggregates in Alzheimer's disease contain both 3R and 4R forms of tau deposited as intraneuronal neurofibrillary tangles, but astrocytic tau pathology is rarely detected.⁶⁹ However, astrocyte reactivity is a well observed phenomenon in Alzheimer's disease⁷⁰ and is therefore thought to play a central role in disease pathogenesis. Studies examining the effect of any aggregated species, including oligomers and fibrils, from Alzheimer's disease patients on astrocytes have yet to be conducted but will provide important insights.

Astrocytes exhibit a range of morphological phenotypes across different tauopathies including tufted astrocytes in PSP, astrocytic

plaques in CBD, thorn shaped astrocytes, granular fuzzy astrocytes in ageing-related tau astrogliopathy (ARTAG), ramified astrocytes in PiD and astrocytes with globular inclusions in globular glial tauopathy. One of the first studies to note paired helical filament or phospho-tau S396/S404 positive tau pathology in aged mice demonstrated that astrocytes may be involved in the uptake and spread of tau pathology.⁷¹ Interestingly, oligodendrocytes are also known to form tau inclusions in the form of coiled bodies and occasionally globular inclusions.⁷² One study demonstrated that tau strains from CBD and PSP post-mortem material could seed specifically in oligodendrocytes and astrocytes. Additionally, glial tau seeding was achieved in non-transgenic mice, thereby implicating glial tau pathology in the progression of tauopathies.⁷³ The same group reported that astrocytic tau pathology from CBD patients and tufted astrocyte pathology from PSP patients could be transmitted via their unique tau strains. Oligodendroglial tau seeding and spreading could propagate along white matter tracts in a neuronal tau knock-down (KD) mouse model whereas astrocytic tau pathology could not. These findings suggest that oligodendroglial tau pathology potentially has a distinct pathogenic impact when compared to astrocytic pathology.⁷⁴ Furthermore, astroglial tau pathology from patients with ARTAG can be seeded to wild-type mice in neuronal and glial populations in the brain,⁷² and CBD astrocytic plaques and tufted astrocytes from PSP patients can also be transmitted via unique strains maintaining a glial cell type specificity for propagation.⁷⁵ Together these data show that some tau ‘strains’ are more likely to affect glia than others and further implicate a nuanced glial involvement in tauopathies.

The effects of tau inclusions on the health of astrocytes are currently unknown and represent an interesting research avenue for future studies. As glial cells typically do not secrete or express much tau protein, the deposition of tau pathology in glial cells is an unusual phenomenon that may be explained by the uptake of neuron-derived monomeric and aggregated tau from the extracellular space. Uptake of extracellular tau by astrocytes cells can occur via many mechanisms⁷⁶ with the uptake of fibrillar and aggregated species occurring mainly via the lysosomal pathway^{77,78} and heparan sulphate proteoglycans (HSPGs),^{77–79} and monomeric tau uptake via an unknown HSPG independent mechanism.⁸⁰ Indeed, a recent paper identified low density lipoprotein receptor-related protein 1 (LRP1) as a novel receptor that mediates tau uptake *in vitro* and *in vivo* models of tau spread with a significant role in astrocytic (as well as neuronal) tau uptake.⁷⁹ In the early stages of disease, this uptake may represent a protective mechanism becoming pathological at a later stage when cellular clearance mechanisms may become overwhelmed. Furthermore, recent data suggest that filamentous tau aggregates can activate integrin signalling in astrocytes where seeds can be taken up and induce further astrocytic activation to release a variety of pro-inflammatory chemokines and cytokines constituting a deleterious reactive astrocyte phenotype.⁸¹ Recent data showed that astrocytes were found to rapidly accumulate toxic tau oligomers in an APP-dependent manner whereby intracellular calcium signalling and calcium-dependent release of gliotransmitters were disrupted. This subsequently induced synaptic dysfunction in neighbouring neurons suggesting that astrocytes play a major role in Alzheimer’s disease pathology by exacerbating synaptic toxicity via the reduction of gliotransmitters.⁸² It should also be noted that astrocytic tau inclusions have been found in aged mice independent of dementia suggesting that astrocytes may also have a role in internalizing and clearing tau.^{83–85} However, further studies of astrocyte activity during the early and late stages of aggregation in *in vitro* co-culture models and *in vivo* models will shed more light on the temporal role of astrocytes in the pathogenesis in the tauopathies.

Synucleinopathies

Parkinson’s disease, dementia with Lewy bodies (DLB) and multiple system atrophy (MSA) constitute a group of neurodegenerative conditions known as synucleinopathies, pathologically represented by the presence of aggregated α -synuclein protein. α -Synuclein is the main component of a pathological hallmark of synucleinopathies known as Lewy bodies and Lewy-like inclusions, found in the neurons of patients with Parkinson’s disease and DLB, and in the astrocytes and oligodendrocytes of patients with MSA.^{86–88} Indeed, α -synuclein inclusions in oligodendrocytes termed glial cytoplasmic inclusions (GCIs) are considered the pathological hallmark of MSA specifically, raising the hypothesis that glial cytoplasmic inclusions drive pathogenesis in this condition. Astrocytic α -synuclein pathology is less common than oligodendrocyte GCIs⁸⁹ and is prevalent at later stages of Parkinson’s disease and DLB.⁹⁰ Indeed, astrocytic α -synuclein is also found in 40% of MSA patients, predominantly in subpial and periventricular regions.⁹¹ These cell types have low levels of expression of α -synuclein and the mechanistic basis for the development of these inclusions is not completely understood.

α -Synuclein is a predominantly synaptic protein that is secreted from neuronal cells by exocytosis.⁹² Oligomeric forms of the protein were found to be elevated in CSF and blood plasma of Parkinson’s disease patients.⁹³ One of the first papers to report the effects of α -synuclein on astrocytes showed that aggregated α -synuclein can be exocytosed from neurons and taken up by astrocytes, forming inclusions and causing an upregulation of inflammatory response genes.⁹⁴ Subsequent studies demonstrated that various recombinant forms of α -synuclein could induce microglial and astrocyte reactivity, but toll-like receptor 4 (TLR4) was not required for uptake in astrocytes, unlike microglia.^{95,96} Another study demonstrated the uptake of recombinant α -synuclein into cortical neurons, astrocytes and fibroblasts, and reported mitochondrial dysfunction in these cell types.⁹⁷ Using purified α -synuclein aggregates from human Parkinson’s disease brain tissue, high content imaging and microfluidics on primary rat cortical neurons and astrocytes, one group has shown that astrocytes can more readily uptake α -synuclein aggregates when compared to neurons via endocytosis. It was also seen that these aggregates could readily transfer between these cell types, resulting in neuronal death if transferred from astrocytes to neurons,⁹⁸ thereby demonstrating a gain of toxic astrocyte function. Loria et al.⁹⁹ interestingly demonstrated that this spread and propagation of α -synuclein fibrils was cell type dependent. They demonstrate that transfer of fibrillar alpha synuclein is most optimal from astrocyte to astrocyte and neuron to astrocyte but less efficient from astrocyte to neuron. Furthermore, astrocytes were able to more efficiently degrade this fibrillar alpha synuclein than neurons providing further evidence to support the hypothesis that astrocytes can capture and degrade pathological α -synuclein as a neuroprotective response.⁹⁹ Another study demonstrated the accumulation of α -synuclein oligomers in human embryonic stem cell derived astrocytes, where failure of the lysosomal digestion pathway to process excess oligomers resulted in inclusion formation in the trans Golgi network, resulting in ER swelling and mitochondrial abnormalities. This study also demonstrated that upon oligomer accumulation, astrocytes formed tunnelling nanotubes to secrete accumulated oligomers to neighbouring healthy astrocytes. These data suggest a potential early protective mechanism and role for astrocytes in the transfer of α -synuclein in the disease process.⁸

Another study introducing monomeric, oligomeric and fibrillar forms of synuclein to primary rat astrocyte cultures described changes akin to reactive transformation such as morphological changes, and an increase in intracellular ROS, cytokine release and GFAP expression.⁹ Here, co-culture experiments revealed




Figure 1 Diagrammatic representation of hypothetical astrocytic involvement in prion-like mechanisms of early- and late-stage neurodegenerative disease. The early stage represents an increased spread, uptake and clearance of aggregates and oligomers from neurons for degradation with continued metabolic and trophic support. Late-stage disease represents a decrease in aggregate and oligomer uptake, degradation and clearance, increased reactive astrocyte transformation, increased aggregate spread and propagation to other neurons and astrocytes and increased oligomer build up resulting in increased neuronal toxicity. Created with BioRender.com.

significant astrocyte-derived cytotoxicity to neurons, with fibrillar α -synuclein-treated astrocytes inducing the highest levels of toxicity. However, oligomeric α -synuclein treatment was the only condition capable of inducing mitochondrial dysfunction and increased hydrogen peroxide production within astrocytes. The levels of TNF- α and cytokine gene expression also varied depending on the species of α -synuclein used. Cumulatively, this study suggested that both oligomeric and fibrillar species potentially contribute to astrocyte mediated toxicity, but through distinct mechanisms and potencies.⁹ Interestingly, different morphological subtypes of α -synuclein inclusion (filamentous and granular) have also been demonstrated within primary astrocyte cultures derived from mutant (A53T or A30P) human α -synuclein transgenic mice exposed to MSA brain homogenate. This study suggested that α -synuclein expression levels determine selective tropism of aggregation.¹⁰⁰

Collectively, these data present a mixed picture of astrocytes playing both a protective and a deleterious role in response to α -synuclein aggregation. Indeed, many of these observations can potentially be attributed to differences in the models used for each study including species (both of cell model and inoculum), astrocyte subtype and duration of treatment. The key to elucidating the role of astrocytes in α -synuclein aggregation induced toxicity lies in further investigation using more physiologically relevant cell types such as human iPSC-derived region-specific astrocytes in co-culture with neurons.^{101–103}

Huntington's disease

Huntington's disease is an autosomal dominant neurodegenerative disease caused by a CAG expansion within the huntingtin gene (*HTT*) that leads to the development of a movement disorder, psychiatric and cognitive symptoms. Translated polyglutamine expansions in the huntingtin protein, if higher than 35 repeats, lead to the aggregation of huntingtin (*HTT*) protein and the development of Huntington's disease. A pathological hallmark of Huntington's disease is the presence of aggregated *HTT* in the form of neuronal intranuclear inclusions located in the striatum. It is thought that such pathology causes the substantial degeneration of medium spiny neurons, but *HTT* aggregation has also been identified in astrocytes, and its role in these cells as part of the disease process is still not clear.^{97,98,104–109} Increasing evidence now suggests that *HTT* also demonstrates prion-like behaviour including the spreading and propagation of *HTT* aggregates in multiple *in vitro* cellular models (reviewed in Bradford et al.¹⁰⁶). However, the role of astrocytes in response to this prion-like behaviour of *HTT* is still not well understood and may provide key insight into the pathogenesis and therapeutic opportunities for Huntington's disease.

Early evidence suggested that glial expression of mutant *HTT* containing polyQ expansion of 115–150 repeats induces its accumulation in glial nuclei in Huntington's disease brains, accompanied by a decrease in glutamate transporters and glutamate uptake in glial cultures and mouse Huntington's disease brains.¹⁰⁵ Wild-type glia were found to be neuroprotective in co-culture models, whereas neurons cultured with mutant Huntington's disease glia were vulnerable to glutamate-induced excitotoxicity.¹¹⁰ The same group developed a transgenic Huntington's disease model with astrocyte-specific expression of the more toxic N-terminal *HTT* fragments (160Q) which displayed an age-related motor phenotype and glutamate transporter deficiency.^{110,111} This mouse model was crossed with another neuronally expressed polyQ *HTT* repeat mouse, resulting in a more severe phenotype and increased vulnerability to excitotoxic stress.¹¹² One group used adeno-associated virus expression of the N-terminal (1–552) fragment of *HTT* with 100Q repeats in primary cortical neurons and found that some astrocytes developed *HTT* inclusions that sequestered clathrin and Golgi complexes, resulting in a significant decrease in secretion of mature brain-derived neurotrophic factor (BDNF), which in turn disrupted neurite development in primary cortical neurons.¹¹³ A more recent study using primate iPSC-derived astrocytes from transgenic mutant *HTT* (mHTT) monkeys also demonstrated the formation of glial *HTT* inclusions which resulted in a reduction in the following parameters: expression of superoxide dismutase 2 (SOD2) (a mitochondrial enzyme that converts superoxide radicals into hydrogen peroxide) and peroxisome proliferator activated receptor γ (PPAR γ) co-activator 1 α (PGC1) (a regulator of mitochondrial biogenesis and oxidative stress), glutamate uptake, 4-aminopyridine (4-AP) response (a non-selective voltage dependent K⁺ channel blocker), and aberrant electrophysiological responses.¹¹⁴ Taken together, these data demonstrate that mHTT aggregates can readily form in

Table 1 Astrocytes are increasingly recognized as playing fundamental roles in a range of neurodegenerative diseases

Common astrocyte-specific pathogenic mechanisms	Proteinopathy (disease-specific protein)					
	Prion disease (PrPsc)	Alzheimer's disease (amyloid- β)	Tauopathies (Tau)	Synucleinopathies (α -synuclein)	Huntington's disease (HTT)	Amyotrophic lateral sclerosis (TDP-43)
Reactive transformation	✓	✓	✓	✓	✓	✓
Increased protein uptake	✓	✓	✓	✓	✓	✓
Accumulation of misfolded protein aggregates/disrupted clearance	✓	✓	✓	✓	✓	✓
Protein transfer/propagation	✓		✓	✓	✓	
Astrocyte-derived toxicity to neurons		✓	✓	✓		
Mitochondrial/metabolic dysfunction		✓		✓		
Disrupted Ca ²⁺ homeostasis		✓	✓			

astrocytic populations and probably influence pathogenesis via alteration of mitochondrial responses to oxidative stress and neuronal activity.

The prion-like transmission of HTT aggregates from neurons to glia was first demonstrated in a *Drosophila* model of Huntington's disease.¹¹⁵ Here, glia were shown to regulate the number of HTT aggregates in neurons through a clearance mechanism mediated via Draper signalling. Draper is the *Drosophila* orthologue of *MEGF10* in mammals, and is an engulfment receptor that regulates the clearance of cellular debris. These data indicate that glia may be contributing to the clearance and transmission of aggregates throughout the CNS and that modulating *MEGF10* signalling may be a potential therapeutic target for reducing spread. Another study using a mouse model with basal ganglia restricted expression of N-terminal HTT fragments in either astrocytes, neurons or both demonstrated that astrocytes are less vulnerable to HTT aggregates. The study also showed that mutant HTT in neurons affected key components of the glial glutamate-glutamine cycle, at least at the mRNA level.¹¹⁶ Some of the mechanistic insights into this increased resistance of astrocytes to mutant HTT aggregates may arise from the increased rate of their degradation in astrocytes compared to neurons, increased levels of proteasome function in clearing soluble K48 linked ubiquitinated HTT,¹¹⁷ increased expression of heat shock proteins such as Hsp70 and C-terminus Hsp70 interacting protein (CHIP) and a reduced expression of HSPB1 protein (a CHIP inhibitory protein) compared to neurons.^{117,118} The evidence for the role of astrocytes and their functional handling of HTT is still in its infancy and requires further investigation; however, these aforementioned data suggest astrocytes may be contributing to disease progression and toxicity, and do appear to play a key role in spreading and propagation of HTT aggregates in the disease process.

TDP-43 proteinopathies

Trans active DNA response binding protein (TARDBP) with a molecular weight of 43 kDa (TDP-43) is a DNA and RNA binding protein. Its abnormal cytoplasmic aggregation is considered the pathological hallmark of ~97% of cases of ALS and ~50% of cases of fronto-temporal lobar degeneration (FTLD-TDP), which have now been reclassified as TDP-43 proteinopathies. Mutations in the TARDBP gene are also causative of a small percentage of ALS and FTLD cases giving TDP-43 a central role in the pathogenesis of ALS and FTLD.^{119–121}

TDP-43 is a predominantly nuclear protein that can dynamically shuttle between the nucleus and the cytoplasm. In ALS and FTLD it becomes mislocalized to the cytoplasm where it is cleaved,

hyperphosphorylated and aggregated. The presence of aggregated TDP-43 is well documented in neurons and is also a common feature in oligodendrocytes.^{122–124} However, the presence of TDP-43 pathology in astrocytes is not well documented and less well studied in ALS and FTLD-TDP. Interestingly, astrocytic TDP-43 pathology is a key feature of Alexander disease, a condition caused by mutations in *GFAP*,¹²⁵ but in this case TDP-43 is not cleaved into C-terminal fragments like in ALS and FTLD-TDP. TDP-43 pathology is also recognized in white matter astrocytes in FTLD-Tau, CBD,¹²⁶ Cockayne syndrome¹²⁷ and brain tumours¹²⁸ suggesting that astroglial TDP-43 pathology does occur in some neurological diseases, but that its pathological relevance is unclear in ALS. *In vitro*, human iPSC-derived astrocytes with a TDP-43 M337V mutation have been reported to show cytoplasmic TDP-43 mislocalization and a cell autonomous survival phenotype.¹²⁹ However, the generalizability of astrocytic cytoplasmic TDP-43 proteinopathy requires further systematic investigation across different model systems.

There is mounting evidence to demonstrate that TDP-43 is also a prion-like protein demonstrating many features of prion-like behaviour,¹³⁰ yet few studies examining the pathological relevance astrocytes in this context have been undertaken. One group used rat neural stem cells differentiated to neurons, astrocytes and oligodendrocytes that virally overexpressed TDP-43 and used proteasome inhibition to induce TDP-43 aggregates in all three cell types. With time lapse imaging they could detect the formation of these aggregates in all three cell types, which was followed by cell death after 72 h and spreading of aggregates to neighbouring cells.¹³¹ This study indeed suggested that the aforementioned cell types are susceptible to pathological TDP-43 protein spread and related toxicity. To begin to address the differences in spread and susceptibility to toxicity between these cell types, we recently demonstrated that control human spinal iPSC-derived motor neurons and astrocytes can seed TDP-43 pathology from serially passaged ALS spinal cord extracts.¹³² We also showed that astrocytes were comparatively resistant to TDP-43 seeding, seeding-induced toxicity and recombinant TDP-43 oligomer-induced toxicity compared to motor neurons. Indeed, upon co-culture with motor neurons we demonstrated that these seeded aggregates can spread between both cell types with a predilection for motor neuron to astrocyte spread. The astrocytes were also neuroprotective to the motor neurons with no deleterious reactive astrocyte changes detected after co-culture.¹³² These data indicate that astrocytes are less vulnerable to TDP-43 protein-mediated toxicity and can—at least on initial exposure—be neuroprotective by clearing aggregated TDP-43 from motor neurons. However, at later disease stages they may be unable to clear aggregates and may participate in the spread and neurotoxic process. The elucidation of the exact

mechanisms of these differential responses of neurons and astrocytes to TDP-43 pathology require further investigation and may highlight novel therapeutic targets to ameliorate protein-mediated toxicity.

Conclusion

Multiple lines of evidence discussed herein implicate astrocytes as key players in proteinopathies in terms of their roles in developing/uptaking, transmitting and clearing pathological protein aggregates in multiple neurodegenerative diseases (Table 1). In addition, they may develop unique responses to protein aggregation where they can display a dual role in mounting initial neuroprotective responses followed by deleterious activity later in the disease process (Fig. 1). The delineation of these exact mechanisms and pathways are exciting areas of research and could lead to the discovery of much needed clinically impactful disease-modifying therapies for these devastating and incurable conditions.

Funding

This work was supported by the Francis Crick Institute which receives its core funding from Cancer Research UK 65 (FC010110), the UK Medical Research Council (FC010110) and the Wellcome Trust (FC010110). R.P. holds an MRC Senior Clinical Fellowship (MR/S006591/1) and a Lister Research Prize Fellowship.

Competing interests

The authors report no competing interests.

References

- Collinge J. Mammalian prions and their wider relevance in neurodegenerative diseases. *Nature*. 2016;539(7628):217–226.
- Andreasen M, Lorenzen N, Otzen D. Interactions between misfolded protein oligomers and membranes: A central topic in neurodegenerative diseases? *Biochim Biophys Acta*. 2015;1848(9):1897–1907.
- Forloni G, Artuso V, La Vitola P, Balducci C. Oligomeropathies and pathogenesis of Alzheimer and Parkinson's diseases. *Mov Disord*. 2016;31(6):771–781.
- Jeffrey M, Goodsir CM, Race RE, Chesebro B. Scrapie-specific neuronal lesions are independent of neuronal PrP expression. *Ann Neurol*. 2004;55(6):781–792.
- Kercher L, Favara C, Chan C-C, Race R, Chesebro B. Differences in scrapie-induced pathology of the retina and brain in transgenic mice that express hamster prion protein in neurons, astrocytes, or multiple cell types. *Am J Pathol*. 2004;165(6):2055–2067.
- Diniz LP, Tortelli V, Matias I, et al. Astrocyte transforming growth factor beta 1 protects synapses against A β oligomers in Alzheimer's disease model. *J Neurosci*. 2017;37(28):6797–6809.
- Ebenezer PJ, Weidner AM, LeVine H 3rd, et al. Neuron specific toxicity of oligomeric amyloid- β : Role for JUN-kinase and oxidative stress. *J Alzheimers Dis*. 2010;22(3):839–848.
- Rostami J, Holmqvist S, Lindström V, et al. Human astrocytes transfer aggregated alpha-synuclein via tunneling nanotubes. *J Neurosci*. 2017;37(49):11835–11853.
- Chavarría C, Rodríguez-Bottero S, Quijano C, Cassina P, Souza JM. Impact of monomeric, oligomeric and fibrillar alpha-synuclein on astrocyte reactivity and toxicity to neurons. *Biochem J*. 2018;475(19):3153–3169.
- Zamanian JL, Xu L, Foo LC, et al. Genomic analysis of reactive astrogliosis. *J Neurosci*. 2012;32(18):6391–6410.
- Liddel SA, Guttenplan KA, Clarke LE, et al. Neurotoxic reactive astrocytes are induced by activated microglia. *Nature*. 2017;541(7638):481–487.
- Anderson MA, Burda JE, Ren Y, et al. Astrocyte scar formation aids central nervous system axon regeneration. *Nature*. 2016;532(7598):195–200.
- Serrano-Pozo A, Muzikansky A, Gómez-Isla T, et al. Differential relationships of reactive astrocytes and microglia to fibrillar amyloid deposits in Alzheimer disease. *J Neuropathol Exp Neurol*. 2013;72(6):462–471.
- Grubman A, Chew G, Ouyang JF, et al. A single-cell atlas of entorhinal cortex from individuals with Alzheimer's disease reveals cell-type-specific gene expression regulation. *Nat Neurosci*. 2019;22(12):2087–2097.
- Mathys H, Davila-Velderrain J, Peng Z, et al. Single-cell transcriptomic analysis of Alzheimer's disease. *Nature*. 2019;570(7761):332–337.
- Zhou Y, Song WM, Andhey PS, et al. Human and mouse single-nucleus transcriptomics reveal TREM2-dependent and TREM2-independent cellular responses in Alzheimer's disease. *Nat Med*. 2020;26(1):131–142.
- Ziff OJ, Taha DM, Crerar H, et al. Reactive astrocytes in ALS display diminished intron retention. *Nucleic Acids Res*. 2021;49(6):3168–3184.
- Ziff OJ, Clarke BE, Taha DM, Crerar H, Luscombe NM, Patani R. Meta-analysis of human and mouse ALS astrocytes reveals multi-omic signatures of inflammatory reactive states. *Genome Res*. 2022;32(1):71–84.
- Taha DM, Clarke BE, Hall CE, et al. Astrocytes display cell autonomous and diverse early reactive states in familial amyotrophic lateral sclerosis. *Brain*. 2022;318(5852):930–936. <https://doi.org/10.1093/brain/awab328>
- Collinge J, Clarke AR. A general model of prion strains and their pathogenicity. *Science*. 2007;318(5852):930–936.
- Huang P, Lian F, Wen Y, Guo C, Lin D. Prion protein oligomer and its neurotoxicity. *Acta Biochim Biophys Sin*. 2013;45(6):442–451.
- Zhou M, Ottenberg G, Sferrazza GF, Lasmézas CI. Highly neurotoxic monomeric α -helical prion protein. *Proc Natl Acad Sci U S A*. 2012;109(8):3113–3118.
- Wu B, McDonald AJ, Markham K, et al. The N-terminus of the prion protein is a toxic effector regulated by the C-terminus. *eLife*. 2017;6:e23473.
- Diedrich JF, Bendheim PE, Kim YS, Carp RI, Haase AT. Scrapie-associated prion protein accumulates in astrocytes during scrapie infection. *Proc Natl Acad Sci U S A*. 1991;88(2):375–379.
- Moser M, Colello RJ, Pott U, Oesch B. Developmental expression of the prion protein gene in glial cells. *Neuron*. 1995;14(3):509–517.
- Raeber AJ, Race RE, Brandner S, et al. Astrocyte-specific expression of hamster prion protein (PrP) renders PrP knockout mice susceptible to hamster scrapie. *EMBO J*. 1997;16(20):6057–6065.
- Sarasa R, Martínez A, Monleón E, et al. Involvement of astrocytes in transmissible spongiform encephalopathies: A confocal microscopy study. *Cell Tissue Res*. 2012;350(1):127–134.
- Hernández RS, Sarasa R, Toledano A, Badiola JJ, Monzón M. Morphological approach to assess the involvement of astrocytes in prion propagation. *Cell Tissue Res*. 2014;358(1):57–63.
- Yamasaki T, Suzuki A, Hasebe R, Horiuchi M. Flow cytometric detection of PrP in neurons and glial cells from prion-infected mouse brains. *J Virol*. 2018;92(1):e01457-17.
- Ye X, Scallet AC, Kascsak RJ, Carp RI. Astrocytosis and amyloid deposition in scrapie-infected hamsters. *Brain Res*. 1998;809(2):277–287.

31. Cronier S, Laude H, Peyrin J-M. Prions can infect primary cultured neurons and astrocytes and promote neuronal cell death. *Proc Natl Acad Sci U S A*. 2004;101(33):12271–12276.
32. Choi YP, Head MW, Ironside JW, Priola SA. Uptake and degradation of protease-sensitive and -resistant forms of abnormal human prion protein aggregates by human astrocytes. *Am J Pathol*. 2014;184(12):3299–3307.
33. Hollister JR, Lee KS, Dorward DW, Baron GS. Efficient uptake and dissemination of scrapie prion protein by astrocytes and fibroblasts from adult hamster brain. *PLoS One*. 2015;10(1):e0115351.
34. Victoria GS, Arkhipenko A, Zhu S, Syan S, Zurzolo C. Astrocyte-to-neuron intercellular prion transfer is mediated by cell-cell contact. *Sci Rep*. 2016;6:2076.
35. Parchi P, Castellani R, Capellari S, et al. Molecular basis of phenotypic variability in sporadic Creutzfeldt-Jakob disease. *Ann Neurol*. 1996;39(6):767–778.
36. Krejciova Z, Alibhai J, Zhao C, et al. Human stem cell-derived astrocytes replicate human prions in a genotype-dependent manner. *J Exp Med*. 2017;214(12):3481–3495.
37. Watts JC, Prusiner SB. β -Amyloid Prions and the Pathobiology of Alzheimer's Disease. *Cold Spring Harb Perspect Med*. 2018;8(5):a023507.
38. Ding Y, Zhao J, Zhang X, et al. Amyloid beta oligomers target to extracellular and intracellular neuronal synaptic proteins in Alzheimer's disease. *Front Neurol*. 2019;10(1140):1140.
39. Olsson TT, Klementieva O, Gouras GK. Prion-like seeding and nucleation of intracellular amyloid- β . *Neurobiol Dis*. 2018;113:1–10.
40. Umeda T, Ramser EM, Yamashita M, et al. Intracellular amyloid β oligomers impair organelle transport and induce dendritic spine loss in primary neurons. *Acta Neuropathol Commun*. 2015;3:51.
41. Nath S, Agholme L, Kurudenkandy FR, Granseth B, Marcusson J, Hallbeck M. Spreading of neurodegenerative pathology via neuron-to-neuron transmission of β -amyloid. *J Neurosci*. 2012;32(26):8767–8777.
42. Matsunaga W, Shirokawa T, Isobe K. Specific uptake of Abeta1-40 in rat brain occurs in astrocyte, but not in microglia. *Neurosci Lett*. 2003;342(1-2):129–131.
43. Nagele RG, D'Andrea MR, Lee H, Venkataraman V, Wang H-Y. Astrocytes accumulate A beta 42 and give rise to astrocytic amyloid plaques in Alzheimer disease brains. *Brain Res*. 2003;971(2):197–209.
44. Wyss-Coray T, Loike JD, Brionne TC, et al. Adult mouse astrocytes degrade amyloid-beta in vitro and in situ. *Nat Med*. 2003;9(4):453–457.
45. Alarcón R, Fuenzalida C, Santibáñez M, von Bernhardi R. Expression of scavenger receptors in glial cells. Comparing the adhesion of astrocytes and microglia from neonatal rats to surface-bound beta-amyloid. *J Biol Chem*. 2005;280(34):30406–30415.
46. Shankar GM, Li S, Mehta TH, et al. Amyloid-beta protein dimers isolated directly from Alzheimer's brains impair synaptic plasticity and memory. *Nat Med*. 2008;14(8):837–842.
47. Cerf E, Gustot A, Goormaghtigh E, Ruyschaert J-M, Raussens V. High ability of apolipoprotein E4 to stabilize amyloid- β peptide oligomers, the pathological entities responsible for Alzheimer's disease. *FASEB J*. 2011;25(5):1585–1595.
48. Lesné SE. Breaking the code of amyloid- β oligomers. *Int J Cell Biol*. 2013;2013:950783.
49. Müller-Schiffmann A, Herring A, Abdel-Hafiz L, et al. Amyloid- β dimers in the absence of plaque pathology impair learning and synaptic plasticity. *Brain*. 2016;139(Pt 2):509–525.
50. Ono K. The oligomer hypothesis in α -synucleinopathy. *Neurochem Res*. 2017;42(12):3362–3371.
51. Rodríguez-Arellano JJ, Parpura V, Zorec R, Verkhratsky A. Astrocytes in physiological aging and Alzheimer's disease. *Neuroscience*. 2016;323:170–182.
52. White JA, Manelli AM, Holmberg KH, Van Eldik LJ, Ladu MJ. Differential effects of oligomeric and fibrillar amyloid-beta 1-42 on astrocyte-mediated inflammation. *Neurobiol Dis*. 2005;18(3):459–465.
53. Riddell DR, Zhou H, Atchison K, et al. Impact of apolipoprotein E (ApoE) polymorphism on brain ApoE levels. *J Neurosci*. 2008;28(45):11445–11453.
54. Manelli AM, Bulfinch LC, Sullivan PM, LaDu MJ. Abeta42 neurotoxicity in primary co-cultures: Effect of apoE isoform and Abeta conformation. *Neurobiol Aging*. 2007;28(8):1139–1147.
55. Perez JL, Carrero I, Gonzalo P, et al. Soluble oligomeric forms of beta-amyloid (Abeta) peptide stimulate Abeta production via astrogliosis in the rat brain. *Exp Neurol*. 2010;223(2):410–421.
56. Hou L, Liu Y, Wang X, et al. The effects of amyloid- β 42 oligomer on the proliferation and activation of astrocytes in vitro. *In Vitro Cell Dev Biol Anim*. 2011;47(8):573–580.
57. Wang W, Hou T-T, Jia L-F, Wu Q-Q, Quan M-N, Jia J-P. Toxic amyloid- β oligomers induced self-replication in astrocytes triggering neuronal injury. *EBioMedicine*. 2019;42:174–187.
58. Alberdi E, Wyssenbach A, Alberdi M, et al. Ca(2+)-dependent endoplasmic reticulum stress correlates with astrogliosis in oligomeric amyloid β -treated astrocytes and in a model of Alzheimer's disease. *Aging Cell*. 2013;12(2):292–302.
59. Allaman I, Gavillet M, Bélanger M, et al. Amyloid-beta aggregates cause alterations of astrocytic metabolic phenotype: Impact on neuronal viability. *J Neurosci*. 2010;30(9):3326–3338.
60. Angelova PR, Abramov AY. Interaction of neurons and astrocytes underlies the mechanism of A β -induced neurotoxicity. *Biochem Soc Trans*. 2014;42(5):1286–1290.
61. Muñoz Y, Paula-Lima AC, Núñez MT. Reactive oxygen species released from astrocytes treated with amyloid beta oligomers elicit neuronal calcium signals that decrease phospho-Ser727-STAT3 nuclear content. *Free Radic Biol Med*. 2018;117:132–144.
62. Shrivastava AN, Kowalewski JM, Renner M, et al. β -amyloid and ATP-induced diffusional trapping of astrocyte and neuronal metabotropic glutamate type-5 receptors. *Glia*. 2013;61(10):1673–1686.
63. Zhang H, Su Y-J, Zhou W-W, et al. Activated scavenger receptor A promotes glial internalization of a β . *PLoS One*. 2014;9(4):e94197.
64. Pitt J, Wilcox KC, Tortelli V, et al. Neuroprotective astrocyte-derived insulin/insulin-like growth factor 1 stimulates endocytic processing and extracellular release of neuron-bound A β oligomers. *Mol Biol Cell*. 2017;28(20):2623–2636.
65. Reid MJ, Beltran-Lobo P, Johnson L, Perez-Nievas BG, Noble W. Astrocytes in tauopathies. *Front Neurol*. 2020;11:572850.
66. Binder LI, Frankfurter A, Rebhun LI. The distribution of tau in the mammalian central nervous system. *J Cell Biol*. 1985;101(4):1371–1378.
67. Zhang Y, Chen K, Sloan SA, et al. An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J Neurosci*. 2014;34(36):11929–11947.
68. Clavaguera F, Duyckaerts C, Haik S. Prion-like properties of Tau assemblies. *Curr Opin Neurobiol*. 2020;61:49–57.
69. Garwood CJ, Ratcliffe LE, Simpson JE, Heath PR, Ince PG, Wharton SB. Review: Astrocytes in Alzheimer's disease and other age-associated dementias: A supporting player with a central role. *Neuropathol Appl Neurobiol*. 2017;43(4):281–298.
70. Nicoll JAR, Wilkinson D, Holmes C, Steart P, Markham H, Weller RO. Neuropathology of human Alzheimer disease after immunization with amyloid-beta peptide: A case report. *Nat Med*. 2003;9(4):448–452.

71. de Calignon A, de Calignon A, Polydoro M, et al. Propagation of tau pathology in a model of early Alzheimer's disease. *Neuron*. 2012;76(2):461.
72. Ferrer I, García MA, González IL, et al. Aging-related tau astroglial pathology (ARTAG): not only tau phosphorylation in astrocytes. *Brain Pathol*. 2018;28(6):965–985.
73. Narasimhan S, Guo JL, Changolkar L, et al. Pathological tau strains from human brains recapitulate the diversity of tauopathies in nontransgenic mouse brain. *J Neurosci*. 2017;37(47):11406–11423.
74. Narasimhan S, Changolkar L, Riddle DM, et al. Human tau pathology transmits glial tau aggregates in the absence of neuronal tau. *J Exp Med*. 2020;217(2):e20190783.
75. He Z, McBride JD, Xu H, et al. Transmission of tauopathy strains is independent of their isoform composition. *Nat Commun*. 2020;11(1):7.
76. Leyns CEG, Holtzman DM. Glial contributions to neurodegeneration in tauopathies. *Mol Neurodegener*. 2017;12(1):50.
77. Holmes BB, DeVos SL, Kfoury N, et al. Heparan sulfate proteoglycans mediate internalization and propagation of specific proteopathic seeds. *Proc Natl Acad Sci U S A*. 2013;110(33):E3138–E3147.
78. Martini-Stoica H, Cole AL, Swartzlander DB, et al. TFEB enhances astroglial uptake of extracellular tau species and reduces tau spreading. *J Exp Med*. 2018;215(9):2355–2377.
79. Rauch JN, Chen JJ, Sorum AW, et al. Tau Internalization is Regulated by 6-O Sulfation on Heparan Sulfate Proteoglycans (HSPGs). *Sci Rep*. 2018;8(1):6382.
80. Perea JR, López E, Díez-Ballesteros JC, Ávila J, Hernández F, Bolós M. Extracellular Monomeric Tau Is Internalized by Astrocytes. *Front Neurosci*. 2019;13:442.
81. Wang P, Ye Y. Filamentous recombinant human Tau activates primary astrocytes via an integrin receptor complex. *Nat Commun*. 2021;12(1):95.
82. Piacentini R, Li Puma DD, Mainardi M, et al. Reduced gliotransmitter release from astrocytes mediates tau-induced synaptic dysfunction in cultured hippocampal neurons. *Glia*. 2017;65(8):1302–1316.
83. Kovacs GG, Astroglia T. New perspectives. *Front Aging Neurosci*. 2020;12:96.
84. Schultz C, Ghebremedhin E, Del Tredici K, Rüb U, Braak H. High prevalence of thorn-shaped astrocytes in the aged human medial temporal lobe. *Neurobiol Aging*. 2004;25(3):397–405.
85. Lace G, Ince PG, Brayne C, et al. Mesial temporal astrocyte tau pathology in the MRC-CFAS ageing brain cohort. *Dement Geriatr Cogn Disord*. 2012;34(1):15–24.
86. Tu PH, Galvin JE, Baba M, et al. Glial cytoplasmic inclusions in white matter oligodendrocytes of multiple system atrophy brains contain insoluble alpha-synuclein. *Ann Neurol*. 1998;44(3):415–422.
87. Wakabayashi K, Hayashi S, Yoshimoto M, Kudo H, Takahashi H. NACP/ α -synuclein-positive filamentous inclusions in astrocytes and oligodendrocytes of Parkinson's disease brains. *Acta Neuropathol*. 2000;99(1):14–20.
88. Terada S, Ishizu H, Yokota O, et al. Glial involvement in diffuse Lewy body disease. *Acta Neuropathol*. 2003;105(2):163–169.
89. Wenning GK, Jellinger KA. The role of alpha-synuclein and tau in neurodegenerative movement disorders. *Curr Opin Neurol*. 2005;18(4):357–362.
90. Braak H, Sastre M, Del Tredici K. Development of alpha-synuclein immunoreactive astrocytes in the forebrain parallels stages of intraneuronal pathology in sporadic Parkinson's disease. *Acta Neuropathol*. 2007;114(3):231–241.
91. Nakamura K, Mori F, Kon T, et al. Accumulation of phosphorylated α -synuclein in subpial and periventricular astrocytes in multiple system atrophy of long duration. *Neuropathology*. 2016;36(2):157–167.
92. Emmanouilidou E, Melachroinou K, Roumeliotis T, et al. Cell-produced α -synuclein is secreted in a calcium-dependent manner by exosomes and impacts neuronal survival. *J Neurosci*. 2010;30(20):6838–6851.
93. El-Agnaf OMA, Salem SA, Paleologou KE, et al. Detection of oligomeric forms of alpha-synuclein protein in human plasma as a potential biomarker for Parkinson's disease. *FASEB J*. 2006;20(3):419–425.
94. Lee H-J, Kim C, Lee S-J. Alpha-synuclein stimulation of astrocytes: Potential role for neuroinflammation and neuroprotection. *Oxid Med Cell Longev*. 2010;3(4):283–287.
95. Fellner L, Irschick R, Schanda K, et al. Toll-like receptor 4 is required for α -synuclein dependent activation of microglia and astroglia. *Glia*. 2013;61(3):349–360.
96. Rannikko EH, Weber SS, Kahle PJ. Exogenous α -synuclein induces toll-like receptor 4 dependent inflammatory responses in astrocytes. *BMC Neurosci*. 2015;16:57.
97. Braidy N, Gai W-P, Xu YH, et al. Uptake and mitochondrial dysfunction of alpha-synuclein in human astrocytes, cortical neurons and fibroblasts. *Transl Neurodegener*. 2013;2(1):20.
98. Cavaliere F, Cerf L, Dehay B, et al. In vitro α -synuclein neurotoxicity and spreading among neurons and astrocytes using Lewy body extracts from Parkinson disease brains. *Neurobiol Dis*. 2017;103:101–112.
99. Loria F, Vargas JY, Bousset L, et al. α -Synuclein transfer between neurons and astrocytes indicates that astrocytes play a role in degradation rather than in spreading. *Acta Neuropathol*. 2017;134(5):789–808.
100. Krejciova Z, Carlson GA, Giles K, Prusiner SB. Replication of multiple system atrophy prions in primary astrocyte cultures from transgenic mice expressing human α -synuclein. *Acta Neuropathol Commun*. 2019;7(1):81.
101. Krencik R, Weick JP, Liu Y, Zhang Z-J, Zhang S-C. Specification of transplantable astroglial subtypes from human pluripotent stem cells. *Nat Biotechnol*. 2011;29(6):528–534.
102. Bradley RA, Shireman J, McFalls C, et al. Regionally specified human pluripotent stem cell-derived astrocytes exhibit different molecular signatures and functional properties. *Development*. 2019;146(13):dev170910.
103. Clarke BE, Taha DM, Ziff OJ, et al. Human stem cell-derived astrocytes exhibit region-specific heterogeneity in their secretory profiles. *Brain*. 2020;143(10):e85.
104. Masnata M, Cicchetti F. The evidence for the spread and seeding capacities of the mutant huntingtin protein in systems and their therapeutic implications. *Front Neurosci*. 2017;11:647.
105. Shin J-Y, Fang Z-H, Yu Z-X, Wang C-E, Li S-H, Li X-J. Expression of mutant huntingtin in glial cells contributes to neuronal excitotoxicity. *J Cell Biol*. 2005;171(6):1001–1012.
106. Bradford J, Shin J-Y, Roberts M, Wang C-E, Li X-J, Li S. Expression of mutant huntingtin in mouse brain astrocytes causes age-dependent neurological symptoms. *Proc Natl Acad Sci U S A*. 2009;106(52):22480–22485.
107. Bradford J, Shin J-Y, Roberts M, et al. Mutant huntingtin in glial cells exacerbates neurological symptoms of Huntington disease mice. *J Biol Chem*. 2010;285(14):10653–10661.
108. Wang L, Lin F, Wang J, et al. Expression of mutant N-terminal huntingtin fragment (htt552-100Q) in astrocytes suppresses the secretion of BDNF. *Brain Res*. 2012;1449:69–82.
109. Singhrao SK, Thomas P, Wood JD, et al. Huntingtin protein colocalizes with lesions of neurodegenerative diseases: An investigation in Huntington's, Alzheimer's, and Pick's diseases. *Exp Neurol*. 1998;150(2):213–222.

110. Shin JY, Fang ZH, Yu ZX, Wang CE, Li SH, Li XJ. Expression of mutant huntingtin in glial cells contributes to neuronal excitotoxicity. *J Cell Biol.* 2005;171(6):1001–1012.
111. Bradford J, Shin JY, Roberts M, Wang CE, Li XJ, Li S. Expression of mutant huntingtin in mouse brain astrocytes causes age-dependent neurological symptoms. *Proc Natl Acad Sci U S A.* 2009;106(52):22480–22485.
112. Faideau M, Kim J, Cormier K, et al. In vivo expression of polyglutamine-expanded huntingtin by mouse striatal astrocytes impairs glutamate transport: a correlation with Huntington's disease subjects. *Hum Mol Genet.* 2010;19(15):3053–3067.
113. Jansen AHP, van Hal M, den Kelder IC Op, et al. Frequency of nuclear mutant huntingtin inclusion formation in neurons and glia is cell-type-specific. *Glia.* 2017;65(1):50–61.
114. Cho IK, Yang B, Forest C, Qian L, Chan AWS. Amelioration of Huntington's disease phenotype in astrocytes derived from iPSC-derived neural progenitor cells of Huntington's disease monkeys. *PLoS One.* 2019;14(3):e0214156.
115. Pearce MMP, Spartz EJ, Hong W, Luo L, Kopito RR. Prion-like transmission of neuronal huntingtin aggregates to phagocytic glia in the *Drosophila* brain. *Nat Commun.* 2015;6(1):6768.
116. Meunier C, Merienne N, Jollé C, Déglon N, Pellerin L. Astrocytes are key but indirect contributors to the development of the symptomatology and pathophysiology of Huntington's disease. *Glia.* 2016;64(11):1841–1856.
117. Zhao T, Hong Y, Li S, Li X-J. Compartment-dependent degradation of mutant huntingtin accounts for its preferential accumulation in neuronal processes. *J Neurosci.* 2016;36(32):8317–8328.
118. Zhao T, Hong Y, Yin P, Li S, Li X-J. Differential HspBP1 expression accounts for the greater vulnerability of neurons than astrocytes to misfolded proteins. *Proc Natl Acad Sci U S A.* 2017;114(37):E7803–E7811.
119. Kabashi E, Valdmanis PN, Dion P, et al. TARDBP mutations in individuals with sporadic and familial amyotrophic lateral sclerosis. *Nat Genet.* 2008;40(5):572–574.
120. Rutherford NJ, Zhang Y-J, Baker M, et al. Novel mutations in TARDBP (TDP-43) in patients with familial amyotrophic lateral sclerosis. *PLoS Genet.* 2008;4(9):e1000193.
121. Sreedharan J, Blair IP, Tripathi VB, et al. TDP-43 mutations in familial and sporadic amyotrophic lateral sclerosis. *Science.* 2008;319(5870):1668–1672.
122. Neumann M, Kwong LK, Truax AC, et al. TDP-43-positive white matter pathology in frontotemporal lobar degeneration with ubiquitin-positive inclusions. *J Neuropathol Exp Neurol.* 2007;66(3):177–183.
123. Nishihira Y, Tan C-F, Onodera O, et al. Sporadic amyotrophic lateral sclerosis: Two pathological patterns shown by analysis of distribution of TDP-43-immunoreactive neuronal and glial cytoplasmic inclusions. *Acta Neuropathol.* 2008;116(2):169–182.
124. Zhang Y-J, Xu Y-F, Cook C, et al. Aberrant cleavage of TDP-43 enhances aggregation and cellular toxicity. *Proc Natl Acad Sci U S A.* 2009;106(18):7607–7612.
125. Walker AK, Daniels CML, Goldman JE, Trojanowski JQ, Lee VM-Y, Messing A. Astrocytic TDP-43 pathology in Alexander disease. *J Neurosci.* 2014;34(19):6448–6458.
126. Uryu K, Nakashima-Yasuda H, Forman MS, et al. Concomitant TAR-DNA-binding protein 43 pathology is present in Alzheimer disease and corticobasal degeneration but not in other tauopathies. *J Neuropathol Exp Neurol.* 2008;67(6):555–564.
127. Sakurai A, Makioka K, Fukuda T, Takatama M, Okamoto K. Accumulation of phosphorylated TDP-43 in the CNS of a patient with Cockayne syndrome. *Neuropathology.* 2013;33(6):673–677.
128. Lee EB, Lee VM-Y, Trojanowski JQ, Neumann M. TDP-43 immunoreactivity in anoxic, ischemic and neoplastic lesions of the central nervous system. *Acta Neuropathol.* 2008;115(3):305–311.
129. Serio A, Bilican B, Barmada SJ, et al. Astrocyte pathology and the absence of non-cell autonomy in an induced pluripotent stem cell model of TDP-43 proteinopathy. *Proc Natl Acad Sci U S A.* 2013;110(12):4697–4702.
130. Nonaka T, Hasegawa M. Prion-like properties of assembled TDP-43. *Curr Opin Neurobiol.* 2020;61:23–28.
131. Ishii T, Kawakami E, Endo K, Misawa H, Watabe K. Formation and spreading of TDP-43 aggregates in cultured neuronal and glial cells demonstrated by time-lapse imaging. *PLoS ONE.* 2017;12(6):e0179375.
132. Smethurst P, Risse E, Tyzack GE, et al. Distinct responses of neurons and astrocytes to TDP-43 proteinopathy in amyotrophic lateral sclerosis. *Brain.* 2020;143(2):430–440.