



Acute and Subacute Toxicity Assessment of Oxyclozanide in Wistar Rats

Weiwei Wang^{1,2,3}, Zhen Dong^{1,2,3}, Jili Zhang^{1,2,3}, Xuzheng Zhou^{1,2,3}, Xiaojuan Wei^{1,2,3}, Fusheng Cheng^{1,2,3}, Bing Li^{1,2,3} and Jiyu Zhang^{1,2,3*}

¹ Key Laboratory of New Animal Drug Project of Gansu Province, Lanzhou, China, ² Key Laboratory of Veterinary Pharmaceutical Development, Ministry of Agriculture, Lanzhou, China, ³ Lanzhou Institute of Husbandry and Pharmaceutical Sciences, Chinese Academy of Agricultural Sciences, Lanzhou, China

OPEN ACCESS

Edited by:

Abdurrahman Aksoy,
Ondokuz Mayıs University, Turkey

Reviewed by:

Ayhan Filazi,
Ankara University, Turkey
Cengiz Gokbulut,
Balıkesir University, Turkey

*Correspondence:

Jiyu Zhang
infzy@sina.com

Specialty section:

This article was submitted to
Veterinary Pharmacology and
Toxicology,
a section of the journal
Frontiers in Veterinary Science

Received: 02 July 2019

Accepted: 16 August 2019

Published: 06 September 2019

Citation:

Wang W, Dong Z, Zhang J, Zhou X,
Wei X, Cheng F, Li B and Zhang J
(2019) Acute and Subacute Toxicity
Assessment of Oxyclozanide in Wistar
Rats. *Front. Vet. Sci.* 6:294.
doi: 10.3389/fvets.2019.00294

Oxyclozanide is an effective anthelmintic and has shown good properties in other ways including anti-adenovirus, anti-biofilm, antifungal, and antibacterial activity. This study aimed to investigate the acute and subacute 28-days repeated dose oral toxicity of an oxyclozanide suspension in Wistar rats. A high oral lethal dose (LD₅₀) of 3,707 mg/kg was observed in the acute toxicity test. During the 28-days time period, no obvious adverse effects or death were detected. Histopathological changes were observed in the heart, liver, and kidney of animals treated with high dose of oxyclozanide. Based on the hematological parameters, there were no statistical differences between the oxyclozanide-treated group and the control group. For biochemistry assays, ALP, AST, GLU, TBIL, GLO, TG, BUN, UA, LDH, and CK were statistically changed in the treatment groups. These data suggested that the LD₅₀ of oxyclozanide was ~3,707 mg/kg body weight (BW), and the lowest observed adverse effect level (LOAEL) of oxyclozanide was at a dose of 74 mg/kg in rats.

Keywords: oxyclozanide, acute, subacute, toxicity, rats

INTRODUCTION

Infectious parasitic diseases have become a serious threat to animal health and productivity in developing countries (1–3). Among them, fascioliasis, as a zoonotic disease, is a food-borne parasitic disease, which is transmitted to certain hosts, humans, and other herbivorous mammals through contaminated water or green vegetables (4). It is also considered to be one of the most serious neglected tropical diseases. The pathogens of fascioliasis are *Fasciola hepatica* as well as *Fasciola gigantica*. *F. hepatica* is an important tissue parasite, which can cause serious pathological damage to the host and have a great impact on animal husbandry. It is estimated that the annual loss of global animal production caused by *F. hepatica* infection is more than \$3.2 billion. In addition, *F. hepatica* is considered to be a new infectious disease in human beings. The World Health Organization estimates that 180 million people are at risk of infection and 2.4 million are infected with fascioliasis (5).

Since no vaccine has been successfully developed, the effective strategy for controlling fascioliasis depends mainly on the use of pharmaceuticals (fasciolicide). Oxyclozanide was developed in 1966 (6) and has achieved an effective effect in subsequent clinical applications (7, 8). In clinical trials, oxyclozanide has been found to be effective not only for the treatment of *F. hepatica* infection but also for other tissue parasites, such as *Hymenolepis* sp. (9) and *Paramphistomum* sp. (10).

Compared with other fasciolicides, such as closantel, albendazole, and clorsulon, oxytetracycline is the most effective drug in ruminants naturally infected with *F. hepatica* (11–13). According to Rana's recent review, oxytetracycline was effective on mature fluke, especially for adults over 14 weeks old (14). Additionally, a resistance of triclabendazole and albendazole in sheep, cattle, and water buffalos was reported widely (15–19). Fortunately, oxytetracycline resistance has still not been reported in animals and has a very good killing effect on *F. hepatica* resistant to triclabendazole (20). Oxytetracycline has also good pharmacokinetic characteristics, including rapid absorption and long elimination of half-life (10, 21, 22).

In recent years, oxytetracycline has still been explored and researched. The researchers found that oxytetracycline showed anti-adenovirus activity and almost no cytotoxicity at low concentration and proved its mechanism as a specific target to inhibit the transcription of early adenovirus gene E1A (23). Maiden and co-workers defined oxytetracycline as an anti-biofilm agent that enhances the activity of aminoglycosides and tetracycline against *Pseudomonas aeruginosa* biofilm by reducing membrane potential, penetrating cells, and enhancing accumulation of tobramycin in biofilm (24). Oxytetracycline also showed excellent antifungal activity and antibacterial activity at low concentrations (25, 26). It reported that the uncoupling of mitochondria could reduce the formation of intestinal polyps and the metastasis of colon cancer cells (27).

Those studies have attracted much more attention for the application of oxytetracycline. Thus, the toxic evaluation of oxytetracycline for clinic use seems to be important and need to be further studied. Although the study of oxytetracycline has already begun broadly in the 1960s, the toxicity assessment is still less known. The acute and 28-day repeated oral toxicity of oxytetracycline were carried out to supply the toxicity of oxytetracycline and to provide guidance for phase II clinical trials. In this study, we evaluated the acute and subacute toxicity of oxytetracycline in rats for oral administration.

MATERIALS AND METHODS

Ethics Statement

All animal studies were performed according to the US National Institutes of Health Guidelines for the Care and Use of Laboratory Animals and approved by the Institutional Animal Care and Use Committee of Lanzhou Institute of Husbandry and Pharmaceutical Science of Chinese Academy of Agricultural Sciences. The animals were examined and adapted to the new environmental conditions for a week before the formal experiment.

Oxytetracycline Suspension

3,3',5,5',6-Pentachloro-2,2'-dihydroxybenzamide (oxytetracycline) is a white crystalline solid, non-hygroscopic, and virtually insoluble in water (7). Oxytetracycline was purchased from Jiangsu Quality Horizons Pharmtech Incorporation. The purity of the drug was 98.76%, which was assessed by high-performance liquid chromatography (HPLC, Agilent, USA) in our laboratory. For the physical properties of the drug, it was mixed with 0.2%

sodium carboxymethyl cellulose solution. The concentrations of oxytetracycline suspension were 250.00, 184.50, 136.50, 101.25, and 75.00 mg/mL. For the acute study, 20 rats were divided into five groups (two males, two females) in pre-experiments. The high-dose group was 5,000 mg/kg and the low-dose group was 1,000 mg/kg. After dosing, according to Veterinary Drug Research (28), rats were observed and recorded for 7 days. Animal death was recorded. In the formula, "a" referred to LD₀, "b" referred to LD₁₀₀. N (group number), and r (ratio of dosage from each group) were calculated according to Koch's Method ($N = b/a$; $r = \lg - 1[(\lg b - \lg a)/(N - 1)]$). Based on the results of pre-experiments, 60 rats were divided into six groups. The treated group received oxytetracycline at the dosage of 5,000, 3,690, 2,730, 2,025, and 1,500 mg/kg BW/day. For the subacute study, according to Veterinary Drug Research (28), one-tenth to one-fourth of LD₅₀ is the high dose daily administration for 28 days. In the 28-day test, one-tenth is chosen. In the medium-dose group, one-half of the high dose is medium dose. In the low-dose group, one half of medium dose is low dose. The concentrations of oxytetracycline suspension were 55.0, 27.8, and 11.1 mg/mL.

Animals

Healthy male and female Wistar rats (100–150 g) were purchased from the Laboratory Animal Center of Lanzhou Veterinary Research Institute [Certification No. SCXK (G) 2015-001]. The toxicity test was performed strictly with the method of Compilation of Technical guidelines for Veterinary Drug Research (28) and was also referred to Organization for Economic Cooperation and Development (OECD) Guideline No. 407 (29). All animals were housed in plastic cages and kept separately according to sex (temperature $22 \pm 2^\circ\text{C}$, relative humidity $55 \pm 10\%$, and 12-h light/dark cycle). All rats were allowed to access water and food freely.

Acute Toxicity Experiment

Sixty rats (8 weeks old) were used in this experiment and were randomly assigned into six groups (five males and five females). The control group received 0.2% carboxymethyl cellulose sodium solution in the same volume, while the treated group received oxytetracycline at the dosage of 5,000, 3,690, 2,730, 2,025, and 1,500 mg/kg BW/day. After 7-day acclimatization, rats were treated once. The general clinic observations will start after rats were given drugs 4 h later and then this record will last for at least 7 days every 8 h.

Subacute Toxicity Experiment

Ninety-six rats (6 weeks old) were randomly assigned into four groups (12 male and 12 female rats in each group). The obtained median lethal dose (LD₅₀) of oxytetracycline was 3,707 mg/kg BW/day in rats. The four groups were divided into: high dose (370 mg/kg BW/day), medium dose (185 mg/kg BW/day), low dose (74 mg/kg BW/day), and control (0.2% carboxymethyl cellulose sodium solution in same volume). Rats were administered once daily by gavage with oxytetracycline suspension at three doses as above (370, 185, and 74 mg/kg BW/day) at 2:00–3:00 p.m. throughout the experiment for 28 continuous days. Hold the animal with the left hand and the

syringe with the right hand. Keep the head and neck of the rat in a straight line, so that the needle can enter the mouth easily. In Baoding rats, the head must be fixed well to avoid the head twisting at will and affect the operation of gavage. The intragastric needle enters from the angulus oris of the animal to make the mouth and esophagus align. When the intragastric needle enters about 5 cm deep, gently push the syringe to make the drug enter the rat body slowly. If there is no excessive struggle, we can try to inject drugs slowly. If the resistance is small, we can inject all drugs. In the case of rat struggle, the intragastric needle should be pulled out, reserved and operated again. The animals' general clinic observations, body weight, morbidity, and mortality were recorded daily during administration. Additionally, measurements of food consumption were recorded weekly.

Hematological Index

Blood samples were collected and placed into tubes containing EDTA-K₂ for the hematological analyses. At the end of the drug administration period, hematological index of rats was analyzed. Standard hematological and biochemistry tests were used to determine the hematological parameters, enzymes, substrates, and products of metabolism. The following indicators were red blood cell count (RBC), white blood cell count (WBC), lymphocyte count (LYM), monocyte count (MON), granulocyte count (GRAN), hemoglobin (HGB), hematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular hemoglobin (MCH), mean corpuscular hemoglobin concentration (MCHC), red blood cell distribution width (RDW), platelet count (PLT), platelet distribution width (PDW), mean platelet volume (MPV), and thrombocytocrit (PCT).

Blood Biochemical Index

Blood samples were collected, centrifuged at 4,000 rpm for 10 min at 4°C and stored at -20°C. At the end of the drug administration period, blood biochemical index of rats in each group was analyzed. The main biochemical indicators were alanine aminotransferase (ALT), aspartate aminotransferase (AST), alkaline phosphatase (ALP), urea nitrogen (BUN), total bilirubin (TBIL), direct bilirubin (DBIL), creatinine (Cr), total protein (TP), blood sugar (Glu), serum albumin (Alb), total protein (TP), total cholesterol (TCH) and triglyceride (TG), globulin (GLO), total bile acid (TBA), glucose (GLU), TG, total cholesterol (TC), creatine kinase (CK), lactate dehydrogenase (LDH), and uric acid (UA).

Necropsy

After 28-day drug administration, treated and untreated rats were euthanized and were then necropsied. Necropsies were conducted carefully on all the rats, which either died or survived during experiments. Tissues like heart, liver, spleen, lungs, kidney, brain, reproductive organs (tests, ovary and uterus), prostate, gland, adrenal gland, white adipose tissue, thyroid, thymus, stomach, urinary bladder, intestine, pancreas, bones, and peripheral nerves were examined for the macroscopic morphology and then removed quickly, washed twice with ice-cold saline, weighed, and kept in AAF fixation solution (Solarbio,

Beijing, China) for at least 48 h. In addition, the relative weight of each organ (viscera/body ratio) was calculated based on the animal's body weights according to the following formula: organ weight/body weight on sacrifice day × 100, as described previously (30).

Histopathological Examination

Tissues were dehydrated in ethyl alcohol and treated with ethyl salicylate. After the tissue was embedded in paraffin and sectioned at the 5-μm thickness, sections were dyed with hematoxylin and eosin (H&E) according to the universally accepted procedure for light microscopy.

Statistical Analysis

In the toxicity study, data were expressed as mean value and standard deviations (mean ± SD). Statistical analyses containing body weight, organ weight, blood biochemical, and biochemistry data were performed by one-way analysis of variance (ANOVA) and Levene's test for homogeneity of variance with SPSS Statistics 19.0 (IBM, Chicago, USA), and the graph was drawn by using GraphPad Prism 5.0 (GraphPad Prism, San Diego, USA). A *p* < 0.05 was considered statistically significant.

RESULTS

Acute Toxicity

In the acute test, the toxic symptoms that were observed in rats included depression, rough hair, and inappetence in Groups 1, 2, 3, and 4. The lips, limbs, and tails of the dead rats were cyanotic. In addition, individual rats have swelling spleen, bleeding spots in the liver, and swelling stomachs in the high-dose group. Most of the animals died after administration during the first 12 h. The number of deaths during acute toxicity is shown in **Table 1**. Particularly, the death was mainly half male and half female, which illustrated that death has no relation to sex. Therefore, LD₅₀ of oxyclozanide was 3,707 mg/kg BW/day for experimental rats in both sexes (95 confidence interval, 3,148–4,365 mg/kg BW/day), which was calculated by the Koch's method (31) (**Table 1**).


Subacute Toxicity

Morphological Alteration

No treatment-related deaths occurred during the 28-day test. Clinic general observations showed that there was no apparent alteration in general signs like posture, gait, and response to handling, but slight depression and diarrhea were detected

TABLE 1 | The death of rats during the acute toxicity study.

Group	Number dosed	Dosage (mg/kg)	Mortality rates (%)	Deaths (n)	Survivals (n)
1	10	5,000	100%	10	0
2	10	3,690	40%	4	6
3	10	2,730	20%	2	8
4	10	2,025	10%	1	9
5	10	1,500	0%	0	10



in the high-dose group during administration. Furthermore, oxyclozanide hasn't changed the skin, fur colors, eyes, mucous membrane, the occurrence of secretions and excretions, motor activities, and autonomic activity.

Assessment of Food Consumption and Body Weight

Food consumption was recorded but no statistical alternation in food consumption (data not shown), illustrating that the oxyclozanide has no obvious effect on food consumption.

The mean values of the body weight gain in the high-dose group and the medium-dose group were lower than those in the control group ($p < 0.05$ for male rats in the medium-dose group and female rats in the high-dose group vs. their representative controls). However, no statistical changes ($p > 0.05$) were observed between the low-dose group and the control group (Figure 1).

Hematology Index

All the relative blood parameters, including Lym, Mon, WBC, HBC, and PCT, were not statistically changed between the oxyclozanide-treated group and the control group, except MCH levels in both male and female rats, and MCHC, RDW, and PCT levels in female rats vs. their respective controls (Table 2). In particular, the statistical findings occurred in the low-dose group. These statistical changes were still in the range of historical controls. Deep reasons of differences need to be further studied.

Biochemical Analyses

Compared with the control group, biochemical analyses showed some statistical differences ($p < 0.05$) between oxyclozanide treatment and control group (Table 3). For instance, the levels of ALB, UA, LDH, and CK were statistically increased and GLU were decreased ($p < 0.01$ and $p < 0.05$, respectively) in the medium- and/or high-dose groups as compared with their respective controls only in male rats. In contrast, TBIL, GLO, and TG levels were statistically decreased ($p < 0.01$) in the medium- and/or high-dose groups as compared with their respective controls only in female rats. The levels of AST, ALP, and BUN were statistically elevated in the medium- and/or high-dose group vs. their respective

controls in both male and female rats ($p < 0.01$ and $p < 0.05$, respectively).

Effect of Oxyclozanide on Relative Organ Weight

During the 28-day experimental period, the mean of relative organ/body ratio (organ weight/body weight \times 100%) was statistically decreased in the lung in male rats and increased in the liver in female animals after using high dose of oxyclozanide ($p < 0.05$; Table 3). No statistical ($p > 0.05$) changes in other organs were observed between oxyclozanide treatment and control group (Table 4).

Macroscopic Gross Necropsy

Macroscopic observations showed that no lesions or abnormalities in the architecture of various organs except swelling of kidney occurred more than half in the high-dose group in both sexes. The observations indicated that there was no inflammation, hypertrophy, edema, and atrophy in organ macroscopic examination.

Histopathological Analyses

The histopathological changes could be observed in the heart, liver, kidney, and duodenum of rats treated with the high-dose drug (Figures 2A–F) and changes are summarized in Table 5. For instance, myocardial fiber degeneration, myocardial necrosis, and interstitial hydrops could be detected in cardiac tissues having received high or medium dose of oxyclozanide. Glomerulonephritis and granular degeneration of tubular epithelium were shown in renal histology in the high-dose and reversible high-dose groups. Additionally, steatosis and granular degeneration of hepatocytes were observed in the liver in the high-dose group. Compared with the control group, there was any alteration in the brain, lung, and spleen in the treatment groups (Figures 2C–H). Moreover, the macroscopic structure of testis, ovary, and uterus was normal in the oxyclozanide treatment groups as compared with the control group (Figures 2G–J).

TABLE 2 | Hematological values of rats in the oxyclozanide-treated group and the control group.

	Historical controls range	0 mg/kg (control)	Oxyclozanide doses (mg/kg)		
			74 mg/kg (low dose)	185 mg/kg (medium dose)	370 mg/kg (high dose)
MALE					
WBC	4–6	5.28 ± 2.12	4.00 ± 1.73	5.47 ± 2.44	5.13 ± 1.22
LYM	2–5	2.87 ± 1.70	3.50 ± 1.22	4.10 ± 2.01	3.88 ± 0.96
Mon	0.05–0.10	0.05 ± 0.071	0.06 ± 0.074	0.08 ± 0.071	0.08 ± 0.044
Gran	0.5–1.0	0.54 ± 0.46	0.63 ± 0.43	0.80 ± 0.48	0.98 ± 0.48
RBC	6–8	7.04 ± 0.38	7.29 ± 0.51	7.03 ± 0.48	6.93 ± 0.45
HGB	120–140	137.55 ± 5.47	128.60 ± 22.83	135.38 ± 10.21	136.17 ± 10.53
HCT	35–45	41.57 ± 1.50	43.14 ± 3.10	41.26 ± 3.02	41.18 ± 3.11
MCV	55–65	59.16 ± 1.91	59.24 ± 0.97	58.75 ± 1.88	59.47 ± 2.32
MCH	17–21	19.51 ± 0.48	17.37 ± 3.01**	19.18 ± 0.46	19.58 ± 0.67
MCHC	270–340	330.36 ± 4.39	293.67 ± 50.54	327.50 ± 7.44	330.17 ± 7.98
RDW	10–15	11.60 ± 2.21	10.84 ± 1.69	12.75 ± 1.84	12.60 ± 1.81
PLT	800–1,000	860.18 ± 153.11	939.44 ± 154.04	956.38 ± 68.66	759.40 ± 144.87
MPV	5–7	6.38 ± 0.25	6.28 ± 0.31	6.09 ± 0.50	5.95 ± 0.66
PDW	15–17	16.19 ± 0.18	16.29 ± 0.32	16.13 ± 0.25	16.22 ± 0.43
PCT	0.4–0.8	0.53 ± 0.079	0.56 ± 0.073	0.58 ± 0.052	0.58 ± 0.090
FEMALE					
WBC	4–6	4.14 ± 2.16	5.76 ± 1.63	4.66 ± 1.98	5.01 ± 1.36
LYM	3–5	3.46 ± 1.55	4.64 ± 0.612	4.25 ± 1.09	4.11 ± 1.14
Mon	0.05–0.10	0.09 ± 0.074	0.10 ± 0.076	0.08 ± 0.067	0.06 ± 0.052
Gran	0.6–1.1	0.86 ± 0.45	0.90 ± 0.41	1.00 ± 0.42	0.92 ± 0.20
RBC	6–8	7.45 ± 0.21	7.21 ± 1.31	7.26 ± 1.47	7.23 ± 0.35
HGB	120–140	148.45 ± 7.85	126.38 ± 28.23	136.80 ± 38.84	143.88 ± 6.94
HCT	35–45	44.50 ± 1.22	42.80 ± 7.84	42.99 ± 8.63	43.70 ± 2.07
MCV	60 ± 6	59.86 ± 1.73	53.05 ± 19.12	59.30 ± 1.00	60.55 ± 2.37
MCH	17–21	19.89 ± 0.81	17.53 ± 2.35*	18.44 ± 3.49	19.88 ± 0.69
MCHC	270–340	333.00 ± 15.50	296.13 ± 41.19*	311.78 ± 58.95	328.88 ± 4.12
RDW	10–15	14.92 ± 0.45	13.80 ± 1.16**	14.78 ± 0.55	14.41 ± 0.77
PLT	900–1,100	1015.82 ± 91.63	970.38 ± 215.43	921.38 ± 192.32	953.33 ± 155.58
MPV	5–7	6.25 ± 0.22	6.33 ± 0.26	6.46 ± 0.69	6.18 ± 0.37
PDW	15–17	16.07 ± 0.13	16.13 ± 0.18	16.50 ± 1.15	16.10 ± 0.41
PCT	0.4–0.8	0.63 ± 0.056	0.55 ± 0.132*	0.49 ± 0.174	0.50 ± 0.197

p* < 0.05 vs. control group, *p* < 0.01 vs. control group.

DISCUSSION

An acute oral LD₅₀ of oxyclozanide in rats was determined to be 3,707 mg/kg. Thus, the design of our current sub-chronic oral toxicity study was based on the LD₅₀ dose, and three different doses (370, 175, and 74 mg/kg) representing high, medium, and low dose, respectively, were examined in order to determine a no observed adverse effect level in rats. To our best knowledge, to date, there is no systemic research about subacute toxicity of oxyclozanide to support clinic drug safety. Therefore, in the current study, we performed a series of experiments to study the toxicity and safety of oxyclozanide in Wistar rats.

During the 28-day oral administration of oxyclozanide, no deaths in rats were observed. The results of clinic symptoms

showed no obvious changes in general behavior, mucosa color, and posture. However, minor depression and diarrhea were seen in the high and medium dose of oxyclozanide groups only at the first several days of drug administration. This result suggested that the impact of oxyclozanide on depression and diarrhea was short, not permanent. In Walley's research, the toxic signs of oxyclozanide in sheep and cattle are similar to that in rats. Side effects, including diarrhea, loss of appetite, depression, and weight loss, can occur when the dose is as high as 30 mg/kg or higher in sheep and cattle. Weight loss was dramatically reduced and death occurs at a dosage of 60 mg/kg in those animals (7). Additionally, duodenal inflammation was found in animals, which was consistent with our results, indicating that diarrhea symptoms could disturb intestinal function, which is consistent with EMEA's (European

TABLE 3 | Liver function related to biochemical profiles.

	Historical controls range	0 mg/kg (control)	Oxyclozanide doses		
			74 mg/kg (low dose)	185 mg/kg (medium dose)	370 mg/kg (high dose)
MALE					
TBIL	0.3–0.7	0.52 ± 0.23	0.54 ± 0.29	0.39 ± 0.19	0.35 ± 0.17
DBIL	0.8–1.4	1.20 ± 0.55	0.97 ± 0.32	1.01 ± 0.22	1.32 ± 0.51
TP	30–50	41.94 ± 15.15	36.17 ± 11.22	41.83 ± 9.38	49.97 ± 8.13
ALB	18–26	19.45 ± 6.61	18.30 ± 5.53	19.63 ± 4.04	24.36 ± 3.48*
GLO	16–26	22.45 ± 8.63	17.87 ± 5.73	22.20 ± 5.57	25.62 ± 4.90
ALT	25–35	34.88 ± 12.47	27.35 ± 9.97	34.62 ± 9.68	32.45 ± 5.23
AST	95–150	98.43 ± 33.86	132.14 ± 35.72	158.11 ± 48.63**	199.45 ± 32.55**
ALP	120–200	131.71 ± 25.22	135.00 ± 26.93	197.00 ± 46.29*	224.00 ± 81.28**
TG	0.5–0.7	0.62 ± 0.18	0.63 ± 0.17	0.68 ± 0.33	0.56 ± 0.16
TC	1.0–2.0	1.52 ± 0.47	1.34 ± 0.45	1.70 ± 0.36	1.70 ± 0.36
GLU	4–8	6.64 ± 1.50	6.00 ± 1.15	5.49 ± 1.45	4.95 ± 1.09**
BUN	5–9	5.49 ± 1.34	7.01 ± 1.98	7.17 ± 1.33	8.76 ± 2.99**
CR	20–30	24.89 ± 17.67	21.05 ± 7.06	25.22 ± 5.19	29.46 ± 8.73
UA	0.04–0.06	0.046 ± 0.012	0.045 ± 0.017	0.065 ± 0.021	0.080 ± 0.048*
LDH	1,200–2,000	1320.30 ± 469.50	1582.11 ± 303.31	1864.00 ± 433.37**	1954.77 ± 236.24**
CK	1,100–1,700	1188.72 ± 488.39	1210.48 ± 279.84	1499.38 ± 405.02	1628.94 ± 312.20**
FEMALE					
TBIL	0.4–0.8	0.69 ± 0.27	0.45 ± 0.26*	0.49 ± 0.14	0.27 ± 0.12**
DBIL	1–2	1.32 ± 0.42	1.27 ± 0.47	1.46 ± 0.49	1.48 ± 0.43
TP	40–48	45.85 ± 9.98	42.24 ± 10.40	45.40 ± 13.24	40.32 ± 4.21
ALB	20–26	23.33 ± 4.65	21.60 ± 5.10	22.49 ± 6.09	25.87 ± 3.98
GLO	20–30	22.52 ± 5.34	20.65 ± 5.36	22.91 ± 7.30	28.97 ± 6.49**
ALT	25–35	26.11 ± 7.17	25.64 ± 10.56	30.68 ± 11.99	33.92 ± 6.11
AST	130–180	137.10 ± 31.18	155.70 ± 39.79	177.71 ± 49.51*	206.60 ± 25.16**
ALP	90–140	97.82 ± 31.68	122.00 ± 30.99	146.17 ± 23.81**	190.67 ± 21.56**
TG	0.5–1.0	0.76 ± 0.25	0.56 ± 0.14*	0.54 ± 0.17*	0.45 ± 0.06**
TC	1.0–2.0	1.54 ± 0.29	1.59 ± 0.35	1.64 ± 0.43	1.73 ± 0.22
GLU	4–8	5.73 ± 1.33	5.59 ± 0.68	5.86 ± 2.01	4.62 ± 0.61
BUN	5–10	5.75 ± 2.14	8.30 ± 1.94**	7.95 ± 1.58*	9.63 ± 1.59**
CR	25–30	27.02 ± 6.23	25.96 ± 6.94	26.28 ± 6.42	28.80 ± 5.45
UA	0.05–0.07	0.061 ± 0.016	0.056 ± 0.011	0.071 ± 0.034	0.063 ± 0.020
LDH	1,200–1,800	1605.75 ± 278.48	1802.50 ± 342.99	1840.88 ± 343.80	1895.00 ± 195.69
CK	1,100–1,700	1306.00 ± 309.40	1547.52 ± 369.63	1595.46 ± 323.62	1503.43 ± 192.21

***p* < 0.01 vs. control group, **p* < 0.05 vs. control group.

Medicines Agency) report (21). In our study, myocardial fibers of rats were damaged, suggesting that oxyclozanide toxicity may cause some damage to the heart. Similarly, blood spots were found in the endocardium of cattle and sheep (7).

The biochemical analyses showed that AST was different between oxyclozanide-treated and untreated control groups. It is known that AST and ALT are released into the serum after relative cells and tissues were damaged (32). In most toxicological studies, the increase of ALT is generally considered to be a practical and specific enzyme indicator of hepatocyte injury. The

amount of serum AST may increase during hepatocyte injury. This increase is non-specific because of the high activity of AST in muscle. In liver injury, the AST is often parallel to serum ALT activity. In general, the elevation in AST concentration is related to the number of affected hepatocytes and does not reflect the severity or reversibility of the lesion on a pathological basis. The ALT representing a sensitive indicator in liver damage was not statistically altered during the 28-day study. Thus, the observed AST changes in this study could be associated with some of the other histological changes observed or may be contributed to by multiple tissues. Additionally, oxyclozanide

TABLE 4 | Effect of oxyclozanide on the body and relative organ weight.

	Historical controls range	Oxyclozanide doses			
		0 mg/kg (control)	74 mg/kg (low dose)	185 mg/kg (medium dose)	370 mg/kg (high dose)
MALE					
Initial BW		136.91 ± 8.76	141.08 ± 11.28	147.93 ± 10.05	149.83 ± 7.06
28 days BW		263.01 ± 13.57	259.50 ± 23.02	237.61 ± 19.23	269.23 ± 12.24
Heart	0.3–0.5	0.40 ± 0.055	0.35 ± 0.068	0.35 ± 0.068	0.33 ± 0.061
Liver	3–4	3.60 ± 0.420	3.51 ± 0.612	3.39 ± 0.423	3.46 ± 0.592
Spleen	0.2–0.4	0.26 ± 0.045	0.26 ± 0.063	0.25 ± 0.053	0.24 ± 0.042
Lung	0.5–0.8	0.61 ± 0.097	0.61 ± 0.133	0.65 ± 0.189	0.46 ± 0.128*
Kidney	0.3–0.5	0.38 ± 0.055	0.39 ± 0.095	0.39 ± 0.050	0.39 ± 0.068
Adrenal gland	0.02–0.04	0.02 ± 0.007	0.03 ± 0.012	0.02 ± 0.004	0.02 ± 0.004
Brain	0.6–0.8	0.68 ± 0.109	0.70 ± 0.100	0.71 ± 0.102	0.70 ± 0.123
FEMALE					
Initial BW		124.53 ± 9.66	129.67 ± 9.33	127.09 ± 7.46	129.31 ± 10.42
28 days BW		188.79 ± 14.95	191.37 ± 13.95	182.33 ± 10.15	181.31 ± 7.43
Heart	0.3–0.5	0.42 ± 0.068	0.41 ± 0.080	0.36 ± 0.076	0.35 ± 0.071
Liver	3–4	3.30 ± 0.601	3.51 ± 0.616	3.49 ± 0.586	4.00 ± 1.057*
Spleen	0.2–0.4	0.29 ± 0.079	0.28 ± 0.047	0.27 ± 0.044	0.32 ± 0.053
Lung	0.5–0.8	0.70 ± 0.183	0.80 ± 0.186	0.70 ± 0.138	0.74 ± 0.129
Kidney	0.3–0.5	0.42 ± 0.089	0.43 ± 0.077	0.41 ± 0.074	0.45 ± 0.087
Adrenal gland	0.02–0.04	0.03 ± 0.009	0.04 ± 0.009	0.03 ± 0.009	0.03 ± 0.007
Brain	0.6–1.0	0.94 ± 0.147	0.89 ± 0.115	0.94 ± 0.132	0.95 ± 0.163

BW, body weight. * $p < 0.01$ vs. control group.

was reported to cause ALP increase and toxic effects on the liver in experimental rats (7). The increase of ALP activity is an indication of cholestasis, which may be secondary to intrahepatic bile duct obstruction or extrahepatic obstruction caused by hepatocyte swelling. Drug-induced hepatocytes necrosis can also increase the activity of ALP (33). In this test, the level of ALP in the high- and medium-dose groups was statistically elevated ($p < 0.01$). These results indicate that oxyclozanide may bring liver toxicity. It reported that the level of TBIL, a breakdown product of normal haem catabolism, is an indicator for estimating liver function and cholestasis (34). Pathological reductions in TBIL are common in anemia caused by chronic glomerulonephritis (35) or aplastic anemia (36), but are not common in toxicological studies and do not have sufficient hematological changes to support this viewpoint. BUN is a kind of marker for assessing renal function in clinic diagnosis (32). During the study, the level of BUN increased in the high-dose oxyclozanide group in both males and females, which indicates that there was damage for functional nephrons because of inflammation or toxin productions in the kidney. Moreover, the level of UA was increased in males in the high-dose group, which may imply the oxyclozanide has an influence on renal function (32). Compared with the control values, the levels of LDH and CK were elevated in the treated group, which suggest that there were muscle damages, especially in liver and heart (32). The liver and kidney are important and sensitive organs to drug actions

(37). Specifically, the liver is an organ for the metabolism of cholesterol and glucose and the kidney was the main organ for excretion (38). The statistical elevation ($p < 0.05$ or $p < 0.01$) of BUN levels and the histopathology results with alternation of functional nephrons suggested that oxyclozanide could damage renal function.

During the histopathological examination of cardiac tissues, we detected some histopathological findings as evidence of toxicity like fiber degeneration, necrosis, and interstitial hydrops, which indicated that there were toxic damages after drug administration. The histopathological study also showed that glomerulonephritis and granular degeneration of tubular epithelium have alternated in renal function, which implied that oxyclozanide produced toxicity in the renal system. Steatosis and granular degeneration were found in the liver tissue of rats after administration, suggesting that there were toxic lesions in the liver. In the previous report, during the 90-day subchronic oral toxicity study, the brain had vacuolization phenotype in a high-dose oxyclozanide group in rats and dogs (21). However, our histopathological studies showed no pathological changes in brains between the treatment and the controls. It suggests that oxyclozanide doesn't show damage to the nervous system under subacute exposure. Besides, the histology of duodenum showed that there was catarrhal enteritis in the normal architecture resulting from diarrhea after administration.




FIGURE 2 | Histopathological analyses of the liver (A) and heart (B) in high-dose rats with/without oxyclozanide treatment. The arrows indicate that there are granular degeneration and fatty degeneration of liver cells in the liver (A) and myocardial fiber degeneration in the heart (B). Histopathological analyses of the kidney (C) and lung (D) in high-dose rats with/without oxyclozanide treatment. The arrows indicate that there are granular degeneration in renal tubular epithelial cells in the kidney (C) and normal in the lung (D). Histopathological analyses of the spleen (E) and duodenum (F) in high-dose rats with/without oxyclozanide treatment. The arrows indicate that they are normal in the spleen (E) and catarrhal enteritis in the duodenum (F). Histopathological analyses of the brain (G) and testis (H) in high-dose rats with/without oxyclozanide treatment. The arrows indicate that it is normal in the brain (G) and testis (H). Histopathological analyses of the ovary (I) and uterus (J) in high-dose rats with/without oxyclozanide treatment. The arrows indicate that it is normal in the ovary (I) and uterus (J).

The uncoupling effect of oxidative phosphorylation from drugs had a pharmacological effect on parasite. As a famous drug of salicylanilides, niclosamide has been considered to have an excellent safety range for a very long time. LD₅₀ of niclosamide was 2,500 mg/kg in rats (39) and 1,000 mg/kg in mice (40). The maximum tolerable dose of niclosamide was 2,000 mg/kg during the 28-day toxicity study of oral administration in rats. Moreover, many researchers have confirmed that niclosamide brought developmental and genetic toxicity to animals (41–44). However, there was still no related report on oxyclozanide.

CONCLUSION

The LD₅₀ of oxyclozanide was 3,707 mg/kg BW for experimental rats in both sex in the acute study. During the 28-day administration, differences in the histopathological, hematological, and biochemical assays were found at the 185- and 370-mg/kg dose groups vs. the untreated control group. These results suggest that oxyclozanide could cause damage in the liver, kidney, and heart. According to those results, the LOAEL level was 74 mg/kg for rats during

TABLE 5 | Histopathology results of organs in rats treated with oxyclozanide for 28 days.

Sex/organ	Changes	Dose			
		Control	Low	Medium	High
MALE					
Liver	Granular degeneration	0/12	2/12 ^c	7/12 ^b	10/12 ^a
	Fatty degeneration	0/12	0/12	1/12 ^a	4/12 ^b
Heart	Myocardial fiber degeneration	0/12	1/12 ^c	5/12 ^b	10/12 ^a
Kidney	Granular degeneration	0/12	3/12 ^c	7/12 ^b	10/12 ^a
Duodenum	Catarrhal enteritis	0/12	0/12	2/12 ^c	8/12 ^b
FEMALE					
Liver	Granular degeneration	0/12	2/12 ^c	6/12 ^b	10/12 ^a
	Fatty degeneration	0/12	0/12	0/12	3/12 ^b
Heart	Myocardial fiber degeneration	0/12	1/12 ^c	4/12 ^b	9/12 ^a
Kidney	Granular degeneration	0/12	2/12 ^c	5/12 ^b	9/12 ^a
Duodenum	Catarrhal enteritis	0/10	0/12	4/12 ^c	8/12 ^b

^aSevere, ^bModerate, ^cMild.
n = 12 photos for each group.

the 28-day toxicity study, which provides basis for clinic use of oxyclozanide and for determining a reasonable safe dose.

DATA AVAILABILITY

The datasets generated for this study are available on request to the corresponding author.

REFERENCES

- Ali F, Hussain R, Qayyum A, Gul ST, Iqbal Z, Hassan MF. Milk somatic cell counts and some hemato-biochemical changes in sub-clinical mastitic dromedary she-camels (*Camelus dromedarius*). *Pak Vet J.* (2016) 36:405–8.
- Elhaig MM, Selim A, Mahmoud MM, El-Gayar EK. Molecular confirmation of *Trypanosoma evansi* and *Babesia bigemina* in cattle from lower Egypt. *Pak Vet J.* (2016) 36:409–14.
- Qayyum A, Khan JA, Hussain R, Avais M, Ahmad N, Khan MS. Investigation of milk and blood serum biochemical profile as an indicator of sub-clinical mastitis in cholistani cattle. *Pak Vet J.* (2016) 36:275–9.
- Mas-Coma S, Funatsu IR, Bargues MD. *Fasciola hepatica* and lymnaeid snails occurring at very high altitude in South America. *Parasitology.* (2001) 123:S115–27. doi: 10.1017/S0031182001008034
- Mulatu H, Addis M. Study on the prevalence and risk factors of fasciolosis in small ruminants in and around Hirna town, Ethiopia. *Glob Vet.* (2011) 7:497–501.
- Broome AW, Jones WG. A new drug for the treatment of fascioliasis in sheep and cattle. *Nature.* (1966) 210:744–5. doi: 10.1038/210744a0
- Walley JK. Oxyclozanide (3,3',5,5',6-pentachloro-2, 2'-dihydroxybenzanilide-6Zanil[®]) in the treatment of the liver fluke *Fasciola hepatica* in sheep and cattle. *Vet Rec.* (1966) 78:267. doi: 10.1136/vr.78.8.267
- Jones EH. Fascioliasis and oxyclozanide. *Vet Rec.* (1966) 79:716–7.
- Hopkins CA, Grant PM, Stallard H. The effect of oxyclozanide on *Hymenolepis microstoma*, and *H. diminuta*. *Parasitology.* (1973) 66:11. doi: 10.1017/S0031182000045285
- Sanabria R, Moreno L, Alvarez L, Lanusse C, Romero J. Efficacy of oxyclozanide against adult *Paramphistomum leydeni* in naturally infected sheep. *Vet Parasitol.* (2014) 206:277–81. doi: 10.1016/j.vetpar.2014.09.022
- Arafa WM, Shokeir KM, Khateib AM. Comparing an *in vivo* egg reduction test and *in vitro* egg hatching assay for different anthelmintics against *Fasciola* species, in cattle. *Vet Parasitol.* (2015) 214:152–8. doi: 10.1016/j.vetpar.2015.09.023
- Coles GC, Stafford KA. Activity of oxyclozanide, nitroxylin, clorsulon and albendazole against adult triclabendazole-resistant *Fasciola hepatica*. *Vet Rec.* (2001) 148:723–4. doi: 10.1136/vr.148.23.723
- Ahmad M, Khan MN, Sajid MS, Muhammad G, Qudoos A, Rizwan HM. Prevalence, economic analysis and chemotherapeutic control of small ruminant fasciolosis in the Sargodha district of Punjab, Pakistan. *Vet Ital.* (2017) 53:47–53. doi: 10.12834/VetIt.114.316.6
- Rana MAA, Roohi N, Khan MA. Fascioliasis in cattle—a review. *J Anim Plant. Sci.* (2014) 24:668–75.
- Moll L, Gaasenbeek CPH, Vellema P, Borgsteede FHM. Resistance of *Fasciola hepatica* against triclabendazole in cattle and sheep in the Netherlands. *Vet Parasitol.* (2000) 91:153–8. doi: 10.1016/S0304-4017(00)00267-3
- Álvarez-Sánchez MA, Mainar-Jaime RC, Pérez-García J, Rojo-Vázquez FA. Resistance of *Fasciola hepatica* to triclabendazole and albendazole in sheep in Spain. *Vet Rec.* (2006) 159:424–5. doi: 10.1136/vr.159.13.424
- Olaechea F, Lovera V, Larroza M, Raffo F, Cabrera R. Resistance of *Fasciola hepatica* against triclabendazole in cattle in Patagonia (Argentina). *Vet Parasitol.* (2011) 178:364–6. doi: 10.1016/j.vetpar.2010.12.047
- Ortiz P, Scarcella S, Cerna C, Rosales C, Cabrera M, Guzmán M, et al. Resistance of *Fasciola hepatica* against triclabendazole in cattle in Cajamarca (Peru): a clinical trial and an *in vivo* efficacy test in sheep. *Vet Parasitol.* (2013) 195:118–21. doi: 10.1016/j.vetpar.2013.01.001

ETHICS STATEMENT

The animal study was reviewed and approved by Institutional Animal Care and Use Committee of Lanzhou Institute of Husbandry and Pharmaceutical Science of Chinese Academy of Agricultural Sciences.

AUTHOR CONTRIBUTIONS

JiyZ, WW, and ZD conceived and designed the experiments. XZ, XW, ZD, and WW performed the experiments. ZD, JiLZ, and BL analyzed the data. ZD, FC, and WW interpreted the references and provided background information. WW and ZD wrote the paper.

FUNDING

This work was supported by the Research on the Key Technologies of the Creation and Industrialization of New Special Chemical Drugs for Animals (grant number 2015BAD11B01-01) and Drug Development and Clinical Drug Use Posts of National Beef Yak Industry Technical System (grant number CARS-37).

ACKNOWLEDGMENTS

The authors appreciate Prof. Ning Jia (Gansu Agriculture University), Research Assistant Zhengying Qiu, and Guibo Wang for their generous support on the histopathological assays.

19. Venturina VM, Alejandro MAF, Baltazar CP, Abes NS, Mingala CN. Evidence of *Fasciola* spp. resistance to albendazole, triclabendazole and bromofenofos in water buffaloes (*Bubalus bubalis*). *Ann Parasitol.* (2015) 61:283–9. doi: 10.17420/ap6104.20
20. Richards R. The efficacy of triclabendazol and other anthelmintics against *Fasciola hepatica* in controlled studies in cattle. *Vet Rec.* (1990) 126:213.
21. EMEA. *Committee for Medicinal Products for Veterinary Use Oxytetracycline (Extrapolation to All Ruminants) Summary Report (3)*. European Agency for the Evaluation of Medicinal Products (2004). p. 1–6. Available online at: https://www.ema.europa.eu/documents/mrl-report/oxytetracycline-extrapolation-all-ruminants-summary-report-3-committee-veterinary-medicinal-products_en.pdf
22. Gokbulut C, Yalinkilinc HS, Aksit D, Veneziano V. Comparative pharmacokinetics of levamisole–oxytetracycline combination in sheep and goats following per os administration. *Can J Vet Res.* (2014) 78:316–20.
23. Marrugal-Lorenzo JA, Serna-Gallego A, Berastegui-Cabrera J, Pachón J, Sánchez-Céspedes J. Repositioning salicylanilide anthelmintic drugs to treat adenovirus infections. *Sci Rep.* (2019) 9:17. doi: 10.1038/s41598-018-37290-3
24. Maiden MM, Zachos MP, Waters CM. The ionophore oxytetracycline enhances tobramycin killing of *Pseudomonas aeruginosa* biofilms by permeabilizing cells and depolarizing the membrane potential. *J Antimicrob Chemother.* (2019) 74:894–906. doi: 10.1093/jac/dky545
25. Pic E, Burgain A, Sellam A. Repurposing the anthelmintic salicylanilide oxytetracycline against susceptible and clinical resistant *Candida albicans* strains. *Med Mycol.* (2019) 57:387–390. doi: 10.1093/mmy/myy027
26. Rajamuthiah R, Fuchs BB, Conery AL, Kim W, Jayamani E, Kwon B, et al. Repurposing salicylanilide anthelmintic drugs to combat drug resistant *Staphylococcus aureus*. *PLOS ONE.* (2015) 10:e0124595. doi: 10.1371/journal.pone.0124595
27. Alasadi A, Chen M, Swapna GVT, Tao H, Guo J, Collantes J, et al. Effect of mitochondrial uncouplers niclosamide ethanolamine (NEN) and oxytetracycline on hepatic metastasis of colon cancer. *Cell Death Dis.* (2018) 9:215. doi: 10.1038/s41419-017-0092-6
28. Veterinary Drug Assessment Center of the Ministry of Agriculture of the People's Republic of China. *Compilation of Technical Guidelines for Veterinary Drug Research*. Beijing: Chemical Industry Press (2012).
29. OECD 407. *Guideline for the Testing of Chemicals Test No. 407: Repeated Dose 28-Day Oral Toxicity Study in Rodents*. Paris (2008).
30. Lancaster PA, Carstens GE, Ribeiro FRB, Davis ME, Lyons JG, Welsh TH. Effects of divergent selection for serum insulin-like growth factor-I concentration on performance, feed efficiency, and ultrasound measures of carcass composition traits in Angus bulls and heifers. *J Anim Sci.* (2008) 86:2862–71. doi: 10.2527/jas.2008-1083
31. Litchfield JTJ, Wilcoxon F. A simplified method of evaluating dose–effect experiments. *J Pharmacol Exp Ther.* (1949) 96:99–113.
32. Haschek WM, Rousseaux CG. *Fundamentals of Toxicologic Pathology*. New York, NY: Academic Press (2010).
33. Foster JR. Spontaneous and drug-induced hepatic pathology of the laboratory beagle dog, the cynomolgus macaque and the marmoset. *Toxicol Pathol.* (2005) 33:63–74. doi: 10.1080/01926230590890196
34. Vitek L. The role of bilirubin in diabetes, metabolic syndrome, and cardiovascular diseases. *Front Pharmacol.* (2012) 3:1–7. doi: 10.3389/fphar.2012.00055
35. Loge JB, Lange RD, Moore CV. Characterization of the anemia associated with chronic renal insufficiency. *Am J Med.* (1958) 24:4–18. doi: 10.1016/0002-9343(58)90357-7
36. Rubin E, Gottlieb C, Vogel P. Syndrome of hepatitis and aplastic anemia. *Am J Med.* (1968) 45:88–97. doi: 10.1016/0002-9343(68)90010-7
37. Lin Q, Jia Z, Xu X, Xu S, Han T, Gao Y, et al. Sub-chronic toxicity study of arecae semen aqueous extract in Wistar rats. *J Ethnopharmacol.* (2018) 215:176–83. doi: 10.1016/j.jep.2017.08.031
38. Ezeja MI, Anaga AO, Asuzu IU. Acute and sub-chronic toxicity profile of methanol leaf extract of *Gouania longipetala* in rats. *J Ethnopharmacol.* (2014) 151:1155–64. doi: 10.1016/j.jep.2013.12.034
39. Hartley D, Kidd H. *The Agrochemicals Handbook*. Nottingham: The Royal Society of Chemistry (1983).
40. Perkow W, Ploss H. Wirksubstanzen der pflanzenschutz und schadlingsbekämpfungsmittel. *Anzeiger Schädlingskunde Pflanzenschutz Umweltschutz.* (1996) 69:10. doi: 10.1007/BF01905859
41. Huq F. Molecular modelling analysis of the metabolism of aniracetam. *J Pharmacol Toxicol.* (2003) 1:317–27. doi: 10.3923/jpt.2006.317.327
42. Tucker JD, Auletta A, Cimino MC, Dearfield KL, Jacobson-Kram D, Tice RR, et al. Sister-chromatid exchange: second report of the Gene-Tox program. *Mutat Res Rev Genet Toxicol.* (1993) 297:101–80. doi: 10.1016/0165-1110(93)90001-4
43. Ostrosky-Wegman P, García G, Montero R, Pérez Romero B, Alvarez Chacón R, Cortinas de Nava C. Susceptibility to genotoxic effects of niclosamide in human peripheral lymphocytes exposed *in vitro* and *in vivo*. *Mutat Res Lett.* (1986) 173:81–7. doi: 10.1016/0165-7992(86)90015-1
44. Abreu FC, Goulart MO, Brett AM. Detection of the damage caused to DNA by niclosamide using an electrochemical DNA-biosensor. *Biosens Bioelectron.* (2002) 17:913–9. doi: 10.1016/S0956-5663(02)00082-9

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Wang, Dong, Zhang, Zhou, Wei, Cheng, Li and Zhang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.