

Research Article

Wide Distribution of Virulence Genes among *Enterococcus faecium* and *Enterococcus faecalis* Clinical Isolates

Sara Soheili,¹ Sobhan Ghafourian,^{1,2} Zamberi Sekawi,¹ Vasanthakumari Neela,¹ Nourkhoda Sadeghifard,² Ramliza Ramli,³ and Rukman Awang Hamat¹

¹ Department of Medical Microbiology and Parasitology, Faculty of Medicine and Health Sciences, Universiti Putra Malaysia, 43400 Serdang, Malaysia

² Clinical Microbiology Research Center, Ilam University of Medical Sciences, Ilam, Iran

³ Department of Medical Microbiology and Immunology, Medical Faculty, Universiti Kebangsaan Malaysia Medical Center, Cheras, 56000 Kuala Lumpur, Malaysia

Correspondence should be addressed to Rukman Awang Hamat; rukman@upm.edu.my

Received 2 April 2014; Revised 13 June 2014; Accepted 19 June 2014; Published 15 July 2014

Academic Editor: Livia Visai

Copyright © 2014 Sara Soheili et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Enterococcus, a Gram-positive facultative anaerobic cocci belonging to the lactic acid bacteria of the phylum Firmicutes, is known to be able to resist a wide range of hostile conditions such as different pH levels, high concentration of NaCl (6.5%), and the extended temperatures between 5°C and 65°C. Despite being the third most common nosocomial pathogen, our understanding on its virulence factors is still poorly understood. The current study was aimed to determine the prevalence of different virulence genes in *Enterococcus faecalis* and *Enterococcus faecium*. For this purpose, 79 clinical isolates of Malaysian enterococci were evaluated for the presence of virulence genes. *pilB*, *fms8*, *efaAfm*, and *sgrA* genes are prevalent in all clinical isolates. In conclusion, the pathogenicity of *E. faecalis* and *E. faecium* could be associated with different virulence factors and these genes are widely distributed among the enterococcal species.

1. Introduction

Enterococci are considered to be part of normal gut microbiota in both humans and animals and capable to survive in a diverse range of harsh conditions. Among them only two, that is, *Enterococcus faecium* and *Enterococcus faecalis*, are now being increasingly recognized to be involved in human infections such as bacteremia, endocarditis, urinary tract infections, and surgical site infections [1, 2]. This could be explained by their inherent resistance to various antibiotics with greater adaptability in hospital environments and has high genetic diversity as well as the presence of various virulence factors [3–5].

These opportunistic bacteria possess various virulence factors including enterococcal surface protein (Esp) and aggregation substance (Agg) which could enhance the colonization process in the host and binding to the host

epithelium, respectively [6]. Others such as cytolyisin [7], enterolysin A [8], gelatinase [9], hyaluronidase, Zn-metalloendopeptidase, enhanced expression of pheromone (Eep) [10], and adhesion-associated protein EfaA (*E. faecalis* endocarditis antigen A) [11] have been reported to be among the most important virulence factors.

Surprisingly through NCBI database analysis, there are some similarities observed in terms of their virulence determinants among enterococci. For instance, *pilA* in *E. faecium* has 99% similarity with the cell wall surface anchor family protein of *Enterococcus* sp. GMD4E (Accession: EKA01662). The *pilB* of *E. faecium* has 74% similarity with the cell wall surface anchor protein of *E. faecalis* B318 (Accession: ETU21897). Meanwhile, the *ecbA* in *E. faecium* has 92% similarity with the cell wall surface anchor protein of *E. faecalis* (Accession: WP_010708790) and 91% with Cna protein B-type domain protein *E. faecalis* (Accession: WP_002417581)

TABLE 1: Characteristics of different primers used in the current study.

Primer name	Sequence (5'-3')	Size (bp)	Accession number
<i>pilA</i>	Forward: AAAACGCCACCAGAGAAGGT Reverse: CATTGGCGCAATCACAACCA	459	ENA ACI49671
<i>pilB</i>	Forward: GATACCCAGCTGACGGCTTT Reverse: GGTACTGCCGAAAACGAAGC	959	EFF34880
<i>fms8</i>	Forward: AGACGAGCAGATGAACAGCC Reverse: CCCGTCAATCGTCTACTGT	765	YP_006376887
<i>efaAfm</i>	Forward: AAAAGGCAAGCGACGCAGAT Reverse: AGGTCTAGCCAAGCATGAGG	199	FJ609170
<i>sgrA</i>	Forward: CTGATCGGATTGTTTATGA Reverse: AATAAACTTCCCCAATAACTT	150	AGS75503
<i>ecbA</i>	Forward: GGAGTGAGGCTTTTAAACCAGA Reverse: GGAAACAGGGTACTTTG	182	AGS75851
<i>hyl</i>	Forward: CCCTGGACACATGAAATGCG Reverse: AGCATCGGCCGTTGATAGAC	605	AF544400
<i>scm</i>	Forward: GTTTACTAGTCTAGTTGC Reverse: TCTGTACTGTCGCTTGTGTC	1015	YP_006377279

(these regions are part of a collagen binding MSCRAMM, *ecbA*).

The cell wall adhesion (*efaAfm*) in *E. faecium* has 66% similarity with endocarditis specific antigen of *E. faecalis* T8 (Accession: EEU25576). With regard to the collagen binding MSCRAMM *fms8* in *E. faecium*, 57% similarity has been observed with the collagen binding domain protein of *E. faecalis* (Accession: WP_016632409). The blast analysis for the collagen binding MSCRAMM *scm* in *E. faecium* has revealed 99% similarity with the collagen-binding MSCRAMM *scm* (*Fms10*) in *Enterococcus* sp. GMD5E. Finally, 37% similarity of the cell wall anchored protein *sgrA* in *E. faecium* has been detected when compared to the surface adhesion protein in *Enterococcus* sp. C1 (Accession: WP_008378446). These findings could be related to the notion that horizontal transfer of both resistance and virulence determinants is very common among enterococci [12]. Thus, this study was conducted to investigate the distribution of the diverse virulence factors and specify the dominant virulence genes among *E. faecalis* and *E. faecium* clinical isolates. Based on the similarity results and also the possibility of horizontal virulence gene transfer the *pilA*, *pilB*, *hyl*, *ecbA*, *scm*, *fms8*, *efaAfm*, and *sgrA* genes were chosen for this study.

2. Material and Methods

2.1. Bacterial Isolates. A total of 79 clinical isolates of *E. faecalis* (50 isolates) and *E. faecium* (29 isolates) were identified during May 2009 to March 2010 from a tertiary teaching hospital. Nonrepetitive isolates were collected from these samples such as urine, blood, pus, vaginal, and sterile body fluid.

2.2. DNA Extraction of Enterococcus. The DNA was extracted using the DNA extraction kit (Gene ALL, South Korea) according to the manufacturer's instructions.

2.3. Evaluation of Different Virulence Genes. All isolates were subjected to the amplified *pilA*, *pilB*, *hyl*, *ecbA*, *scm*, *fms8*, *efaAfm*, and *sgrA* genes, using specific primers as listed in Table 1. These virulence genes are most strongly associated with clinical lineages of *E. faecium*; however, not much data has been reported among *E. faecalis*. The PCR amplification was carried out in a DNA thermocycler (Bio-Rad) using the amplification parameters as follows: initial denaturation at 95°C for 2 minutes, followed by 35 cycles of denaturation at 95°C for 20 seconds, annealing at 58°C for 10 seconds, and extension at 72°C for 20 seconds, with a final extension at 72°C for 5 minutes. All PCR products were analyzed by 1% agarose gel electrophoresis.

2.4. Purification of Virulence Genes from Agarose Gel. PCR products of each virulence gene were run on 1% (w/v) molecular grade agarose gel (Sigma, USA), using a Bio-Rad mini-gel electrophoresis system at 80 V for 70 minutes. The DNA was purified using DNA purification kit (Gene ALL, South Korea) according to the manufacturer's instructions.

2.5. Sequence Analysis. The PCR products were purified from gel agarose and then the purified products were sequenced by Sigma Company (Singapore). The results of DNA sequencing were run in Chromas Lite program to analyze the similarity to the sequenced gene in GenBank library.

3. Results and Discussion

A total of 79 *E. faecalis* (50 isolates) and *E. faecium* (29 isolates) were analyzed for the presence of different virulence genes including *pilA*, *pilB*, *hyl*, *ecbA*, *scm*, *fms8*, *efaAfm*, and *sgrA*. The analysis showed different prevalence of virulence genes in *Enterococcus* which ranged from 35.4% to 100%. The *pilB*, *fms8*, *efaAfm*, and *sgrA* were identified as the dominant virulence genes in all isolates. The second most prevalent virulence gene, *scm*, was found in 92.4% of the isolates

TABLE 2: Distribution of different virulence factors in *E. faecium* and *E. faecalis*.

Virulence genes	<i>E. faecalis</i> (n = 50)	<i>E. faecium</i> (n = 29)	Total (N = 79)
<i>pilB</i>	50 (100%)	29 (100%)	79 (100%)
<i>fms8</i>	50 (100%)	29 (100%)	79 (100%)
<i>efaAfm</i>	50 (100%)	29 (100%)	79 (100%)
<i>sgrA</i>	50 (100%)	29 (100%)	79 (100%)
<i>scm</i>	44 (88%)	29 (100%)	73 (92.4%)
<i>ecbA</i>	35 (70%)	29 (100%)	64 (81%)
<i>pilA</i>	29 (58%)	29 (100%)	58 (73.4%)
<i>hyl</i>	4 (8%)	24 (82.7%)	28 (35.4%)

(n = 73/79). The *ecbA* was determined as the third most common prevalent virulence gene with 81% (n = 64/79) frequency. The *pilA* showed 73.4% (n = 58/79) prevalence in clinical isolates of *Enterococcus*. The lowest prevalence detected was *hyl* with the prevalence of 35.4% (n = 28/79). The prevalence of different virulence genes is shown in Figure 1. With regard to both enterococcal species, the analysis showed that all selected virulence genes were positive among *E. faecium* isolates. The exception is *hyl* where it was detected with 82.7% prevalence rate. Among *E. faecalis* isolates, *pilB*, *fms8*, *efaAfm*, and *sgrA* genes were detected in all isolates, followed by *scm* (88%), *ecbA* (70%), and *pilA* (58%). The least prevalence was *hyl* which was detected in only 4 isolates (8%). Table 2 shows the distribution of different virulent determinants among *E. faecium* and *E. faecalis* clinical isolates. We believe that this is the first report on the prevalence of selected virulence genes among *E. faecalis* which could be demonstrated by the possibility of horizontal gene transfer among *E. faecium* and *E. faecalis*.

Pili which are also known as fimbriae have been detected in Gram-positive bacteria [13–15]. This surface organelle is responsible for endocarditis and biofilm formation in Gram-positive bacteria [16], and mediates attachment to human epithelium and skin [17] and confers resistance against macrophages [18].

So far, limited data has been documented on the prevalence of pili in *E. faecium*. In our study, high prevalence rate of *pilB* in clinical *E. faecium* and *E. faecalis* isolates (100% for each) has been observed. In comparison with *pilB*, 21 (26.6%) clinical *E. faecalis* isolates were negative for the presence of *pilA*. Figures 2 and 3 show the presence of *pilA* and *pilB* genes in representative isolates. Hendrickx et al. [19] have demonstrated the presence of *pilA* and *pilB* in clinical isolates of *E. faecium*. Their analysis showed that *pilA* could not be expressed in broth condition and the best temperature for their expression was 37°C. This could probably be a reason as to why we could not detect *pilA* in some of our clinical *E. faecalis* isolates. Another possible reason could be explained by the lack of horizontal gene transfer among *E. faecalis* isolates.

Nowadays, it is obvious that in pathogenic bacteria several proteins have evolved to adhere to and invade into the host cells and subsequently escape and resist host defense




FIGURE 1: Prevalence of different virulence genes in clinical *E. faecium* and *E. faecalis* isolates. The *pilB*, *fms8*, *efaAfm*, and *sgrA* were identified in all clinical isolates.

[20]. MSCRAMMs which are known as microbial surface components recognizing adhesive matrix molecules have an important role for cell adhesion and involve in pathogenicity of bacteria [21]. *In silico* analysis has revealed a family of genes that encode MSCRAMM-like proteins in *Enterococcus* such as *ebp* (endocarditis and biofilm-associated pilus of *E. faecalis*) operon, *ace* (adhesion of collagen from *E. faecalis*), and *acm* (adhesion of collagen from *E. faecium*) [22–24].

The role of collagen-binding MSCRAMM *acm* (*Fms8*) is to bind to the collagen types I and IV. These types of collagen are also an important antigen involved in human during endocarditis. In the current research, high prevalence rate of *fms8* (100% for each) was detected among *E. faecium* and *E. faecalis* clinical isolates and could play an important role in pathogenicity of both enterococcal species. Figure 4 shows the presence of *fms8* genes in representative isolates.

The exact role of *efaAfm* is unknown although it is believed to be involved in cell wall adherence. The *efaAfm* gene was only found in *E. faecium* isolates [25]. A study by Barbosa et al. [26] has demonstrated 27% of isolates harbored the *efaAfm* gene while the current study showed all isolates were positive for *efaAfm*. The presence of *efaAfm* gene among representative isolates is shown in Figure 5. Our finding corroborates with Martin et al. [27] where they have demonstrated all *E. faecium* isolates carrying *efaAfm* virulence gene.

The pathogenic *E. faecium* is enriched with two *orf2351* and *orf2430* genes which encode the *sgrA* and *ecbA* LPXTG-like cell wall anchored proteins, respectively. *sgrA* is a surface adhesion that can bind to the extracellular matrix molecules nidogen 1 and nidogen 2 and is also involved in biofilm formation. Meanwhile, *ecbA* binds to the collagen type V and




FIGURE 2: PCR results of *pilA*. 1 = marker = 100 bp; 2-16 = *pilA* = 495 bp; 17 = negative control.



FIGURE 3: PCR results of *pilB*. 1, 10 = marker = 100 bp; 2-8 = *pilB* = 959 bp; 9 = negative control.




FIGURE 5: PCR results of *efaAfm*. 1, 8 = marker = 100 bp; 2-6 = *efaAfm* = 199 bp; 7 = negative control.




FIGURE 4: PCR results of *fms8*. 1 = marker = 100 bp; 2-8 = *fms8* = 765 bp; 9 = negative control.




FIGURE 6: PCR results of *sgrA*. 1 = marker = 100 bp; 2-11 = *sgrA* = 150 bp; 12 = negative control.

fibrinogen. Both *ecbA* and *sgrA* were reported to be prevalent in clinical strains of *E. faecium* [28]. However, our study demonstrated that *sgrA* was more prevalent than *ecbA* (100% versus 81%). The representative isolates for the presence of *sgrA* and *ecbA* genes are shown in Figures 6 and 7.

In addition, *hyl* virulence gene encodes for a putative glycosyl hydrolase that is considered as a plasmid harboring gene which has colonized the gastrointestinal tracts of mice subsequently caused an increase of pathogenicity of *E. faecium* in experimental peritonitis [29]. In a study by Panesso et al. [30] that subjected 32 hospitals in Colombia, Ecuador, Peru, and Venezuela, the results revealed that 23% of *E. faecium* strains carried the *hyl* virulence gene. Our results

are in accordance with the previous study by Panesso et al. [30]. The *hyl* gene among representative isolates is shown in Figure 8. In a research by Worth et al. [31] among Australian haematology patients, only 2.1% of isolates showed positive reaction to *hyl*. A study by Vanckerckhoven et al. [32] revealed 71% of VRE and 29% of vancomycin sensitive *Enterococcus* (VSE) harbored *hyl* virulence gene. Since *hyl* is not prevalent in all clinical isolates in our study, we believe that this gene could have little role in pathogenicity of *Enterococcus* in comparison with other prevalent virulence genes.



FIGURE 7: PCR results of *ecbA*. 1 = marker = 100 bp; 2–11 = *ecbA* = 182 bp; 12 = negative control.




FIGURE 8: PCR results of *hyl*. 11 = marker = 100 bp; 2–9 = *hyl* = 605 bp; 1, 10 = negative control.

Scm (second collagen adhesion of *E. faecium*) binds to collagen type V and fibrinogen and it is commonly distributed among clinical and nonclinical isolates of *E. faecium*. *Scm* was first described in *E. faecium* in 2008 [33] but limited information on the prevalence of *scm* in *E. faecium* has been reported. Nonetheless, we documented a high prevalence of *scm* in clinical isolates of *E. faecium* (92.4%). Isolates representing for the presence of *scm* gene are shown in Figure 9.

4. Conclusion

The wide distribution of several virulence genes that is *pilB*, *fms8*, *efaAfm*, and *sgrA* in *E. faecalis* and *E. faecium* clinical isolates could give a clue that these virulence genes might play an important role in the pathogenicity of both enterococcal species. This could also be explained that the horizontal virulence gene transfer is common among the clinical isolates.

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.




FIGURE 9: PCR results of *scm*. 7 = marker = 100 bp; 1–5 = *scm* = 1015 bp; 6 = negative control.

Acknowledgment

This study was supported by Universiti Putra Malaysia, Research University Grant Scheme (RUGS), with Project no. 04-20-12-1766RU.

References

- [1] V. Shankar, A. S. Baghdayan, M. M. Huycke, G. Lindahl, and M. S. Gilmore, "Infection-derived *Enterococcus faecalis* strains are enriched in *esp*, a gene encoding a novel surface protein," *Infection and Immunity*, vol. 67, no. 1, pp. 193–200, 1999.
- [2] I. G. Sava, E. Heikens, and J. Huebner, "Pathogenesis and immunity in enterococcal infections," *Clinical Microbiology and Infection*, vol. 16, no. 6, pp. 533–540, 2010.
- [3] E. N. Vergis, N. Shankar, J. W. Chow et al., "Association between the presence of enterococcal virulence factors gelatinase, hemolysin, and enterococcal surface protein and mortality among patients with bacteremia due to *Enterococcus faecalis*," *Clinical Infectious Diseases*, vol. 35, no. 5, pp. 570–575, 2002.
- [4] J. Top, R. Willems, and M. Bonten, "Emergence of CC17 *Enterococcus faecium*: from commensal to hospital-adapted pathogen," *FEMS Immunology and Medical Microbiology*, vol. 52, no. 3, pp. 297–308, 2008.
- [5] P. L. Weng, R. Ramli, M. N. Shamsudin, Y.-K. Cheah, and R. A. Hamat, "High genetic diversity of *Enterococcus faecium* and *Enterococcus faecalis* clinical isolates by pulsed-field gel electrophoresis and multilocus sequence typing from a hospital in Malaysia," *BioMed Research International*, vol. 2013, Article ID 938937, 6 pages, 2013.
- [6] A. P. Johnson, "The pathogenicity of enterococci," *Journal of Antimicrobial Chemotherapy*, vol. 33, no. 6, pp. 1083–1089, 1994.
- [7] Y. Ike, D. B. Clewell, R. A. Segarra, and M. S. Gilmore, "Genetic analysis of the pAD1 hemolysin/bacteriocin determinant in *Enterococcus faecalis*: Tn917 insertion mutagenesis and cloning," *Journal of Bacteriology*, vol. 172, no. 1, pp. 155–163, 1990.
- [8] T. Nilsen, I. F. Nes, and H. Holo, "Enterolysin A, a cell wall-degrading bacteriocin from *Enterococcus faecalis* LMG 2333,"

- Applied and Environmental Microbiology*, vol. 69, no. 5, pp. 2975–2984, 2003.
- [9] Y. A. Su, M. C. Sulavik, P. He et al., “Nucleotide sequence of the gelatinase gene (gelE) from *Enterococcus faecalis* subsp. *liquefaciens*,” *Infection and Immunity*, vol. 59, no. 1, pp. 415–420, 1991.
- [10] F. Y. An, M. C. Sulavik, and D. B. Clewell, “Identification and characterization of a determinant (eep) on the *Enterococcus faecalis* chromosome that is involved in production of the peptide sex pheromone cAD1,” *Journal of Bacteriology*, vol. 181, no. 19, pp. 5915–5921, 1999.
- [11] A. M. Lowe, P. A. Lambert, and A. W. Smith, “Cloning of an *Enterococcus faecalis* endocarditis antigen: homology with adhesins from some oral streptococci,” *Infection and Immunity*, vol. 63, no. 2, pp. 703–706, 1995.
- [12] T. J. Eaton and M. J. Gasson, “Molecular Screening of Enterococcus Virulence Determinants and Potential for Genetic Exchange between Food and Medical Isolates,” *Applied and Environmental Microbiology*, vol. 67, no. 4, pp. 1628–1635, 2001.
- [13] P. Lauer, C. D. Rinaudo, M. Soriani et al., “Microbiology: Genome analysis reveals pili in group B streptococcus,” *Science*, vol. 309, no. 5731, p. 105, 2005.
- [14] M. K. Yeung and P. A. Ragsdale, “Synthesis and function of *Actinomyces naeslundii* T14V type 1 fimbriae require the expression of additional fimbria-associated genes,” *Infection and Immunity*, vol. 65, no. 7, pp. 2629–2639, 1997.
- [15] M. K. Yeung, J. A. Donkersloot, J. O. Cisar, and P. A. Ragsdale, “Identification of a gene involved in assembly of *Actinomyces naeslundii* T14V type 2 fimbriae,” *Infection and Immunity*, vol. 66, no. 4, pp. 1482–1491, 1998.
- [16] S. R. Nallapareddy and B. E. Murray, “Ligand-signaled upregulation of *Enterococcus faecalis* ace transcription, a mechanism for modulating host-*E. faecalis* interaction,” *Infection and Immunity*, vol. 74, no. 9, pp. 4982–4989, 2006.
- [17] E. L. Abbot, W. D. Smith, G. P. S. Siou et al., “Pili mediate specific adhesion of *Streptococcus pyogenes* to human tonsil and skin,” *Cellular Microbiology*, vol. 9, no. 7, pp. 1822–1833, 2007.
- [18] H. C. Maisey, D. Quach, M. E. Hensler et al., “A group B streptococcal pilus protein promotes phagocyte resistance and systemic virulence,” *The FASEB Journal*, vol. 22, no. 6, pp. 1715–1724, 2008.
- [19] A. P. A. Hendrickx, M. J. M. Bonten, M. van Luit-Asbroek, C. M. E. Schapendonk, A. H. M. Kragten, and R. J. L. Willems, “Expression of two distinct types of pili by a hospital-acquired *Enterococcus faecium* isolate,” *Microbiology*, vol. 154, no. 10, pp. 3212–3223, 2008.
- [20] J. Pizarro-Cerdá and P. Cossart, “Bacterial adhesion and entry into host cells,” *Cell*, vol. 124, no. 4, pp. 715–727, 2006.
- [21] J. M. Patti, B. L. Allen, M. J. McGavin, and M. Hook, “MSCRAMM-mediated adherence of microorganisms to host tissues,” *Annual Review of Microbiology*, vol. 48, pp. 585–617, 1994.
- [22] M. G. Bowden, W. Chen, J. Singvall et al., “Identification and preliminary characterization of cell-wall-anchored proteins of *Staphylococcus epidermidis*,” *Microbiology*, vol. 151, no. 5, pp. 1453–1464, 2005.
- [23] S. R. Nallapareddy, X. Qin, G. M. Weinstock, M. Hook, and B. E. Murray, “*Enterococcus faecalis* adhesin, ace, mediates attachment to extracellular matrix proteins collagen type IV and laminin as well as collagen type I,” *Infection and Immunity*, vol. 68, no. 9, pp. 5218–5224, 2000.
- [24] Y. Xu, X. Liang, Y. Chen, T. M. Koehler, and M. Höök, “Identification and biochemical characterization of two novel collagen binding MSCRAMMs of *Bacillus anthracis*,” *Journal of Biological Chemistry*, vol. 279, no. 50, pp. 51760–51768, 2004.
- [25] M. Martín, J. Gutiérrez, R. Criado, C. Herranz, L. M. Cintas, and P. E. Hernández, “Genes encoding bacteriocins and their expression and potential virulence factors of enterococci isolated from wood pigeons (*Columba palumbus*),” *Journal of Food Protection*, vol. 69, no. 3, pp. 520–531, 2006.
- [26] J. Barbosa, V. Ferreira, and P. Teixeira, “Antibiotic susceptibility of enterococci isolated from traditional fermented meat products,” *Food Microbiology*, vol. 26, no. 5, pp. 527–532, 2009.
- [27] B. Martin, M. Garriga, M. Hugas, and T. Aymerich, “Genetic diversity and safety aspects of enterococci from slightly fermented sausages,” *Journal of Applied Microbiology*, vol. 98, no. 5, pp. 1177–1190, 2005.
- [28] A. P. A. Hendrickx, M. Van Luit-Asbroek, C. M. E. Schapendonk et al., “SgrA, a nidogen-binding LPXTG surface adhesin implicated in biofilm formation, and EcbA, a collagen binding MSCRAMM, are two novel adhesins of hospital-acquired *Enterococcus faecium*,” *Infection and Immunity*, vol. 77, no. 11, pp. 5097–5106, 2009.
- [29] C. A. Arias, D. Panesso, K. V. Singh, L. B. Rice, and B. E. Murray, “Cotransfer of antibiotic resistance genes and a hylEfm-containing virulence plasmid in *Enterococcus faecium*,” *Antimicrobial Agents and Chemotherapy*, vol. 53, no. 10, pp. 4240–4246, 2009.
- [30] D. Panesso, J. Reyes, S. Rincón et al., “Molecular epidemiology of vancomycin-resistant *Enterococcus faecium*: a prospective, multicenter study in South American hospitals,” *Journal of Clinical Microbiology*, vol. 48, no. 5, pp. 1562–1569, 2010.
- [31] L. J. Worth, M. A. Slavin, V. Vankerckhoven, H. Goossens, E. A. Grabsch, and K. A. Thursky, “Virulence determinants in vancomycin-resistant *Enterococcus faecium* vanB: clonal distribution, prevalence and significance of esp and hyl in Australian patients with haematological disorders,” *Journal of Hospital Infection*, vol. 68, no. 2, pp. 137–144, 2008.
- [32] V. Vankerckhoven, T. Van Autgaerden, C. Vael et al., “Development of a multiplex PCR for the detection of *asaI*, *gelE*, *cylA*, *esp*, and *hyl* genes in enterococci and survey for virulence determinants among European hospital isolates of *Enterococcus faecium*,” *Journal of Clinical Microbiology*, vol. 42, no. 10, pp. 4473–4479, 2004.
- [33] J. Sillanpää, S. R. Nallapareddy, V. P. Prakash et al., “Identification and phenotypic characterization of a second collagen adhesin, Scm, and genome-based identification and analysis of 13 other predicted MSCRAMMs, including four distinct pilus loci, in *Enterococcus faecium*,” *Microbiology*, vol. 154, no. 10, pp. 3199–3211, 2008.