

Original Article
Preventive Medicine,
Occupation &
Environmental Medicine



Spatial and Temporal Trends of Number of Deaths Attributable to Ambient PM_{2.5} in the Korea

Changwoo Han ¹, Soontae Kim ², Youn-Hee Lim ^{3,4}, Hyun-Joo Bae ⁵,
and Yun-Chul Hong ^{1,3,4}

¹Department of Preventive Medicine, Seoul National University College of Medicine, Seoul, Korea

²Department of Environmental and Safety Engineering, Ajou University, Suwon, Korea

³Institute of Environmental Medicine, Seoul National University Medical Research Center, Seoul, Korea

⁴Environmental Health Center, Seoul National University College of Medicine, Seoul, Korea

⁵Korea Environmental Institute, Seoul, Korea



Received: Jan 9, 2018

Accepted: May 10, 2018

Address for Correspondence:

Yun-Chul Hong, MD, PhD

Department of Preventive Medicine, Seoul National University College of Medicine, 130 Daehak-ro, Jongno-gu, Seoul 03080, Korea.
E-mail: ychong1@snu.ac.kr


© 2018 The Korean Academy of Medical Sciences.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID iDs

Changwoo Han 

<https://orcid.org/0000-0002-5127-4680>

Soontae Kim 


<https://orcid.org/0000-0002-1198-934X>

Youn-Hee Lim 

<https://orcid.org/0000-0002-1290-5814>

Hyun-Joo Bae 

<https://orcid.org/0000-0003-2723-5013>

Yun-Chul Hong 

<https://orcid.org/0000-0001-9010-7271>

Funding

This research was supported by the National Strategic Project-Fine particle of the National Research Foundation of Korea (NRF), funded by the Ministry of Science and ICT (MSIT), the Ministry of Environment (ME),

ABSTRACT

Background: We aimed to evaluate the spatial and temporal trends of the health burden attributable to particulate matter less than 2.5 µm in diameter (PM_{2.5}) in the metropolitan cities and provinces of the Korea.

Methods: We used modeled PM_{2.5} concentration data for the basic administrative levels, comprising the cities and the provinces of Korea, the corresponding annual population census data for each level, and the age and cause specific mortality data. We applied cause-specific integrated exposure-response functions to calculate the premature mortality attributable to ambient PM_{2.5} for four disease end points (ischemic heart disease [IHD], chronic obstructive pulmonary disease [COPD], lung cancer [LC], and cerebrovascular disease [stroke]) for the year 2015. Moreover, the temporal trends of the health burden from 2006 to 2015 were assessed.

Results: The annual average PM_{2.5} concentration for Korea was 24.4 µg/m³, and 11,924 premature deaths were attributable to PM_{2.5} exposure in 2015. By simulating the reduction in the annual mean values of PM_{2.5} to 10 µg/m³, about 8,539 premature deaths were preventable. There was spatial variation in mortality burden attributable to PM_{2.5} across the sub-national regions of Korea. In particular, the high burden was concentrated at Seoul and Gyeonggi province due to the high population density. However, decreasing trends were noted for most of the metropolitan cities and provinces of Korea since 2006.

Conclusion: Our findings show that further actions to improve air quality in Korea would substantially improve the health burden due to particulate matter.

Keywords: Particulate Matter; Health Burden; Mortality; Korea

INTRODUCTION

Particulate matter 2.5 µm or less in diameter (PM_{2.5}) is becoming a global health threat. Previous systematic reviews of epidemiologic studies revealed causal relationships between PM_{2.5} exposure and mortality outcomes such as ischemic heart disease, chronic obstructive pulmonary disease, lung cancer, and cerebrovascular disease.¹ According to a recent report from the World Health Organization (WHO), a global estimation of about 3 million deaths

and the Ministry of Health and Welfare (MOHW) (#2017M3D8A1092008 and #2017M3D8A1092009).

Disclosure

The authors have no potential conflicts of interest to disclose.

Author Contributions

Conceptualization: Han C, Hong YC. Data curation: Han C, Kim S. Formal analysis: Han C. Investigation: Han C, Lim YH, Bae HJ, Hong YC. Writing - original draft: Han C, Hong YC. Writing - review & editing: Han C, Kim S, Lim YH, Bae HJ, Hong YC.

and 84 million disability-adjusted life years (DALYs) were attributable to the ambient PM_{2.5} exposure in 2012.² The Global Burden of Disease (GBD) group reported that ambient PM_{2.5} is the fifth ranking mortality risk factor, which accounts for the 4.2 million deaths and 103 million DALYs in 2015.³ Due to increasing number of studies reporting possible associations between ambient particulate matter exposure and diverse health outcomes, total disease burden from particulate matter exposure might be higher than the current estimation indicate.⁴

There is a growing public interest regarding PM_{2.5} in the Republic of Korea. Among the Organization for Economic Co-operation and Development (OECD) countries, Korea has the second highest population weighted annual PM_{2.5} exposure levels with 29 µg/m³ in 2015.⁵ Decreasing trends of the annual particulate matter concentrations has been flattened in recent years, and days with high concentrations of particulate matter has been increased in Korea.^{6,7} To meet the increasing public demands, the government of Korea commenced a sub-national level forecasting system on particulate matter 10 µm or less in diameter (PM₁₀) and PM_{2.5} from February 2014 and January 2015, respectively. The government unveiled special management measures to deal with PM_{2.5} in June 2016 and, as part of the process, eight old coal-fired power plants (30 years or older) were shut down for 1 month in June 2017.

Recent health impact assessment of air pollution from the global levels of estimation indicates that 11,523 and 18,148 deaths were attributable to ambient PM_{2.5} exposure in Korea for 2012 and 2015, respectively.^{2,8} However, no study has been conducted at the sub-national levels to evaluate the spatial distribution and the temporal trends of the health burden across Korea.

Therefore, in this study, we conducted a sub-national level analysis to estimate the health burden attributable to PM_{2.5} exposure and evaluated the temporal trends of the health burden in Korea. We used the latest publicly available population structure and mortality data with modeled PM_{2.5} concentration covering all the sub-national regions of the country from 2006 to 2015.

METHODS

Study design

Korea is composed of eight metropolitan cities and nine provinces. Each metropolitan city is composed of 'gu' while provinces are composed of 'si' and 'gun,' which are the basic administration units of Korea. To calculate the number of deaths attributable to PM_{2.5} in the sub-national level, we need following three information: 1) annual PM_{2.5} concentrations, 2) the population structure, 3) age and disease specific mortality rates. For this study, annual PM_{2.5} concentration and the population structure data were collected in basic administration unit levels, while the mortality data were collected in metropolitan cities or the province level.

PM_{2.5} exposure data

We used a Weather Research and Forecast (WRF, version 3.3.1) - Sparse Matrix Operator Kernel Emission (SMOKE, version 3.1) - Community Multiscale Air Quality (CMAQ, version 4.7.1) system to prepare daily PM_{2.5} concentration from 2006 to 2015 in the basic administrative unit levels. Details of the modeling methodologies and validation results are presented in a previous research articles.^{9,10} In brief, we utilized the modeling system to

estimate formation, transport, and removal of $PM_{2.5}$. We used the CMAQ with the AERO5 aerosol module and the Statewide Air Pollution Research Center version 99 to simulate air quality in Korea. We conducted meteorology simulation using WRF with the National Center for Environmental Protection FNL (Final) data as the initial field. The CMAQ-ready meteorological inputs were prepared using the Meteorology Chemistry Interface Processor version 3.6. For anthropogenic emissions, the Korean National Emissions Inventory named Clean Air Policy Support System 2010 was processed with SMOKE. Biogenic emissions were estimated with the Model of Emissions of Gases and Aerosols from Nature (MEGAN). We adopted two modeling domains with horizontal resolutions of 27-km and 9-km. The coarse modeling domain covers the Northeast Asia including China, Japan, and Korea. Simulation results from the coarse modeling domain was used to derive the boundary conditions for the 9-km modeling domain. Gridded hourly $PM_{2.5}$ concentrations from CMAQ simulations for the fine model domain were then resampled for each administration in Korea based on a GIS shape file. During the process, hourly $PM_{2.5}$ observation data were nudged on the modeling field to minimize model uncertainties.

Population structure and the mortality rate data

To calculate the health burden attributable to $PM_{2.5}$, we used the 2015 population census data, which are the results of the complete enumeration survey. Owing to the enumeration surveys being conducted every five years, we used the mid-year population of the resident registration data from 2006 to 2015 to estimate the temporal trends of the health burden. The number of the population for different age groups (25–29, 30–34, 35–39, 40–44, 45–49, 50–54, 55–59, 60–64, 65–69, 70–74, 75–79, and 80–) in each administrative unit ('si,' 'gun,' and 'gu') were collected from the Korean Statistical Information Service website.¹¹

Mortality data of each city and province were collected from the Korean Statistical Information Service MicroData Integrated Service homepage.¹² We selected the following four disease endpoints (corresponding International Classification of Disease, 10th revision code): ischemic heart disease (IHD, I20–I25), chronic obstructive pulmonary disease (COPD, J40–J44), lung cancer (LC, C33–C34), and cerebrovascular disease (stroke, I60–I69) to calculate the health burden attributable to $PM_{2.5}$ exposure. Disease outcomes were selected based on the epidemiological studies indicating their robust causal relationship with $PM_{2.5}$ exposure.^{1,13} Age-specific mortality rate (/100,000 persons) was calculated for the city or province levels by dividing the number of yearly age groups, and cause-specific mortality rate with the corresponding mid-year population. We calculated the city or province level mortality rates because public deaths registration data were unavailable in basic administration unit levels.

Statistical analysis

We applied cause-specific integrated exposure-response (IER) functions developed for the Global Burden of Disease 2010 and 2013 to estimate the premature mortality attributable to ambient $PM_{2.5}$.¹³ In previous health impact assessment studies, researchers used diverse concentration-mortality functions based on epidemiological cohort studies to estimate the health burden attributable to $PM_{2.5}$. However, most of the previous studies were conducted in Western regions where $PM_{2.5}$ concentrations were relatively low, resulting in uncertainties regarding concentration-mortality functions at higher $PM_{2.5}$ concentration. Therefore, researchers integrated the relative risk (RR) values of diverse sources of $PM_{2.5}$ such as the second-hand tobacco smoking, active tobacco smoking, and the household air pollution from burning solid fuels to estimate the concentration-mortality functions in higher $PM_{2.5}$

concentration. The IER functions have been widely used by the WHO, GBD groups, and other researchers for estimating the health burden attributable to PM_{2.5}.^{2,3,14}

Based on previous research article which provided parameters for the calculated IER functions, we calculated the mean values of RR and their 95% confidence intervals for PM_{2.5} concentrations ranging from 0–150 µg/m³ in 0.1 µg/m³ steps.¹³ The lower and upper bounds of counterfactual concentration with theoretical minimum risk, which indicates no health benefits for reducing PM_{2.5} concentration below certain concentration, were defined as 5.8 and 8.8 µg/m³, respectively.

In previous global PM_{2.5} health impact assessment studies, the basic unit for the calculation of the health burden attributable to PM_{2.5} was 10 km × 10 km grid cell. They used the global spatially resolved estimates of annual average ambient PM_{2.5} concentrations in 10 km × 10 km resolutions, the global gridded population data within the same resolution, the regional level mortality, and demographic data to calculate the mortality impacts of PM_{2.5}. Detailed calculation equations were as follows:

$$M_{i,j} = P_i \times \hat{I}_{j,k} \times (RR_j(C_i) - 1)$$

$$\hat{I}_{j,k} = \frac{I_{j,k}}{RR_{j,k}}$$

$$\overline{RR}_{j,k} = \frac{\sum_{i=1}^N P_i \times RR_j(C_i)}{\sum_{i=1}^N P_i}$$

Where $M_{i,j}$ is premature mortality for grid cell i (for disease j) located in region k (consisting of N sub-grid cells), P_i is the population in grid cell i , $I_{j,k}$ is the average mortality rate for disease j in region k , $RR_j(C_i)$ is the IER RR value for the disease j at PM_{2.5} concentration of C (for grid cell i).¹⁴ To estimate the health benefits ($\Delta M_{i,j}$, change in mortality) of achieving the WHO PM_{2.5} air quality guideline values of 10 µg/m³, the following equation was used:

$$\Delta M_{i,j} = P_i \times \hat{I}_{j,k} \times \{RR_j(C_i^*) - RR_j(10)\}$$

This approach could be very useful for calculating and comparing country level PM_{2.5} health burden by using the global level estimates (population, PM_{2.5} concentration, mortality) in the same unit resolution (10 km × 10 km grid and regional level). However, up to date demographic and mortality data of each country are often not used for the analysis and sub-national level estimates for most of the countries were not available. In addition, 10 km × 10 km grid cell may not clearly reflect the regional differences in the age structure and the mortality rates in various metropolitan cities and provinces of a single country. Besides, the demographic and mortality data are organized in administrative units in most of the countries and thus have to be converted into 10 km × 10 km grid cell levels.

Therefore, we used the same equations but applied minor changes regarding the basic calculation units. Our basic analysis units were 'si,' 'gun,' and 'gu,' the basic administrative units for the metropolitan cities and provinces of Korea. For example, for the calculation of premature mortality in metropolitan cities, the above equation was modified as follows: where $M_{i,j}$ is premature mortality for 'gu' i (for disease j) located in metropolitan city k (consisting of N number of 'gu'), P_i is the population in 'gu' i , I_k is the average mortality

rate in metropolitan city *k*. We applied the same concept for calculating the premature mortality in major provinces by using the population and PM_{2.5} concentration data in 'si' and 'gun' levels. The 95% confidence intervals for the attributable mortality and the changes in mortality was estimated by applying 2.5% and 97.5% RR values of IER model for certain PM_{2.5} exposure values.

To account for the lag effect of PM_{2.5} exposure, we used annual levels of PM_{2.5} concentration for 2006 (the earliest year with modelled PM_{2.5} data) to calculate number of deaths attributable to ambient PM_{2.5} for 2015 as suggested in the previous article.¹⁵ In addition, we averaged PM_{2.5} concentration from year 2006 to 2015 and estimated the number of deaths attributable to ambient PM_{2.5} for 2015 as a sensitivity analysis. We additionally evaluated the annual temporal trends of the health burden from 2006 to 2015 in metropolitan cities and provinces of Korea by using yearly PM_{2.5} concentration, mortality, and population structure data. SAS version 9.4 (SAS Institute Inc., Cary, NC, USA) and R statistical software (version 3.1.3; R Foundation for Statistical Computing, Vienna, Austria) were used in our analysis.

Ethics statements

The Institutional Review Board of the Seoul National University Hospital, Korea, exempted this study from review because we used publicly available population level data from the Korean Statistical Information Service (IRB No. E-1710-004-889).

RESULTS

The annual average PM_{2.5} concentration and the population weighted PM_{2.5} concentration for 2015 in Korea were 24.4 µg/m³ and 25.1 µg/m³, respectively. **Fig. 1** shows the spatial variation in annual PM_{2.5} concentrations across major metropolitan cities and provinces of Korea. The sub-national regions with the highest PM_{2.5} concentrations (over 30 µg/m³) are located in Incheon metropolitan city (Dong-, Nam-, Namdong-, and Bupyeong-gu) and Gyeonggi (Bucheon-si), Chungcheongnam (Nonsan-si), and Jeollabuk (Jeonju-, Iksan-, Gimje- and Wanju-si) provinces (**Supplementary Table 1**).




Fig. 1. Mean concentration of PM_{2.5} and population density by basic administrative unit of the Korea in year 2015. **(A)** PM_{2.5} concentration and **(B)** population density. PM_{2.5} = particulate matter less than 2.5 µm in diameter.

The total 11,924 deaths were attributable to PM_{2.5} exposure in 2015 in Korea. Specifically, 3,303 deaths due to IHD, 637 deaths due to COPD, 2,338 deaths due to LC, and 5,646 deaths due to stroke were attributable to PM_{2.5} exposure in 2015 (Table 1). Fig. 2 and Table 1 shows the spatial variation of PM_{2.5} health burden across the sub-national regions. Particularly, deaths attributable to PM_{2.5} exposure were concentrated around Seoul (1,763 deaths), Busan (947 deaths), and Gyeonggi province (2,352 deaths). The number of attributable deaths due to PM_{2.5} according to basic administration level of metropolitan cities and province are summarized at Supplementary Table 1.

General association between annual mean PM_{2.5} concentration and age-standardized mortality rate due to PM_{2.5} exposure in Korea for year 2015 are described in Fig. 3. There was positive correlation between age-standardized mortality rate and annual mean PM_{2.5} concentration. After age-standardization (per 100,000 population), mortality rate attributable to PM_{2.5} exposure was high in Sejong, Daegu, and Incheon metropolitan cities and Jeollabuk and Chungcheongbuk provinces.

By simulating the reduction in the annual mean values of PM_{2.5} according to the WHO guideline value (10 µg/m³), about 8,539 deaths could be averted in Korea (Table 2). By conducting a 9-year time lag analysis with the annual PM_{2.5} concentration of 2006, the total of 13,856 deaths were attributable to PM_{2.5} exposure in 2015 (Supplementary Table 2). In the sensitivity analysis, with an average annual PM_{2.5} concentration from 2006 to 2015, the total 12,808 deaths were attributable to PM_{2.5} exposure in 2015 (Supplementary Table 3).

Most of the metropolitan cities and provinces in Korea showed decreasing or stable temporal trends of the mortality burden attributable to PM_{2.5} from 2006 to 2015 (Fig. 4, Supplementary Tables 4 and 5). Compared to 2006, Seoul, Busan, and Incheon metropolitan cities and Gyeonggi province showed marked decrease in number of attributable deaths due to PM_{2.5} in 2015. However, compared to 2014, Jeollanam and Jeollabuk provinces and Gwangju metropolitan city showed increase in number of attributable deaths due to PM_{2.5} in 2015.



Fig. 2. Spatial distribution of the health burden attributable to ambient PM_{2.5} exposure in basic administrative units of the Korea.
PM_{2.5} = particulate matter less than 2.5 µm in diameter.

Table 1. Health burden attributable to ambient PM_{2.5} exposure in major cities and provinces of the Republic of Korea

Cities and provinces	Population ^a	PM _{2.5} (µg/m ³) ^b	Number of deaths attributable to PM _{2.5}					Total	Age-standardized rate (per 100,000)
			Ischemic heart disease	Chronic obstructive pulmonary disease	Lung cancer	Stroke			
Metropolitan cities									
Seoul	7,189,442	24.2	487.7 (457.9–517.5)	60.9 (52.8–69)	367.4 (320.7–414.1)	847.4 (743.6–951.2)	1,763.4 (1,645.5–1,881.4)	19.3	
Busan	2,557,993	25.9	314.1 (287.9–340.2)	48.7 (40.5–56.9)	169.6 (142.1–197.2)	415.1 (349.2–480.9)	947.4 (871–1,023.9)	24.9	
Daegu	1,769,795	24.9	217.1 (192.5–241.8)	26.8 (20.3–33.3)	119.1 (91.4–146.7)	309 (238.6–379.4)	671.9 (592.1–751.8)	27.4	
Incheon	2,045,392	27.6	63.9 (54.8–73.1)	17.5 (12.6–22.4)	68.8 (50.1–87.4)	159.2 (117–201.4)	309.4 (262.2–356.7)	26.0	
Gwangju	1,028,181	24.1	185.2 (163–207.3)	24.2 (18.9–29.6)	131 (103.2–158.9)	316.3 (253.4–379.2)	656.8 (584.3–729.2)	25.8	
Daejeon	1,060,749	27.2	88.7 (76.5–100.9)	18.8 (13.2–24.4)	60.5 (43.1–77.9)	174.2 (125.8–222.6)	342.2 (289–395.3)	22.9	
Ulsan	809,987	24.6	56.6 (48.9–64.4)	15.3 (10.7–19.8)	38.4 (27.5–49.4)	112 (80.9–143)	222.3 (188.2–256.4)	25.9	
Sejong	134,303	27.7	11.5 (7.9–15.1)	3 (1.2–4.8)	10.2 (4.3–16.2)	24.5 (11–38)	49.2 (33.9–64.5)	28.4	
Provinces									
Gyeonggi	8,571,943	25.0	678.2 (644.9–711.4)	110.1 (98.7–121.6)	459.8 (414.3–505.2)	1,104.2 (998.6–1,209.8)	2,352.3 (2,232.1–2,472.6)	21.9	
Gangwon	1,104,304	20.6	119.2 (106.7–131.7)	29 (22.7–35.3)	82.1 (64.7–99.6)	212.3 (165.6–259)	442.7 (390.9–494.5)	22.5	
Chungcheongbuk	966,579	26.4	88.8 (80.3–97.4)	23.7 (19.2–28.2)	89.9 (73.5–106.4)	201 (165.3–236.6)	403.4 (363–443.9)	24.8	
Chungcheongnam	1,471,674	25.8	144.4 (132.2–156.6)	41.6 (34.7–48.5)	125.3 (105.2–145.3)	266.6 (224.9–308.4)	577.9 (529.5–626.2)	22.3	
Jeollabuk	1,313,542	29.1	130.1 (116.1–144.1)	37.4 (30.8–44)	130.1 (107.5–152.8)	340.5 (286.6–394.5)	638.2 (577.7–698.8)	26.1	
Jeollanam	1,323,030	22.4	181 (167.6–194.5)	44.9 (37.8–52)	121.1 (102.5–139.7)	286.5 (244.4–328.6)	633.6 (585.1–682.1)	22.3	
Gyeongsangbuk	1,960,926	21.6	248.5 (230.4–266.6)	68.9 (58.1–79.6)	167.2 (141.7–192.7)	400.7 (340.4–460.9)	885.3 (816.6–954)	24.4	
Gyeongsangnam	2,364,910	25.0	268.9 (249–288.9)	62.5 (53.3–71.8)	182.2 (155.8–208.7)	449.6 (385.5–513.8)	963.4 (890.6–1,036.2)	25.9	
Jeju	418,612	13.6	18.5 (13.2–23.9)	3.6 (1.2–5.9)	15.1 (5.8–24.3)	27.2 (13.3–41.2)	64.4 (46.6–82.1)	10	
Republic of Korea	36,091,362	24.4	3,302.6 (3,227.9–3,377.4)	636.8 (608.2–665.5)	2,338 (2,234.9–2,441)	5,646.3 (5,404.9–5,887.7)	11,923.7 (11,649.3–12,198.1)	22.9	

PM_{2.5} = particulate matter less than 2.5 µm in diameter.

^aPopulation census data of year 2015 (age over 25); ^bAnnual mean PM_{2.5} concentration at year 2015.




Fig. 3. Association between annual mean PM_{2.5} concentration and age-standardized mortality rate due to PM_{2.5} exposure in Korea for year 2015. PM_{2.5} = particulate matter less than 2.5 µm in diameter.

Table 2. Health benefits for meeting WHO PM_{2.5} air quality guideline in major cities and provinces of the Republic of Korea

Cities and provinces	PM _{2.5} (µg/m ³) ^a	Number of deaths averted by meeting WHO PM _{2.5} guideline (10 µg/m ³)				
		Ischemic heart disease	Chronic obstructive pulmonary disease	Lung cancer	Stroke	Total
Metropolitan cities						
Seoul	24.2	253.3 (220.8–285.8)	43.1 (40.3–46)	271.9 (253.9–289.9)	684.2 (653.9–714.4)	1,252.5 (1,204.5–1,300.5)
Busan	25.9	168.9 (142.6–195.3)	35.4 (32.7–38.2)	128.7 (118.6–138.9)	342.3 (324.5–360.1)	675.5 (642–708.9)
Daegu	24.9	114.3 (87.8–140.7)	19.2 (17–21.4)	89.1 (78.6–99.5)	252.1 (232.6–271.7)	474.7 (440–509.3)
Incheon	27.6	35.3 (26.4–44.1)	13 (11.3–14.6)	53 (46.4–59.7)	133.1 (122.2–144)	234.3 (218.6–249.9)
Gwangju	24.1	104.4 (84.4–124.3)	18.3 (16.6–19.9)	102.7 (93.2–112.2)	266.7 (250.3–283)	492 (464.5–519.5)
Daejeon	27.2	45.9 (32.5–59.3)	13.3 (11.3–15.3)	44.7 (37.9–51.4)	140.7 (126.6–154.8)	244.6 (223.8–265.3)
Ulsan	24.6	29.7 (21.3–38.1)	10.9 (9.3–12.5)	28.7 (24.5–32.8)	91 (82.3–99.7)	160.3 (147.4–173.2)
Sejong	27.7	6.3 (2.9–9.8)	2.2 (1.6–2.8)	7.9 (5.8–10)	20.4 (16.8–24)	36.9 (31.4–42.3)
Provinces						
Gyeonggi	25.0	358.2 (323.2–393.1)	79.1 (75.2–83)	344.5 (327.3–361.6)	901.5 (871.7–931.2)	1,683.2 (1,634–1,732.3)
Gangwon	20.6	58.3 (45.6–70.9)	19.7 (17.5–21.9)	58.4 (51.8–65.1)	166.5 (153.6–179.4)	302.8 (283.4–322.3)
Chungcheongbuk	26.4	48.4 (40–56.7)	17.4 (15.9–18.9)	68.8 (62.9–74.7)	166.9 (157.5–176.4)	301.5 (287.5–315.6)
Chungcheongnam	25.8	77.1 (65.2–89)	30.1 (27.8–32.5)	94.8 (87.4–102.1)	219.6 (208.3–230.9)	421.6 (403.4–439.8)
Jeollabuk	29.1	74.5 (63.1–85.9)	28.5 (26.5–30.5)	103.2 (95.7–110.7)	289.6 (275.6–303.7)	495.8 (476.1–515.5)
Jeollanam	22.4	89.6 (74.7–104.5)	30.7 (28.1–33.3)	86.8 (79.3–94.4)	226.3 (213.1–239.5)	433.4 (412–454.9)
Gyeongsangbuk	21.6	123.9 (104.2–143.5)	47.3 (43.5–51.2)	120.4 (110.3–130.4)	317.4 (299.7–335.2)	609 (580.4–637.6)
Gyeongsangnam	25.0	142.2 (122.3–162)	45 (41.9–48.1)	137 (127.2–146.7)	368.4 (351.1–385.7)	692.5 (664.3–720.8)
Jeju	13.6	5.4 (–1.1–12)	1.6 (0.4–2.7)	7.1 (1.6–12.5)	14.1 (5.3–23)	28.2 (15.9–40.6)
Republic of Korea	24.4	1,735.6 (1,658.4–1,812.8)	454.8 (445–464.7)	1,747.5 (1,708.7–1,786.3)	4,600.9 (4,532.9–4,668.9)	8,538.8 (8,428.4–8,649.2)

WHO = World Health Organization, PM_{2.5} = particulate matter less than 2.5 µm in diameter.

^aAnnual mean PM_{2.5} concentration at year 2015.




Fig. 4. Temporal trends of the health burden attributable to ambient $PM_{2.5}$ exposure in the Korea. **(A)** Metropolitan cities and **(B)** provinces. $PM_{2.5}$ = particulate matter less than 2.5 μm in diameter.

DISCUSSION

By using $PM_{2.5}$ modeling data, IER functions, and up to date publicly available population and mortality data, we evaluated the health burden attributable to $PM_{2.5}$ in the metropolitan cities and provinces of Korea. The total 11,924 deaths were attributable to $PM_{2.5}$ in Korea in 2015. There was also decreasing trends of the health burden in most of the cities and provinces when compared to 2006. As expected, deaths burden attributable to $PM_{2.5}$ was elevated in the basic administrative units where $PM_{2.5}$ concentration and population density are high.

After age-standardization (per 100,000 population), mortality rate attributable to $PM_{2.5}$ was high in Sejong, Daegu, and Incheon metropolitan cities and Jeollabuk and Chungcheongbuk provinces. Although number of deaths attributable to $PM_{2.5}$ was highest in Seoul metropolitan city and Gyeonggi province, age-standardized rate become relatively low in these two sub-regions. We believe this results comes from high population density of Seoul and Gyeonggi, but relatively low age and disease specific mortality rates compared to other sub-regions of Korea.

According to 2016 WHO report, a total of 11,523 deaths and 253,512 DALYs were attributable to PM_{2.5} exposure in Korea for 2012.² These results are comparable to our estimation of 11,872 deaths for 2012. In the recent GBD 2015 study, premature deaths attributable to PM_{2.5} exposure in Korea in 2005, 2010, and 2015, were 13,094, 13,777, and 18,148 deaths, showing increasing trends in the three most recent five-yearly estimates.⁸ However, in our study, the number of deaths attributable to PM_{2.5} generally decreased since 2006 in Korea, even though there were increasing patterns in 2013 and 2014. This discrepancy might have resulted from the use of different PM_{2.5} exposure data, mortality data, and IER functions between this study and that used in the 2015 GBD study. For example, although we used publicly available IER functions of GBD 2010 and 2013 in our study, there was a recent update of the functions with changes in the theoretical minimum risk exposure levels (2.4–5.9 µg/m³) and the model fitting algorithm, which resulted in a higher estimation of RR values compared to the previous GBD studies.³ Therefore, there are possibilities of underestimation in our study compared to recent 2015 GBD study. However, our study aim was to assess the regional and temporal variations in mortality burden due to PM_{2.5} exposure in Korea and the above changes of RR values may not affect our trend analysis results seriously.

Therefore, we believe that the major discrepancy between our trend analysis and those of the GBD 2015 results may come from the use of different PM_{2.5} exposure data. In the GBD 2015 study, annual mean population weighted exposure levels of PM_{2.5} increased from 25.18 µg/m³ (2010) to 28.66 µg/m³ (2015) in Korea.⁸ In our modeling data, the annual mean concentration and the population weighted PM_{2.5} levels were 26.85 µg/m³ and 28.74 µg/m³ for 2010, and 24.47 µg/m³ and 25.13 µg/m³ for 2015, respectively. Because the annual PM_{2.5} exposure data of 2015 used in our study shows about 3 µg/m³ lower levels compared to PM_{2.5} exposure data used in GBD studies, this may result in lower values of number of attributable deaths for year 2015. However, recent country wide monitoring data shows similar results to our PM_{2.5} exposure modelling data. The annual report of air quality in Korea indicated decreasing trends of annual PM₁₀ concentration from 2006 to 2015.^{6,7} In addition, the only available nation-wide PM_{2.5} ground monitoring data of 2015 (Seoul: 23, Busan: 26, Daegu: 26, Incheon: 29, Gwangju: 26, Daejeon: 28, Ulsan: 25, all measured in µg/m³) showed comparable results to that of the PM_{2.5} modeling data used in our study (Seoul: 24.2, Busan: 25.9, Daegu: 24.9, Incheon: 27.6, Gwangju: 24.1, Daejeon: 27.2, Ulsan: 24.6 all measured in µg/m³). Therefore, we believe our modeling data is reasonable for the health impact assessment study of PM_{2.5} in the sub-national levels and our study results may contribute additional spatial and temporal information to the global level estimations. However, long term (from 2004 to 2013) particulate matter measurement data from one monitoring site in Seoul showed complicated annual PM_{2.5} concentration patterns, despite their overall decreasing pattern.¹⁶ Therefore, we believe our PM_{2.5} modeling data should be further validated using satellite data and in the future, by a nation-wide ground monitoring data of PM_{2.5}.

There are few studies which evaluated the health impacts of particulate matter in Korea. By using modelled PM_{2.5} data, a total number of 15,346 (4,498–26,242) deaths were estimated to be attributable to PM_{2.5} exposure in Seoul, Incheon and Gyeonggi provinces in 2010.¹⁷ Another city level health impact assessment study estimated that about 12,000 deaths can be averted by reducing PM_{2.5} concentrations in the seven metropolitan cities of Korea according to the WHO guideline values of 10 µg/m³.¹⁸ However, these previous results could be overestimated due to the use of the linear exposure mortality functions and by incorporating diverse disease outcomes rather than the four specific disease outcomes (IHD, LC, COPD, and stroke) analyzed in our study and in GBD studies. Another study estimated number of

attributable deaths due to PM_{2.5} exposure by using exposure data of GBD 2013 and linear exposure mortality functions.¹⁹ They estimated that about 16,871 deaths were attributable to PM_{2.5} exposure in year 2013 by LC, IHD, and stroke.

In our study, deaths burden attributable to ambient PM_{2.5} exposure has generally decreased from 2006 to 2015. However, about 52% and 98% of the Korean population are still living in conditions with an annual concentration level of PM_{2.5} greater than the WHO interim Target 2 (25 µg/m³) and Target 3 (15 µg/m³) recommendations with our data. There were 43% and 79% of days in Korea, which exceeded WHO interim Target 2 and 3 recommendations in year 2015. Furthermore, nearly all of the total Korean population were living in conditions where annual concentration level of PM_{2.5} are above 10 µg/m³ and 92% of days were above 10 µg/m³ in 2015. By decreasing the annual PM_{2.5} concentrations to 10 µg/m³, about 8,500 deaths in 2015 can be averted. Therefore, despite the decreasing patterns of the PM_{2.5} attributable deaths in our analysis, there is great need for improvement regarding PM_{2.5} in Korea.

Significant decrease in the mortality burden in Seoul, Incheon, and Gyeonggi provinces, which are Seoul Metropolitan Areas (SMA), are worthy of note. In December 2003, the Special Act on the improvement of air quality in SMA was legislated.²⁰ Based on the Special Act, the first phase of the air quality improvement plan for the SMA (2005–2014) aimed to reduce the annual PM₁₀ and nitric oxide concentrations to 40 µg/m³ and 22 ppb respectively by regulating the total amount of the emissions in the workplace, supplying low-emission vehicles, and strengthening the gas emission management regulations. As a result, ground monitored annual PM₁₀ concentration in SMA showed decreasing patterns from 2005 to 2015.^{6,7} Decreasing patterns of the annual PM_{2.5} modeling data and attributable deaths in our study in SMA also indicate the effects of the Special Act in some aspects.

In December 2013, the Ministry of Environment, the Republic of Korea announced the implement of policy guidelines named the Comprehensive countermeasures against particulate matter.^{21,22} The main measures were, implementation of nationwide forecasting and warning system on particulate matter, enhancing international level cooperation with neighboring countries, and commencing the second phase (2015–2024) of the air quality improvement plan for the SMA. In June 2016, the government unveiled the emission reduction plan and Special Measures on Air Quality, to tackle particulate matter issues, with the intention to reduce the average annual concentration of PM_{2.5} to be comparable to that of European city in 2021 (20 µg/m³) and 2026 (18 µg/m³).²³ However, current governmental measures have not considered regional disparities and temporal trends of the health burden attributable to PM_{2.5} exposure. Therefore, we believe our study findings can enable politicians and local government officers to recognize the current sub-national states of the health burden and to estimate potential health benefits by reducing particulate matter concentration in sub-national levels.

Several conventional limitations of PM_{2.5} health-impact assessment studies are known, and particularly, issues regarding the use of IER functions was related to our study. First, IER functions only provide cause specific RR values for IHD, LC, COPD, and stroke for adults.¹³ Due to the increasing number of studies reporting the possible association between air pollution exposure and diverse disease outcomes, our study may underestimate the true health burden attributable to PM_{2.5} exposure. Second, because IER functions do not provide the source-specific RR, health impact assessment studies using IER functions rely on the concentration of the particulate matter. Therefore, the different components and sources of the PM_{2.5} at the sub-national regions were not considered in our analysis. However, evidence of different

toxicities from the sources and the components of particulate matters are still lacking.^{24,25} Third, the RR estimates of large-scale cohort studies used to model IER functions were based on studies conducted in western countries where the annual PM_{2.5} values are relatively low, compared to Korea. Although IER functions are believed to provide reasonable estimates of RR values at higher PM_{2.5} concentrations,³ large scale cohort studies evaluating PM_{2.5} exposure and mortality outcomes in regions with high levels of PM_{2.5} concentrations are needed to generate more precise estimates for the calculation of the health burden in polluted countries. Therefore, cohort studies evaluating the association between chronic PM_{2.5} exposure and mortality outcomes in Korea are needed in the future. Finally, for trend analysis of HIA, we used yearly PM_{2.5} concentration, demographic, and mortality data. Therefore, results of the trend analysis should be interpreted with caution due to the lag effects of particulate exposure on several chronic disease outcomes evaluated in this study. In addition, number of attributable deaths due to PM_{2.5} exposure depends on not only annual PM_{2.5} concentration, but also population structure and age and disease specific mortality data. Therefore, improvements on medical quality and system of Korea during 2006–2015 can also affect the trend analysis results.

In conclusion, about 11,924 premature deaths were attributable to PM_{2.5} exposure in 2015 in the Republic of Korea. Our findings show that further actions to improve air quality in Korea would substantially improve health burden. Due to the spatial variations attributable to PM_{2.5}, future particulate matter resolution policies should be reflective of the city and province level health-impact assessment results.

SUPPLEMENTARY MATERIALS

Supplementary Table 1

Health burden attributable to ambient PM_{2.5} exposure in basic administration units of the Republic of Korea

[Click here to view](#)

Supplementary Table 2

Health burden attributable to ambient PM_{2.5} exposure in major cities and provinces of the Republic of Korea (analysis with annual PM_{2.5} exposure of 2006)

[Click here to view](#)

Supplementary Table 3

Health burden attributable to ambient PM_{2.5} exposure in major cities and provinces of the Republic of Korea (analysis with annual average PM_{2.5} exposure from 2006 to 2015)

[Click here to view](#)

Supplementary Table 4

Temporal trends of the health burden attributable to ambient PM_{2.5} exposure in metropolitan cities of the Republic of Korea from 2006 to 2015

[Click here to view](#)

Supplementary Table 5

Temporal trends of the health burden attributable to ambient PM_{2.5} exposure in provinces of the Republic of Korea from 2006 to 2015

[Click here to view](#)

REFERENCES

1. Lim SS, Vos T, Flaxman AD, Danaei G, Shibuya K, Adair-Rohani H, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380(9859):2224–60. [PUBMED](#) | [CROSSREF](#)
2. World Health Organization. *Ambient Air Pollution: a Global Assessment of Exposure and Burden of Disease*. Geneva: World Health Organization; 2016.
3. Cohen AJ, Brauer M, Burnett R, Anderson HR, Frostad J, Estep K, et al. Estimates and 25-year trends of the global burden of disease attributable to ambient air pollution: an analysis of data from the Global Burden of Diseases Study 2015. *Lancet* 2017;389(10082):1907–18. [PUBMED](#) | [CROSSREF](#)
4. Han C, Lim YH, Jung K, Hong YC. Association between ambient particulate matter and disorders of vestibular function. *Environ Res* 2017;155:242–8. [PUBMED](#) | [CROSSREF](#)
5. Health Effect Institute. *State of Global Air 2017*. Boston, MA: Health Effect Institute; 2017.
6. Kim HC, Kim S, Kim BU, Jin CS, Hong S, Park R, et al. Recent increase of surface particulate matter concentrations in the Seoul metropolitan area, Korea. *Sci Rep* 2017;7(1):4710. [PUBMED](#) | [CROSSREF](#)
7. Hong YD. *Annual Report of Air Quality in Korea 2015*. Sejong: Ministry of Environment; 2016.
8. Global Burden of Disease Study. *Global Burden of Disease Study 2015 (GBD 2015) Results*. Seattle, WA: Institute for Health Metrics and Evaluation; 2016.
9. Kim BU, Bae C, Kim HC, Kim E, Kim S. Spatially and chemically resolved source apportionment analysis: case study of high particulate matter event. *Atmos Environ* 2017;162:55–70. [CROSSREF](#)
10. Kim HC, Kim E, Bae C, Cho JH, Kim BU, Kim S. Regional contributions to particulate matter concentration in the Seoul metropolitan area, South Korea: seasonal variation and sensitivity to meteorology and emissions inventory. *Atmos Chem Phys* 2017;17(17):10315–32. [CROSSREF](#)
11. Statistical database. http://kosiskr/statisticsList/statisticsList_01Listjsp?vwcd=MT_ZTITLE&parentId=A#SubCont. Accessed November 27, 2017.
12. MicroData integrated service website. https://mdiskostatgokr/extract/extSimpleSurvSearchdo?curMenuNo=UI_POR_P1200. Accessed November 1, 2017.
13. Burnett RT, Pope CA 3rd, Ezzati M, Olives C, Lim SS, Mehta S, et al. An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure. *Environ Health Perspect* 2014;122(4):397–403. [PUBMED](#)
14. Apte JS, Marshall JD, Cohen AJ, Brauer M. Addressing global mortality from ambient PM_{2.5}. *Environ Sci Technol* 2015;49(13):8057–66. [PUBMED](#) | [CROSSREF](#)
15. Lo WC, Shie RH, Chan CC, Lin HH. Burden of disease attributable to ambient fine particulate matter exposure in Taiwan. *J Formos Med Assoc* 2017;116(1):32–40. [PUBMED](#) | [CROSSREF](#)
16. Ahmed E, Kim KH, Shon ZH, Song SK. Long-term trend of airborne particulate matter in Seoul, Korea from 2004 to 2013. *Atmos Environ* 2015;101:125–33. [CROSSREF](#)
17. Leem JH, Kim ST, Kim HC. Public-health impact of outdoor air pollution for 2(nd) air pollution management policy in Seoul metropolitan area, Korea. *Ann Occup Environ Med* 2015;27(1):7. [PUBMED](#) | [CROSSREF](#)

18. Yorifuji T, Bae S, Kashima S, Tsuda T, Doi H, Honda Y, et al. Health impact assessment of PM10 and PM2.5 in 27 Southeast and East Asian cities. *J Occup Environ Med* 2015;57(7):751-6.
[PUBMED](#) | [CROSSREF](#)
19. Cheong HK. *Development of Standardized Methodology and Assessment of National Environmental Burden of Disease in Korea. Final Report for Environmental Health Action Program*. Sejong: Ministry of Environment; 2016.
20. Chae Y. Co-benefit analysis of an air quality management plan and greenhouse gas reduction strategies in the Seoul metropolitan area. *Environ Sci Policy* 2010;13(3):205-16.
[CROSSREF](#)
21. Ministry of Environment. *Comprehensive Measures against Fine Particles*. Sejong: Ministry of Environment; 2013.
22. Shim C. *Policy Measures for Mitigating Fine Particle Pollution in Korea and Suggestions for Expediting International Dialogue in East Asia*. Tokyo: JICA Research Institute; 2017.
23. Ministry of Environment. *Special Measures on Air Quality*. Sejong: Ministry of Environment; 2016.
24. Stanek LW, Sacks JD, Dutton SJ, Dubois JJ. Attributing health effects to apportioned components and sources of particulate matter: an evaluation of collective results. *Atmos Environ* 2011;45(32):5655-63.
[CROSSREF](#)
25. Lippmann M, Chen LC, Gordon T, Ito K, Thurston GD. National Particle Component Toxicity (NPACT) Initiative: integrated epidemiologic and toxicologic studies of the health effects of particulate matter components. *Res Rep Health Eff Inst* 2013;(177):5-13.
[PUBMED](#)