

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

1*H*-Benzotriazole–4-hydroxybenzoic acid (1/1)A. Thirunavukkarasu,^a A. Silambarasan,^a G. Chakkaravarthi,^{b*} R. Mohankumar^c and P. R. Umarani^{d*}^aDepartment of Physics, Presidency College, Chennai 600 005, India, ^bDepartment of Physics, CPCL Polytechnic College, Chennai 600 068, India, ^cDepartment of Physics, Presidency College, Chennai 600 005, India, and ^dKunthavai Naacchiyaar Govt. Arts College (W), Thanjavur 613 007, India

Correspondence e-mail: chakkaravarthi_2005@yahoo.com, kan_uma6@yahoo.com


Received 23 September 2013; accepted 29 September 2013

Key indicators: single-crystal X-ray study; *T* = 295 K; mean $\sigma(\text{C}-\text{C})$ = 0.003 Å; *R* factor = 0.031; *wR* factor = 0.074; data-to-parameter ratio = 12.6.

The asymmetric unit of the title compound, C₆H₅N₃·C₇H₆O₃, comprises independent benzotriazole and 4-hydroxybenzoic acid molecules. The dihedral angle between the benzene ring and the benzotriazole ring system is 15.18 (7)°. The mean plane of the carboxyl group is twisted at an angle of 18.55 (1)° with respect to the benzene ring. The crystal structure is stabilized by weak intermolecular N—H···N, O—H···N, O—H···O and C—H···O interactions, forming a three-dimensional network.

Related literature

For biological activities of benzotriazole derivatives, see: Dubey *et al.* (2011); Gaikwad, *et al.* (2012). For reported structures, see: Sieroń (2007); Sudhahar *et al.* (2013); Yang *et al.* (2010).



Experimental

Crystal data

C₆H₅N₃·C₇H₆O₃*M_r* = 257.25Orthorhombic, *Pna*2₁*a* = 17.3634 (13) Å*b* = 11.4669 (9) Å*c* = 6.0818 (4) Å*V* = 1210.91 (15) Å³*Z* = 4Mo *K*α radiation μ = 0.10 mm⁻¹*T* = 295 K

0.30 × 0.26 × 0.24 mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Absorption correction: multi-scan

(SADABS; Sheldrick, 1996)

T_{min} = 0.970, *T_{max}* = 0.976

6611 measured reflections

2195 independent reflections

1925 reflections with *I* > 2σ(*I*)*R_{int}* = 0.033

Refinement

 $R[F^2 > 2\sigma(F^2)]$ = 0.031 $wR(F^2)$ = 0.074*S* = 1.04

2195 reflections

174 parameters

1 restraint

H-atom parameters constrained

 $\Delta\rho_{\text{max}}$ = 0.15 e Å⁻³ $\Delta\rho_{\text{min}}$ = -0.11 e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
N1—H1···N2 ⁱ	0.86	2.20	2.982 (2)	152
O2—H2A···N3 ⁱⁱ	0.82	1.87	2.6817 (19)	169
O3—H3A···O1 ⁱⁱⁱ	0.82	1.88	2.6912 (17)	171
C13—H13···O3 ^{iv}	0.93	2.49	3.373 (2)	159

Symmetry codes: (i) $-x, -y + 2, z + \frac{1}{2}$; (ii) $x + \frac{1}{2}, -y + \frac{3}{2}, z + 1$; (iii) $-x + \frac{1}{2}, y + \frac{1}{2}, z - \frac{1}{2}$; (iv) $x, y, z + 1$.

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97*.

The authors thanks SAIF, IIT, Madras, for data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: VM2200).

References

- Bruker (2004). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Dubey, A., Srivastava, S. K. & Srivastava, S. D. (2011). *Bioorg. Med. Chem. Lett.* **21**, 569–573.
- Gaikwad, N. D., Patil, S. V. & Bodade, V. D. (2012). *Bioorg. Med. Chem. Lett.* **22**, 3449–3454.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Sieroń, L. (2007). *Acta Cryst.* **E63**, o2089–o2090.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Sudhahar, S., Krishnakumar, M., Sornamurthy, B. M., Chakkaravarthi, G. & Mohankumar, R. (2013). *Acta Cryst.* **E69**, o279.
- Yang, Y. X., Li, K., Wang, Y. J. & Li, Q. (2010). *Beijing Shifan Dax. Xue. Zir. Kex. (J. B. Norm. Univ.)*, **46**, 160–165.

supplementary materials

Acta Cryst. (2013). E69, o1605 [doi:10.1107/S1600536813026767]

1H-Benzotriazole–4-hydroxybenzoic acid (1/1)

A. Thirunavukkarasu, A. Silambarasan, G. Chakkaravarthi, R. Mohankumar and P. R. Umarani

1. Comment

Benzotriazole derivatives exhibit numerous essential bioactivities, especially in antitubercular and antimicrobial (Dubey *et al.*, 2011; Gaikwad *et al.*, 2012) activities. We herewith report the crystal structure of the title compound (I) (Fig.1). The geometric parameters are comparable with reported structures (Sieroń, 2007; Sudhakar *et al.*, 2013; Yang *et al.*, 2010).

The benzene ring (C1-C6) is planar, with the maximum deviation of 0.010 (2) Å. The dihedral angle between the benzene ring and benzotriazole ring system is 15.18 (7)°. The mean plane of carboxyl group is twisted at an angle of 18.55 (1)° with the benzene ring. The crystal structure is stabilized by weak intermolecular N—H···N, O—H···N, O—H···O, C—H···O (Table 1 & Fig. 2) interactions to form a three dimensional network.

2. Experimental


Benzotriazole (C₆H₅N₃, 1.1913 g) and p-hydroxy benzoic acid (C₇H₆O₃, 1.3812 g) were mixed in equimolar ratio in methanol and the prepared solution was allowed for slow evaporation at room temperature. Good quality crystals suitable for X-ray intensity data collection were collected in a period of 10 days.

3. Refinement


H atoms were positioned geometrically and refined using riding model with C-H = 0.93 Å and U_{iso}(H) = 1.2U_{eq}(C) for CH, N-H = 0.86 Å and U_{iso}(H) = 1.2U_{eq}(C) for NH, O-H = 0.82 Å and U_{iso}(H) = 1.5U_{eq}(C) for OH.

Computing details

Data collection: *APEX2* (Bruker, 2004); cell refinement: *SAINT* (Bruker, 2004); data reduction: *SAINT* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008).

**Figure 1**

The molecular structure of (I), with atom labels and 30% probability displacement ellipsoids for non-H atoms.

**Figure 2**

The packing of (I), viewed down *c* axis. Intermolecular hydrogen bonds are shown as dashed lines. H atoms not involved in hydrogen bonding have been omitted.

1H-Benzotriazole-4-hydroxybenzoic acid (1/1)

Crystal data

C₆H₅N₃·C₇H₆O₃

M_r = 257.25

Orthorhombic, *Pna*2₁

Hall symbol: P 2c -2n

a = 17.3634 (13) Å

b = 11.4669 (9) Å

c = 6.0818 (4) Å

V = 1210.91 (15) Å³

Z = 4

F(000) = 536

D_x = 1.411 Mg m⁻³

Mo *Kα* radiation, λ = 0.71073 Å

Cell parameters from 2578 reflections

θ = 2.1–25.2°

μ = 0.10 mm⁻¹

T = 295 K

Block, colourless

0.30 × 0.26 × 0.24 mm

Data collection

Bruker Kappa APEXII CCD

diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

ω and φ scan

Absorption correction: multi-scan

(*SADABS*; Sheldrick, 1996)

T_{min} = 0.970, *T_{max}* = 0.976

6611 measured reflections

2195 independent reflections

1925 reflections with *I* > 2σ(*I*)

R_{int} = 0.033

θ_{max} = 26.8°, θ_{min} = 2.1°

h = -20→22

k = -13→14

l = -7→4

Refinement

Refinement on *F*²

Least-squares matrix: full

R[*F*² > 2σ(*F*²)] = 0.031

wR(*F*²) = 0.074

S = 1.04

2195 reflections

174 parameters

1 restraint

Primary atom site location: structure-invariant

direct methods

Secondary atom site location: difference Fourier

map

Hydrogen site location: inferred from

neighbouring sites

H-atom parameters constrained

w = 1/[σ²(*F_o*²) + (0.0362*P*)² + 0.0577*P*]

where *P* = (*F_o*² + 2*F_c*²)/3

(Δ/σ)_{max} < 0.001

Δρ_{max} = 0.15 e Å⁻³

Δρ_{min} = -0.11 e Å⁻³

Extinction correction: *SHELXL97* (Sheldrick,

2008), *F_c** = *kF_c*[1 + 0.001*xF_c*²λ³/sin(2θ)]^{-1/4}

Extinction coefficient: 0.025 (2)

Special details

Geometry. All esds (except the esd in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell esds are taken into account individually in the estimation of esds in distances, angles and torsion angles; correlations between esds in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell esds is used for estimating esds involving l.s. planes.

Refinement. Refinement of *F*² against ALL reflections. The weighted *R*-factor *wR* and goodness of fit *S* are based on *F*², conventional *R*-factors *R* are based on *F*, with *F* set to zero for negative *F*². The threshold expression of *F*² > 2σ(*F*²) is used only for calculating *R*-factors(gt) etc. and is not relevant to the choice of reflections for refinement. *R*-factors based on *F*² are statistically about twice as large as those based on *F*, and *R*-factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	<i>U_{iso}</i> */ <i>U_{eq}</i>
C7	0.37669 (8)	0.74471 (13)	0.7594 (3)	0.0363 (4)
C4	0.32444 (9)	0.79794 (14)	0.5984 (3)	0.0353 (4)
C3	0.30244 (10)	0.73727 (15)	0.4112 (3)	0.0457 (4)

H3	0.3204	0.6617	0.3895	0.055*
C2	0.25482 (10)	0.78675 (16)	0.2581 (3)	0.0492 (5)
H2	0.2416	0.7452	0.1323	0.059*
C1	0.22618 (9)	0.89793 (15)	0.2887 (3)	0.0385 (4)
C6	0.24605 (9)	0.95891 (15)	0.4762 (3)	0.0417 (4)
H6	0.2261	1.0331	0.4997	0.050*
C5	0.29535 (10)	0.91033 (14)	0.6286 (3)	0.0419 (4)
H5	0.3093	0.9528	0.7527	0.050*
N2	-0.00946 (8)	0.88952 (13)	0.2498 (3)	0.0477 (4)
C9	0.03383 (9)	0.72630 (14)	0.3824 (3)	0.0345 (4)
C8	0.05616 (10)	0.81341 (14)	0.5280 (3)	0.0366 (4)
C13	0.09730 (10)	0.78968 (17)	0.7199 (3)	0.0485 (5)
H13	0.1117	0.8480	0.8178	0.058*
C12	0.11490 (10)	0.67520 (17)	0.7544 (4)	0.0539 (5)
H12	0.1419	0.6551	0.8809	0.065*
C11	0.09398 (12)	0.58653 (16)	0.6071 (3)	0.0521 (5)
H11	0.1083	0.5101	0.6375	0.062*
C10	0.05347 (10)	0.60939 (14)	0.4216 (3)	0.0459 (4)
H10	0.0393	0.5504	0.3248	0.055*
N3	-0.00650 (8)	0.77773 (12)	0.2146 (2)	0.0418 (3)
N1	0.02792 (8)	0.91193 (12)	0.4377 (3)	0.0456 (4)
H1	0.0332	0.9803	0.4935	0.055*
O1	0.38942 (6)	0.63969 (10)	0.7712 (2)	0.0472 (3)
O2	0.40973 (7)	0.81933 (10)	0.8925 (2)	0.0504 (3)
H2A	0.4380	0.7843	0.9779	0.076*
O3	0.17907 (8)	0.94246 (11)	0.13231 (19)	0.0531 (4)
H3A	0.1627	1.0060	0.1729	0.080*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
C7	0.0366 (8)	0.0325 (9)	0.0398 (9)	-0.0021 (6)	0.0031 (7)	-0.0020 (9)
C4	0.0346 (8)	0.0338 (8)	0.0376 (9)	-0.0022 (6)	0.0022 (7)	-0.0032 (7)
C3	0.0509 (10)	0.0377 (9)	0.0485 (11)	0.0083 (7)	-0.0051 (9)	-0.0119 (10)
C2	0.0575 (10)	0.0467 (10)	0.0434 (11)	0.0084 (8)	-0.0093 (10)	-0.0186 (9)
C1	0.0391 (8)	0.0405 (9)	0.0359 (9)	0.0005 (7)	-0.0001 (7)	-0.0029 (8)
C6	0.0480 (9)	0.0303 (8)	0.0468 (10)	0.0029 (7)	-0.0041 (8)	-0.0082 (7)
C5	0.0463 (9)	0.0375 (9)	0.0418 (10)	-0.0014 (7)	-0.0062 (8)	-0.0101 (8)
N2	0.0591 (9)	0.0351 (8)	0.0488 (9)	-0.0029 (6)	-0.0070 (8)	0.0017 (7)
C9	0.0379 (8)	0.0318 (8)	0.0336 (9)	-0.0030 (6)	-0.0001 (7)	-0.0033 (7)
C8	0.0397 (9)	0.0324 (9)	0.0376 (9)	-0.0030 (7)	0.0033 (7)	-0.0053 (7)
C13	0.0527 (10)	0.0538 (12)	0.0391 (10)	-0.0031 (8)	-0.0053 (8)	-0.0128 (10)
C12	0.0563 (11)	0.0610 (12)	0.0445 (11)	0.0052 (9)	-0.0124 (10)	0.0009 (11)
C11	0.0582 (11)	0.0405 (11)	0.0574 (12)	0.0063 (8)	-0.0091 (10)	0.0023 (10)
C10	0.0524 (10)	0.0326 (9)	0.0528 (12)	-0.0006 (7)	-0.0056 (10)	-0.0074 (9)
N3	0.0519 (8)	0.0330 (8)	0.0406 (8)	-0.0022 (6)	-0.0058 (7)	-0.0004 (7)
N1	0.0587 (9)	0.0295 (7)	0.0486 (9)	-0.0043 (6)	-0.0009 (8)	-0.0086 (7)
O1	0.0533 (7)	0.0323 (7)	0.0560 (8)	0.0004 (5)	-0.0065 (6)	-0.0008 (7)
O2	0.0630 (7)	0.0339 (7)	0.0542 (8)	0.0015 (5)	-0.0221 (7)	-0.0036 (6)
O3	0.0654 (8)	0.0509 (8)	0.0430 (7)	0.0145 (6)	-0.0147 (6)	-0.0089 (6)

Geometric parameters (Å, °)

C7—O1	1.2265 (18)	C9—N3	1.371 (2)
C7—O2	1.3102 (19)	C9—C8	1.390 (2)
C7—C4	1.468 (2)	C9—C10	1.404 (2)
C4—C3	1.388 (2)	C8—N1	1.348 (2)
C4—C5	1.396 (2)	C8—C13	1.395 (3)
C3—C2	1.369 (2)	C13—C12	1.364 (3)
C3—H3	0.9300	C13—H13	0.9300
C2—C1	1.381 (2)	C12—C11	1.403 (3)
C2—H2	0.9300	C12—H12	0.9300
C1—O3	1.3545 (19)	C11—C10	1.355 (3)
C1—C6	1.381 (2)	C11—H11	0.9300
C6—C5	1.380 (2)	C10—H10	0.9300
C6—H6	0.9300	N1—H1	0.8600
C5—H5	0.9300	O2—H2A	0.8200
N2—N3	1.3008 (19)	O3—H3A	0.8200
N2—N1	1.339 (2)		
O1—C7—O2	121.74 (15)	N3—C9—C10	131.39 (15)
O1—C7—C4	123.97 (15)	C8—C9—C10	120.69 (16)
O2—C7—C4	114.28 (13)	N1—C8—C9	103.95 (15)
C3—C4—C5	118.12 (15)	N1—C8—C13	133.65 (17)
C3—C4—C7	120.59 (15)	C9—C8—C13	122.39 (16)
C5—C4—C7	121.30 (15)	C12—C13—C8	115.53 (17)
C2—C3—C4	121.11 (16)	C12—C13—H13	122.2
C2—C3—H3	119.4	C8—C13—H13	122.2
C4—C3—H3	119.4	C13—C12—C11	122.75 (19)
C3—C2—C1	120.55 (16)	C13—C12—H12	118.6
C3—C2—H2	119.7	C11—C12—H12	118.6
C1—C2—H2	119.7	C10—C11—C12	121.73 (17)
O3—C1—C2	118.07 (15)	C10—C11—H11	119.1
O3—C1—C6	122.67 (15)	C12—C11—H11	119.1
C2—C1—C6	119.26 (16)	C11—C10—C9	116.89 (16)
C5—C6—C1	120.36 (15)	C11—C10—H10	121.6
C5—C6—H6	119.8	C9—C10—H10	121.6
C1—C6—H6	119.8	N2—N3—C9	108.75 (14)
C6—C5—C4	120.56 (15)	N2—N1—C8	111.29 (14)
C6—C5—H5	119.7	N2—N1—H1	124.4
C4—C5—H5	119.7	C8—N1—H1	124.4
N3—N2—N1	108.09 (14)	C7—O2—H2A	109.5
N3—C9—C8	107.91 (14)	C1—O3—H3A	109.5
O1—C7—C4—C3	-18.0 (2)	N3—C9—C8—C13	178.84 (16)
O2—C7—C4—C3	161.82 (15)	C10—C9—C8—C13	-1.3 (3)
O1—C7—C4—C5	161.86 (16)	N1—C8—C13—C12	179.32 (19)
O2—C7—C4—C5	-18.3 (2)	C9—C8—C13—C12	0.6 (3)
C5—C4—C3—C2	1.3 (3)	C8—C13—C12—C11	0.6 (3)
C7—C4—C3—C2	-178.86 (17)	C13—C12—C11—C10	-1.2 (3)
C4—C3—C2—C1	-1.3 (3)	C12—C11—C10—C9	0.5 (3)

C3—C2—C1—O3	-179.75 (16)	N3—C9—C10—C11	-179.47 (17)
C3—C2—C1—C6	-0.1 (3)	C8—C9—C10—C11	0.6 (2)
O3—C1—C6—C5	-178.87 (16)	N1—N2—N3—C9	-0.20 (18)
C2—C1—C6—C5	1.5 (3)	C8—C9—N3—N2	0.25 (18)
C1—C6—C5—C4	-1.5 (3)	C10—C9—N3—N2	-179.65 (18)
C3—C4—C5—C6	0.1 (2)	N3—N2—N1—C8	0.09 (19)
C7—C4—C5—C6	-179.74 (16)	C9—C8—N1—N2	0.06 (18)
N3—C9—C8—N1	-0.18 (17)	C13—C8—N1—N2	-178.79 (18)
C10—C9—C8—N1	179.72 (16)		

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
N1—H1 \cdots N2 ⁱ	0.86	2.20	2.982 (2)	152
O2—H2A \cdots N3 ⁱⁱ	0.82	1.87	2.6817 (19)	169
O3—H3A \cdots O1 ⁱⁱⁱ	0.82	1.88	2.6912 (17)	171
C13—H13 \cdots O3 ^{iv}	0.93	2.49	3.373 (2)	159

Symmetry codes: (i) $-x, -y+2, z+1/2$; (ii) $x+1/2, -y+3/2, z+1$; (iii) $-x+1/2, y+1/2, z-1/2$; (iv) $x, y, z+1$.