

# Postrenal acute kidney injury from a sliding inguinal hernia


Rosemarie Vincent MD, Shannon A. Fraser MD MSc, Evan G. Wong MD MPH

■ Cite as: *CMAJ* 2022 August 22;194:E1112. doi: 10.1503/cmaj.211805

A 92-year-old man with chronic kidney disease (baseline creatinine 120 µmol/L) developed acute kidney injury (highest creatinine 422 µmol/L) while hospitalized following his third myocardial infarction. He had no pain or lower urinary tract symptoms, but had noticed a bulge in his left inguinal area for the previous 2 years. Physical examination showed that the patient was frail, with a large, partially reducible, left inguinal mass. His acute kidney injury did not resolve after intravenous hydration. We ordered pelvic ultrasonography, which showed moderate-to-severe bilateral hydronephrosis and a poorly visualized left testicle. We inserted a Foley catheter and consulted a urologist. We ordered computed tomography to look for intrinsic and extrinsic causes of the hydronephrosis, which showed that most of the bladder had herniated through a left inguinal hernia. It also showed bilateral hydronephrosis and hydroureter (Figure 1).

We attempted a mechanical reduction of the bladder hernia, which was not successful. We offered the patient and his family surgical repair or more conservative management. Because of his high surgical risk, they chose nonoperative management. We inserted bilateral nephrostomy tubes, and the acute kidney injury improved. We subsequently removed the tubes and discharged the patient with an indwelling Foley catheter. He did not have further complications related to his hernia or his urinary system during the 6 months after discharge.

Bladder hernias are uncommon, accounting for 1%–4% of inguinal hernias.<sup>1</sup> Only 7%–10% are diagnosed preoperatively.<sup>1,2</sup> Inguinal bladder hernias presenting as obstructive uropathy are infrequent, but should be considered in patients with an inguinal mass and concomitant acute kidney injury based on history and physical examination.<sup>2</sup> The diagnosis can be confirmed with ultrasonography, computed tomography or cystography.<sup>1</sup> Patients who are at low-to-moderate operative risk are usually offered surgery, but decompression with transient nephrostomy tubes and a permanent bladder catheter is an alternative for patients who have a high operative risk.



**Figure 1:** Computed tomography scans of the abdomen and pelvis of a 92-year-old man with acute kidney injury. (A) Sagittal view shows most of the bladder herniating through a left inguinal hernia (white arrow). Foley catheter is visible within the small intra-abdominal portion of the bladder (red arrow). (B) Axial view shows bilateral hydronephrosis (green arrows) and hydroureter (blue arrow).

## References

1. Elkbuli A, Narvel RI, McKenney M, et al. Inguinal bladder hernia: a case report and literature review. *Int J Surg Case Rep* 2019;58:208-11.
2. Branchu B, Renard Y, Larre S, et al. Diagnosis and treatment of inguinal hernia of the bladder: a systematic review of the past 10 years. *Turk J Urol* 2018;44:384-8.

**Competing interests:** Rosemarie Vincent reports grants from McGill University, outside the submitted work. No other competing interests were declared.

This article has been peer reviewed.

The authors have obtained patient consent.

**Affiliations:** Faculty of Medicine (Vincent); Division of General Surgery (Fraser, Wong), Department of Surgery, McGill University, Montréal, Que.

**Content licence:** This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY-NC-ND 4.0) licence, which permits use, distribution and reproduction in any medium, provided that the original publication is properly cited, the use is noncommercial (i.e., research or educational use), and no modifications or adaptations are made. See: <https://creativecommons.org/licenses/by-nc-nd/4.0/>

**Correspondence to:** Evan Wong, [evan.wong@mcgill.ca](mailto:evan.wong@mcgill.ca)