

RESEARCH

Open Access



An explanatory model of depressive symptoms from anxiety, post-traumatic stress, somatic symptoms, and symptom perception: the potential role of inflammatory markers in hospitalized COVID-19 patients

David Villarreal-Zegarra^{1,2} , Rubí Paredes-Angeles² , Nikol Mayo-Puchoc² , Ana L. Vilela-Estada² , Anthony Copez-Lonzoy^{3,2,4} and Jeff Huarcaya-Victoria^{5,6*}

Abstract

Background: The context of the COVID-19 pandemic has harmed the mental health of the population, increasing the incidence of mental health problems such as depression, especially in those who have had COVID-19. Our study puts forward an explanatory model of depressive symptoms based on subjective psychological factors in those hospitalized for COVID-19 with and without biological markers (i.e., inflammatory markers). Therefore, we aim to evaluate the hypotheses proposed in the model to predict the presence of depressive symptoms.

Method: We conducted a cross-sectional study, using a simple random sampling. Data from 277 hospitalized patients with COVID-19 in Lima-Peru, were collected to assess mental health variables (i.e., depressive, anxiety, post-traumatic stress, and somatic symptoms), self-perception of COVID-19 related symptoms, and neutrophil/lymphocyte ratio (NLR) such as inflammatory marker. We performed a structural equation modeling analysis to evaluate a predictive model of depressive symptoms.

Results: The results showed a prevalence of depressive symptoms (11.2%), anxiety symptoms (7.9%), somatic symptoms (2.2%), and symptoms of post-traumatic stress (6.1%) in the overall sample. No association was found between the prevalence of these mental health problems among individuals with and without severe inflammatory response. The mental health indicators with the highest prevalence were sleep problems (48%), low energy (47.7%), nervousness (48.77%), worry (47.7%), irritability (43.7%) and back pain (52%) in the overall sample. The model proposed to explain depressive symptoms was able to explain more than 83.7% of the variance and presented good goodness-of-fit indices. Also, a different performance between the proposed model was found between those with and without severe inflammatory response. This difference was mainly found in the relationship between anxiety and post-traumatic stress symptoms, and between the perception of COVID-19 related symptoms and somatic symptoms.

*Correspondence: jeff.huarcaya@upsjb.edu.pe

⁶ Departamento de Psiquiatría, Servicio de Psiquiatría de Adultos, Unidad de Psiquiatría de Enlace, Hospital Nacional Guillermo Almenara Irigoyen, Lima, Perú

Full list of author information is available at the end of the article



Conclusions: Results demonstrated that our model of mental health variables may explain depressive symptoms in hospitalized patients of COVID-19 from a third-level hospital in Peru. In the model, perception of symptoms influences somatic symptoms, which impact both anxiety symptoms and symptoms of post-traumatic stress. Thus, anxiety symptoms could directly influence depressive symptoms or through symptoms of post-traumatic stress. Our findings could be useful to decision-makers for the prevention of depression, used to inform the creation of screening tools (i.e., perception of symptoms, somatic and anxiety symptoms) to identify vulnerable patients to depression.

Keywords: COVID-19, Inflammation, Depression, Anxiety, Post-traumatic stress, Peru

Background

Several studies have reported that COVID-19 patients had experienced various mental health problems (i.e., depression, anxiety, and post-traumatic symptoms) [1–3]. Systematic reviews have identified a high prevalence of depressive symptoms (52%), anxiety symptoms (47%) [4], and symptoms of post-traumatic stress (26.9%) [5] as a result of COVID-19. The evidence suggests that individuals who contracted COVID-19 suffered a negative impact on their mental health, however, this impact was greater in individuals who were hospitalized for COVID-19 [6]. In this way, patients who were hospitalized had a greater negative impact on their mental health due to various clinical factors such as demographics (i.e. sex, age, and proceeding outside of the capital), clinical (i.e. self-perception of the severity of COVID-19, the persistence of COVID-19 symptoms, a history of psychiatric treatment, and history of a family member infected by COVID-19), immune factors (i.e. neutrophil–lymphocyte index greater of 6.5), and psychosocial characteristics (i.e. isolation or quarantine, fear of COVID-19, being discriminate because of COVID) [7–10].

There are two mechanisms of action that may explain the presence of mental health problems in hospitalized patients with COVID-19: biological and psychological responses. Although the neuropsychiatric complications of COVID-19 are under study, there is evidence that inflammatory markers may cause mental health problems, such as depression. An invasion of SARS-CoV-2 to the respiratory tract could induce an acute respiratory syndrome with consequent release of proinflammatory cytokines such as IL-1 β and IL-6. Consequently, a systemic immune response in the form of a “cytokine storm” is produced [11]. Moreover, studies have reported these cytokines have increased in various psychiatric disorders (i.e., schizophrenia, depression, and post-traumatic stress) [12]. The relationship between elevated cytokine levels in COVID-19 and mental health problems could indicate that immune/inflammatory pathways are one of the possible mechanisms involved in mental health problems in this infection [12].

The neutrophil–lymphocyte ratio (NLR) is an inexpensive marker calculated through a complete blood count. Its pathogenic role has been studied in a wide variety of diseases [13–16]. Thus, elevated NLR has been related to an increase of cytokines and C-reactive protein (CRP). Elevated NLR levels are also associated with a state of chronic inflammation. Recent meta-analyses have documented the relevance of NLR in psychiatric diseases such as schizophrenia [14], and mood disorders [17]. If the measurement of cytokine performance is not possible, we can indirectly assess an increase in cytokines through an elevation of NLR.

From a psychological point of view, patients hospitalized with COVID-19 experience physical discomfort related to the COVID-19 symptoms themselves and other somatic symptoms, which can lead to stress-induced mental health problems [18, 19]. As a novel and life-threatening disease, COVID-19 can cause fear and stress in patients, especially for those being treated in the isolation ward. Also, the uncertainty regarding the consequences of the infection during the hospitalization may intensify patients’ experiences of panic [20]. Systematic reviews with meta-analysis evidence that the isolation, physical discomfort, and adverse effects of treatment may increase sensitivity among patients around symptoms of the infection, which could lead to worsening of mental health [18, 21–23]. Additionally, stress and anxiety could cause depressive symptoms [24].

There are studies that separately evaluate the relationship between mental health in hospitalized COVID-19 patients with psychological factors and biological markers. However, few studies investigate the impact of both factors on mental health problems (i.e., depression) in COVID-19 patients. Therefore, we conducted a study to explain the presence of one of the most prevalent mental health problems (i.e., depressive symptoms) from subjective psychological factors (i.e., somatic symptoms, anxiety symptoms, and symptoms of post-traumatic stress) in individuals hospitalized for COVID-19 with and without biological markers (i.e., inflammatory markers) (see Fig. 1).




Fig. 1 Model tested using structural equation modeling (SEM)

Perception of the severity of COVID-19 symptoms influences somatic symptoms (hypothesis 1) (see Fig. 1)

Patients' self-perception of the severity of illness (i.e., reducing or increasing symptoms) is related to the severity of mental health problems. A possible explanation is that concerns about their illness or condition add to their psychological burden [25]. Furthermore, the perception of symptoms related to COVID-19 (i.e., fever, cough, trouble breathing) are closely connected to somatic symptoms (i.e., headache, feeling tired, etc.). Therefore, it is also related to other mental health problems.

Somatic symptoms influence anxiety symptoms (hypothesis 2) and symptoms of post-traumatic stress (hypothesis 3) in patients hospitalized for COVID-19 (see Fig. 1)

It has been evidenced that the prevalence of somatic symptoms is significantly related to psychological outcomes (i.e., anxiety and post-traumatic stress). Evidence shows a high prevalence of moderate or severe anxiety during the COVID-19 pandemic among the general public. In many cases, a common anxiety-induced comorbidity was somatization [26]. In contrast, PTSD is a severe psychological consequence when a person experiences a stressful event as highly traumatic [27]. Indeed, longitudinal studies about PTSD in trauma survivors reported that symptoms of post-traumatic stress establish a more consistent relationship with somatic symptoms over time [28, 29].

Anxiety symptoms influence post-traumatic stress symptoms (hypothesis 4) in patients hospitalized for COVID-19 (see Fig. 1)

There is much evidence to support the triad of fear, anxiety, and stress. This triad is a fear-induced sequence of responses (being hospitalized for COVID-19) that leads to an anxiety response. This in turn leads to symptoms

of post-traumatic stress. Firstly, fear may increase sympathetic nervous system arousal and induce defensive or escape behavior in the face for specific and real threatening stimulus. Anxiety is like fear as an emotional reaction, but unlike fear, the source of threat is unclear. Thus, it is associated with preventive behaviors such as avoidance [30]. Moreover, fear and anxiety are related to the amygdala, which recruits and expresses the memory of these emotions in both animals and humans [31].

The fear caused by COVID-19 can be implicated in mental health problems (i.e., insomnia, increased alcohol and tobacco use, anxiety, among others), such as high infection and death rates, strict public health measures, etc. [7]. In many cases, the excessive exposure to anxiety behaviors triggers post-traumatic stress disorder (PTSD) [27].

Symptoms of post-traumatic stress and anxiety symptoms influence the presence of depressive symptoms (hypothesis 5 and 6) in patients hospitalized for COVID-19 (see Fig. 1)

The evidence on the relationship between anxiety, depression, and post-traumatic stress is abundant. Studies have identified that anxiety and fear of being hospitalized for COVID-19 can generate a state of acute stress in individuals [32, 33]. Acute stress and symptoms of post-traumatic stress often trigger different mental health problems such as depression in hospitalized patients [20, 34]. Therefore, it is hypothesized that PTSD symptoms precede depressive symptoms (hypothesis 6: symptoms of post-traumatic stress influences depressive symptoms). On the other hand, there is ample evidence that both anxiety and depression are closely related [35, 36] especially in COVID-19 pandemic [37]. Previous studies confirmed this hypothesis, where anxiety, post-traumatic stress, and depression were closely related to each other. Also, anxiety had the greatest influence on the prevalence of depressive

symptoms [24, 38] (hypothesis 5: anxiety symptoms influence depressive symptoms).

While studies about mental health in COVID-19 are in ascendant progress, there is a lack of clarity about the functioning of these variables and their subsequent impact on mental health. Therefore, the present study proposes to evaluate these hypotheses to predict the presence of depressive symptoms from subjective psychological factors (i.e., somatic symptoms, anxiety symptoms, symptoms of post-traumatic stress) in those hospitalized for COVID-19 with and without biological markers (i.e., inflammatory markers).

Methods

Study design

The study design was a cross-sectional investigation.

Participants

We used secondary data from Mental Health in COVID-2019 Survivors from a General Hospital in Peru: Sociodemographic, Clinical, and Inflammatory Variable Associations [39]. Participants were individuals with COVID-19 who were discharged from the “Hospital Nacional Guillermo Almenara Irigoyen” between March and September 2020 in Lima, Peru. Inclusion criteria included the following: 18 years or older, having been assessed at their admission and release from the hospital. Participants were excluded as follows: individuals who had missing data in the variables of interest (i.e., anxiety symptoms, depressive symptoms, somatic symptoms, post-traumatic stress symptoms, and NLR) and demographic variables (i.e., age, sex, civil status, degree of education, employment status, partners, relatives with COVID-19).

We calculated sample size with the Epidat v44.2 program (Dirección Xeral de Saúde Pública da Consellería de Sanidade, Galicia, España) and the sample was selected by a simple random sampling from a total number of 1190 participants.

Setting

The data of this second study were collected for the HNGAI from September to November of 2020. “Hospital Nacional Guillermo Almenara Irigoyen” is classified as a third-level specialized health institute in 2015, and it is the second-largest hospital in the “Seguridad Social de Salud del Perú” (ESSALUD).

During the COVID-19 pandemic, the health system was focused on the care of COVID-19 patients. Thus, third-level hospitals were responsible for providing hospital beds from their different specialties to these patients due to the high demand for care. COVID-19 was diagnosed by serological and molecular tests.

Variables and measurement instruments

Depressive symptoms

The Patient Health Questionnaire-9 (PHQ-9) is a self-reporting instrument developed to identify possible causes or measure the severity of depressive symptoms within the last two weeks [40]. It is based on the 9 criteria diagnostics from the Diagnostic Statistical Manual of Mental Disorders, five editions [DSM-5]. The items are scored on a four-point scale, ranging from 0 (“not at all”) to 3 (“nearly every day”). Total scores range from 0 to 27 with severity levels of minimal (score 0 to 4), mild (score 5 to 9), moderate (score 10 to 14), moderate-severe (score 15 to 19), and severe (score 20 to 27) depressive symptoms. Also, the screening cut-off point of 10 or more is considered as the presence of clinically relevant depressive symptoms [41, 42]. The Spanish version of the PHQ-9 conducted in Peru, developed by Villarreal-Zegarra [43], showed good psychometric properties. In this study, the scale displayed good levels of reliability (Cronbach’s $\alpha = 0.88$) and validity (See details in Supplementary material 1).

Anxiety symptoms

The General Anxiety Disorder-7 scale (GAD-7), is a self-report scale that assesses the presence or severity of generalized anxiety disorder (GAD) during the 2 weeks before self-application. The items reflect the most prominent diagnostic features of the DSM-5 symptoms criteria for GAD. Response options are scored on a four-point scale: 0 (“not at all”), 1 (“several days”), 2 (“more than half the days”), and 3 (“nearly every day”). Total scores range from 0 to 21 and are categorized as follows: minimal (score 0 to 4), mild (score 5 to 9), moderate (score 10 to 14) and severe levels of anxiety symptoms (score 15 to 21) [44]. In addition, the GAD-7 has a cut-off range of 10 points or more to identify the presence of clinically relevant anxiety symptoms [45–47]. The scale has been translated into Spanish and validated by García-Campayo et al. [48]. In the present study, GAD-7 scale had adequate levels of internal consistency (Cronbach’s $\alpha = 0.87$) and validity (See details in Supplementary material 1).

Somatic symptoms

The Patient Health Questionnaire-15 (PHQ-15) is a scale derived from the full PHQ. It measures 15 somatic symptoms that entail more than 90% of the physical complaints during the past 4 weeks. Items are based on the most prevalent DSM-IV somatization disorder somatic symptoms. The items has three-type Likert response options: 0 (“Not bothered at all”), 1 (“Bothered a little”), and 2 (“Bothered a lot”). Total scores range from 0 to 30 with severity levels of minimal (score 1 to 4), low (score 5 to 9), medium (score 10 to 14), and high (score 15 to

30) somatic symptoms [49, 50]. It presents a cut-off point of 15 to consider clinically significant somatization. The PHQ-15 scale has been translated and validated into Spanish by Ros [51].

Due to differences in some samples in terms of the specific factors: one, two, three [52–54] and four factors [50, 55, 56], we conducted a sub-analysis to assess the psychometric properties of the PHQ-15 through factor analysis and reliability (See the Supplementary material 1). As a result, instead of using the PHQ-15 scale, we decided to use a version of 12 items (PHQ-12), which showed good psychometric properties (Cronbach's $\alpha = 0.83$).

Symptoms of post-traumatic stress

The Impact of Events Scale-Revised (IES-R) is a self-report scale that measures the degree of suffering caused by a life event, described as a form of subjective stress during the past 7 days. The IES-R has 22 items scored with five-point scale, ranging from: 0 ("not at all") to 4 ("extremely"). It is categorized in three dimensions: a) Intrusion dimension (e.g. intrusive distressing thoughts, nightmares, feelings, and images), which items are 1, 2, 3, 6, 9, 14, 16, and 20, b) avoidance dimension (e.g. avoidance of feelings, situations or ideas), which items are 5, 7, 8, 11, 12, 13, 17, and 22, and c) hyperarousal dimension (e.g. anger, hypervigilance, irritability, difficulty concentrating), which items are 4, 10, 15, 18, 19, and 21 [57]. The total scores reflect severity levels of distress symptoms as follows: normal (score 0 to 8), mild (score 9–25), moderate (score 26 to 43), and severe (score 44 to 88) [58]. Moreover, the scale presents a cut-off point of 33 or more that entails clinically relevant symptoms of post-traumatic stress [59]. The scale has been translated and validated in an into Spanish [60]. In the current study, IES-R total score had adequate levels of internal consistency (Cronbach's $\alpha = 0.95$) and validity (See Supplementary material 1).

Perception of symptoms

Self-perception of COVID-19 related symptoms was assessed through two questions in Spanish. The first question asked about how many symptoms the person self-reported at the time of admission to hospitalization and the second question asked about the number of symptoms the person reported at the time of assessment with the psychological instruments. To determine the self-perception of symptoms, the difference between these two questions was assessed. It was expected that if the number of symptoms increased the person self-perceived that his or her illness worsened (positive values) and if the number of symptoms decreased the person perceived that his or her health status improved (negative values). The symptoms were fever, fatigue, myalgia,

cough, dyspnea, odynophagia, rhinorrhea, diarrhea, nausea or vomiting, anosmia, ageusia, headache, dizziness, ataxia, and convulsions.

Neutrophil–lymphocyte ratio (NLR)

The neutrophil–lymphocyte ratio (NLR) was obtained from the patients' complete blood counts on admission. It consists of the ratio between the neutrophil count and the lymphocyte count. The NLR was categorized into < 6.5 and ≥ 6.5 . This cutoff point was chosen considering its ability to predict mortality in patients with COVID-19 [61].

Sociodemographic variables

Information was provided on age, sex, civil status, and employment status. The following questions were also queried: 1) Do you belong to any religion? 2) Have you had a family member with COVID-19? 3) Has any member of your family passed away due to COVID-19? 4) Do you have a previous psychiatric diagnosis? and 5) Have you had a previous psychological treatment?

Data analysis

Descriptive and prevalence

A descriptive analysis of participants was conducted. The prevalence of depressive symptoms (PHQ-9 > 10 or more) [41, 42], anxiety symptoms (GAD-7 > 10 or more) [45, 46], symptoms of post-traumatic stress (IES-R > 33 or more) [59]). We performed a differentiated analysis of the symptoms and indicators of the PHQ-9, GAD-7, IES-R and PHQ-12. The results were stratified based on those with high neutrophil counts (NLR ≥ 6.5), indicating risk of mortality in patients with COVID-19 [61].

Relation between variables

Spearman's Correlation was used to measure the degree of association between variables. We categorized the size of the correlation coefficient as follows: a large ($r > 0.70$), moderate ($r > 0.50$), or small ($r > 0.30$) ratio [62].

Structural regression model

Due the data were non-normal (i.e., categorical indicators), we used a structural regression model with the weighted least squares means and variance adjusted (WLSML) estimator [62]. A polychronic correlation matrix for the nature of the items was also used. Four goodness-of-fit indices were evaluated the proposed model for hospitalized participants with high and low NLR: Comparative Fit Index (CFI), Tucker-Lewis's index (TLI), standardized root mean square residual (SRMR), and root mean square error of approximation (RMSEA). Also, its points cohort is as follows: a) CFI and TLI > 0.95 or more, b) SRMR and RMSEA < 0.08 or flew [63, 64].

In addition, we evaluate the R2 of the outcome variable (depressive symptoms) to determine how much variance explains the proposed model [65].

To ensure sufficient statistical power to perform structural regression analysis, it was considered necessary to have at least 200 participants in total [65]. In addition, to maintain the internal validity of the results, we sought to ensure that exposed and unexposed cases with the inflammatory response (i.e., NRL) had similar sizes.

Statistical software

All analyses were done in R studio version 4.1.1, with the packages “lavaan”, “semTools” and “semPlot” [66].

Results

General characteristics and prevalence

From 319 patients with a diagnosis of COVID-19, we excluded 42 of them as they did not record NLR measurements. Thus, we analyzed data from 277 participants (86.8% of the total number of patients). The average age was 54.2 (± 14.9) years, and most patients were men (61.4%). Two-hundred and twenty-five (81.2%) had at least one family member with COVID-19, and eighty-five (30.7%) had at least one relative die by COVID-19. Most participants did not have a psychiatric diagnosis (93.1%) and did not receive psychological treatment (91%) prior to their COVID-19 infection. Regarding the prevalence of mental health problems, 11.2% was the overall prevalence of depressive symptoms, 7.9% for anxiety symptoms, 2.2% for somatic symptoms, and 6.1% for post-traumatic stress symptoms.

The 48.7% of the participants had a severe inflammatory response, the analysis differentiated by those with and without severe response can be seen in Table 1. In addition, an association was found between age and sex with severe inflammatory response ($p < 0.05$).

Regarding the prevalence of the overall sample, we observed a prevalence in sleep problems (48%) and low energy (47.7%) as depression indicators in the overall sample. Nervousness (48.77%), worry (47.7%) and irritability (43.7%) were the most prevalent indicators of anxiety. Back pain (52%) and trouble sleeping (46.6%) were the most common somatic symptoms. Similarities were observed among inflammatory responses. Sleeping problems (>45.9%) and low energy (>46.5%) were the most common depressive indicators in both groups with and without an inflammatory response. Likewise, while worry (>46.7%) and nervousness (>48.6%) were prevalent in both groups, irritability (47.4%) was higher in patients with severe inflammatory responses than those without inflammatory responses. Differences in somatic symptoms were observed between groups. Back pain (>50%) and trouble sleeping (>46.5%) were the most prevalent

indicators in both samples. Pain in arms and legs (48.1%), feeling tired (43.7%), and shortness of breath indicators (34.8%) were higher in the group with inflammatory responses in comparison with the other group. The clinical indicators for each of the mental health problems by the group are summarized in Fig. 2 and detailed in Supplementary material 2.

Relationship between variables

The findings in Table 2 indicate that correlations between scores for depression, anxiety and somatization symptoms were high for overall participants ($r > 0.70$, $p < 0.05$). A moderate relationship was also observed between symptoms of post-traumatic stress with depression, anxiety and somatic symptoms ($r > 0.50$, $p < 0.05$). A small relationship was found between the perception of COVID-19 symptoms with the other variables, in the group of all participants.

A differential analysis on the strength of the correlation between participants with and without severe inflammatory response was conducted. Among participants without inflammatory response the relationship between post-traumatic stress and symptom perception was small and significant ($r > 0.20$, $p < 0.05$). However, in the group of inflammatory responders, this same correlation was not significant. The strength of the relationship between the other variables did not change.

Structural regression model

The general model that includes individuals with and without severe inflammatory response identified an optimal overall fit, reaching adequate goodness-of-fit indices in all the indexes evaluated (see Table 3). The proposed model explains 85% of the variance of depressive symptoms.

In the general model (see Fig. 3A), the perception of symptoms influenced somatic symptoms ($\beta = 0.223$, $p < 0.05$). In addition, somatic symptoms influenced anxiety symptoms ($\beta = 0.922$, $p < 0.05$) and post-traumatic stress symptoms ($\beta = 0.623$, $p < 0.05$). A non-significant relationship was found between anxiety and post-traumatic stress symptoms ($\beta = 0.205$, $p = 0.262$) and stress-post traumatic stress with depressive symptoms ($\beta = 0.026$, $p = 0.727$). Finally, the relationship between anxiety and depressive symptoms was high and significant ($\beta = 0.902$, $p < 0.05$).

When analyzing separately the overall performance of the models for participants with and without inflammatory response, both models presented adequate goodness-of-fit indices and explained more than 83% of the variance of depressive symptoms. However, the SRMR values are high, possibly due to the small sample size ($n < 200$) (see Table 3).

Table 1 Socio-demographic characteristics ($n = 277$)

		Overall		With severe inflammatory response ($n = 135$)		Without severe inflammatory response ($n = 142$)		p
		n	%	n	%	n	%	
Age (in years)	20 a 39	53	19.1%	15	11.1%	38	26.8%	0.004
	40 a 59	119	43.0%	65	48.1%	54	38%	
	60 a 94	105	37.9%	55	40.7%	50	35.2%	
Sex	Men	170	61.4%	99	73.3%	71	50%	< 0.001
	Women	107	38.6%	36	26.7%	71	50%	
Civil status	Single	34	12.3%	15	11.1%	19	13.4%	0.505
	Married	195	70.4%	99	73.3%	96	67.6%	
	Divorced	18	6.5%	6	4.4%	12	8.5%	
	Widowed	30	10.8%	15	11.1%	15	10.6%	
Employment status	Unemployed	120	43.3%	55	40.7%	65	45.8%	0.469
	Employed	157	56.7%	80	59.3%	77	54.3%	
Do you belong to any religion?	No	21	7.6%	11	8.1%	10	7%	0.904
	Yes	256	92.4%	124	91.9%	132	93%	
Have you had a family member with COVID-19?	No	52	18.8%	27	20%	25	17.6%	0.722
	Yes	225	81.2%	108	80%	117	82.4%	
Has any member of your family passed away due to COVID-19?	No	192	69.3%	91	67.4%	101	71.1%	0.589
	Yes	85	30.7%	44	32.6%	41	28.9%	
Previous psychiatric diagnosis	No	258	93.1%	128	94.8%	130	91.5%	0.403
	Yes	19	6.9%	7	5.2%	12	8.5%	
Previous psychological treatment	No	252	91.0%	124	91.9%	128	90.1%	0.774
	Yes	25	9.0%	11	8.1%	14	9.9%	
Depressive symptoms	No	246	88.8%	116	85.9%	130	91.5%	0.196
	Yes	31	11.2%	19	14.1%	2	8.5%	
Anxiety symptoms	No	255	92.1%	121	89.6%	134	94.4%	0.217
	Yes	22	7.9%	14	10.4%	8	5.6%	
Somatic symptoms	No	271	97.8%	132	97.8%	139	97.9%	0.919
	Yes	6	2.2%	3	2.2%	3	2.1%	
Symptoms of post-traumatic stress	No	260	93.9%	126	93.3%	134	94.4%	0.914
	Yes	17	6.1%	9	6.7%	8	5.6%	

The p -value is obtained from the chi-square analysis


The specific assessment of the relationships of the proposed models identified different performances for participants with a severe inflammatory response (see Fig. 3B) and for participants without a severe inflammatory response (see Fig. 3C). The relationship between perception of COVID-19 symptoms and somatic symptoms was found to be significant for this group of participants without severe inflammatory response ($\beta = 0.289$, $p < 0.05$), but the relationship was not significant in the group of participants with a severe inflammatory response ($\beta = 0.289$, $p = 0.117$). The relationship between symptoms of post-traumatic stress and anxiety symptoms in the group without severe inflammatory response was direct, significant, and high ($\beta = 0.684$, $p < 0.05$). However, the relationship was inverted and not significant

among participants with a severe inflammatory response ($\beta = -0.277$, $p = 0.531$). A non-significant relationship was found between symptoms of post-traumatic stress and depressive symptoms in both groups (i.e., with and without a severe inflammatory response).

Discussion

Main findings and significance of the results

There is extensive discussion on the role of biological and psychological variables in the occurrence of depressive symptoms in patients with COVID-19. Previous studies found higher values in inflammatory markers (i.e., NLR) in patients with depression, compared with the non-depression groups [25, 61]. Our research presents evidence that a model depicting the subjective perception



of symptoms (psychosomatic and COVID-19 symptoms) with anxiety and post-traumatic stress symptoms explains the presence of depressive symptoms in patients with and without severe inflammatory response in NLR who were hospitalized for COVID-19.

We used the anxiety-PTSD-depression triad as a base model and added the variables related to the subjective perception of symptoms. Our model explained 85% of depressive symptoms in the overall sample, indicating that several mood disorders occur simultaneously prior to the development of depression [67].

The model also showed that psychosomatic symptoms and anxiety have the most influence in the occurrence of depressive symptoms. The impact of these variables may be explained by biological perspectives. For instance, an increase of anxiety is associated with presence of somatic symptoms, such as headaches and shoulder and limb pain [67]. Besides, the relationship between psychosomatic symptoms and symptoms of post-traumatic stress

represents a relevant effect, which can be explained due to somatic symptoms being more prevalent during periods of stress [68].

Contrasting findings with existing literature

Prevalence and indicators

In our study, the most prevalent mental health problems were anxiety (11.2%), depression (7.1%) and PTSD (6.1%). Similarly, a previous study conducted in Peru reported the prevalence for depression (12.1%), anxiety (8.4%), and PTSD (10.5%) in health care-workers during the pandemic [24]. The prevalence reported in systematic reviews indicated higher values than our study showed for depression (45%), anxiety (47%), and PTSD (16 to 22.6%) [4, 69, 70]. These differences among researchers could be explained due to the different socio-demographic compositions, different study designs, and measurement instruments used, which may influence the degree of prevalence. Even so, results evidenced that the

Table 2 Correlations between depressive symptoms, anxiety symptoms, somatic symptoms, symptoms of post-traumatic stress, and perception of physical symptom (n = 277)

Group	Variable	1	2	3	4	4.1	4.2	4.3	5
Overall (n = 277)	1. Depressive symptoms	1							
	2. Anxiety symptoms	0.77*	1						
	3. Somatic symptoms	0.73*	0.71*	1					
	4. Symptoms of post-traumatic stress disorder	0.64*	0.65*	0.67*	1				
	4.1 Intrusion	0.63*	0.63*	0.65*	0.93*	1			
With severe inflam- matory response (NLR ≥ 6.5) (n = 135)	4.2 Avoidance	0.52*	0.57*	0.59*	0.92*	0.82*	1		
	4.3 Hyperarousal	0.68*	0.65*	0.69*	0.90*	0.81*	0.74*	1	
	5. Perception of physical symptoms	0.17*	0.17*	0.22*	0.11	0.13*	0.10	0.13*	1
	1. Depressive symptoms	1							
	2. Anxiety symptoms	0.79*	1						
Without severe inflam- matory response (NLR < 6.5) (n = 142)	3. Somatic symptoms	0.72*	0.70*	1					
	4. Symptoms of post-traumatic stress disorder	0.62*	0.64*	0.65*	1				
	4.1 Intrusion	0.62*	0.67*	0.64*	0.92*	1			
	4.2 Avoidance	0.45*	0.54*	0.56*	0.91*	0.81*	1		
	4.3 Hyperarousal	0.68*	0.63*	0.69*	0.91*	0.83*	0.73*	1	
	5. Perception of physical symptoms	0.18*	0.08*	0.15*	0.02*	0.03	-0.01	0.05	1

The relationship was assessed using Spearman's coefficient

* p < 0.05: significant correlation between variables

Table 3 Goodness-of-fit indices of the structural regression model

Model	n	χ ²	χ ² /df	CFI	TLI	RMSEA [90% CI]	SRMR	R ²
Overall	277	1963.3	1.61	0.984	0.984	0.047 [0.043—0.051]	0.085	0.850
With severe inflam- matory response	135	1667.8	1.37	0.971	0.971	0.053 [0.046—0.059]	0.109	0.837
Without severe inflammatory response	142	1552.1	1.28	0.960	0.959	0.044 [0.037—0.051]	0.128	0.898


χ² chi-square, CFI comparative fit index, TLI Tucker-Lewis's index, RMSEA root mean square error of approximation, CI confidence intervals, SRMR standardized root mean square. R² coefficient of determination, Degrees of freedom = 1,216

patients with COVID-19 present several mental health problems at the same time. This could mean that, during the treatment, patients may develop multiple related psychiatric diseases which form a mutually influential symptom network, in turn, influencing their recovery [71].


Sleep problems were one of the most frequent indicators of mental health problems. This relationship was previously reported in a systematic review, in which sleep problems were associated with higher levels of mental

health problems (i.e., anxiety and depression) among mental health care-workers, general population, and COVID-19 patients [72]. Also, this indicator has been found to be prevalent in healthcare workers even before COVID-19 [72]. A possible explanation for the prevalence of sleep problems could be fear of COVID-19. Due to worries about the disease, patients may struggle with sleep and consequently develop insomnia [73]. Moreover, if the individual cannot manage the fear for a specific

A (Overall)



B (With severe inflammatory response)



C (Without severe inflammatory response)




Fig. 3 Path Analysis. Note: **A** Overall participant. **B** Participants with severe inflammatory response. **C** Participants without severe inflammatory response. The model was estimated with the WLSMV method. Values in red are not significant. * $p < 0.05$

time, they may experience mental health problems (e.g., depression, anxiety) [74]. We also found that low energy had the second highest, which is a comorbid symptom of many psychiatric problems in patients with COVID-19 [75]. In addition, low energy could be caused by lack of sleep or one of the other mental health problems such as depression.

The main indicators related to anxiety were nervousness, irritability, and worry. This finding was also reported in other studies in individuals with COVID-19, with the most common symptoms of anxiety being insomnia, irritability, restlessness, and excessive worrying [76]. As previously mentioned, being hospitalized, patients may experience fear of COVID-19 and worries related to their health, family, financial issues, and environmental conditions (e.g., isolation, uncertainty about the evolution of the disease). In our study, somatic complaints (i.e., backache, arms, and legs pain, and feeling exhausted) were most prevalent. This may be due to similarities between somatic complaints and physiological symptoms of COVID-19.

Structural equation model and relationship between variables

Previous studies of predicted models identified an impact negative of COVID-19 on mental health problems (i.e., depression, anxiety, stress, fear of COVID-19, among others). Two studies proposed models in which fear of COVID is significantly and positively related to depression, anxiety, and insomnia [77, 78]. Moreover, epidemiological studies reported the SARS-CoV-2 virus could lead to systems immune changes, which in turn could reflect in mental health problems. These psychiatric outcomes can be influenced by other factors as well (i.e., biological, factor social isolation, adverse effects of treatments, etc.) [12, 79]. However, a small number of studies has proposed predictive models about the relationship of biological responses with these mental clinical problems. For example, a study, using SEM, proposed a model that cortisol (as an indicator of Hypothalamic pituitary adrenal) predicts depression, which predicts circulating pro-inflammatory cytokines (IL-2, IL-6, TNF- α) In patients diagnosed with chronic fatigue syndrome (CFS) [80]. Another study explored the relationship between biological factors (i.e., sex, disease duration, self-perceived illness severity, and inflammatory markers) and mental health status in inpatients with COVID-19. In the SEM, inflammatory markers (i.e., NLR, IL-1 β as observed variables) and mental health (i.e., insomnia, depression, and anxiety as observed variables) were set as latent variables. Results indicated the inflammatory markers had a significant and direct effect on mental health. Moreover, the

disease duration and inflammatory markers indirectly influenced mental health, through self-perceived illness severity as a mediator [25]. These findings suggest that inflammatory responses could be related to psychological disorders.

Following this hypothesis, studies have found a heterogeneous influence of NLR on psychological mental problems. First, one study, using regression analysis, demonstrated the influence of NLR markers on both the prevalence of depression and anxiety in Chinese patients with gastric cancer [81]. In contradiction to this finding, a multi-linear regression study showed a weak association between inflammatory biomarkers and depression in a three-month cohort of stroke patients [82]. As similar immune responses exist in both COVID-19 infection and mood disorders, they may share biological response as well. Both states induce the production of abnormal levels of cytokines, chemokines, and other inflammatory mediators [83], showing a hyperinflammatory state [84]. While patients with mental health problems showed high levels of biomarkers [17], a meta-analysis, with 16 studies, evidenced higher counts of biomarkers (i.e. IL-6, CRP, PCT, among others) in severe cases of COVID-19 [85].

Another interesting result was the high influence of anxiety on depression in all three models. This finding is in accordance with other studies which evidenced that anxiety symptoms had a direct and significant relationship with depression. One study that proposed a model of the triad fear-anxiety-stress in the development of depression symptoms in pandemic disease symptoms in health workers, indicated that the fear of COVID-19, anxiety and post-traumatic symptoms explains depression symptoms. The SEM demonstrated that anxiety was the most influential variable in depression symptoms in comparison with post-traumatic stress [24]. Preceding COVID-19, researchers have shown that anxiety may contribute directly or as mediating variables in depression. These results show how the different variables (i.e., stress, self-esteem, stressful negative events) influences depression, where increases in anxiety may lead to increases in depression [86, 87]. In sum, these findings suggest that the role of anxiety in the occurrence of depressive symptoms is significant and is even maintained in the COVID-19 pandemic. Anxiety is a common adaptive response against threatening situations, which could be increased due to factors such as stress or fear and could trigger prolonged anxiety. Thus, pathological anxiety can affect functioning in the daily routine of patients, which in turn may cause or be comorbid with other mental disorders such as depression [88].

Another result was the influence of the PTSD variable on depression. Our results demonstrated that

PTSD symptoms do not present a significant influence on depression in hospitalized patients with and without severe inflammatory markers. This finding might be related to the PTSD symptoms changes over time. Other studies have found the different prevalence of PTSD symptoms in each stage of COVID-19 disease (i.e., recovering from COVID-19 infection, being quarantined) [89, 90]. Likewise, another reason could be the similarity between our variables. There are studies that report a high association between PTSD and somatic symptoms. Findings support the fact that somatic symptoms may be related to the patient's psychophysiological dysregulation and lead to psychological symptoms (e.g., PTSD) [91, 92].

Implications in public health and making decisions

Our findings provide a theoretical model, which may help establish policies to prevent depression among inpatients with COVID-19. Specifically, the model revealed that somatic and anxiety symptoms are the most relevant predictors to develop depression. Health workers could employ screening measures for anxiety and somatic symptoms to prioritize the care of patients with high levels in these conditions, and thus avoid possible cases of depressive symptoms. It is a necessity because Peru is one of the countries that reported worse mental health levels in the world during the pandemic [93]. Also, the prevalence of depression in 2020 was five times higher than previous years [94].

Interventions to reduce symptoms of anxiety, fear and worry in hospitalized patients could prevent subsequent cases of mental illness [95]. For example, telephone-based interventions also have been useful to reduce symptoms of anxiety and depression, providing psychological support, information about the process of the disease and promoting a sense of emotional stability [96, 97]. Thus, the implementation of telephones during hospitalizations could be a strategy to prevent psychological problems in hospital isolated patient so and could be a facility for patients to have access to make calls or send messages to their relatives.

Strengths and limitations

This study has limitations that should be mentioned. First, some patients did not have inflammatory markers recorded, so they were excluded. This exclusion could lead to an information bias. Second, NLR was evaluated as the only inflammatory measure. However, using inflammatory markers to assess inflammatory response is not considered a gold standard. Therefore, doing so may have caused errors when grouping participants into those with and without a severe inflammatory response. Third, this study has a cross-sectional design, thus we cannot infer causality in the interpretation of the findings. Fourth, we employed

self-reported measures, which may have been influenced by social desirability or memory bias. Fifth, the data includes a single hospital in a Peruvian city. Therefore, results should not be generalized to other cities or contexts. Sixth, we used a validated scale such as the IES-R to measure PTSD, however, IES-R does not include the entire concept of PTSD. The Diagnostic and Statistical Manual of Mental Disorders (DSM-5) considers four dimensions and the IES-R only assesses three of these dimensions. This could imply a partial evaluation of the symptoms of PTSD. Finally, other confounding variables were not considered, such as fear of COVID-19 [24] and coping [98]. Thus, it is possible that the model is partial or influenced by other variables.

On the other hand, our study has three main strengths. This investigation presents a larger sample compared to previous studies evaluating hospitalized patients [25, 99]. We also employed structural equation modelling, which allowed us to assess several variables simultaneously. Moreover, to our knowledge, this is the first study that provides a framework of biological and psychological variables that explain depressive symptoms as an outcome in the context of the COVID-19.

Conclusions and recommendations

Results demonstrated that our model of mental health variables may explain depression in hospitalized patients of COVID-19 from a third-level hospital in Peru. In the model, perception of symptoms influences somatic symptoms, which influence both anxiety symptoms and symptoms of post-traumatic stress. Thus, anxiety symptoms could directly influence depressive symptoms or through PTSD symptoms. Additionally, our model was found to have a good overall fit and explained more than 83% of the depressive symptoms.

Regarding clinical indicators, patients presented a high prevalence of depression, anxiety, and psychosomatic indicators. Our findings could be useful to decision-makers for the prevention of depression, such as to encourage the use of screening tools (i.e., perception of symptoms, somatic symptoms, anxiety) that may sooner identify patients vulnerable to depression.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12888-022-04277-4>.

Additional file 1. Supplementary material 1. Psychometric properties of the scales used (n=277).

Additional file 2. Supplementary material 2. Values of the prevalence of clinical indicators of depression, anxiety and psychosomatic symptoms (n=277).

Acknowledgements

The authors acknowledge the contribution of Jessica Barreto, Angela Podestà, Claudia Castillo, Rosa Guija and Lucía Aire, for their dedicated work in the data collection.

Authors' contributions

David Villarreal-Zegarra: Formal Analysis, Methodology, Supervision, Validation, Writing – Original version, Approval of the final version. Rubí Paredes: Conceptualization, Formal Analysis, Methodology, Validation, Writing – Original version, Approval of the final version. Nikol Mayo: Conceptualization, Methodology, Validation, Writing – Original version, Approval of the final version. Anthony Copez-Lonzoy: Methodology, Validation, Writing – Original version, Approval of the final version. Ana L. Vilela-Estrada: Methodology, Validation, Writing – Original version, Approval of the final version. Jeff Huarcaya-Victoria: Conceptualization, Methodology, Validation, Writing – Review & Editing, Approval of the final version.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Availability of data and materials

The datasets generated and analysed during the current study are not publicly available due their containing information that could compromise the privacy of research participants but are available from the corresponding author on reasonable request.

Declarations

Ethics of approval and consent to participate

This study has been approved by the Institutional Review Board (IRB) for COVID-19 of the "Seguro Social del Perú" (EsSalud). All methods were carried out in accordance with relevant guidelines and regulations. The guidelines of the Helsinki Declaration were followed. Each participant provided signed virtual informed consent. The data collected were codified and duly protected to guarantee the confidentiality of the information and the results of the patients.

Consent for publication

Not applicable.

Competing interests

The authors report no conflict of interest when conducting the study, analyzing the data, or writing the manuscript.

Author details

¹Escuela de Medicina, Universidad César Vallejo, Trujillo, Peru. ²Instituto Peruano de Orientación Psicológica, Lima, Peru. ³Unidad de Investigación en Bibliometría, Universidad San Ignacio de Loyola, Lima, Peru. ⁴PSYCOPERU Peruvian Research Institute of Educational and Social Psychology, Lima, Peru. ⁵Escuela Profesional de Medicina Humana, Universidad Privada San Juan Bautista, Filial Ica, Peru. ⁶Departamento de Psiquiatría, Servicio de Psiquiatría de Adultos, Unidad de Psiquiatría de Enlace, Hospital Nacional Guillermo Almendra Irigoyen, Lima, Perú.

Received: 11 April 2022 Accepted: 5 August 2022

Published online: 10 October 2022

References

- Alamri HS, Mousa WF, Algarni A, Megahid SF, Al Bshabshe A, Alshehri NN, et al. Mental health of COVID-19 patients-a cross-sectional survey in Saudi Arabia. *Int J Environ Res Public Health*. 2021;18. <https://doi.org/10.3390/ijerph18094758>.
- Ismail F, Bizario JCS, Battagin T, Zaramella B, Leal FE, Torales J, et al. Post-infection depressive, anxiety and post-traumatic stress symptoms: a prospective cohort study in patients with mild COVID-19. *Prog Neuropsychopharmacol Biol Psychiatry*. 2021;111: 110341. <https://doi.org/10.1016/j.pnpb.2021.110341>.
- Li T, Sun S, Liu B, Wang J, Zhang Y, Gong C, et al. Prevalence and risk factors for anxiety and depression in patients with COVID-19 in Wuhan. *China Psychosom Med*. 2021;83:368–72. <https://doi.org/10.1097/PSY.0000000000000934>.
- Deng J, Zhou F, Hou W, Silver Z, Wong CY, Chang O, et al. The prevalence of depression, anxiety, and sleep disturbances in COVID-19 patients: a meta-analysis. *Ann NY Acad Sci*. 2021;1486:90–111. <https://doi.org/10.1111/nyas.14506>.
- Yuan K, Gong Y-M, Liu L, Sun Y-K, Tian S-S, Wang Y-J, et al. Prevalence of posttraumatic stress disorder after infectious disease pandemics in the twenty-first century, including COVID-19: a meta-analysis and systematic review. *Mol Psychiatry*. 2021. <https://doi.org/10.1038/s41380-021-01036-x>.
- Mohammadian Khonsari N, Shafiee G, Zandifar A, Mohammad Poornami S, Ejtahed H-S, Asayesh H, et al. Comparison of psychological symptoms between infected and non-infected COVID-19 health care workers. *BMC Psychiatry*. 2021;21:170. <https://doi.org/10.1186/s12888-021-03173-7>.
- Huarcaya-Victoria J, Villarreal-Zegarra D, Podestà A, Luna-Cuadros MA. Psychometric Properties of a Spanish Version of the Fear of COVID-19 scale in general population of Lima, Peru. *Int J Ment Health Addict*. 2020. <https://doi.org/10.1007/s11469-020-00354-5> [cited 22 May 2021].
- Huarcaya-Victoria J, Barreto J, Aire L, Podestà A, Caqui M, Guija-Igreda R, et al. Mental health in COVID-19 survivors from a general hospital: association with sociodemographic, clinical, and inflammatory variables. *In Review*; 2021. <https://doi.org/10.21203/rs.3.rs-146200/v1>.
- Kontoangelos K, Economou M, Papageorgiou C. Mental health effects of COVID-19 Pandemia: a review of clinical and psychological traits. *Psychiatry Investig*. 2020;17:491–505. <https://doi.org/10.30773/pi.2020.0161>.
- Wu T, Jia X, Shi H, Niu J, Yin X, Xie J, et al. Prevalence of mental health problems during the COVID-19 pandemic: a systematic review and meta-analysis. *J Affect Disord*. 2021;281:91–8. <https://doi.org/10.1016/j.jad.2020.11.117>.
- Conti P, Ronconi G, Caraffa A, Gallenga C, Ross R, Frydas I, et al. Induction of pro-inflammatory cytokines (IL-1 and IL-6) and lung inflammation by Coronavirus-19 (COVI-19 or SARS-CoV-2): anti-inflammatory strategies. *J Biol Regul Homeost Agents*. 2020;34:327–31. <https://doi.org/10.23812/CONTI-E>.
- Raony Í, de Figueiredo CS, Pandolfo P, Giestal-de-Araujo E, Oliveira-Silva Bomfim P, Savino W. Psycho-neuroendocrine-immune interactions in COVID-19: potential impacts on mental health. *Front Immunol*. 2020;11. <https://doi.org/10.3389/fimmu.2020.01170>.
- Beltrán BE, Paredes S, Cotrina E, Sotomayor EM, Castillo JJ. The impact of the neutrophil:lymphocyte ratio in response and survival of patients with de novo diffuse large B-cell lymphoma. *Leuk Res*. 2018;67:82–5. <https://doi.org/10.1016/j.leukres.2018.02.011>.
- Karageorgiou V, Milas GP, Michopoulos I. Neutrophil-to-lymphocyte ratio in schizophrenia: a systematic review and meta-analysis. *Schizophr Res*. 2019;206:4–12. <https://doi.org/10.1016/j.schres.2018.12.017>.
- Seropian IM, Romeo FJ, Pizarro R, Vulcano NO, Posatini RA, Marenchino RG, et al. Neutrophil-to-lymphocyte ratio and platelet-to-lymphocyte ratio as predictors of survival after heart transplantation. *ESC Heart Fail*. 2018;5:149–56. <https://doi.org/10.1002/ehf2.12199>.
- Yoshitomi R, Nakayama M, Sakoh T, Fukui A, Katafuchi E, Seki M, et al. High neutrophil/lymphocyte ratio is associated with poor renal outcomes in Japanese patients with chronic kidney disease. *Ren Fail*. 2019;41:238–43. <https://doi.org/10.1080/0886022X.2019.1595645>.
- Mazza MG, Lucchi S, Tringali AGM, Rossetti A, Botti ER, Clerici M. Neutrophil/lymphocyte ratio and platelet/lymphocyte ratio in mood disorders: a meta-analysis. *Prog Neuropsychopharmacol Biol Psychiatry*. 2018;84:229–36. <https://doi.org/10.1016/j.pnpb.2018.03.012>.
- Wang M, Hu C, Zhao Q, Feng R, Wang Q, Cai H, et al. Acute psychological impact on COVID-19 patients in Hubei: a multicenter observational study. *Transl Psychiatry*. 2021;11:1–9. <https://doi.org/10.1038/s41398-021-01259-0>.
- Willis C, Chalder T. Concern for Covid-19 cough, fever and impact on mental health. What about risk of Somatic Symptom Disorder? *J Ment Health*. 2021;0:1–5. <https://doi.org/10.1080/09638237.2021.1875418>.
- Xiang Y-T, Yang Y, Li W, Zhang L, Zhang Q, Cheung T, et al. Timely mental health care for the 2019 novel coronavirus outbreak is urgently needed. *Lancet Psychiatry*. 2020;7:228–9. [https://doi.org/10.1016/S2215-0366\(20\)30046-8](https://doi.org/10.1016/S2215-0366(20)30046-8).

21. Abad C, Fearday A, Safdar N. Adverse effects of isolation in hospitalised patients: a systematic review. *J Hosp Infect.* 2010;76:97–102. <https://doi.org/10.1016/j.jhin.2010.04.027>.
22. Kang E, Lee SY, Kim MS, Jung H, Kim KH, Kim K-N, et al. The psychological burden of COVID-19 stigma: evaluation of the mental health of isolated Mild condition COVID-19 Patients. *J Korean Med Sci.* 2021;36. <https://doi.org/10.3346/jkms.2021.36.e33>.
23. Purrussell E, Gould D, Chudleigh J. Impact of isolation on hospitalised patients who are infectious: systematic review with meta-analysis. *BMJ Open.* 2020;10:e030371. <https://doi.org/10.1136/bmjopen-2019-030371>.
24. Villarreal-Zegarra D, Copez-Lonzoy A, Vilela-Estrada A, Huarcaya-Victoria J. Depression, post-traumatic stress, Anxiety, and Fear of COVID-19 in the general population and health-care workers: prevalence, relationship, and explicative Model in Peru. 2021. <https://doi.org/10.21203/rs.3.rs-151028/v1>.
25. Hu Y, Chen Y, Zheng Y, You C, Tan J, Hu L, et al. Factors related to mental health of inpatients with COVID-19 in Wuhan. *China Brain Behav Immun.* 2020;89:587–93. <https://doi.org/10.1016/j.cbbi.2020.07.016>.
26. Shangguan F, Quan X, Qian W, Zhou C, Zhang C, Zhang XY, et al. Prevalence and correlates of somatization in anxiety individuals in a Chinese online crisis intervention during COVID-19 epidemic. *J Affect Disord.* 2020;277:436–42. <https://doi.org/10.1016/j.jad.2020.08.035>.
27. Longo MS de C, Vilete LMP, Figueira I, Quintana MI, Mello MF, Bressan RA, et al. Comorbidity in post-traumatic stress disorder: a population-based study from the two largest cities in Brazil. *J Affect Disord.* 2020;263:715–721. <https://doi.org/10.1016/j.jad.2019.11.051>.
28. McAndrew LM, Lu S-E, Phillips LA, Maestro K, Quigley KS. Mutual maintenance of PTSD and physical symptoms for Veterans returning from deployment. *Eur J Psychotraumatology.* 2019;10:1608717. <https://doi.org/10.1080/20008198.2019.1608717>.
29. Stensland SØ, Thoresen S, Jensen T, Wentzel-Larsen T, Dyb G. Early pain and other somatic symptoms predict posttraumatic stress reactions in survivors of terrorist attacks: the longitudinal utøya cohort study. *J Trauma Stress.* 2020;33:1060–70. <https://doi.org/10.1002/jts.22562>.
30. Leeuw M, Goossens MEJB, Linton SJ, Crombez G, Boersma K, Vlaeyen JWS. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med.* 2007;30:77–94. <https://doi.org/10.1007/s10865-006-9085-0>.
31. Gonzalez P, Martinez KG. The role of stress and fear on the development of psychopathology. *Psychiatr Clin North Am.* 2014;37:535–46. <https://doi.org/10.1016/j.psc.2014.08.010>.
32. Beck K, Vincent A, Becker C, Keller A, Cam H, Schaefer R, et al. Prevalence and factors associated with psychological burden in COVID-19 patients and their relatives: a prospective observational cohort study. *PLoS ONE.* 2021;16:e0250590. <https://doi.org/10.1371/journal.pone.0250590>.
33. Guo Q, Zheng Y, Shi J, Wang J, Li G, Li C, et al. Immediate psychological distress in quarantined patients with COVID-19 and its association with peripheral inflammation: A mixed-method study. *Brain Behav Immun.* 2020;88:17–27. <https://doi.org/10.1016/j.cbbi.2020.05.038>.
34. Bo H-X, Li W, Yang Y, Wang Y, Zhang Q, Cheung T, et al. Posttraumatic stress symptoms and attitude toward crisis mental health services among clinically stable patients with COVID-19 in China. *Psychol Med.* 2021;51:1052–3. <https://doi.org/10.1017/S0033291720000999>.
35. Jacobson NC, Newman MG. Anxiety and depression as bidirectional risk factors for one another: a meta-analysis of longitudinal studies. *Psychol Bull.* 2017;143:1155–200. <https://doi.org/10.1037/bul0000111>.
36. Silk JS, Price RB, Rosen D, Ryan ND, Forbes EE, Siegle GJ, et al. A longitudinal follow-up study examining adolescent depressive symptoms as a function of prior anxiety treatment. *J Am Acad Child Adolesc Psychiatry.* 2019;58:359–67. <https://doi.org/10.1016/j.jaac.2018.10.012>.
37. Choi EPH, Hui BPH, Wan EYF. Depression and Anxiety in Hong Kong during COVID-19. *Int J Environ Res Public Health.* 2020;17. <https://doi.org/10.3390/ijerph17103740>.
38. Ni MY, Jiang C, Cheng KK, Zhang W, Gilman SE, Lam TH, et al. Stress across the life course and depression in a rapidly developing population: the Guangzhou Biobank Cohort Study. *Int J Geriatr Psychiatry.* 2016;31:629–37. <https://doi.org/10.1002/gps.4370>.
39. Huarcaya-Victoria J, Barreto J, Aire L, Podestá A, Caqui M, Guija-Igreda R, et al. Mental Health in COVID-2019 Survivors from a General Hospital in Peru: Sociodemographic, Clinical, and Inflammatory Variable Associations. *Int J Ment Health Addict.* 2021;1–22. <https://doi.org/10.1007/s11469-021-00659-z>.
40. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001;16:606–13. <https://doi.org/10.1046/j.1525-1497.2001.016009606.x>.
41. Levis B, Benedetti A, Thombs BD, DEPRESSion Screening Data (DEPRESSD) Collaboration. Accuracy of Patient Health Questionnaire-9 (PHQ-9) for screening to detect major depression: individual participant data meta-analysis. *BMJ.* 2019;365:11476. <https://doi.org/10.1136/bmj11476>.
42. Manea L, Gilbody S, McMillan D. A diagnostic meta-analysis of the patient health questionnaire-9 (PHQ-9) algorithm scoring method as a screen for depression. *Gen Hosp Psychiatry.* 2015;37:67–75. <https://doi.org/10.1016/j.genhosppsych.2014.09.009>.
43. Villarreal-Zegarra D, Copez-Lonzoy A, Bernabé-Ortiz A, Melendez-Torres GJ, Bazo-Alvarez JC. Valid group comparisons can be made with the Patient Health Questionnaire (PHQ-9): a measurement invariance study across groups by demographic characteristics. *PLoS ONE.* 2019;14:e0221717. <https://doi.org/10.1371/journal.pone.0221717>.
44. Spitzer RL, Kroenke K, Williams JBW, Löwe B. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med.* 2006;166:1092–7. <https://doi.org/10.1001/archinte.166.10.1092>.
45. Plummer F, Manea L, Trepel D, McMillan D. Screening for anxiety disorders with the GAD-7 and GAD-2: a systematic review and diagnostic metaanalysis. *Gen Hosp Psychiatry.* 2016;39:24–31. <https://doi.org/10.1016/j.genhosppsych.2015.11.005>.
46. Löwe B, Decker O, Müller S, Brähler E, Schellberg D, Herzog W, et al. Validation and standardization of the generalized anxiety disorder screener (GAD-7) in the general population. *Med Care.* 2008;46:266–74. <https://doi.org/10.1097/MLR.0b013e318160d093>.
47. Kroenke K, Spitzer RL, Williams JBW, Monahan PO, Löwe B. Anxiety disorders in primary care: prevalence, impairment, comorbidity, and detection. *Ann Intern Med.* 2007;146:317–25. <https://doi.org/10.7326/0003-4819-146-5-200703060-00004>.
48. García-Campayo J, Zamorano E, Ruiz MA, Pardo A, Pérez-Páramo M, López-Gómez V, et al. Cultural adaptation into Spanish of the generalized anxiety disorder-7 (GAD-7) scale as a screening tool. *Health Qual Life Outcomes.* 2010;8:8. <https://doi.org/10.1186/1477-7525-8-8>.
49. Kroenke K, Spitzer RL, Williams JBW. The PHQ-15: validity of a new measure for evaluating the severity of somatic symptoms. *Psychosom Med.* 2002;64:258–66.
50. Witthöft M, Hiller W, Loch N, Jasper F. The latent structure of medically unexplained symptoms and its relation to functional somatic syndromes. *Int J Behav Med.* 2013;20:172–83. <https://doi.org/10.1007/s12529-012-9237-2>.
51. Ros Montalbán S, Comas Vives A, Garcia-García M. Validation of the Spanish version of the PHQ-15 questionnaire for the evaluation of physical symptoms in patients with depression and/or anxiety disorders: DEPRE-SOMA study. *Actas Esp Psiquiatr.* 2010;38:345–57.
52. Liao S-C, Huang W-L, Ma H-M, Lee M-T, Chen T-T, Chen I-M, et al. The relation between the patient health questionnaire-15 and DSM somatic diagnoses. *BMC Psychiatry.* 2016;16. <https://doi.org/10.1186/s12888-016-1068-2>.
53. Zhang L, Fritzsche K, Liu Y, Wang J, Huang M, Wang Y, et al. Validation of the Chinese version of the PHQ-15 in a tertiary hospital. *BMC Psychiatry.* 2016;16:89. <https://doi.org/10.1186/s12888-016-0798-5>.
54. Grover S, Sahoo S, Chakrabarti S, Avasthi A. Anxiety and somatic symptoms among elderly patients with depression. *Asian J Psychiatry.* 2019;41:66–72. <https://doi.org/10.1016/j.ajp.2018.07.009>.
55. Zhou Y, Xu J, Rief W. Are comparisons of mental disorders between Chinese and German students possible? An examination of measurement invariance for the PHQ-15, PHQ-9 and GAD-7. *BMC Psychiatry.* 2020;20:480. <https://doi.org/10.1186/s12888-020-02859-8>.
56. Walentynowicz M, Witthöft M, Raes F, Van Diest I, Van den Bergh O. Sensory and affective components of symptom perception: A psychometric approach. *J Exp Psychopathol.* 2018;9:jep.059716. <https://doi.org/10.5127/jep.059716>.
57. Weiss DS. The Impact of Event Scale: Revised. In: Wilson JP, Tang CS, editors. *Cross-Cultural Assessment of Psychological Trauma and PTSD*. Boston, MA: Springer US; 2007. pp. 219–238. https://doi.org/10.1007/978-0-387-70990-1_10.

58. Guest R, Tran Y, Gopinath B, Cameron ID, Craig A. Prevalence and psychometric screening for the detection of major depressive disorder and post-traumatic stress disorder in adults injured in a motor vehicle crash who are engaged in compensation. *BMC Psychol*. 2018;6:4. <https://doi.org/10.1186/s40359-018-0216-5>.
59. Creamer M, Bell R, Failla S. Psychometric properties of the impact of event scale - revised. *Behav Res Ther*. 2003;41:1489–96. <https://doi.org/10.1016/j.brat.2003.07.010>.
60. Caamaño WL, Fuentes MD, González BL, Melipillán AR, Sepúlveda CM, Valenzuela GE. Adaptación y validación de la versión chilena de la escala de impacto de evento-revisada (EIE-R). *Rev Médica Chile*. 2011;139:1163–8. <https://doi.org/10.4067/S0034-98872011000900008>.
61. Li X, Liu C, Mao Z, Xiao M, Wang L, Qi S, et al. Predictive values of neutrophil-to-lymphocyte ratio on disease severity and mortality in COVID-19 patients: a systematic review and meta-analysis. *Crit Care Lond Engl*. 2020;24:647. <https://doi.org/10.1186/s13054-020-03374-8>.
62. Akoglu H. User's guide to correlation coefficients. *Turk J Emerg Med*. 2018;18:91–3. <https://doi.org/10.1016/j.tjem.2018.08.001>.
63. Brown TA. Confirmatory factor analysis for applied research. New York, NY, US: The Guilford Press; 2006. pp. xiii, 475.
64. Iacobucci D. Structural equations modeling: Fit Indices, sample size, and advanced topics. *J Consum Psychol*. 2010;20:90–8. <https://doi.org/10.1016/j.jcps.2009.09.003>.
65. Hair JF, Anderson RE, Tatham RL, Black WC. Análisis multivariante. 2004. Available: <https://dialnet.unirioja.es/servlet/libro?codigo=320227>.
66. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2021. Available: <https://www.R-project.org/>.
67. Wong JY-H, Fong DY-T, Chan KK-W. Anxiety and insomnia as modifiable risk factors for somatic symptoms in Chinese: a general population-based study. *Qual Life Res*. 2015;24: 2493–2498. <https://doi.org/10.1007/s11136-015-0984-9>.
68. Basant MM. Revision Notes in Psychiatry. 2014; 204.
69. Sun L, Yi B, Pan X, Wu L, Shang Z, Jia Y, et al. PTSD Symptoms and sleep quality of covid-19 patients during hospitalization: an observational study from two centers. *Nat Sci Sleep*. 2021;13:1519–31. <https://doi.org/10.2147/NSS.S317618>.
70. Nagarajan R, Krishnamoorthy Y, Basavarachar V, Dakshinamoorthy R. Prevalence of post-traumatic stress disorder among survivors of severe COVID-19 infections: a systematic review and meta-analysis. *J Affect Disord*. 2022;299:52–9. <https://doi.org/10.1016/j.jad.2021.11.040>.
71. Chen Y, Huang X, Zhang C, An Y, Liang Y, Yang Y, et al. Prevalence and predictors of posttraumatic stress disorder, depression and anxiety among hospitalized patients with coronavirus disease 2019 in China. *BMC Psychiatry*. 2021;21:80. <https://doi.org/10.1186/s12888-021-03076-7>.
72. Villarreal-Zegarra D, Torres-Puente R, Alfaro SO, Al-kassab-Córdova A, de Castro JR, Mezones-Holguín E. Spanish version of Jenkins Sleep Scale in physicians and nurses: psychometric properties from a Peruvian nationally representative sample. *J Psychosom Res*. 2022; 110759. <https://doi.org/10.1016/j.jpsychores.2022.110759>.
73. Lin C-Y, Broström A, Griffiths MD, Pakpour AH. Investigating mediated effects of fear of COVID-19 and COVID-19 misunderstanding in the association between problematic social media use, psychological distress, and insomnia. *Internet Interv*. 2020;21: 100345. <https://doi.org/10.1016/j.invent.2020.100345>.
74. Kuriyama K, Soshi T, Kim Y. Sleep deprivation facilitates extinction of implicit fear generalization and physiological response to fear. *Biol Psychiatry*. 2010;68:991–8. <https://doi.org/10.1016/j.biopsych.2010.08.015>.
75. Zou S, Liu Z-H, Yan X, Wang H, Li Y, Xu X, et al. Prevalence and correlates of fatigue and its association with quality of life among clinically stable older psychiatric patients during the COVID-19 outbreak: a cross-sectional study. *Glob Health*. 2020;16:119. <https://doi.org/10.1186/s12992-020-00644-6>.
76. Medeiros G, Beach S. Exacerbation of anxiety symptoms in the setting of COVID-19 pandemic: An overview and clinically-useful recommendations. 2021;48: 69–70. <https://doi.org/10.15761/0101-60830000000281>.
77. Caycho-Rodríguez T, Tomás JM, Vilca LW, Carbajal-León C, Cervigni M, Gallegos M, et al. Socio-Demographic Variables, Fear of COVID-19, Anxiety, and Depression: Prevalence, Relationships and Explanatory Model in the General Population of Seven Latin American Countries. *Front Psychol*. 2021;12. Available: <https://www.frontiersin.org/article/10.3389/fpsyg.2021.695989>.
78. Vilca LW, Chávez BV, Fernández YS, Caycho-Rodríguez T, White M. Impact of the fear of catching COVID-19 on mental health in undergraduate students: A Predictive Model for anxiety, depression, and insomnia. *Curr Psychol N B NJ*. 2022; 1–8. <https://doi.org/10.1007/s12144-021-02542-5>.
79. Mazza MG, De Lorenzo R, Conte C, Poletti S, Vai B, Bollettini I, et al. Anxiety and depression in COVID-19 survivors: Role of inflammatory and clinical predictors. *Brain Behav Immun*. 2020;89:594–600. <https://doi.org/10.1016/j.bbi.2020.07.037>.
80. Milrad SF, Hall DL, Jutagir DR, Lattie EG, Czaja SJ, Perdomo DM, et al. Depression, evening salivary cortisol and inflammation in chronic fatigue syndrome: a psychoneuroendocrinological structural regression model. *Int J Psychophysiol*. 2018;131:124–30. <https://doi.org/10.1016/j.ijpsycho.2017.09.009>.
81. Xu L, Pan Q, Lin R. Prevalence rate and influencing factors of preoperative anxiety and depression in gastric cancer patients in China: Preliminary study. *J Int Med Res*. 2016;44:377–88. <https://doi.org/10.1177/0300060515616722>.
82. Liegey JS, Sagnier S, Debruxelles S, Poli M, Olindo S, Renou P, et al. Influence of inflammatory status in the acute phase of stroke on post-stroke depression. *Rev Neurol (Paris)*. 2021;177:941–6. <https://doi.org/10.1016/j.neurol.2020.11.005>.
83. Najjar S, Pearlman DM, Alper K, Najjar A, Devinsky O. Neuroinflammation and psychiatric illness. *J Neuroinflammation*. 2013;10:43. <https://doi.org/10.1186/1742-2094-10-43>.
84. Chakrabarti S. Mental health in hospitalised COVID 19 Patients in quarantine during second wave in a South Indian private teaching hospital. *J Multidiscip Healthc*. 2021;14:2777–89. <https://doi.org/10.2147/JMDH.S330812>.
85. Zeng F, Huang Y, Guo Y, Yin M, Chen X, Xiao L, et al. Association of inflammatory markers with the severity of COVID-19: A meta-analysis. *Int J Infect Dis*. 2020;96:467–74. <https://doi.org/10.1016/j.ijid.2020.05.055>.
86. Anyan F, Worsley L, Hjemdal O. Anxiety symptoms mediate the relationship between exposure to stressful negative life events and depressive symptoms: a conditional process modelling of the protective effects of resilience. *Asian J Psychiatry*. 2017;29:41–8. <https://doi.org/10.1016/j.ajp.2017.04.019>.
87. Nima AA, Rosenberg P, Archer T, Garcia D. Anxiety, affect, self-esteem, and stress: mediation and moderation effects on depression. *PLoS ONE*. 2013;8:e73265. <https://doi.org/10.1371/journal.pone.0073265>.
88. Kessler RC, Wang PS. The descriptive epidemiology of commonly occurring mental disorders in the United States. *Annu Rev Public Health*. 2008;29:115–29. <https://doi.org/10.1146/annurev.publhealth.29.020907.090847>.
89. Janiri D, Carfi A, Kotzalis GD, Bernabei R, Landi F, Sani G, et al. Posttraumatic stress disorder in patients after severe COVID-19 infection. *JAMA Psychiat*. 2021;78:567–9. <https://doi.org/10.1001/jamapsychiatry.2021.0109>.
90. Ripon RK, Mim SS, Puente AE, Hossain S, Babor MdMH, Sohan SA, et al. COVID-19: psychological effects on a COVID-19 quarantined population in Bangladesh. *Heliyon*. 2020;6: e05481. <https://doi.org/10.1016/j.heliyon.2020.e05481>.
91. Ketcheson Felicia, St C, King Lisa, Don R. Influence of PTSD and MDD on somatic symptoms in treatment-seeking military members and Veterans. *J Mil Veteran Fam Health*. 2018 [cited 9 Feb 2022]. <https://doi.org/10.3138/jmvfh.2017-0029>.
92. McFarlane ACAO, Graham K. The ambivalence about accepting the prevalence somatic symptoms in PTSD: Is PTSD a somatic disorder? *J Psychiatr Res*. 2021;143:388–94. <https://doi.org/10.1016/j.jpsychores.2021.09.030>.
93. Ipsos. One year of COVID-19. 2021. Available: https://www.ipsos.com/sites/default/files/ct/news/documents/2021-04/wef_-_expectations_about_when_life_will_return_to_pre-covid_normal_final.pdf.
94. Antiporta DA, Cutipé YL, Mendoza M, Celentano DD, Stuart EA, Bruni A. Depressive symptoms among Peruvian adult residents amidst a National Lockdown during the COVID-19 pandemic. *BMC Psychiatry*. 2021;21:111. <https://doi.org/10.1186/s12888-021-03107-3>.
95. Kim J, Seo YE, Sung HK, Park HY, Han MH, Lee SH. Predictors of the Development of mental disorders in Hospitalized COVID-19 Patients without Previous psychiatric history: a single-center retrospective study in South

- Korea. *Int J Environ Res Public Health*. 2022;19:1092. <https://doi.org/10.3390/ijerph19031092>.
96. Kim J-W, Stewart R, Kang S-J, Jung S-I, Kim S-W, Kim J-M. Telephone based Interventions for Psychological Problems in Hospital Isolated Patients with COVID-19. *Clin Psychopharmacol Neurosci*. 2020;18:616–20. <https://doi.org/10.9758/cpn.2020.18.4.616>.
97. Tausch A, e Souza RO, Viciano CM, Cayetano C, Barbosa J, Hennis AJ. Strengthening mental health responses to COVID-19 in the Americas: A health policy analysis and recommendations. *Lancet Reg Health - Am*. 2022;5: 100118. <https://doi.org/10.1016/j.lana.2021.100118>.
98. Freyhofer S, Ziegler N, de Jong EM, Schippers MC. Depression and anxiety in times of COVID-19: how coping strategies and loneliness relate to mental health outcomes and academic performance. *Front Psychol*. 2021;12:682684. <https://doi.org/10.3389/fpsyg.2021.682684>.
99. Ouanes S, Al-Amin H, Hussein NB, Khan F, Al Shahrani A, David P, et al. Physical and psychosocial well-being of hospitalized and non-hospitalized patients with covid-19 compared to the general population in Qatar. *Front Psychiatry*. 2021;12:792058. <https://doi.org/10.3389/fpsyg.2021.792058>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

