

SCIENTIFIC REPORTS



OPEN

A novel recombinant 6A β 15-THc-C chimeric vaccine (rCV02) mitigates Alzheimer's disease-like pathology, cognitive decline and synaptic loss in aged 3 \times Tg-AD mice

Received: 30 January 2016

Accepted: 13 May 2016

Published: 03 June 2016

Yun-Zhou Yu^{1,*}, Si Liu^{1,*}, Hai-Chao Wang^{1,2}, Dan-Yang Shi¹, Qing Xu², Xiao-Wei Zhou¹, Zhi-Wei Sun¹ & Pei-Tang Huang¹

Alzheimer's disease (AD) is a neurodegenerative disorder that impairs memory and cognition. Targeting amyloid- β (A β) may be currently the most promising immunotherapeutic strategy for AD. In this study, a recombinant chimeric 6A β 15-THc-C immunogen was formulated with alum adjuvant as a novel A β B-cell epitope candidate vaccine (rCV02) for AD. We examined its efficacy in preventing the cognitive deficit and synaptic impairment in 3 \times Tg-AD mice. Using a toxin-derived carrier protein, the rCV02 vaccine elicited robust A β -specific antibodies that markedly reduced AD-like pathology and improved behavioral performance in 3 \times Tg-AD mice. Along with the behavioral improvement in aged 3 \times Tg-AD mice, rCV02 significantly decreased calpain activation concurrent with reduced soluble A β or oligomeric forms of A β , probably by preventing dynamin 1 and PSD-95 degradation. Our data support the hypothesis that reducing A β levels in rCV02-immunized AD mice increases the levels of presynaptic dynamin 1 and postsynaptic PSD-95 allowing functional recovery of cognition. In conclusion, this novel and highly immunogenic rCV02 shows promise as a new candidate prophylactic vaccine for AD and may be useful for generating rapid and strong A β -specific antibodies in AD patients with pre-existing memory Th cells generated after immunization with conventional tetanus toxoid vaccine.

Alzheimer's disease (AD) is characterized by senile plaques (SPs) and neurofibrillary tangles (NFTs). The onset and progression of AD is thought to be caused by the production and accumulation of excessive amyloid- β (A β) in the brain, which results in amyloid plaque deposition as a defining pathological hallmark, and ultimately leads to neuron loss, cognitive decline and brain atrophy^{1,2}. Human A β -directed active and passive immunization can effectively clear the cerebral A β load in various AD mouse models³⁻⁵ and human AD patients⁶⁻⁹. Furthermore, immunotherapeutic reduction of A β in the brain ameliorates AD-like behavioral symptoms in AD model mice and, in humans, immunotherapy with a monoclonal antibody directed at the mid-region of A β (Solanezumab) has also shown some beneficial cognitive effects in mildly affected AD patients¹⁰. Therefore, the removal or lowering of A β from the brain in patients with very early AD pathology or even in presymptomatic subjects could be an effective therapeutic measure; obviously, a safe active vaccine might be beneficial for such preventive treatments of AD^{11,12}.

Synapse loss occurs early in AD and accompanies A β accumulation; therefore, these characteristics are considered the best neuropathological correlates of cognitive decline¹³⁻¹⁶. Some therapeutic strategies for AD attenuate synaptic dysfunction and improve cognitive behavior in AD models¹⁷⁻²⁴. Given the remarkable recovery of cognition in AD models of targeted-A β immunotherapy, it is necessary to determine the molecular correlations associated with improvement. A recombinant chimeric 6A β 15-THc-C immunogen developed as protein vaccine for AD generated a robust anti-A β 42 antibody response, and attenuated A β pathology and cognitive deficits in the PDAPP^{V7171} mouse model²⁵. However, the potential of this treatment to rescue synaptic dysfunction in preclinical

¹Beijing Institute of Biotechnology, Beijing 100071, China. ²Institute of Life Science and Biotechnology, Beijing Jiaotong University, Beijing 100044, China. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to Y.-Z.Y. (email: yunzhouyu@163.com)




Figure 1. rCV02 induces strong Th2-polarized A β -specific humoral immune responses and activates proliferation of THc-specific T cells in 3 \times Tg-AD mice. (A) Titer of anti-A β 42 antibodies in 3 \times Tg-AD mice immunized with rCV02. The titer was determined for the indicated IgM and IgG isotypes (IgG1, IgG2a, IgG2b, and IgG3). (B) T cell proliferation in rCV02-immunized mice. Splenocytes were harvested from rCV02-immunized and control mice and restimulated *in vitro* with 10 μ g/mL A β 42 and THc. Cytokine production from splenocytes was used as a surrogate marker of Th1 (IFN- γ ; (E)) and Th2 (IL-4; (D)) bias in the immune response to rCV02. IL-4 and IFN- γ levels were measured by ELISA. Data represent the mean \pm SD (n = 8). Statistically significant differences were determined by Student's *t*-test. * $p < 0.05$, ** $p < 0.01$, compared with the control group (3 \times Tg-AD).

models of AD remains to be clarified. In this study, this recombinant chimeric 6A β 15-THc-C immunogen was formulated with alum adjuvant as a novel A β B-cell epitope candidate vaccine (rCV02). We performed a comprehensive evaluation of its efficacy for the prevention of the cognitive deficit and synaptic impairment in 3 \times Tg-AD mice. Furthermore, we sought to determine the molecular correlations between the recovery of cognition and the improvement of synaptic functions. Moreover, the immune mechanism associated with rCV02 vaccination with the aid of a toxin-derived carrier was defined in 3 \times Tg-AD mice.

Results

The immunogenicity of rCV02 in 3 \times Tg-AD mice. To evaluate the immune response to the rCV02, the humoral and cellular immune responses were analyzed in experimental and control 3 \times Tg-AD mice. As shown in Fig. 1A, high levels of A β -specific IgG antibodies were induced in the rCV02-immunized mice following multiple immunizations (2, 3, or 4). Lymphocyte proliferative responses showed that rCV02 induced THc-specific responses, but not A β -specific T cell immunity in immunized mice (Fig. 1B, $p < 0.05$) and significantly increased IL-4 (Th2) or IFN- γ (Th1) cytokine levels induced by THc but not by A β 42 (Fig. 1C,D, $p < 0.01$). Furthermore, immunization with rCV02 resulted mainly in IgG1 and IgG2b isotypes, which are associated with non-inflammatory Th2 responses. These results indicate that the rCV02 can induce robust anti-inflammatory Th2-polarized immune responses without activating self-reactive A β 42-specific T cells.

Having demonstrated that rCV02 elicited a robust immune response, we tested its immunogenicity in prophylactic and therapeutic treatment models (supplementary Figure S1). As a prophylactic treatment, immunization with rCV02 generated a strong A β -specific antibody response after two immunizations; all antibody titers were further increased following a booster (Fig. 2A). The antibody levels remained high over a 7-month period after the last booster (five immunizations).

As a therapeutic treatment, rCV02 generated a relatively low antibody response after the initial three immunizations, while the antibody response was markedly augmented by two boosters. However, the high level of A β -specific IgG antibodies could not be sustained over a 4-month period after the last booster (Fig. 2B).

Immunization with rCV02 restores cognitive memory in aged 3 \times Tg-AD mice. The spatial learning and memory abilities of 3 \times Tg-AD and wild-type (C57/BL6) mice were evaluated using the Morris water maze test. In the prophylactic treatment, the latency to reach the platform for the rCV02-immunized AD mice was significantly decreased on days 6–7 of training (Fig. 3A, $p < 0.01$ and 0.001, compared to the control AD mice) as age-matched non-transgenic C57/BL6. Notably, a decrease trend in this latency was similar to a group of non-transgenic mice of the same age. In the probe trials for spatial memory ability, significantly shorter initial latency to reach the platform and significant increased number of correct platform location crosses were observed in the rCV02-immunized AD mice when compared to the control AD mice ($p < 0.05$, Fig. 3B,C).




Figure 2. Duration of rCV02-induced immunity in 3 \times Tg-AD mice following prophylactic or therapeutic administration. Serum samples from 3 \times Tg-AD mice immunized with rCV02 were collected at the time-points indicated (Supplementary Figure S1) and the anti-A β 42 IgG titers were analyzed by ELISA. Serum samples from individual mice were assayed, and the geometric mean titer (GMT) was calculated for each group ($n = 8$). The serum antibody titer was monitored for 14 months for the prophylactic treatment (A) and 7 months for the therapeutic treatment (B). The x-axis indicates the age of the mice at the time of the blood sampling (months). The solid arrows indicate the time of rCV02 vaccine administration.

In the therapeutic treatment, the rCV02-immunized AD mice also showed significantly shorter escape latencies on days 6–7 compared with the control AD mice ($p < 0.05$ and $p < 0.01$, Fig. 3D). In spatial memory testing, the rCV02-immunized AD mice had a significantly shorter initial latency to reach the platform, and a decrease in the number of correct platform location crosses was observed, while the effect was not statistically significant (Fig. 3E,F). These data indicate that immunization with rCV02 generates effective prophylactic or therapeutic effects on the cognitive memory of aged 3 \times Tg-AD mice and restores their cognitive ability.

Immunization with rCV02 mitigates the development of AD-like pathology in aged 3 \times Tg-AD mice.

To better understand the recovery of cognition in the rCV02-immunized mice, we sought to determine the molecular correlations associated with improvement. We first assessed whether rCV02 mitigated AD-like pathology in the brains of vaccinated 3 \times Tg-AD mice (Figs 4–6). In both the prophylactic and therapeutic setting, immunization with rCV02 significantly reduced the number of visible plaques and the total percentage of the brain area occupied by plaques (Fig. 4A–C). Compared to the control AD mice, the A β plaque load in the hippocampus of the rCV02-immunized mice was reduced by 68.6% and 65.1% in the prophylactic and therapeutic models, respectively (Fig. 4D). Prophylactic treatment significantly reduced the levels of insoluble and soluble A β 42 peptide in the brains of immunized mice compared to control mice (Fig. 5A). We also detected a significant reduction in soluble A β 42 concentrations following therapeutic treatment (Fig. 5C).

The A β oligomeric species present in soluble fractions of brain homogenates of vaccinated and non-vaccinated 3 \times Tg-AD mice were also detected by Western blot analysis. Levels of 6E10 monoclonal antibody-reactive A β oligomers, primarily hexamers and nonamers (28 and 42 kDa, respectively), were significantly reduced by prophylactic and therapeutic administration of rCV02 (Fig. 6, $p < 0.05$). Thus, A β oligomers appeared to decrease in immunized mice in parallel with changes in plaque load and cerebral A β levels.

Moreover, we assessed the tau pathology in the brains of immunized 3 \times Tg-AD mice by immunohistochemistry and ELISA. A significant reduction in the number of HT-7 positive neurons of brains was observed in the brains of immunized mice compared to controls, which indicated that both prophylactic and therapeutic administration of rCV02 decreased total tau accumulation ($p < 0.01$, Supplementary Figure S2). The levels of insoluble and soluble total tau were also significantly reduced in aged 3 \times Tg-AD mice following prophylactic immunization (Fig. 5B), while it was not significantly reduced following therapeutic




Figure 3. Vaccination with rCV02 improves cognitive performance in aged 3 × Tg-AD mice following prophylactic and therapeutic administration. 3 × Tg-AD mice were immunized with rCV02 at 3 months (prophylactic; panels A–C) or 12 months (therapeutic; panels D–F) of age and subjected to the Morris water maze test at 19 months and 21 months of age, respectively. Animals were tested daily for 7 days. (A,D) The mean escape latency (time) to reach the platform on days 1–7 of the water test during the training trials. (B,E) The mean number of platform location crosses during the probe trial after the last training trial. (C,F) Initial latency (time) to reach the platform location in probe trials after the last training trial. Data represent the mean ± SD for each group (n = 6–8). Statistically significant differences were determined by ANOVA. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, compared with the control group (3 × Tg-AD).




Figure 4. Vaccination with rCV02 significantly reduced the A β plaques in brains in aged 3 \times Tg-AD mice. 3 \times Tg-AD mice were immunized with rCV02 at 3 months (prophylactic) or 12 months (therapeutic) of age. The brains were collected for evaluation at 19 months and 21 months of age, respectively. (A) Representative images showing the distribution of A β plaques (6E10 antibody staining) in the hippocampus. Scale bar, 300 μ m. (B–D) Quantification of A β plaques in the hippocampus. Vaccination with rCV02 significantly reduced the plaque count (B), percentage of brain area occupied by plaques (C), and A β load (D). Data represent the mean \pm SD for each group (n = 8). Statistically significant differences were determined by Student's *t*-test. ***p* < 0.01, ****p* < 0.001, compared with the control group (3 \times Tg-AD).

immunization (Fig. 5D). Thus, our results indicated that immunization with rCV02 markedly reduced the levels of A β and tau in the brains of aged 3 \times Tg-AD mice, and improved behavioral performance.




Figure 5. Vaccination with rCV02 reduced the level of Aβ42 and tau in the brains of 3 × Tg-AD mice. 3 × Tg-AD mice were immunized with rCV02 at 3 months (prophylactic; panels A and B) or 12 months (therapeutic; panels C,D) of age. The brains were collected for evaluation at 19 months and 21 months of age, respectively. (A,C) Soluble and insoluble Aβ42 levels in the brain were measured by sandwich ELISA. (B,D) Soluble and insoluble tau levels in the brain were measured by ELISA. Data represent the mean ± SE for each group (n = 8). Statistically significant differences were determined by Student's *t*-test. **p* < 0.05, ***p* < 0.01, ****p* < 0.001, compared with the control group (3 × Tg-AD).




Figure 6. rCV02 reduced the levels of soluble Aβ oligomeric species in the brains of 3 × Tg-AD mice. 3 × Tg-AD mice were immunized with rCV02 at 3 months (prophylactic; panel A) or 12 months (therapeutic; panels B) of age. The brains were collected for evaluation at 19 months and 21 months of age, respectively. The 6-, 9- and 12-mers (28, 42 and 56 kDa) of Aβ in soluble fractions of brain homogenates were detected by Western blot analysis using the 6E10 monoclonal antibody; representative images are shown (lanes 1–3, non-vaccinated control 3 × Tg-AD; lanes 4–6, rCV02-vaccinated 3 × Tg-AD). Densitometry calculations were performed using Image J and normalized to β-actin. Data represent the mean ± SD for each group. Statistically significant differences were determined by Student's *t*-test. **p* < 0.05, ***p* < 0.01, compared with the control group (3 × Tg-AD).




Figure 7. rCV02 decreased calpain activation in the brains of 3 × Tg-AD mice. 3 × Tg-AD mice were immunized with rCV02 at 3 months (prophylactic; panel A) or 12 months (therapeutic; panels B) of age. The brains were collected for evaluation at 19-months and 21-months of age, respectively. The spectrin content in the soluble fractions of the brain homogenates were detected by Western blot analysis; representative images are shown (lanes 1–3, non-vaccinated control 3 × Tg-AD; lanes 4–6, rCV01-vaccinated 3 × Tg-AD; lanes 6–9, non-vaccinated control C57/BL6). The graphs show the spectrin 150/240 kDa ratio in control and treated mice. Data represent the mean ± SE for each group. Statistically significant differences were determined by ANOVA. ** $p < 0.01$, compared with the control group (3 × Tg-AD).

Immunization with rCV02 decreases calpain activation and prevents the loss of synaptic proteins in aged 3 × Tg-AD mice. A β accumulation has been shown to induce calpain activation^{26–30}, which may lead to the degradation of dynamin 1, a protein involved in synaptic vesicle release, and postsynaptic density protein PSD-95, which is crucial for synapse maturation and plasticity^{31–33}. Given the remarkable recovery of cognition in the rCV02-immunized aged 3 × Tg-AD mice with reduced AD-like pathology, we further explored the molecular mechanisms associated with this improvement. We first determined whether administration of rCV02 decreased calpain activation in 3 × Tg-AD mice. Spectrin degradation is highly sensitive to calpain activation and considered an excellent marker of the activity of this protease^{26,29}. Western blot analysis of the immunoreactivity of soluble fractions of brain homogenates with a specific spectrin antibody showed a significant increase in the spectrin 150/240 kDa ratio in the brains of aged 3 × Tg-AD mice compared to age-matched C57/BL6 mice. Prophylactic or therapeutic administration of rCV02 prevented the increase in the spectrin 150/240 ratio in aged 3 × Tg-AD mice (Fig. 7A,B), which indicated that immunization with rCV02 decreases calpain activation in aged 3 × Tg-AD mice.

To examine the potential of rCV02 immunization to prevent the loss of synaptic proteins or to mitigate synaptic impairment, we assessed the levels of dynamin 1 and PSD-95 in the brains of mice. Compared to the age-matched C57/BL6 controls, the non-immunized 3 × Tg-AD mice had significantly reduced levels of both synaptic proteins in the brain. However, prophylactic or therapeutic administration of rCV02 prevented the significant decrease of dynamin 1 (Fig. 8A,B) and PSD-95 (Fig. 8C,D) in aged 3 × Tg-AD mice. Thus, along with the behavioral improvement, immunization with rCV02 significantly decreased calpain activation concurrent with reduced soluble A β or oligomeric forms of A β in aged 3 × Tg-AD mice. This effect may prevent dynamin 1 and PSD-95 degradation.

Immune mechanism underlying the use of a tetanus toxin fragment (THc-C) of rCV02 as carrier protein. The novel recombinant 6A β 15-THc-C chimeric vaccine (rCV02) was reconstructed by fusion of 6A β 15 with THc-C, which had been proposed as a carrier or molecular adjuvant^{34,35}. Here, the immune mechanism of THc-C of rCV02 as carrier protein was further defined in 3 × Tg-AD mice. As shown in Fig. 9A,B, immunization with rCV02 induced strong A β -specific humoral immune responses and activated THc-specific Th cell proliferation ($p < 0.05$, compared to 6A β 15 or control groups), but did not activate A β -specific Th cells in mice. In contrast, immunization with 6A β 15 or 6A β 15 + THc induced only very mild A β -specific antibody responses ($p < 0.001$, compared to rCV02). Thus, the presence of the foreign toxin fragment as a carrier protein in the rCV02 vaccine was critical for generating strong A β -specific antibodies without activating self-reactive A β 42-specific T cells.

We also evaluated the capacity of rCV02 to elicit a rapid and strong antibody response in THc-immunized old 3 × Tg-AD mice (Fig. 9C,D). Two or three boosters with rCV02 induced robust anti-A β antibody responses only in THc-immunized old 3 × Tg-AD mice with THc-specific pre-existing memory Th cells. Multiple immunizations with rCV02 still generated a relatively low antibody response in control mice. Moreover, a stronger Th cell response specific to THc was also detected in this group of mice with pre-existing memory Th cells compared with the responses generated in control mice (Fig. 9D, $p < 0.05$). Thus, these data indicated that immunization with rCV02 could activate pre-existing THc-specific memory Th cells, leading to rapid and strong THc-specific cellular responses and anti-A β antibody responses. Taken together, rCV02 induced strong A β -specific humoral




Figure 8. rCV02 increased the levels of synaptic proteins in the brains of $3 \times$ Tg-AD mice. $3 \times$ Tg-AD mice were immunized with rCV02 at 3 months (prophyllactic; panel **A** and **C**) or 12 months (therapeutic; panels **B** and **D**) of age. The brains were collected for evaluation at 19 months and 21 months of age, respectively. Levels of dynamamin 1 (**A,B**) and PSD-95 (**C,D**) in the soluble fractions of the brain homogenates were detected by Western blot analysis; representative images are shown (lanes 1–3, non-vaccinated control $3 \times$ Tg-AD; lanes, 4–6 rCV01-vaccinated $3 \times$ Tg-AD; lanes 6–9, non-vaccinated control C57/BL6). The graphs show the levels of dynamamin 1 (**A, B**) and PSD-95 (**C,D**) in control and treated mice. Data represent the mean \pm SE for each group. Statistically significant differences were determined by ANOVA. * $p < 0.05$, ** $p < 0.01$, compared with the control group ($3 \times$ Tg-AD).

immunity via the help provided by the foreign Th cells specific to the Th epitopes of THc-C and generated a rapid and strong A β -specific antibody response in aged $3 \times$ Tg-AD mice with pre-existing memory Th cells.

Discussion

The objective of this study was to undertake a comprehensive evaluation of the immunopotency of a novel clinical grade epitope immunogen 6A β 15–THc-C formulated with alum adjuvant (rCV02) in $3 \times$ Tg-AD mice and to determine the subsequent effects on A β -related pathologies and cognition. The recombinant chimeric 6A β 15–THc-C immunogen was developed as a novel A β B-cell epitope candidate vaccine composed of hexavalent foldable A β 1–15 fused with a tetanus toxin fragment (THc-C) as carrier protein or molecular adjuvant. This chimeric vaccine was designed to circumvent the problem of A β 42-specific T cell autoreactivity and to overcome tolerance induction to self-antigens. In this study, immunization with rCV02 induced a strong Th2-polarized anti-A β antibody response in $3 \times$ Tg-AD mice via the help of a toxin-derived carrier protein without the induction of auto-immune A β 42-specific T cell responses. Both prophylactic and therapeutic administration of rCV02 mitigated Alzheimer's disease-like pathology, cognitive decline and synaptic impairment in aged $3 \times$ Tg-AD mice.

A potential problem of AD immunotherapy highlighted in previous studies is that a reduction of insoluble A β or A β deposits may lead to increased levels of the soluble forms of this peptide³⁶, especially the most neurotoxic oligomers, which will impair synaptic and cognitive function^{16,37–39}. In a previous study, the antibodies elicited




Figure 9. Immune mechanism underlying the effects of the tetanus toxin fragment (THc-C) of rCV02 as a carrier protein in 3 × Tg-AD mice. Titer of anti-Aβ42 antibodies (A) and T cell proliferation (B) were analyzed in 3 × Tg-AD mice immunized with different vaccines. The rCV02 vaccine induced strong Aβ-specific humoral immunity via help from foreign Th cells specific to the Th epitopes of THc-C. Titer of anti-Aβ 42 antibodies (C) and T cell proliferation (D) were also analyzed in THc-immunized aged 3 × Tg-AD mice. Strong humoral and cellular immune responses to rCV02 were generated in THc-immunized aged 3 × Tg-AD mice. Data represent the mean ± SD (n = 8). Statistically significant differences were determined by ANOVA or Student's *t*-test. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, compared with 6Aβ15 or the control group.

by 6Aβ15-THc-C immunization bound to Aβ oligomers and inhibited Aβ42 oligomer-mediated neurotoxicity *in vitro*²⁵. This indicated the potential of based-rCV02 vaccination to mediate therapeutic effects in 3 × Tg-AD mice *in vivo*. Here, we show that prophylactic immunization with rCV02 reduces not only Aβ plaques, but also soluble Aβ peptide, including Aβ nonamers and hexamers. These findings are consistent with those of previous reports^{24,40–42}. The therapeutic effects of immunization with rCV02 in reducing the soluble Aβ are very important because the elicited anti-Aβ antibodies will clear or neutralize the oligomeric Aβ, producing beneficial cognitive effects.

Remarkably, in our study prophylactic immunization significantly reduced the total levels of insoluble and soluble tau in the brain in aged 3 × Tg-AD mice. While Aβ accumulation may be the primary event in AD pathogenesis and accelerate phosphorylation of tau or NFT formation^{43,44}, tau pathology involved in synaptic loss also plays an important role in disease progression^{12,45,46}. Previous studies have shown that Aβ immunotherapy decreases Aβ and tau pathology in 3 × Tg-AD mice^{42,47–50}, which is consistent with the results presented here. Coincidentally, a decrease in soluble tau levels has been reported in the CSF of immunized patients⁶ and in the brain of immunized 3 × Tg-AD mice⁴⁹. Therefore, our results suggest that the rCV02-mediated reduction in Aβ may facilitate the clearance of total tau-related pathology. Furthermore, previous reports indicate that a reduction in both soluble Aβ and tau is necessary or beneficial in synergistically ameliorating cognitive impairments^{49–50}.

In AD, cognitive decline is associated with synapse loss, which precedes neuron death. The neurotoxic effects of Aβ on the central synapses have been described previously and are reflected in the decrease levels of some synaptic proteins^{14,16,51}. Indeed, the levels of dynamin 1 and PSD-95 in unvaccinated 3 × Tg-AD mice were significantly reduced compared to those in age-matched non-transgenic controls (C57/BL6). It is well established in animal AD models that Aβ-induced synaptic injury or neuronal loss can be prevented by calpain inhibitors^{28,29} or Aβ-directed immunotherapy^{17,19,20,24}. Therefore, we hypothesized that immunization with rCV02 could decrease calpain activation and further protect synaptic function. In the current study, calpain activation was significantly decreased in the brains of immunized-rCV02 3 × Tg-AD mice. Along with the decreased calpain activity and Aβ levels, there were obviously increased levels of dynamin 1 and PSD-95. Thus, it is likely that the decreased calpain activity contributed to preventing degradation of dynamin 1 and PSD-95. These results were suggestive of a remarkable association between the levels of Aβ, cognitive function, and synaptic function in 3 × Tg-AD mice.

Taken together, these findings demonstrate that selective reduction of soluble A β or oligomers by immunotherapy with rCV02 is sufficient to prevent the loss of the synaptic proteins, allowing functional recovery of cognition. Since synaptic dysfunction is an important phenotypic manifestation of AD^{14,20,22,38}, it can be speculated that there may be a molecular link to AD pathogenesis that will allow dynamin 1 and PSD-95 to be useful as biomarkers for evaluating prospective treatments for AD.

In the design of the rCV02 vaccine, we selected the immunodominant B-cell epitope peptide of A β 42 as the immunogen and constructed a hexavalent foldable A β 1–15, with each peptide sequence separated by a GS small linker (6A β 15). This novel immunogen for AD was designed to mimic the assembly states of A β 42 and improve immunogenicity over that of short A β 1–15 peptide. We also replaced the T cell epitopes of A β 42 with the C fragment of TeNT (THc-C) with the aim of activating non-self pre-existing memory Th cells in the general human population with a conventional tetanus toxoid (TT) vaccine. Data from this study demonstrate that immunization with rCV02 produced robust anti-A β 42 antibody responses while breaking immune self-tolerance to the A β 42 self-antigen in 3 \times Tg-AD mice. It should be noted that the enhanced A β -specific antibody response was observed only when the 6A β 15 antigen was fused directly to the THc-C carrier protein. Thus, it can be speculated that the strong immunogenicity of rCV02 can be attributed to the fusion construct, in which the THc-C carrier protein acts as an efficient molecular adjuvant. Moreover, in presence of THc-specific memory Th cells, rCV02 mediated strong activation of pre-existing memory Th cells specific to the Th epitopes of this vaccine and led to a rapid and robust production of antibodies specific to the B-cell epitope (A β 1–15) of the same vaccine as previously reported⁵². Therefore, the immunogenic carrier protein THc-C in rCV02, containing strong universal CD4⁺T cell epitopes^{34,35}, can provide T cell help and activate pre-existing THc-specific memory Th cells for A β -specific antibody production.

The prophylactic treatment of 3-month-old 3 \times Tg-AD mice showed a trend toward the generation and maintenance of relatively stable and adequate antibody levels over a 7-month period after the last vaccination. The therapeutic treatment of 12-month-old 3 \times Tg-AD mice generated a relatively low antibody response and there was an obvious reduction in A β -specific antibody levels after the last vaccination. Lower levels of A β and tau and greater cognitive benefits were observed in the prophylactic model, which is consistent with previous reports of the correlation of high titers of therapeutic anti-A β antibodies with reduced A β pathology and improved cognitive ability in preclinical and clinical trials^{6,8}. The post-vaccination production of A β -specific antibodies over time may be considered a pharmacological response of the immune system to the vaccine. The generation of robust immunity was more difficult due to old age-associated hyporesponsiveness in elderly 3 \times Tg-AD mice. However, rCV02 immunization overcame hyporesponsiveness in 12-month-old 3 \times Tg-AD mice with THc-specific memory T cells and generated a rapid and strong A β -specific humoral response. Therefore, the immunological mechanism of action of rCV02 indicate this vaccine would be highly beneficial for inducing therapeutically potent anti-A β antibody responses in future clinical trials of middle-aged or elderly AD patients with pre-existing memory.

To avoid stimulating adverse immune responses, second-generation AD vaccines targeting the N-terminal epitopes of A β -specific B cells have been developed and are being tested in current clinical trials^{5,8,12}. In addition, universal CD4⁺ T cell epitopes have been identified within the C fragment of TeNT and other proteins, and their use as carrier proteins has been tested in both mice and humans^{35,53}. Elan and Wyeth designed an AD vaccine ACC-001, in which an N-terminal sequence of A β (A β 1–6) was conjugated to diphtheria toxin (DT) to provide foreign Th epitopes to overcome A β -associated hyporesponsiveness for A β -specific antibody production. MER5101 is also a similar A β 1–15:DT conjugate vaccine for AD⁵⁴. A recombinant protein vaccine (Lu AF20513) composed of two foreign Th cell epitopes from TT, P30, and P2 and three copies of the B-cell epitopes of A β 42 (A β 1–12) is currently being tested in phase I trials⁵². Lu AF20513 is expressed in *E. coli* and is found primarily in the inclusion bodies, while in this study a novel recombinant chimeric 6A β 15-THc-C antigen expressed in *E. coli* (BL21) in a fully soluble form was constructed and developed as the rCV02 vaccine for AD. Unlike the current ACC-001 or other vaccines in which an N-terminal A β sequence is conjugated to DT or other carriers, this type of recombinant protein vaccine carries the advantages of expected safety as well as ease of construction and large-scale production in a chemically homogeneous form. Moreover, these two recombinant protein vaccines may represent an effective and safe form of active immunotherapy that may overcome the A β and old age-associated hyporesponsiveness via the help of foreign Th cell epitopes from TT⁵².

In summary, we have comprehensively characterized the immunogenicity, efficacy, and mechanism of action of rCV02 in both prophylactic and therapeutic 3 \times Tg-AD mouse models. Our findings indicate the promise of rCV02 as a novel candidate vaccine for AD as well as the potential for inducing potent anti-A β antibody responses in AD patients with pre-existing memory Th cells specific to TT. After completion of preclinical safety and toxicity studies, human clinical trials of the rCV02 as a new prophylactic vaccine for AD will be initiated.

Methods

Preparation of rCV02. In this study, a novel recombinant chimeric 6A β 15-THc-C immunogen was expressed in *Escherichia coli* (BL21) in a fully soluble form²⁵ and purified through a series of chromatographic methods and buffer exchanges to yield the final formulation. We used the recombinant chimeric 6A β 15-THc-C immunogen as a novel A β B-cell epitope vaccine (rCV02) containing a hexavalent foldable A β 1–15 (6A β 15) fused to a tetanus toxin fragment (THc-C) as a carrier protein or molecular adjuvant. The vaccine for AD was formulated in a Th2-biased aluminum hydroxide adjuvant (Brenntag Biosector, Frederikssund, Denmark).

Immunization of 3 \times Tg-AD mice with rCV02. A colony of 3 \times Tg-AD homozygous mice harboring the human APP_{Swe}, PS1_{M146V} and Tau_{P301L} mutations were generated from breeding pairs obtained from Jackson Lab (Bar Harbor, ME, USA). All 3 \times Tg-AD mice were housed in a temperature- and light-cycle controlled animal facility at the Beijing Laboratory Animal Center (Beijing, China). All animal procedures were conducted with

the approval of the Beijing Institute of Biotechnology Institutional Animal Care and Use Committee and the methods were carried out in accordance with the approved guidelines. Eight $3 \times$ Tg-AD mice (aged 2 months, 4 females and 4 males) were immunized with $5 \mu\text{g}$ purified 6A β 15-THc-C immunogen formulated with 0.2% (w/w) AlhydrogelTM (Brenntag Biosector). Four injections were administered at 2-wk intervals ($100 \mu\text{l}$ /injection). PBS formulated with the adjuvant was used as a negative control.

To assess the protective effects of rCV02 in $3 \times$ Tg-AD mice, we established prophylactic (Supplementary Fig. S1A) and therapeutic (Supplementary Fig. S1B) models. The prophylactic group consisted of 8 mice (4 females and 4 males) that were aged 3 months at the time of immunization and did not have obvious behavioral impairment or amyloid plaques. The prophylactic group was immunized intramuscularly (i.m.) four times with $5 \mu\text{g}$ of 6A β 15-THc-C (rCV02) at 1-month intervals, with a final booster immunization administered 5 months after the fourth dose. The therapeutic group consisted of 8 mice (4 females and 4 males) that were aged 12 months at the time of immunization and had the typical behavioral impairment and amyloid plaque deposition. The therapeutic group was immunized i.m. three times with rCV02 at 1-month intervals, and received two booster immunization (the first 2 months after the third dose and the second 3 months later). Eight $3 \times$ Tg-AD mice (4 females and 4 males) receiving only vehicle was used as a negative control.

We performed two experiments in $3 \times$ Tg-AD mice to further assess the mechanism of the immune response associated with rCV02 vaccination. In one experiment, groups of eight mice (aged 3 months) were immunized three times at 1-month intervals with $5 \mu\text{g}$ of 6A β 15-THc-C (rCV02), 6A β 15²⁵, THc of tetanus toxin⁵⁵, or 6A β 15 + THc (a mixed vaccine composed of 6A β 15 and THc) formulated with AlhydrogelTM 0.2% (w/w). In another experiment, 3-month-old mice were prime-immunized twice at 1-month intervals with $5 \mu\text{g}$ of THc formulated with 0.2% (w/w) AlhydrogelTM. Control mice received the aluminum hydroxide adjuvant only. The 12-month-old mice from the THc-primed and control groups were boosted with 1, 2 or 3 doses of rCV02 at 1-month intervals after a resting period of 8 months. The humoral and cellular immune responses were analyzed in experimental and control mice.

Detection of antibody responses. ELISAs were used to determine the levels of anti-A β 42 antibodies in sera from mice in the different treatment groups as previously described²⁵. Briefly, the sera were serially diluted (from 100-fold) before testing. The total IgG and the IgG isotype antibody responses were detected using horseradish peroxidase (HRP)-conjugated anti-mouse antibodies (Santa Cruz Biotechnology, Inc., Santa Cruz, CA, USA) at a dilution of 1:2,000. After incubation with the secondary antibody, reactivity was visualized by adding $100 \mu\text{l}$ of citrate buffer (pH 5.0) containing 0.04% (w/v) o-phenylenediamine and 0.02% (v/v) hydrogen peroxide for 10 min at room temperature. The reaction was stopped by the addition of $50 \mu\text{l}$ of 2 M H₂SO₄, and the absorbance was read at 492 nm using a Thermo Labsystems microplate reader (Franklin, MA, USA). Serum samples from individual mice were assayed, and the geometric mean titer (GMT) was calculated for each group ($n = 8$).

Lymphocyte proliferative responses and evaluation of cytokine levels. Cell suspensions of splenocytes from immunized mice were prepared and treated as previously described²⁵. The proliferative responses of splenocytes were determined using the cell counting kit-8 (CCK-8; Dojindo Molecular Technologies, Kumamoto, Japan) according to the manufacturer's protocol. The levels of mouse IL-4 and IFN- γ produced by the splenocytes were assayed using commercial ELISA kits according to the manufacturer's protocol (Bender Medsystems, Vienna, Austria). Data are presented as means \pm SD of $n = 8$ samples.

Behavioral test. The Morris water maze was used for behavioral tests as previously described²⁵. Briefly, the $3 \times$ Tg-AD mice aged 19 months and 21 months that had received five vaccinations were trained to swim in the Morris water maze apparatus. Age-matched non-transgenic C57/BL6 mice were used as controls ($n = 6$, 3 females and 3 males). For each trial, the mouse was released into the tank at one of four designated start locations and allowed to find and escape onto the hidden platform. During the learning period each mouse was subjected to a daily four-trial session for seven consecutive days. The escape latency was recorded as the time from being put into the water to climbing the escape platform during training. Retention (probe trial) of spatial learning was assessed 24 h after the last training trial. The platform was removed from the pool, and each mouse was subjected to one 60 s swim probe trial. The parameters measured during the probe trial were the initial latency to reach the platform location and number of platform location crosses.

Immunohistochemistry and semi-quantitative image analysis. After the Morris water maze test, brains were removed from the $3 \times$ Tg-AD mice aged 19 months and 21 months and divided sagittally along the interhemispheric fissure; half of the brain was used for immunohistochemical analysis. Immunohistochemical analysis and quantification of the stained sections were conducted as previously described^{25,56}. The A β deposits were detected using the anti-human A β monoclonal antibody 6E10 (1:1,000, Covance, Emeryville, CA, USA) and general tau pathology was detected with HT7 (1:40, Thermo scientific, Waltham, MA, USA), recognizing epitopes 159–163.

Determination of A β and tau levels in soluble/insoluble fractions. The other half of the brain collected following the Morris water maze test was homogenized in Tris-buffered saline (TBS) buffer containing a protease inhibitor cocktail (Roche, Mannheim, Germany). The samples were ultrasonicated and then centrifuged at $100,000 \times g$ for 2 h at 4 °C. The supernatant was collected to detect soluble A β 42 or tau. The sediment was treated with 5 M guanidine buffer to solubilize the A β or tau. Biosource ELISA kits (Invitrogen, Carlsbad, CA, USA) were used to detect A β 42 and total tau levels in the brain extracts. A β levels were calculated as ng/g brain (wet weight).

Western blot analysis. Western blot analyses were performed to determine the levels of A β oligomers, dynamin 1, spectrin, and PSD-95 in the soluble fractions of the brain homogenates from immunized, control (3 \times Tg-AD), and wild-type (C57/BL6) mice. Protein samples in TBS buffer were separated by SDS-PAGE (10–12% gels), and transferred onto a PVDF membrane. The membrane was blocked with 10% non-fat milk in 20 mM Tris-HCl (pH 7.4) containing 150 mM NaCl and 0.05% Tween 20 (TBS-T). After washing with TBS-T, the membrane was incubated with the following primary antibodies: anti-human A β monoclonal antibody 6E10 (1:1,500), anti-dynamin 1 (C16, 1:2,000, Santa Cruz Biotechnology, Inc.), anti-spectrin (AA6, 1:1,500, Millipore, MA, USA), or anti-PSD-95 (1:500, Invitrogen), or anti- β -actin/tubulin (Sigma, MO, USA) for 1 h at 37 °C. The membrane was incubated with a HRP-conjugated IgG (GIBCO, WA, USA) secondary detection antibody (1: 5,000) for 0.5 h at 37 °C and protein immunoreactivity was visualized using enhanced chemiluminescence reagents (ECL, Pierce, IL, USA). Proteins were quantified by densitometry using the Image J Software and normalized to β -actin/tubulin. The values obtained in untreated controls were considered to represent 100% expression. Values represent the means \pm SE for each group (n = 6–8).

Statistical analysis. All statistical analyses were performed using the SPSS software (version 16.0, Chicago, IL, USA). Statistically significant differences between groups were determined using the one-way analysis of variance (ANOVA) or Student's *t*-test. For all tests, *P*-values < 0.05 were considered to indicate statistical significance.

References

- Gilbert, B. J. The role of amyloid beta in the pathogenesis of Alzheimer's disease. *J Clin Pathol* **66**, 362–366 (2013).
- Pimplikar, S. W. Reassessing the amyloid cascade hypothesis of Alzheimer's disease. *Int J Biochem Cell Biol* **41**, 1261–1268 (2009).
- Wisniewski, T. & Boutajangout, A. Immunotherapeutic approaches for Alzheimer's disease in transgenic mouse models. *Brain Struct Funct* **214**, 201–218 (2010).
- Wisniewski, T. & Sigurdsson, E. M. Murine models of Alzheimer's disease and their use in developing immunotherapies. *Biochim Biophys Acta* **1802**, 847–859 (2010).
- Lannfelt, L., Relkin, N. R. & Siemers, E. R. Amyloid- β -directed immunotherapy for Alzheimer's disease. *J Intern Med* **275**, 284–295 (2014).
- Gilman, S. *et al.* Clinical effects of Abeta immunization (AN1792) in patients with AD in an interrupted trial. *Neurology* **64**, 1553–1562 (2005).
- Lemere, C. A. & Masliah, E. Can Alzheimer disease be prevented by amyloid-beta immunotherapy? *Nat Rev Neurol* **6**, 108–119 (2010).
- Menendez-Gonzalez, M., Perez-Pinera, P., Martinez-Rivera, M., Muniz, A. L. & Vega, J. A. Immunotherapy for Alzheimer's disease: rational basis in ongoing clinical trials. *Curr Pharm Des* **17**, 508–520 (2011).
- Panza, F., Logroscino, G., Imbimbo, B. P. & Solfrizzi, V. Is there still any hope for amyloid-based immunotherapy for Alzheimer's disease? *Curr Opin Psychiatry* **27**, 128–137 (2014).
- Siemers, E. R. *et al.* Phase 3 solanezumab trials: Secondary outcomes in mild Alzheimer's disease patients. *Alzheimers Dement* **12**, 110–120 (2016).
- Carrillo, M. C. *et al.* Can we prevent Alzheimer's disease? Secondary “prevention” trials in Alzheimer's disease. *Alzheimers Dement* **9**, 123–131 (2013).
- Panza, F. *et al.* Amyloid-based immunotherapy for Alzheimer's disease in the time of prevention trials: the way forward. *Expert Rev Clin Immunol* **10**, 405–419 (2014).
- Jacobsen, J. S. *et al.* Early-onset behavioral and synaptic deficits in a mouse model of Alzheimer's disease. *Proc. Natl. Acad. Sci. USA* **103**, 5161–5166 (2006).
- Leuba, G. *et al.* Pathological reorganization of NMDA receptors subunits and postsynaptic protein PSD-95 distribution in Alzheimer's disease. *Behav Brain Res* **11**, 86–96 (2014).
- Pozueta, J., Lefort, R. & Shelanski, M. L. Synaptic changes in Alzheimer's disease and its models. *Neuroscience* **251**, 51–65 (2013).
- Selkoe, D. J. Soluble oligomers of the amyloid beta-protein impair synaptic plasticity and behavior. *Behav Brain Res* **192**, 106–113 (2008).
- Buttini, M. *et al.* Beta-amyloid immunotherapy prevents synaptic degeneration in a mouse model of Alzheimer's disease. *J Neurosci* **25**, 9096–9101 (2005).
- Chen, Y. *et al.* Hsp90 chaperone inhibitor 17-AAG attenuates Abeta-induced synaptic toxicity and memory impairment. *J Neurosci* **34**, 2464–2470 (2014).
- Dorostkar, M. M. *et al.* Immunotherapy alleviates amyloid-associated synaptic pathology in an Alzheimer's disease mouse model. *Brain* **137**, 3319–3326 (2014).
- Hillen, H. *et al.* Generation and therapeutic efficacy of highly oligomer-specific beta-amyloid antibodies. *J Neurosci* **30**, 10369–10379 (2010).
- Inestrosa, N. C. *et al.* Tetrahydroperforin prevents cognitive deficit, Abeta deposition, tau phosphorylation and synaptotoxicity in the APP^{swE}/PSEN1^{DeltaE9} model of Alzheimer's disease: a possible effect on APP processing. *Transl Psychiatry* **1**, e20 (2011).
- Liu, X. *et al.* Long-term treatment with Ginkgo biloba extract EGb 761 improves symptoms and pathology in a transgenic mouse model of Alzheimer's disease. *Brain Behav Immun* **46**, 121–131 (2015).
- Nistico, R., Pignatelli, M., Piccinin, S., Mercuri, N. B. & Collingridge, G. Targeting synaptic dysfunction in Alzheimer's disease therapy. *Mol neurobiol* **46**, 572–587 (2012).
- St-Amour, I. *et al.* IVIg protects the 3 \times Tg-AD mouse model of Alzheimer's disease from memory deficit and Abeta pathology. *J Neuroinflammation* **11**, 54 (2014).
- Yu, Y. Z. *et al.* Strikingly reduced amyloid burden and improved behavioral performance in Alzheimer's disease mice immunized with recombinant chimeric vaccines by hexavalent foldable A β ₁₋₁₅ fused to toxin-derived carrier proteins. *J Alzheimers Dis* **41**, 243–260 (2014).
- Czogalla, A. & Sikorski, A. F. Spectrin and calpain: a ‘target’ and a ‘sniper’ in the pathology of neuronal cells. *Cell Mol Life Sci* **62**, 1913–1924 (2005).
- Getz, G. S. Calpain inhibition as a potential treatment of Alzheimer's disease. *Am J Pathol* **181**, 388–391 (2012).
- Medeiros, R. *et al.* Calpain inhibitor A-705253 mitigates Alzheimer's disease-like pathology and cognitive decline in aged 3 \times TgAD mice. *Am J Pathol* **181**, 616–625 (2012).
- Sinjoanu, R. C. *et al.* The novel calpain inhibitor A-705253 potently inhibits oligomeric beta-amyloid-induced dynamin 1 and tau cleavage in hippocampal neurons. *Neurochem Int* **53**, 79–88 (2008).
- Tu, S., Okamoto, S., Lipton, S. A. & Xu, H. Oligomeric Abeta-induced synaptic dysfunction in Alzheimer's disease. *Mol Neurodegener* **9**, 48 (2014).
- Fa, M., Staniszewski, A., Saeed, F., Francis, Y. I. & Arancio, O. Dynamin 1 is required for memory formation. *Plos one* **9**, e91954, (2014).

32. Savioz, A., Leuba, G. & Vallet, P. G. A framework to understand the variations of PSD-95 expression in brain aging and in Alzheimer's disease. *Ageing Res Rev* **18**, 86–94 (2014).
33. Shao, C. Y., Mirra, S. S., Sait, H. B., Sacktor, T. C. & Sigurdsson, E. M. Postsynaptic degeneration as revealed by PSD-95 reduction occurs after advanced Abeta and tau pathology in transgenic mouse models of Alzheimer's disease. *Acta Neuropathol* **122**, 285–292 (2011).
34. Baraldo, K. *et al.* N19 polypeptide as a carrier for enhanced immunogenicity and protective efficacy of meningococcal conjugate vaccines. *Infect Immun* **72**, 4884–4887 (2004).
35. Diethelm-Okita, B. M., Okita, D. K., Banaszak, L. & Conti-Fine, B. M. Universal epitopes for human CD4+ cells on tetanus and diphtheria toxins. *J Infect Dis* **181**, 1001–1009 (2000).
36. Patton, R. L. *et al.* Amyloid-beta peptide remnants in AN-1792-immunized Alzheimer's disease patients: a biochemical analysis. *Am J Pathol* **169**, 1048–1063 (2006).
37. Sokolow, S. *et al.* AD synapses contain abundant Abeta monomer and multiple soluble oligomers, including a 56-kDa assembly. *Neurobiol Aging* **33**, 1545–1555 (2012).
38. Viola, K. L. & Klein, W. L. Amyloid beta oligomers in Alzheimer's disease pathogenesis, treatment, and diagnosis. *Acta Neuropathol* **129**, 183–206 (2015).
39. Zahs, K. R. & Ashe, K. H. beta-Amyloid oligomers in aging and Alzheimer's disease. *Front Aging Neurosci* **5**, 28 (2013).
40. Gimenez-Llort, L., Rivera-Hernandez, G., Marin-Argany, M., Sanchez-Quesada, J. L. & Villegas, S. Early intervention in the 3 × Tg-AD mice with an amyloid beta-antibody fragment ameliorates first hallmarks of Alzheimer disease. *MAbs* **5**, 665–677 (2013).
41. Movsesyan, N. *et al.* Reducing AD-like pathology in 3 × Tg-AD mouse model by DNA epitope vaccine - a novel immunotherapeutic strategy. *PLoS one* **3**, e2124 (2008).
42. Oddo, S. *et al.* Temporal profile of amyloid-beta (Abeta) oligomerization in an *in vivo* model of Alzheimer disease. A link between Abeta and tau pathology. *J Bio Chem* **281**, 1599–1604 (2006).
43. Chabrier, M. A. *et al.* Soluble abeta promotes wild-type tau pathology *in vivo*. *J Neurosci* **32**, 17345–17350 (2012).
44. Oddo, S. *et al.* Triple-transgenic model of Alzheimer's disease with plaques and tangles: intracellular Abeta and synaptic dysfunction. *Neuron* **39**, 409–421 (2003).
45. Ballatore, C., Lee, V. M. & Trojanowski, J. Q. Tau-mediated neurodegeneration in Alzheimer's disease and related disorders. *Nature reviews. Neuroscience* **8**, 663–672 (2007).
46. Takashima, A. Tau aggregation is a therapeutic target for Alzheimer's disease. *Curr Alzheimer Res* **7**, 665–669 (2010).
47. Minami, S. S. *et al.* Therapeutic versus neuroinflammatory effects of passive immunization is dependent on Abeta/amyloid burden in a transgenic mouse model of Alzheimer's disease. *J Neuroinflammation* **7**, 57 (2010).
48. Oddo, S., Billings, L., Kesslak, J. P., Cribbs, D. H. & LaFerla, F. M. Abeta immunotherapy leads to clearance of early, but not late, hyperphosphorylated tau aggregates via the proteasome. *Neuron* **43**, 321–332 (2004).
49. Oddo, S. *et al.* Reduction of soluble Abeta and tau, but not soluble Abeta alone, ameliorates cognitive decline in transgenic mice with plaques and tangles. *J Bio Chem* **281**, 39413–39423 (2006).
50. Rasool, S., Martinez-Coria, H., Wu, J. W., LaFerla, F. & Glabe, C. G. Systemic vaccination with anti-oligomeric monoclonal antibodies improves cognitive function by reducing Abeta deposition and tau pathology in 3 × Tg-AD mice. *J Neurochem* **126**, 473–482 (2013).
51. Meli, G. *et al.* Conformational targeting of intracellular Abeta oligomers demonstrates their pathological oligomerization inside the endoplasmic reticulum. *Nat Commun* **5**, 3867 (2014).
52. Davtyan, H. *et al.* Immunogenicity, efficacy, safety, and mechanism of action of epitope vaccine (Lu AF20513) for Alzheimer's disease: prelude to a clinical trial. *J Neurosci* **33**, 4923–4934 (2013).
53. Savelyeva, N., King, C. A., Vitetta, E. S. & Stevenson, F. K. Inhibition of a vaccine-induced anti-tumor B cell response by soluble protein antigen in the absence of continuing T cell help. *Proc Natl Acad Sci USA* **102**, 10987–10992 (2005).
54. Liu, B. *et al.* MER5101, a novel Abeta1-15:DT conjugate vaccine, generates a robust anti-Abeta antibody response and attenuates Abeta pathology and cognitive deficits in APP^{swE}/PS1^{DeltaE9} transgenic mice. *J Neurosci* **33**, 7027–7037 (2013).
55. Yu, Y. Z. *et al.* Co-expression of tetanus toxin fragment C in *Escherichia coli* with thioredoxin and its evaluation as an effective subunit vaccine candidate. *Vaccine* **29**, 5978–5985 (2011).
56. Liu, S. *et al.* Co-immunization with DNA and protein mixture: a safe and efficacious immunotherapeutic strategy for Alzheimer's disease in PDAPP mice. *Sci Rep* **5**, 7771 (2015).

Acknowledgements

This work was supported by grants from the Major National Science and Technology projects (2012ZX0901003-001-005), and the Natural Science Foundation of Beijing (7122130). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author Contributions

Y.Z.Y. and Q.X. conceived and designed the experiments. S.L., H.C.W., D.Y.S. and Y.Z.Y. performed the experiments. S.L., H.C.W. and Y.Z.Y. analyzed the data. X.W.Z., Z.W.S. and P.T.H. contributed reagents/materials/analysis tools. S.L. and Y.Z.Y. wrote the paper.

Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Yu, Y.-Z. *et al.* A novel recombinant 6Aβ15-THc-C chimeric vaccine (rCV02) mitigates Alzheimer's disease-like pathology, cognitive decline and synaptic loss in aged 3 × Tg-AD mice. *Sci. Rep.* **6**, 27175; doi: 10.1038/srep27175 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>