

Rescue treatment with eltrombopag in refractory cytopenias after allogeneic stem cell transplantation

Semra Aydin^{ID}, Chiara Dellacasa, Sara Manetta, Luisa Giaccone, Laura Godio, Giorgia Iovino, Benedetto Bruno and Alessandro Busca

Ther Adv Hematol
2020, Vol. 11: 1–10

DOI: 10.1177/
2040620720961910

© The Author(s), 2020.
Article reuse guidelines:
sagepub.com/journals-
permissions

Abstract

Background: Patients with post-transplant cytopenias due to poor graft function or primary engraftment failure show poor prognosis with a high mortality rate mainly because of graft *versus* host disease (GVHD), infection and/or bleeding. Treatment options are scarce and a CD34+ stem cell boost or a second bone marrow transplantation may be required to restore adequate haematopoiesis.

Methods: In the present study patients with primary engraftment failure ($n = 1$) and refractory poor graft function ($n = 11$) were treated with eltrombopag in a single centre. The reason for eltrombopag treatment was trilineage cytopenia in six patients, bilineage cytopenia in three patients and single lineage cytopenia in three patients. Eltrombopag was initiated at a median of 214 (range: 120–877) days after haematopoietic stem cell transplantation (HSCT) and administered for a median time of 114 (range: 12 days to >490) days. In 8/12 patients eltrombopag was introduced at a dose of 75 mg/day and then increased to 150 mg/day after 1 week; 1 patient was given 50 mg eltrombopag per day, and 3 patients received 75 mg daily.

Results: In 10/12 patients eltrombopag significantly enhanced blood count values and patients became transfusion independent. Once stable haematological response was obtained, treatment was tapered until final discontinuation in 9/10 responding patients. No grade 3 or 4 toxicities were observed. At time of last follow up, 3/12 patients were dead, 2 due to disease relapse, 1 due to GVHD and pneumonia. All patients except one maintained their complete response and remain transfusion independent at a median of 858 (range: 429–1119) days.

Conclusion: These preliminary data confirm that eltrombopag is able to rescue multilineage haematopoiesis in patients with treatment-refractory cytopenias after allogeneic HSCT.

Keywords: cytopenia, eltrombopag, haematopoietic stem cell transplantation, poor graft function

Received: 3 June 2020; revised manuscript accepted: 7 September 2020.

Introduction

Poor graft function is a multifactorial complication after allogeneic haematopoietic stem cell transplantation (HSCT), occurring in 7–21% of patients.^{1,2} Multiple pre- and post-transplantation factors have been associated, namely: (a) donor-recipient HLA-disparity; (b) low haematopoietic stem cell (HSC) dose; (c) T-cell

depletion of the donor marrow; (d) intensity of the conditioning regimen; (e) post-transplant immunosuppression; (f) coexisting graft *versus* host disease (GVHD); (g) reactivation of cytomegalovirus (CMV) as well as its treatment with ganciclovir; (h) infections. A small proportion of patients (<3%) fail engraftment after transplant and therefore are classified as primary engraftment failure.³

Correspondence to:
Semra Aydin
A.O.U. Città della
Salute e della Scienza,
Dipartimento di Oncologia,
Ematologia, Corso
Bramante 88, Turin, 10126,
Italy
semay74@hotmail.com

Chiara Dellacasa
Sara Manetta
Luisa Giaccone
Benedetto Bruno
Alessandro Busca
A.O.U. Città della
Salute e della Scienza,
Dipartimento di Oncologia,
SSD Trapianto allogenico
di cellule staminali, Turin,
Italy

Laura Godio
A.O.U. Città della Salute
e della Scienza, Anatomia
Patologica, Turin, and
University of Turin, Italy

Giorgia Iovino
A.O.U. Città della
Salute e della Scienza,
Dipartimento di Oncologia,
Ematologia, Turin, Italy

Several attempts to reconstitute donor haematopoiesis have included CD34+ selected stem cell boost, mesenchymal stem cell transfusion or a second marrow transplantation.⁴⁻⁶ Unfortunately patients achieve unsatisfactory results and their prognosis remains poor with a high mortality rate due to GVHD, infection and/or bleeding, especially in patients with primary engraftment failure.

Recent *in vitro* and *in vivo* data showed that stimulation of the c-MPL-signalling pathway by thrombopoietin (TPO) agonists may have a significant influence on maintenance and self-renewal of HSCs.⁷⁻¹⁰ Recovery of trilineage haematopoiesis after eltrombopag treatment was highly effective in patients with severe aplastic anaemia (SAA) and led to US Food and Drug Administration approval for treatment of SAA, immune thrombocytopenia and hepatitis-C-related thrombocytopenia.¹¹⁻¹⁵

In the setting of HSCT, several authors investigated retrospectively the role of eltrombopag in refractory thrombocytopenia after allogeneic HSCT and reported promising results with significant platelet recovery until achieving transfusion independence.¹⁶⁻¹⁹ On the basis of these results current phase II trials using TPO agonists for prolonged thrombocytopenia after HSCT have been initiated, (NCT01791101, NCT01000051, NCT01927731).

Further, *in vivo* studies have demonstrated that TPO promotes even multilineage haematopoiesis and increases peripheral human platelets and white blood cell counts by recruiting and expanding HSC/haematopoietic progenitor cells from the bone marrow compartment of a NOD/SCID xenotransplant model.^{20,21}

These observations led to the hypothesis that eltrombopag may have a role in haematopoietic recovery in single, bi- or even trilineage cytopenias in patients after allogeneic HSCT as recently reported.²²⁻²⁴ In the present analysis we report a single centre experience with eltrombopag in post-allogeneic transplant patients with refractory single and multilineage cytopenias due to poor graft function or primary engraftment failure.

Patients and methods

Between January 2015 and August 2018 a total of 216 patients underwent allogeneic HSCT at the Stem Cell Transplant Center, AOU Città della

Salute e della Scienza in Turin, Italy. Overall, 12 (5.5%) patients developed refractory single or multilineage cytopenias after transplant due to graft failure ($n=1$) or poor graft function ($n=11$) and were included in the present analysis. Two patients have already been described in a previous study.²² Poor graft function was defined as cytopenia in at least one lineage, namely platelets $<20.000 \times 106/L$, haemoglobin $<90 g/L$, absolute neutrophil count (ANC) $<1000 \times 106/L$ and/or transfusion requirement of blood or platelets beyond day +28 after HSCT together with a full donor chimerism. Patients with severe GVHD, reactivation of CMV or other infections as well as patients receiving myelotoxic treatments at time of eltrombopag treatment were excluded. Three patients with full donor chimerism developed single ($n=1$) and multilineage cytopenia ($n=2$) at days +59, +105 and +131, respectively. They are still receiving eltrombopag after 30, 14 and 15 days, respectively. Interestingly, two of them showed significant improvement in their transfusion requirements. Given the short follow-up period the three patients were not included in the analysis.

Poor graft function was classified as 'primary' if association with obvious clinical events could be excluded.²³ Poor graft function was defined as 'secondary' in the following circumstances: (a) in case of association with immune-mediated events such as GVHD in the previous 4 weeks, or (b) in case of association with infectious complications and/or myelotoxic treatments such as CMV reactivation and anti-CMV treatment in the previous 4 weeks, or other clinically important infections. In the case of missing recovery of the blood count after HSCT the patient was defined as primary engraftment failure.²³

A bone marrow analysis showing a full donor chimerism and excluding disease relapse as well as showing impaired haematopoiesis was performed in all patients before therapy initiation. Before starting eltrombopag treatment and after individual approval by the Hospital Commission for Off-label Use of Drugs for each patient, personal informed consent for off-label treatment was signed by each patient. All patients received oral eltrombopag: in 8/12 patients eltrombopag was initiated at a dose of 75 mg/day and after 1 week was increased to 150 mg/day; 1 patient was given eltrombopag 50 mg/day, and 3 patients received 75 mg daily. Once stable haematological response was obtained, treatment was slowly tapered until final discontinuation.

Complete response or remission (CR) was defined as all of the following: (a) platelet count $>50.000 \times 106/L$; (b) ANC $>1500 \times 106/L$; (c) haemoglobin $>100g/L$ without blood cell transfusion or granulocyte colony-stimulating factor (G-CSF) support.

Treatment toxicities

Eltrombopag was well tolerated by all patients with no evidence of cataract, thrombosis or any other grade 3/4 toxicities, only grade 1–2 toxicities were observed. No endogenous TPO levels were measured in this patient cohort. A total of 11 patients started eltrombopag therapy at 75 mg daily in order to assess the initial safety profile. After 1 week, in eight patients, the dosage was increased to 150 mg daily, except in four patients. Patient no. 06 developed a grade 2 thrombocytosis up to $50,700 \times 106/L$ when the dose was increased to 150 mg. Eltrombopag was immediately reduced to 75 mg and the platelet count normalized again. Patient no. 08 developed grade 1 nausea after enhancement to 150 mg and subsequently refused the 150 mg dosage. Patient no. 11 started eltrombopag therapy at 75 mg and obtained a CR with this dosage. Patient no. 12 started eltrombopag at 50 mg daily due to moderate liver enzyme enhancement. After a total of 12 days he developed significant increase in liver function tests, therefore eltrombopag was discontinued. He concomitantly developed severe venous occlusive disease.

Statistical analysis

A nonparametric Wilcoxon matched-pairs signed-rank test was used to determine the statistical significance of the single lineage upon eltrombopag treatment in patients with cytopenia considering the single cytopenia line. All statistical tests were two-tailed. Statistical analyses were performed using GraphPad Prism 8.4.1. for Windows (GraphPad Software, San Diego, CA, USA). Unadjusted *p* values less than 0.05 were considered significant.

Results

Transplant characteristics

Patients' baseline transplant characteristics are summarised in Table 1. Indication for allogeneic transplantation was acute leukaemia (including one blast crisis of chronic myeloid leukaemia) in

11 patients and high-risk myelodysplastic syndrome in 1 patient. Median age of recipients and donors was 47.5 (range: 40–66) years and 37 (range: 17–59) years, respectively. The majority of patients (8/12) received grafts from an unrelated donor (8/8 HLA matched in 7 cases), 3 patients from a haploidentical family member and 1 patient from a matched sibling donor. A total of 7/12 patients showed an unfavourable CMV combination (donor negative/patient positive) whereas 5 donor/recipient pairs showed a major ABO incompatibility. All patients received a T-cell depleted graft: in 5/12 patients the haematopoietic stem cell source was bone marrow while 7/12 patients received peripheral blood stem cells. Conditioning regimens were chosen according to standard guidelines on the basis of the underlying disease, the age of the patient and the donor type and are listed in Table 1. A myeloablative conditioning regimen was used in all cases except one and patients received a median of $3.7 \times 108/kg$ (range: 1.77–13.3) total nucleated cells, a median of $6 \times 106/kg$ (range: 1.8–19.2) CD34+ cells and a median of $2.37 \times 108/kg$ (range: 0.25–3.2) CD3+ cells. GVHD prophylaxis included cyclosporine (CSA) and methotrexate (MTX) in matched sibling donor transplants, CSA-MTX and rabbit antithymocyte globulin in unrelated donor transplants and tacrolimus-mycophenolate mofetil combined with post-transplant cyclophosphamide in case of grafts from haploidentical donors.

A total of 11 patients achieved neutrophil recovery $>500 \times 106/L$ after a median of 20.5 (range: 15–33) days and 8 patients achieved platelets $>20.000 \times 106/L$ after a median of 20.5 (range: 13–37) days after transplant.

Cytopenias and treatment characteristics

Cytopenias and treatment characteristics are summarised in Table 2. Six patients developed primary poor graft function while in five patients poor graft function was secondary to GVHD ($n=2$), previous bacterial sepsis ($n=2$) and CMV reactivation requiring therapy up to 1 month before cytopenia appearance ($n=1$). At time of eltrombopag initiation all patients showed no evidence of CMV infection. As part of the routine protocol for patients with refractory cytopenia at our centre, human herpesvirus 6 (HHV6) reactivation was controlled. None of the patients in the present study resulted positive.

Table 1. Patient and transplant characteristics.

No.	Age at HSCT	Disease	Donor	HSCT source	ABO (D/R)	Cond.	TNC × 10 ⁸ /kg	CD34 × 10 ⁶ /kg	CD3 × 10 ⁸ /kg	CMV (D/R)	GVHD prophylaxis	Engraftment ANC >500 (day)	Engraftment PLT >20,000 (day)
01	63	AML	Haplo	BM	0+/B+	MAC	3.18	4.51	0.27	neg/pos	PT/Cy+FK- MMF	+25	NR
02	41	ALL	MUD	PBSC	0+/0+	MAC	8.9	6	2.8	neg/pos	CSA, MTX, ATG	+23	+14
03	47	MDS	MUD	PBSC	0+/A+	MAC	11,14	5.6	2.6	neg/neg	CSA, MTX, ATG	+15	+37
04	47	AML	MUD	PBSC	A-/0+	MAC	8.4	6	2.8	neg/pos	CSA, MTX, ATG	+16	+17
05	50	AML	MUD	PBSC	0+/A-	MAC	13.3	6.48	2.57	neg/pos	CSA, MTX, ATG	+28	+25
06	50	MPAL	MUD	PBSC	A+/0+	MAC	11.7	19.2	3.2	neg/pos	CSA, MTX, ATG	+24	+18
07	42	CML	MUD	PBSC	B+/0+	RIC	3.7	8.71	2.37	neg/pos	CSA, MMF	+19	never below
08	46	AML	Haplo	BM	0+/0+	MAC	7	6.6	0.34	pos/pos	PT/Cy+FK- MMF	+17	+110
09	51	AML	MSD	PBSC	A+/0+	MAC	5.3	4	2.39	pos/pos	CSA-MTX	+14	+13
10	66	AML	MUD	BM	0+/-B-	MAC	2.73	1.8	0.25	pos/pos	CSA, MTX, ATG	+33	NR
11	40	AML	MUD	BM	0-/A-	MAC	3.7	2.3	0.4	neg/pos	CSA, MTX, ATG	+20	+23
12	48	AML	Haplo	BM	B-/A+	MAC	1.77	1.8	0.35	pos/pos	PT/Cy+FK- MMF	+21	NR

ALL, acute lymphoblastic leukaemia; AML, acute myeloid leukaemia; ANC, absolute neutrophil count; ATG, antithymocyte globulin; BM, bone marrow; CML, chronic myeloid leukaemia; CMV, cytomegalovirus; Cond, conditioning regimen; CSA, cyclosporine; D, donor; FK, tacrolimus; GVHD, graft versus host disease; Haplo, haploidentical donor; HSCT, haematopoietic stem cell transplant; MAC, myeloablative conditioning; MDS, myelodysplastic syndrome; MMF, mycophenolate mofetil; MPAL, mixed-phenotype acute leukaemia; MSD, matched sibling donor; MTX, methotrexate; MUD, matched unrelated donor; neg, negative; NR, not reported; PBSC, peripheral blood stem cells; PLT, platelets; pos, positive; PT/Cy, post-transplant cyclophosphamide; R, recipient; RIC, reduced intensity conditioning; TNC, total nuclear cells.

Table 2. Cytopenia and treatment characteristics.

No	Cytopenia origin	Lineage involvement	RBC transfusion	PLT transfusion	BM cellularity	Chimerism (BM)	Previous cytopenia therapy	Eltrombopag start from HSCT (+ days)	Maximum dose (mg)	Therapy duration (days)	State at follow up
01	Primary PGF	PLT, RBC, ANC	yes	yes	20–60%	100%	G-CSF, EPO	148	150	ongoing	alive
02	Secondary PGF	PLT, RBC, ANC	yes	yes	10–70%	100%	G-CSF	191	150	490	alive
03	Secondary PGF	PLT, RBC, ANC	yes	yes	30–80%	100%	G-CSF, EPO, Ig	310	150	63	alive
04	Primary PGF	RBC, ANC	yes	no	10–60	100%	G-CSF, EPO, RTX, PEX	404	150	141	alive
05	Secondary PGF	PLT, ANC, RBC	yes	yes	20–40%	100%	EPO	371	150	341	alive
06	Primary PGF	RBC	yes	no	80%	100%	EPO, RTX, EDX, BRT	877	75	172	alive
07	PEF	PLT, RBC, ANC	yes	yes	10%	4,84%	G-CSF, EPO	132	150	30	alive
08	Primary PGF	PLT	no	yes	70%	100%	only transfusions	186	75	372	alive
09	Primary PGF	RBC, ANC	yes	no	10–50%	100%	G-CSF, EPO, RTX, PEX	543	150	101	alive
10	Secondary PGF	PLT, RBC, ANC	yes	yes	10–70%	100%	G-CSF	140	150	22	dead
11	Primary PGF	PLT	no	yes	5–60%	100%	RTX, Ig	237	75	128	dead
12	Secondary PGF	RBC, PLT	yes	yes	5–20%	100%	G-CSF, EPO	120	50	12	dead

ANC, absolute neutrophil count; BM, bone marrow; BRT, bortezomib; EDX, cyclophosphamide; EPO, erythropoietin; G-CSF, granulocyte colony-stimulating factor; HSCT, haematopoietic stem cell transplant; Ig, immunoglobulin; PEF, primary engraftment failure; PEX, plasma exchange; PGF, poor graft function; PLT, platelets, RBC, red blood cells; RTX, rituximab.




Figure 1. Blood cell count before and after eltrombopag treatment. The analysis included all patients with cytopenia considering the single lineage. (a) Median platelet count was 16,000 (6000–22,000) *versus* 77,000 (50,000–218,000) $\times 10^6/L$ before and after treatment ($p=0.012$). (b) Median haemoglobin was 76 (66–85) *versus* 129 (74–154) g/L ($p=0.008$). (c) Median neutrophil count was 650 (0–990) *versus* 2325 (0–4260) $\times 10^6/L$ ($p=0.019$). Nonparametric Wilcoxon matched-pairs signed-rank test.

The reason for eltrombopag treatment was trilineage cytopenia in six patients, bilineage cytopenia in three patients and single lineage cytopenia in three patients, of whom $n=1$ with isolated anaemia and $n=2$ with isolated thrombocytopenia. All patients required recurrent transfusions of platelet and/or red blood cells. Antibody-mediated cytopenia was investigated at time of pure red cell aplasia diagnosis in three patients with anti-A immunoglobulin (Ig)G and IgM titres (no. 4: IgG 1:128, IgM 1:32; no. 6: IgG 1:2048, IgM 1:256; no. 9: IgG 1:32, IgM 1:4). Two patients (no. 6 and no. 9) did not show significant reduction of anti-A titres before eltrombopag initiation, while one (no. 4) did not determine anti-A titres before eltrombopag initiation. At the time of eltrombopag initiation, the median ferritin level was 3126 (range: 897–6770) ng/ml with 9/12 patients having a ferritin level >2000 ng/ml. Before treatment with eltrombopag patients received erythropoietin ($n=8$), G-CSF ($n=8$), plasma exchange ($n=2$), rituximab ($n=4$), immunoglobulins ($n=2$), bortezomib ($n=1$) and cyclophosphamide ($n=1$). As indicated in Table 2, four patients received rituximab at days +216, +286, +161 and +209 post-transplant, respectively. Two underwent plasma exchange at day +173 and +125, whereas one patient received bortezomib at day +390 after transplant. Patients receiving rituximab therapy were treated after a total of 17, 257, 716 and 195 days, respectively with eltrombopag and all four responded with CR. Nevertheless a long-term effect of rituximab therapy cannot be excluded. Eltrombopag was initiated at a median of 214 (range: 120–877) days after HCST and was administered for a median time of 114 (range: 12– >490) days. In one case

(no. 12) eltrombopag was discontinued after a total of 12 days and not resumed due to the presence of significant increase in liver function tests concomitantly with the development of severe venous occlusive disease. Interestingly, after discontinuation of eltrombopag the platelet count increased up to a maximum of $59,000 \times 10^6/L$, the haemoglobin value rose to a maximum of 113 g/L and the absolute neutrophil count up to a maximum of $1740 \times 10^6/L$. This improvement was temporary; 12.9 months after eltrombopag treatment the patient developed relapse of his acute myeloid leukaemia and died 3 weeks later.

In 10/12 patients eltrombopag significantly increased blood count values and patients became transfusion independent. Among all, the median platelet count of nine patients with persistent thrombocytopenia before eltrombopag treatment was 16,000 (range: 6000–22,000) $\times 10^6/L$ *versus* 77,000 (range: 50,000–218,000) $\times 10^6/L$ after eltrombopag treatment, $p=0.012$ (Figure 1(a)). This maximum response of the platelet count was obtained a median of 66 (range: 44–425) days after eltrombopag initiation. A total of 10 patients with persistent anaemia showed a significant response with a rise in the haemoglobin value from a median value of 76 (range: 66–85) g/L to a median of 129 (range: 74–154) g/L after eltrombopag therapy; this response was obtained after a median of 141 (range: 6–291) days ($p=0.019$) (Figure 1(b)). Finally, a total of eight patients with persistent neutropenia had a median absolute neutrophil count before eltrombopag treatment of 650 (range: 0–990) $\times 10^6/L$ *versus* 2325 (range: 0–4260) $\times 10^6/L$ after a median of 200 (21–379) days, $p=0.008$ (Figure 1(c)).

Two patients did not respond to eltrombopag. The poor graft function of the nonresponding patient no. 10 was defined secondary probably due to a previous CMV reactivation and treatment. Eltrombopag treatment was maintained for up to 22 days at a maximum dose of 150 mg daily, however he developed a significant GVHD and eventually died of pneumonia. The second non-responding patient (no. 7) was classified as a primary engraftment failure, she never recovered haematopoiesis after transplant and was treated with eltrombopag (maximum 150 mg/day) for a total of 30 days. No response was obtained, and she received a second bone marrow transplantation. She is still alive maintaining full donor chimerism and shows no signs of disease relapse.

Overall, at time of last follow up, 9/12 patients are still alive, of whom only 1 patient is still under eltrombopag treatment. In seven patients eltrombopag could be tapered and discontinued. All patients except one maintained their complete response and remain transfusion independent at a median of 858 (range: 429–1119) days. One patient, who initially responded with a CR lost the response 341 days after initiation of the treatment, and eltrombopag was stopped. A total of 3/12 patients were dead at time of last follow up, 2 due to disease relapse and 1 due to GVHD and pneumonia.

Discussion

The receptor for TPO, c-Mpl, is expressed mainly on HSCs. Eltrombopag initiates signal transduction by binding to c-Mpl, consequently Janus kinase 2 phosphorylates tyrosine residues, thereby stimulating the downstream cascade STATs, PI3K, the mitogen-activated protein kinases and extracellular signal-regulated kinases.^{25–27} Thereby, the c-Mpl/TPO signalling pathway is activated and regulates partially HSC cell-cycle progression and renewal.²⁸ Sun *et al.* could previously even demonstrate that eltrombopag induces in umbilical cord blood transplants multilineage responses through the expansion of bone marrow HSCs and haematopoietic progenitor cells.²¹

On the basis of these results, the stimulation of residual HSCs by TPO agonists was postulated and attempted by using eltrombopag in the setting of persistent cytopenia after HSCT by several groups with promising results. Tang *et al.* have treated with eltrombopag a cohort of 12

post-HSCT patients, who responded poorly to standard treatments for secondary poor graft function. A total of 10 (83.3%) patients responded to the treatment; 8 achieved CR while 2 achieved a partial response.²⁴ The 12-month overall survival rate was 83.3% with all CR patients maintaining their response. A further study with a similar cohort size was reported by Marotta *et al.* using eltrombopag as first-line treatment in patients with primary engraftment failure and poor graft function ($n = 13$).²³ Overall, haematological response was reported in 7/13 (53.8%) patients, with 6 patients showing CR (46.2%). In both studies eltrombopag was well tolerated, and no signs of clonal evolution were reported. The dosages used in the study of Marotta *et al.* as well as in the study of Tang *et al.* were quite low (usually <100 mg/day) compared with the 150 mg/day used in aplastic anaemia. In our study the initial dose used was 75 mg and then increased to 150 mg daily in the majority of patients according to tolerance and response.

In the present cohort, a total of 10/12 patients (83.3%) obtained CR achieving transfusion independence. These results were obtained by oral eltrombopag monotherapy in patients that were deeply cytopenic that otherwise might require a CD34+-selected stem cell boost or a second allogeneic HSCT to restore an effective haematopoiesis.

Interestingly, analysis of the bone marrow biopsy before eltrombopag treatment in eight patients revealed the presence of infiltration of cytotoxic T lymphocytes (marked by CD3+ and cytotoxic granule-associated RNA binding protein TIA1) in the range of 10–25%. Whether the infiltration of cytotoxic T cells in the bone marrow may have an impact on poor graft function occurrence is currently unknown. Notably, Nakao *et al.* showed that in patients with aplastic anaemia a T-cell clone markedly inhibited the growth of haematopoietic progenitor cells by direct cell–cell contact and IFN-gamma secretion in response to CD34+ cells.²⁹ They hypothesised that these cytotoxic T cells may even contribute to the pathogenesis of aplastic anaemia.²⁹

Further, iron overload has been shown to have a suppressive effect on the erythroblast differentiation of human CD34+ cells. It increases the level of the intracellular reactive oxygen species (ROS) promoting apoptosis of immature erythroblasts

by suppressing BCL2 gene expression.^{30,31} Kong *et al.* recently reported that elevated ROS levels are associated with increased DNA damage and apoptosis, resulting in exhaustion of CD34+ cells in the bone marrow of patients with poor graft function.³² These studies support the strong association of a high serum ferritin level with poor graft function. Iron-chelation therapy may lead to a significant haematopoietic response in some patients with aplastic anaemia.^{33,34} Similarly, Olivieri *et al.* described a patient with severe aplastic anaemia who was diagnosed with poor graft function after transplantation and achieved complete haematopoietic recovery mainly due to iron-chelation therapy.³⁵

In the present study the median ferritin level before eltrombopag treatment initiation was 3126 (range: 897–6770) ng/ml with 9/12 patients having a ferritin level >2000 ng/ml. This may be attributed to significant blood transfusion support as nearly all were patients with acute leukaemia. Iron chelation was performed in six out of nine patients with severe anaemia, and all except one (no. 10) became blood transfusion independent. On the basis of this preliminary result, a synergistic effect between eltrombopag and iron-chelation therapy may be hypothesised.

At time of last follow up, three patients were dead and one with primary engraftment failure was successfully retransplanted. Of the remaining eight patients seven maintained CR criteria and are transfusion independent. Our study has several limiting factors, namely the retrospective character, the small cohort size and the lack of a control group for the comparison of the drug effect. Defined criteria for eltrombopag initiation and discontinuation were lacking and were determined by the clinical course. A larger prospective study is warranted to confirm our preliminary results.

In conclusion, in the present study, a high overall response rate was achieved with the use of eltrombopag in heavily pretreated patients with post-HSCT cytopenias. These preliminary findings confirm the results from previous groups suggesting that patients with refractory cytopenias following HSCT might benefit from treatment with eltrombopag.

Authors contributions

SA collected data, performed analysis and wrote the manuscript. AB, CD, SM, LG, GI and BB analysed the data and wrote the manuscript. LG performed the bone marrow analysis.

Authors' note

Semra Aydin is also affiliated with Department of Oncology, Hematology, Immuno-oncology and Rheumatology, University Hospital of Bonn, Bonn, Germany.

Conflict of interest statement

AB has been a speaker for Novartis.

Funding

The authors received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Aydin Semra  <https://orcid.org/0000-0001-7393-9690>

References

1. Lee KH, Lee JH, Choi SJ, *et al.* Failure of trilineage blood cell reconstitution after initial neutrophil engraftment in patients undergoing allogeneic hematopoietic cell transplantation - frequency and outcomes. *Bone Marrow Transplant* 2004; 33: 729–734.
2. Woolfrey A and Anasetti C. Allogeneic hematopoietic stem-cell engraftment and graft failure. *Pediatr Transplant* 1999; 3(Suppl. 1): 35–40.
3. Olsson R, Remberger M, Schaffer M, *et al.* Graft failure in the modern era of allogeneic hematopoietic SCT. *Bone Marrow Transplant* 2013; 48: 537–543 *Erratum in: Bone Marrow Transplant* 2013; 48: 616.
4. Ghobadi A, Fiala MA, Ramsingh G, *et al.* Fresh or cryopreserved CD34+-selected mobilized peripheral blood stem and progenitor cells for the treatment of poor graft function after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2017; 23: 1072–1077.
5. Larocca A, Piaggio G, Podestà M, *et al.* Boost of CD34+-selected peripheral blood cells without further conditioning in patients with poor graft function following allogeneic stem cell transplantation. *Haematologica* 2006; 91: 935–940.

6. Liu X, Wu M, Peng Y, *et al.* Improvement in poor graft function after allogeneic hematopoietic stem cell transplantation upon administration of mesenchymal stem cells from third-party donors: a pilot prospective study. *Cell Transplant* 2014; 23: 1087–1098.
7. Zeigler FC, de Sauvage F, Widmer HR, *et al.* In vitro megakaryocytopoietic and thrombopoietic activity of c-mpl ligand (TPO) on purified murine hematopoietic stem cells. *Blood* 1994; 84: 4045–4052.
8. Alexander WS, Roberts AW, Nicola NA, *et al.* Deficiencies in progenitor cells of multiple hematopoietic lineages and defective megakaryocytopoiesis in mice lacking the thrombopoietic receptor c-Mpl. *Blood* 1996; 87: 2162–2170.
9. Qian H, Buza-Vidas N, Hyland CD, *et al.* Critical role of thrombopoietin in maintaining adult quiescent hematopoietic stem cells. *Cell Stem Cell* 2007; 1: 671–684.
10. Chotinantakul K and Leraanansaksiri W. Hematopoietic stem cell development, niches, and signaling pathways. *Bone Marrow Res* 2012; 2012: 270425.
11. Olnes MJ, Scheinberg P, Calvo KR, *et al.* Eltrombopag and improved hematopoiesis in refractory aplastic anemia. *N Engl J Med* 2012; 367: 11–19.
12. Desmond R, Townsley DM, Dumitriu B, *et al.* Eltrombopag restores trilineage hematopoiesis in refractory severe aplastic anemia that can be sustained on discontinuation of drug. *Blood* 2014; 123: 1818–1825.
13. Yang R, Li J, Jin J, *et al.* Multicentre, randomised phase III study of the efficacy and safety of eltrombopag in Chinese patients with chronic immune thrombocytopenia. *Br J Haematol* 2017; 176: 101–110.
14. Bussel JB, Cheng G, Saleh MN, *et al.* Eltrombopag for the treatment of chronic idiopathic thrombocytopenic purpura. *N Engl J Med* 2007; 357: 2237–2247.
15. McHutchison JG, Dusheiko G, Shiffman ML, *et al.* Eltrombopag for thrombocytopenia in patients with cirrhosis associated with hepatitis C. *N Engl J Med* 2007; 357: 2227–2236.
16. Tanaka T, Inamoto Y, Yamashita T, *et al.* Eltrombopag for treatment of thrombocytopenia after allogeneic hematopoietic cell transplantation. *Biol Blood Marrow Transplant* 2016; 22: 919–924.
17. Fu H, Zhang X, Han T, *et al.* Eltrombopag is an effective and safe therapy for refractory thrombocytopenia after haploidentical hematopoietic stem cell transplantation. *Bone Marrow Transplant* 2019; 54: 1310–1318.
18. Yuan C, Boyd AM, Nelson J, *et al.* Eltrombopag for treating thrombocytopenia after allogeneic stem cell transplantation. *Biol Blood Marrow Transplant* 2019; 25: 1320–1324.
19. Mahat U, Rotz SJ and Hanna R. Use of thrombopoietin receptor agonists in prolonged thrombocytopenia after hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 2020; 26: e65–e73
20. Ramsfjell V, Borge OJ, Cui L, *et al.* Thrombopoietin directly and potently stimulates multilineage growth and progenitor cell expansion from primitive (CD34+ CD38-) human bone marrow progenitor cells: distinct and key interactions with the ligands for c-kit and flt3, and inhibitory effects of TGF-beta and TNF-alpha. *J Immunol* 1997; 158: 5169–5177.
21. Sun H, Tsai Y, Nowak I, *et al.* Eltrombopag, a thrombopoietin receptor agonist, enhances human umbilical cord blood hematopoietic stem/progenitor cell expansion and promotes multi-lineage hematopoiesis. *Stem Cell Res* 2012; 9: 77–86.
22. Busca A, Dellacasa C, Giaccone L, *et al.* Eltrombopag for the treatment of refractory pure RBC aplasia after major ABO incompatible hematopoietic stem cell transplantation. *Biol Blood Marrow Transplant* 2018; 24: 1765–1770.
23. Marotta S, Marano L, Ricci P, *et al.* Eltrombopag for post-transplant cytopenias due to poor graft function. *Bone Marrow Transplant* 2019; 54: 1346–1353.
24. Tang C, Chen F, Kong D, *et al.* Successful treatment of secondary poor graft function post allogeneic hematopoietic stem cell transplantation with eltrombopag. *J Hematol Oncol* 2018; 11: 103.
25. Debili N, Wendling F, Katz A, *et al.* The Mpl-ligand or thrombopoietin or megakaryocyte growth and differentiative factor has both direct proliferative and differentiative activities on human megakaryocyte progenitors. *Blood* 1995; 86: 2516–2525.
26. Chou FS and Mulloy JC. The thrombopoietin/MPL pathway in hematopoiesis and leukemogenesis. *J Cell Biochem* 2011; 112: 1491–1498.

27. Tortolani PJ, Johnston JA, Bacon CM, *et al.* Thrombopoietin induces tyrosine phosphorylation and activation of the Janus kinase, JAK2. *Blood* 1995; 85: 3444–3451.
28. Bersenev A, Wu C, Balcerak J, *et al.* Lnk controls mouse hematopoietic stem cell self-renewal and quiescence through direct interactions with JAK2. *J Clin Invest* 2008; 118: 2832–2844.
29. Nakao S, Takami A, Takamatsu H, *et al.* Isolation of a T-cell clone showing HLA-DRB1*0405-restricted cytotoxicity for hematopoietic cells in a patient with aplastic anemia. *Blood* 1997; 89: 3691–3699.
30. Cabantchik ZI, Breuer W, Zanninelli G, *et al.* LPI-labile plasma iron in iron overload. *Best Pract Res Clin Haematol* 2005; 18: 277–287.
31. Taoka K, Kumano K, Nakamura F, *et al.* The effect of iron overload and chelation on erythroid differentiation. *Int J Hematol* 2012; 95: 149–159.
32. Kong Y, Song Y, Hu Y, *et al.* Increased reactive oxygen species and exhaustion of quiescent CD34-positive bone marrow cells may contribute to poor graft function after allotransplants. *Oncotarget* 2016; 7: 30892–30906.
33. Lee JW, Yoon SS, Shen ZX, *et al.* Hematologic responses in patients with aplastic anemia treated with deferasirox: a post hoc analysis from the EPIC study. *Haematologica* 2013; 98: 1045–1048.
34. Visani G, Guiducci B, Giardini C, *et al.* Deferasirox improves hematopoiesis after allogeneic hematopoietic SCT. *Bone Marrow Transplant* 2014; 49: 585–587.
35. Olivieri J, Mancini G, Goteri G, *et al.* Reversal of poor graft function with iron-chelating therapy after allogeneic transplantation for severe aplastic anemia. *Leuk Lymphoma* 2016; 57: 965–968.

Visit SAGE journals online
[journals.sagepub.com/
home/tah](http://journals.sagepub.com/home/tah)

 SAGE journals