

Surgical Repositioning of a Dilacerated Impacted Incisor

¹Vivek Kumar Adlakha, ¹Preetika Chandna, ²Sumir Gandhi, ³Saroj Chopra
³Namita Singh, ⁴Shamsher Singh

¹Senior Lecturer, Department of Pedodontics and Preventive Dentistry, Subharti Dental College, Meerut, Uttar Pradesh, India

²Reader, Department of Oral Surgery, Christian Dental College, Ludhiana, Punjab, India

³Professor, Department of Pedodontics and Preventive Dentistry, Christian Dental College, Ludhiana, Punjab, India

⁴Lecturer, Department of Pedodontics and Preventive Dentistry, Surindra Dental College, Ganganagar, Rajasthan, India

Correspondence: Vivek Kumar Adlakha, Senior Lecturer, Department of Pedodontics and Preventive Dentistry, Subharti Dental College, J-58, Shastri Nagar, Meerut-250004, Uttar Pradesh, India, e-mail: adlakhavivek@gmail.com

ABSTRACT

Dilaceration is one of the most common complications of trauma to deciduous dentition. The possible causes of dilaceration are trauma and developmental disturbances, while some authors suggest an association with some developmental syndromes. Dilaceration can be seen in both the permanent and deciduous dentition. The present case report describes surgical repositioning of a dilacerated impacted maxillary central incisor in a 9-year-old boy.

Keywords : Dental trauma, Impacted, Dilaceration.

INTRODUCTION

Dilaceration refers to an angulation or a sharp curve in the root or crown of a formed tooth. Dilaceration is usually associated with dental trauma to a deciduous predecessor in which the tooth is driven apically into the jaw. The anomaly in tooth happens during the period when the tooth is forming. Trauma results in change of the calcified portion of the tooth and the remaining tooth develops at an angle. The curve or bend in the tooth may occur anywhere along the length of the tooth, depending upon the amount of tooth formed when the injury occurred.¹ The most complicated situation is root dilacerations with the crown in an inverted direction; thus, the tooth is always impacted.² The conventional treatment for a dilacerated incisor is extraction, transplantation or surgical/orthodontic approach.²⁻⁷ This article describes a rare case of dilaceration with the crown in an inverted direction with the palatal aspect of the crown facing the labial side.

CASE REPORT

A 9-year-old boy visited the outpatient clinic of Department of Pedodontics and Preventive Dentistry, Christian Dental College, Ludhiana, India with a chief complaint of missing upper anterior tooth (Fig. 1). Patient gave a history of trauma at the age of 3 to 4 years due to hand pump, which resulted in intrusion of the deciduous maxillary central incisors. The deciduous maxillary central incisors were mobile and were extracted. Intraoral examination revealed missing left maxillary central incisor (tooth no. 21). There was a space

deficiency in the maxillary anterior area due to migration of number 11 into the space that would have been occupied by tooth number 21. The maxillary central incisor, with a dilacerated root and the crown directed upwards, was visible on the radiographs (Figs 2 to 4). The treatment goal was to reopen the incisor space and bring the impacted incisor into its position.

TREATMENT

Orthodontic treatment was carried out to reopen the lost space using a removable Hawley's appliance with finger springs (Fig. 5). Once the lost space was regained, surgical repositioning of the dilacerated incisor was planned. Following presurgical investigations, surgical repositioning



Fig. 1: Preoperative study model shows noneruption of tooth number 21




Fig. 2: Preoperative IOPA radiograph




Fig. 3: Preoperative occlusal radiograph




Fig. 4: Preoperative OPG radiograph

was undertaken under local anesthesia. The impacted incisor was exposed with a buccal flap operation utilizing vertical releasing incisions. The bone surrounding the crown of the tooth was carefully removed and the incisor tooth follicle was gently separated from its bony socket. Special care was taken not to damage the periodontal membrane on



Fig. 5: Hawley's appliance with finger spring for space regaining

the cervical and root portion of the tooth. The incisor was then repositioned in the correct direction but in the semi-erupted position to safeguard marginal bone regeneration. Orthodontic bracket was cemented on the labial surface of the incisors to splint the repositioned incisor. After suturing the flap, the suture was used to splint the incisor lightly with the adjacent teeth. The sutures and splint were removed after one week (Figs 6 to 12).

The repositioned incisor remained vital and responded normally to percussion, mobility and sensitivity testing. The soft tissue, gingival contour and the probing depth were within normal limits. The patient is presently on orthodontic therapy for further positioning and alignment of the teeth (Fig. 13).

DISCUSSION

The present case demonstrated that surgical repositioning of a dilacerated impacted maxillary central incisor may be a realistic treatment option as compared to other conventional techniques like extraction, surgical exposure and orthodontic traction, autotransplantation with premolars or autoallograft tooth transplantation. It is probably because of the high clinical difficulty in surgically positioning the dilacerated impacted incisor tooth; most patients and clinicians probably choose extraction with replacement of prosthesis instead. The present case revealed that dilacerations of the root would not be a great obstacle always if the case had carefully planned procedure.

The various advantages of surgically repositioning the dilacerated impacted incisor are:^{2,8}

1. There is only one surgical site and procedure is done once.
2. Simple orthodontic mechanics; if needed.
3. Impacted tooth used as a donor has a thick periodontal membrane, which is more suitable for reattachment than the narrow periodontal membrane found in erupted, functional tooth.



Fig. 6: During surgical exposure shows the impacted incisor with its crown inverted



Fig. 9: Repositioned tooth number 21 in semierupted state



Fig. 7: Bony socket after removal of tooth number 21



Fig. 10: Sutured flap



Fig. 8: Repositioning of the dilacerated tooth




Fig. 11: Suture splint in place

4. Immediate esthetics: Endodontic treatment was not necessary in this case because of the open apex. There is a good chance of revascularization of replanted teeth with open apex unlike those with a closed apex, where endodontic treatment is planned within a couple of weeks after transplantation.^{5,9}

Andreasen showed that if autotransplantation was carefully done, success rate can be as high as 95%.¹⁰

The impacted tooth was transplanted in a semierupted position; immediate esthetics was achieved. This can be a good compliance incentive for both clinicians and parents.



Fig. 12: Postoperative IOPA radiograph



Fig. 13: Postoperative photograph after 1 month

Surgical repositioning treatment helps in achieving normal periodontal and gingival attachment, and normal dental and skeletal growth.

The treatment expense is reduced since the tooth is positioned in a semi-erupted state. Also, simple mechanics are needed to align the tooth in proper occlusion.² And positioning the tooth in semi-erupted state helps in safeguarding marginal bone regeneration.¹¹

The major disadvantage with surgical repositioning or autotransplantation is injury to the periodontal ligament, resulting in root resorption and/or ankylosis, and injury to the Hertwig's epithelial root sheath causing partial or total arrest of root development.¹⁰⁻¹²

CONCLUSION

Surgical repositioning is a viable treatment option for dilacerated impacted incisor. It is a simple surgical procedure with minimal orthodontic mechanics and can be undertaken safely as an outpatient procedure.

REFERENCES

1. Shafer, WG.; Maynard, KH.; Bernet, ML. A textbook of oral pathology. 4th edition. Philadelphia: WB Saunders Co 1993. p. 40.
2. Tsai TP. Surgical repositioning of an impacted dilacerated incisor in mixed dentition. J Am Dent Assoc 2002 Jan;133(1):61-66.
3. Agnihotri A, Marwah N, Dutta S. Dilacerated unerupted central incisor: A case report. J Indian Soc Pedod Prev Dent 2006 Sep;24(3):152-154.
4. Crawford LB. Impacted maxillary central incisor in mixed dentition treatment. Am J Orthod Dentofacial Orthop 1997 Jul;112(1):1-7.
5. Vermette ME, Kokich VG, Kennedy DB. Uncovering labially impacted teeth: Apically positioned flap and closed-eruption techniques. Angle Orthod 1995;65(1):23-32.
6. Czochrowska EM, Stenvik A, Album B, Zachrisson BU. Autotransplantation of premolars to replace maxillary incisors: A comparison with natural incisors. Am J Orthod Dentofacial Orthop 2000 Dec;118(6):592-600.
7. Lin YT. Treatment of an impacted dilacerated maxillary central incisor. Am J Orthod Dentofacial Orthop 1999 Apr;115(4):406-409.
8. Berglund L, Kurol J, Kvint S. Orthodontic pre-treatment prior to autotransplantation of palatally impacted maxillary canines: Case reports on a new approach. Eur J Orthod 1996 Oct;18(5):449-456.
9. Saad AY, Abdellatif ES. Surgical repositioning of unerupted anterior teeth. J Endod 1996 Jul;22(7):376-379.
10. Andresen, JO. Atlas of replantation and transplantation of teeth. St Louis: Saunders, 1992. p. 16-57.
11. Andreasen, JO.; Petersen, JK.; Laskin, DM. Textbook and color atlas of tooth impactions. Copenhagen, Denmark: Munksgaard 1997. p. 113-124.
12. Azaz B, Steiman Z, Koyoundjisky-Kaye E, Lewin-Epstein J. The sequelae of surgical exposure of unerupted teeth. J Oral Surg 1980 Feb;38(2):121-127.