



BRAZILIAN JOURNAL
OF MEDICAL AND BIOLOGICAL RESEARCH

www.bjournal.com.br

ISSN 1414-431X
Volume 45 (10) 875-994 October 2012

BIOMEDICAL SCIENCES

Braz J Med Biol Res, October 2012, Volume 45(10) 982-987

doi: 10.1590/S0100-879X2012007500091

Behavioral meaningful opioidergic stimulation activates kappa receptor gene expression

E. Teodorov, M.F.R. Ferrari, D.R. Fior-Chadi, R. Camarini and L.F. Felício

The Brazilian Journal of Medical and Biological Research is partially financed by



Ministério da Ciência e Tecnologia




Ministério da Educação



Institutional Sponsors



Explore High - Performance MS Orbitrap Technology In Proteomics & Metabolomics



All the contents of this journal, except where otherwise noted, is licensed under a [Creative Commons Attribution License](https://creativecommons.org/licenses/by-nc/4.0/)

Behavioral meaningful opioidergic stimulation activates kappa receptor gene expression

E. Teodorov¹, M.F.R. Ferrari², D.R. Fior-Chadi³, R. Camarini⁴ and L.F. Felício⁵

¹Centro de Matemática, Computação e Cognição, Universidade Federal do ABC, São Paulo, SP, Brasil

²Departamento de Genética e Biologia Evolutiva, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP, Brasil

³Departamento de Fisiologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, SP, Brasil

⁴Departamento de Farmacologia, Instituto de Ciências Biomédicas, Universidade de São Paulo, São Paulo, SP, Brasil

⁵Departamento de Patologia, Faculdade de Medicina Veterinária e Zootecnia, Universidade de São Paulo, São Paulo, SP, Brasil

Abstract

The periaqueductal gray (PAG) has been reported to be a location for opioid regulation of pain and a potential site for behavioral selection in females. Opioid-mediated behavioral and physiological responses differ according to the activity of opioid receptor subtypes. The present study investigated the effects of the peripheral injection of the kappa-opioid receptor agonist U69593 into the dorsal subcutaneous region of animals on maternal behavior and on *Oprk1* gene activity in the PAG of female rats. Female Wistar rats weighing 200-250 g at the beginning of the study were randomly divided into 2 groups for maternal behavior and gene expression experiments. On day 5, pups were removed at 7:00 am and placed in another home cage that was distant from their mother. Thirty minutes after removing the pups, the dams were treated with U69593 (0.15 mg/kg, sc) or 0.9% saline (up to 1 mL/kg) and after 30 min were evaluated in the maternal behavior test. Latencies in seconds for pup retrieval, grouping, crouching, and full maternal behavior were scored. The results showed that U69593 administration inhibited maternal behavior ($P < 0.05$) because a lower percentage of U69593 group dams showed retrieval of first pup, retrieving all pups, grouping, crouching and displaying full maternal behavior compared to the saline group. Opioid gene expression was evaluated using real-time reverse-transcription polymerase chain reaction (RT-PCR). A single injection of U69593 increased *Oprk1* PAG expression in both virgin ($P < 0.05$) and lactating female rats ($P < 0.01$), with no significant effect on *Oprm1* or *Oprd1* gene activity. Thus, the expression of kappa-opioid receptors in the PAG may be modulated by single opioid receptor stimulation and behavioral meaningful opioidergic transmission in the adult female might occur simultaneously to specific changes in gene expression of kappa-opioid receptor subtype. This is yet another alert for the complex role of the opioid system in female reproduction.

Key words: Real-time reverse-transcription polymerase chain reaction; Opioid receptor; Gene expression; Opioid sensitivity; Kappa opioid receptor

Introduction

Opioidergic stimuli modulate various sensory systems with behavioral consequences for both male and female mammals (1-9). The periaqueductal gray (PAG) is likely to influence a number of brainstem sites that are critically involved in controlling both somatomotor (e.g., mating) and autonomic (e.g., thermoregulation) responses, as well as hypothalamic regions related to the control of generalized arousal and sensorimotor integration (10,11). The PAG is also known to play important roles in the modulation of nociceptive sensory transmission, regulation of the car-

diovascular system (12), vocalization (13), and expression of a variety of behaviors, including defensive (14), sexual (15), maternal (16-18), and feeding behaviors (15,19,20). The PAG occupies a pivotal position in the central nervous system that influences the selection of adaptive behavioral responses. This region receives both limbic information and a vast array of inputs from prefrontal cortical areas (21,22). Previous studies from our laboratory have suggested an integrative role of opioidergic transmission in the PAG in influencing behavioral selection during lactation (6,23-25).

Correspondence: L.F. Felício, Departamento de Patologia, FMVZ, USP, Av. Prof. Dr. Orlando M. de Paiva, 87, 05508-900 São Paulo, SP, Brasil. E-mail: lfelicio@usp.br

Received February 14, 2012. Accepted May 16, 2012. Available online June 1, 2012. Published September 3, 2012.

The role played by the PAG in maternal behavior may be dependent on adequate local expression of opioid-related genes.

Opioid receptors have been the focus of intense molecular, physiological, and pharmacological research. The PAG has a high density of opioid receptors and activation of these metabotropic, G-protein-coupled receptors modulates various aspects of reproductive behavior in females. There are three types of opioid receptors: mu, delta, and kappa (26). Multiple reproductive experiences influence both gene activity and opioid receptor expression in the PAG. A preliminary study suggested that opioidergic stimulation modulates opioid gene expression in the female PAG (27,28).

Our group has shown that the morphine-induced inhibition of maternal behavior is plastic and adaptive and that opioidergic transmission in the rostral PAG plays a role in this process. In addition, it has been demonstrated that both repeated treatment with morphine and reproductive experience have effects on the expression of opioid receptors. The present study further extends this line of research by investigating the hypothesis that one stimulation would modify the activity of a specific receptor gene. Although the major target of morphine and naloxone is *Oprm1*, the acute inhibition of maternal behavior induced by kappa receptor stimulation needs more investigation. Because kappa stimulation is behaviorally meaningful, this study was designed to specifically investigate the dynamics of kappa on the PAG. In order to test for indirect or nonspecific effects, the kappa activity of *Oprm1* and *Oprd1* was also measured.

The present study investigated the effects of U69593, a kappa-opioid receptor agonist, on mu-, kappa-, and delta-opioid receptor *Oprm1*, *Oprk1*, and *Oprd1* gene activity in the PAG of both lactating and virgin female rats. This was performed by evaluating opioid gene expression using real-time reverse-transcription polymerase chain reaction (RT-PCR).

The aim of the present study was to determine whether a single kappa opioidergic stimulation using the kappa agonist U69593 can promote immediate changes in the activity and expression of the kappa-opioid receptor gene in the PAG of female animals. In order to test for indirect effects of such kappa stimulation on other opioid receptor genes, the activity of mu and delta receptor genes was also measured. Both virgin and lactating rats were tested to determine whether the physiological state influences the response to this agonist.

Material and Methods

Animals and housing

Female Wistar rats weighing 200-250 g at the beginning of the study were obtained from Faculdade de Medicina Veterinária, Universidade de São Paulo. The animals were housed in polypropylene cages (32 x 40 x 18 cm) with 3

animals per cage, under conditions of controlled temperature ($22 \pm 2^\circ\text{C}$), a 12/12-h light/dark cycle (lights on at 6:00 am), and free access to food and water during the experimental procedure. All animal procedures were in accordance with the guidelines of the Committee on the Care and Use of Laboratory Animal Resources, National Research Council, USA. We attempted to minimize the number of rats used, and every effort was made to ensure that no rat suffered unnecessarily.

For mating, 2 female rats were placed overnight with 1 sexually experienced male rat. The day on which sperm was observed in the vaginal lavage was designated day 1 of pregnancy. Pregnant rats were housed 2 per cage until day 18 of pregnancy and then individually until the end of the experiments. At parturition, all pups were examined externally, sexed, and weighed, with 8 pups (4 males and 4 females) left with each dam until the maternal behavior test (experiment 1) and gene expression studies (experiment 2).

Maternal tasks

On day 5 of lactation, pups were removed at 7:00 am and placed in another home cage that was distant from their mother. Thirty minutes after removing the pups, the dams were acutely challenged with U69593 ([5- α ,7- α ,8- β]-[+]-*N*-methyl-*N*-{7-(1-pyrrolidinyl)-1-oxaspiro[4.5]dec-8-yl} benzeneacetamide; Sigma Aldrich, USA, 0.15 mg/kg, sc) or saline. Thirty minutes after the injections were given to the dams, all pups were placed back with their mothers, and maternal behavior testing began. Latencies in seconds for pup sniffing, retrieval, grouping, crouching, and full maternal behavior were scored (2,3). Animals were scored as fully maternal if they retrieved all 8 pups to the nest and displayed nursing behavior with their back arched over the pups for 3 consecutive minutes. If animals were not fully maternal after 30 min of continuous observation, they were checked every 15 min up to 60 min and then hourly until full maternal behavior was observed. Events observed after the first 30 min of continuous observation were recorded at the time of first observation (e.g., if full maternal behavior was first observed at 60 min, the full maternal behavior latency was scored as 60 min, or 3600 s). The same criterion was used for all other responses. For all behavioral tests, the observers were blind to the treatment of the test subjects.

Opioid-selective agonist treatment

Sixty virgin and lactating rats were randomly divided into two (experiment 1) and four (experiment 2) groups of 10 animals each. They were treated with U69593 (0.15 mg/kg), saline, or no treatment (i.e., no experimental procedures were conducted on these animals, which were defined as blank). Brains were collected on day 5 of lactation. After 1 h, the animals were decapitated for PAG dissection using a blade dissecting approximately 2 cm² around the coordinate anteroposterior: 6.0; dorsoventral: 4.3; mediolateral: 0.6

coordinates (29). In both experiment, because the results for the saline and no treatment (blank) groups were similar ($P > 0.5$), they are presented as a single group ($N = 10$).

RNA extraction, cDNA construction, and real-time RT-PCR quantification of opioid gene expression

Total RNA was extracted from each tissue sample using TRIZOL reagent (Invitrogen Life Technologies, USA). Thus, immediately after euthanasia by decapitation, the PAG (approximately 40 mg, removed with a scalpel) was suspended in 1 mL ice-cold TRIZOL, and total RNA was extracted according to manufacturer instructions. The final RNA pellets were resuspended in 50 μ L diethyl-pyrocyanate-treated water. The total RNA concentrations were measured spectrophotometrically at 260 nm, and the integrity of RNA samples was analyzed on 1.5% agarose gel (Sigma, USA) containing ethidium bromide (0.5 μ g/mL). Total RNA was then treated with DNase I before further processing and stored at -80°C .

Oligo DT primers (1 μ L) and dNTPs (1 μ L) were added to the total RNA sample (5 μ g) and incubated at 65°C for 5 min; 5X buffer (4 μ L; Superscript II RNase H-reverse transcriptase), DTT (1 M, 2 μ L), and RNaseOUT (1 μ L) were then added, and the mixture was incubated at 42°C for 2 min. Superscript II (1 μ L) was added and the mixture was incubated at 42°C for 50 min. An additional incubation was performed at 70°C for 15 min. One microliter RNaseH was added for the removal of remaining RNA and the mixture was incubated at 37°C for 20 min. All reagents were purchased from Invitrogen Life Technologies.

Real-time RT-PCR was performed using an ABI Prism 7000 Sequence Detection System (Applied Biosystems, USA) with TaqMan Universal Master Mix (part No. 4304437; Applied Biosystems). PCR primers (Applied Biosystems) and the TaqMan probes for *Oprm1*, *Oprd1*, *Oprk1*, and 18S quantification were selected using the Primer Express software (Applied Biosystems), verified by a BLAST search of GenBank, and labeled Rn00565144_m1 for *Oprm1*, Rn00567737_m1 for *Oprk1*, RN00561699_m1 for *Oprd1*, and 4319413E for 18S, which was used as a housekeeping control. The primers were chosen to amplify a 65-bp fragment. The internal TaqMan probe (FAM-3'-TCTGGCA CCTCTCTTT-5'-NFQ) was designed following the general rules outlined by the manufacturer and carried a 5'-reporter

dye [6-carboxy fluorescein (FAM)] and a 3'-non-fluorescent quencher dye (NFQ). The primers and probes were used with 100% efficiency at final concentrations of 0.9 and 0.25 μ M, respectively.

The thermal cycling conditions for cDNA quantification assays were established according to ABI Prism 7000 Sequence Detection System parameters. Relative gene expression data were analyzed by the $2^{-\Delta\Delta\text{CT}}$ method (30).

Statistical analysis

Behavioral data were analyzed by comparing the means using the Student *t*-test. The percents of animals showing each behavior during the 30-min test were compared by the Fisher test. Mean gene expression data were compared by the Student *t*-test. Values of $P < 0.05$ were considered to be statistically significant.

Results

Experiment 1

The 0.15 mg/kg dose of U69593 induced no sedation. Comparison of the saline and U69593 groups revealed that a lower percentage of U69593 group dams showed retrieval of first pup, and retrieving all pups, grouping, crouching and displaying full maternal behavior. Importantly, latencies to sniff the pups did not differ among groups (Table 1). Animals in the U69593 group continued digging in the wood chips and building nests instead of retrieving pups, resulting in longer latencies to retrieve the pups. Such disruptive effects were not observed in the saline group.

Experiment 2

Acute U69593 treatment significantly increased *Oprk1* expression in virgin female rats ($P < 0.05$). This treatment did not induce alterations in *Oprd1* or *Oprm1* gene expression in the PAG of virgin female rats (Figure 1). In lactating female rats, U69593 treatment increased only *Oprk1* expression ($P < 0.01$). No other significant differences were found in opioid receptor gene expression in lactating rats treated with this drug (Figure 1) in the PAG.

Discussion

The PAG belongs to a primary anatomical pathway me-

Table 1. Effects of treatment with a selective agonist kappa-opioid receptor (U69593) on maternal behavior on day 5 of lactation.

Groups	Latency to sniff the 1st pup (s)	Retrieving 1st pup	Retrieving all pups	Grouping	Crouching	Full maternal behavior
Saline	10 \pm 2.8	100	100	100	100	100
U69593 (0.15 mg/kg)	12 \pm 3.4	50*	0*	0*	0*	0*

Data are reported as percentage of animals observed showing the listed behaviors during the first 30 min. Latency to sniff the first pup is reported in seconds (s). * $P < 0.05$ compared to the saline group (Fisher test).




Figure 1. Real-time RT-PCR of 5 µg total RNA extracted from the periaqueductal gray of adult virgin (A, B, C) and lactating female rats (D, E, F) in relation to expression of *Oprm1*, *Oprk1* and *Oprd1*. Data are reported as means ± SEM. *P < 0.05 compared to the blank group (ANOVA).

diating opioid-based analgesia. Morphine administration produces greater antinociception in males than females (31). In reproductive females, physiological activation of one or more PAG opioid receptors interferes with nursing and other aspects of maternal behavior, such as pup retrieval and pup grooming. The physiological role of female PAG opioid receptors, particularly the kappa receptor, has not been well established. The ventrolateral PAG column of the caudal part might be involved in the expression of kyphosis toward pups, which is the arched-back posture that facilitates nursing (17). Maternal aggression and survival strategies involving stressful situations have also been reported to be influenced by opioid receptors (16,17,32). A previous study reported that stimulation of kappa-opioid receptors using different doses of U69593 interferes with maternal behavior (8). Behavioral data reported in the present study show that peripheral injection of 0.15 mg/kg of the kappa agonist inhibits maternal behavior in lactating rats. On the other hand, other investigators administered a kappa 1 agonist (U69596, 0.1 mg/kg) and found no effect on maternal behavior in lactating rats. The larger dose used in the present study may be responsible for this difference. Alternatively, since another drug was used, kinetic differences may have played a role (33).

A previous study from our laboratory has reported the presence of kappa receptor protein in the PAG of both virgin and lactating rats (8). The present study investigated the expression of *Oprm1*, *Oprk1*, and *Oprd1* in the PAG of female rats by evaluating the gene expression

of mu-, kappa-, and delta-opioid receptors in the PAG of animals treated with an acute injection of U69593. The present data suggest that agonist-induced changes in gene expression are similar in both lactating and virgin animals. Treatment with a single peripheral injection of the kappa-opioid agonist U69593 significantly increased kappa receptor gene expression in both lactating and virgin rats. No significant changes were observed for mu or delta receptor genes after this drug challenge. Consistent with its binding affinity for kappa receptors, this action of U69593 on *Oprk1* gene activity suggests a specificity of U69593 for actions on the gene encoding the kappa receptor. In the present study, the behavioral effects of acute kappa opioidergic stimulation in lactating rats were confirmed. In addition, the present data add knowledge to this scenario by demonstrating a specific effect of a kappa agonist on a site that has been functionally implicated in behavioral selection during lactation. Acute stimulation increased *Oprk1* gene activity in both virgin and lactating animals. This may indicate a lack of endocrine modulation of this response. Thus, a specific immediate and likely compensatory gene response to kappa receptor stimulation is described in animals acutely exposed to a low dose of U69593. Such response may play a role in plastic mechanisms, resulting in different responses to similar pharmacological challenges. This hypothesis is to be tested in future studies.

The present study shows that by stimulating a kappa-opioid receptor both inhibition of maternal behavior and enhancement of *Oprk1* are achieved. Although more stud-

ies are necessary to elucidate the functional relationship between these effects, a specific role for the kappa receptor in the context of reproductive behavior now seems feasible. This is yet another alert for the complex role of the opioid system in female reproduction.

References

- Mann PE, Felicio LF, Bridges RS. Investigation into the role of cholecystokinin (CCK) in the induction and maintenance of maternal behavior in rats. *Horm Behav* 1995; 29: 392-406.
- Miranda-Paiva CM, Felicio LF. Differential role of cholecystokinin receptor subtypes in opioid modulation of ongoing maternal behavior. *Pharmacol Biochem Behav* 1999; 64: 165-169.
- Miranda-Paiva CM, Nasello AG, Yin AJ, Felicio LF. Morphine pretreatment increases opioid inhibitory effects on maternal behavior. *Brain Res Bull* 2001; 55: 501-505.
- Miranda-Paiva CM, Ribeiro-Barbosa ER, Canteras NS, Felicio LF. A role for the periaqueductal grey in opioidergic inhibition of maternal behaviour. *Eur J Neurosci* 2003; 18: 667-674.
- Patti CL, Frussa-Filho R, Silva RH, Carvalho RC, Kameda SR, Takatsu-Coleman AL, et al. Behavioral characterization of morphine effects on motor activity in mice. *Pharmacol Biochem Behav* 2005; 81: 923-927.
- Patti CL, Kameda SR, Carvalho RC, Takatsu-Coleman AL, Lopez GB, Niigaki ST, et al. Effects of morphine on the plus-maze discriminative avoidance task: role of state-dependent learning. *Psychopharmacology* 2006; 184: 1-12.
- Yim AJ, Miranda-Paiva CM, Florio JC, Oliveira CA, Nasello AG, Felicio LF. A comparative study of morphine treatment regimen prior to mating and during late pregnancy. *Brain Res Bull* 2006; 68: 384-391.
- Teodorov E, Tomita AT, Banon GP, Gil IG, Bernardi MM, Felicio LF. Behavioral effects of acute stimulation of kappa-opioid receptors during lactation. *Pharmacol Biochem Behav* 2008; 90: 534-539.
- Cruz AM, Maiorka PC, Canteras NS, Sukikara MH, Felicio LF. Morphine treatment during pregnancy modulates behavioral selection in lactating rats. *Physiol Behav* 2010; 101: 40-44.
- Cameron AA, Khan IA, Westlund KN, Cliffer KD, Willis WD. The efferent projections of the periaqueductal gray in the rat: a *Phaseolus vulgaris*-leucoagglutinin study. I. Ascending projections. *J Comp Neurol* 1995; 351: 568-584.
- Cameron AA, Khan IA, Westlund KN, Willis WD. The efferent projections of the periaqueductal gray in the rat: a *Phaseolus vulgaris*-leucoagglutinin study. II. Descending projections. *J Comp Neurol* 1995; 351: 585-601.
- Bandler R, Shipley MT. Columnar organization in the mid-brain periaqueductal gray: modules for emotional expression? *Trends Neurosci* 1994; 17: 379-389.
- Jurgens U. The role of the periaqueductal grey in vocal behaviour. *Behav Brain Res* 1994; 62: 107-117.
- Keay KA, Bandler R. Parallel circuits mediating distinct emotional coping reactions to different types of stress. *Neurosci Biobehav Rev* 2001; 25: 669-678.
- Sakuma Y, Pfaff DW. Facilitation of female reproductive behavior from mesencephalic central gray in the rat. *Am J Physiol* 1979; 237: R278-R284.
- Lonstein JS, Stern JM. Role of the midbrain periaqueductal gray in maternal nurturance and aggression: c-fos and electrolytic lesion studies in lactating rats. *J Neurosci* 1997; 17: 3364-3378.
- Lonstein JS, Stern JM. Site and behavioral specificity of periaqueductal gray lesions on postpartum sexual, maternal, and aggressive behaviors in rats. *Brain Res* 1998; 804: 21-35.
- Moura LM, Canteras NS, Sukikara MH, Felicio LF. Morphine infusions into the rostralateral periaqueductal gray affect maternal behaviors. *Braz J Med Biol Res* 2010; 43: 899-905.
- Carrive P. The periaqueductal gray and defensive behavior: functional representation and neuronal organization. *Behav Brain Res* 1993; 58: 27-47.
- Van Erp AM, Kruk MR, Meelis W, Veening JG. Periaqueductal gray lesions do not affect grooming, induced electrically in the hypothalamic paraventricular area in the rat. *Behav Brain Res* 1993; 59: 95-101.
- Floyd NS, Price JL, Ferry AT, Keay KA, Bandler R. Orbitomedial prefrontal cortical projections to distinct longitudinal columns of the periaqueductal gray in the rat. *J Comp Neurol* 2000; 422: 556-578.
- Mota-Ortiz SR, Sukikara MH, Felicio LF, Canteras NS. Afferent connections to the rostralateral part of the periaqueductal gray: a critical region influencing the motivation drive to hunt and forage. *Neural Plast* 2009; 2009: 612698.
- Sukikara MH, Mota-Ortiz SR, Baldo MV, Felicio LF, Canteras NS. A role for the periaqueductal gray in switching adaptive behavioral responses. *J Neurosci* 2006; 26: 2583-2589.
- Sukikara MH, Platero MD, Canteras NS, Felicio LF. Opiate regulation of behavioral selection during lactation. *Pharmacol Biochem Behav* 2007; 87: 315-320.
- Sukikara MH, Mota-Ortiz SR, Baldo MV, Felicio LF, Canteras NS. The periaqueductal gray and its potential role in maternal behavior inhibition in response to predatory threats. *Behav Brain Res* 2010; 209: 226-233.
- Mansour A, Fox CA, Akil H, Watson SJ. Opioid-receptor mRNA expression in the rat CNS: anatomical and functional implications. *Trends Neurosci* 1995; 18: 22-29.
- Teodorov E, Modena CC, Sukikara MH, Felicio LF. Preliminary study of the effects of morphine treatment on opioid receptor gene expression in brain structures of the female rat. *Neuroscience* 2006; 141: 1225-1231.
- Teodorov E, Bernardi MM, Ferrari MF, Fior-Chadi DR, Felicio LF. Plasticity of opioid receptors in the female periaqueductal gray: multiparity-induced increase in the activity of genes encoding for mu and kappa receptors and a post-translational decrease in delta receptor expression. *J Mol Neurosci* 2011;

Acknowledgments

Research supported by FAPESP (#2010/50415-4) and CNPq grants to L.F. Felício (#478765/04-05) and E. Teodorov (#2004/54696-0).

- 43: 175-181.
29. Paxinos G, Watson C. *The rat brain: in stereotaxic coordinates*. Amsterdam: Elsevier; 2009.
 30. Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 2001; 25: 402-408.
 31. Loyd DR, Murphy AZ. The role of the periaqueductal gray in the modulation of pain in males and females: are the anatomy and physiology really that different? *Neural Plast* 2009; 2009: 462879.
 32. Pavesi E, Enck MC, De Toledo CA, Terenzi MG. Disruption of maternal behaviour by acute conspecific interaction induces selective activation of the lateral periaqueductal grey. *Eur J Neurosci* 2007; 26: 2055-2065.
 33. Mann PE, Kinsley CH, Bridges RS. Opioid receptor subtype involvement in maternal behavior in lactating rats. *Neuroendocrinology* 1991; 53: 487-492.