



## Review Article

# Modern acupuncture-like stimulation methods: a literature review

Min-Ho Jun, Young-Min Kim, Jaeuk U. Kim\*

KM Fundamental Research Division, Korea Institute of Oriental Medicine, Daejeon, Korea

## ARTICLE INFO

## Article history:

Received 12 August 2015

Received in revised form

23 September 2015

Accepted 24 September 2015

Available online 3 October 2015

## Keywords:

acupuncture-like stimulation devices

electrical stimulation

laser stimulation

magnetic stimulation

ultrasonic stimulation

## ABSTRACT

Acupuncture therapy has been proved to be effective for diverse diseases, symptoms, and conditions in numerous clinical trials. The growing popularity of acupuncture therapy has triggered the development of modern acupuncture-like stimulation devices (ASDs), which are equivalent or superior to manual acupuncture with respect to safety, decreased risk of infection, and facilitation of clinical trials. Here, we aim to summarize the research on modern ASDs, with a focus on featured devices undergoing active research and their effectiveness and target symptoms, along with annual publication rates. We searched the popular electronic databases Medline, PubMed, the Cochrane Library, and Web of Science, and analyzed English-language studies on humans. Thereby, a total of 728 studies were identified, of which 195 studies met our inclusion criteria. Electrical stimulators were found to be the earliest and most widely studied devices (133 articles), followed by laser (44 articles), magnetic (16 articles), and ultrasound (2 articles) stimulators. A total of 114 studies used randomized controlled trials, and 109 studies reported therapeutic benefits. The majority of the studies (32%) focused on analgesia and pain-relief effects, followed by effects on brain activity (16%). All types of the reviewed ASDs were associated with increasing annual publication trends; specifically, the annual growth in publications regarding noninvasive stimulation methods was more rapid than that regarding invasive methods. Based on this observation, we anticipate that the noninvasive or minimally invasive ASDs will become more popular in acupuncture therapy.

© 2015 Korea Institute of Oriental Medicine. Published by Elsevier. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

## 1. Introduction

Stimulation of acupoints and meridians has been an important therapeutic modality in traditional Eastern medicine, and it has also become popular in the West, as its clinical effectiveness has been demonstrated through extensive research.

Acupuncture and related modern technologies are increasing in popularity worldwide. According to a 2002 World Health Organization report, acupuncture treatment was shown to be effective in controlled trials of 29 diseases, symptoms, or conditions.<sup>1</sup> However, the safety of acupuncture has engendered controversy with respect to infection, inflammation, and pain management.

\* Corresponding author. KM Fundamental Research Division, Korea Institute of Oriental Medicine, 1672 Yuseongdaero, Yuseong-Gu, Daejeon 305-811, Korea.

E-mail address: [jaeukkim@kiom.re.kr](mailto:jaeukkim@kiom.re.kr) (J.U. Kim).

<http://dx.doi.org/10.1016/j.imr.2015.09.005>

2213-4220/© 2015 Korea Institute of Oriental Medicine. Published by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Clinical effectiveness of acupuncture has widely been studied during the past four decades.<sup>2-6</sup> In addition to the demonstrated effectiveness of traditional acupuncture practices, increased demand has arisen for the development of modern acupuncture-like stimulation devices (ASDs), which are simpler to quantify and standardize and are less dependent on the manipulation techniques of individual clinicians.

The first modern ASD dates back to the early 1950s, which was based on electrical stimulation (ES).<sup>7,8</sup> In addition to its long history, ES is the most extensively studied ASD.<sup>9</sup> Recently, however, several types of ASDs have extensively been studied for their clinical effectiveness and noninferiority to manual acupuncture, including laser stimulation (LS)<sup>10</sup> and magnetic stimulation (MS).<sup>11</sup> In this review, we summarize recent studies of popular ASDs. We first describe the most popular types of ASDs, discuss their clinical effectiveness and target symptoms, and finally, discuss the annual research trends regarding popular ASDs.

## 2. Methods

To analyze the popularity and features of methods for stimulation of acupoints, we searched for studies in the Medline, PubMed, Cochrane Library, and Web of Science electronic databases from their inception to June 2014. First, we searched for studies related to acupuncture or acupoint stimulation, which yielded > 22,000 studies, of which approximately 20,000 were redundant. Among the latter studies, approximately 3000 were related to moxibustion, 1600 to massage (or acupressure), 200 to the cupping method, 5400 to ES, 900 to LS, 700 to MS, and 300 to ultrasound stimulation (US). To narrow the search scope to ASDs, we refined the search to [(acupoint\* or "acupuncture point\*" or meridian\*) and (stimul\* or irritat\* or excit\* or response or respon\* or react\* or reflex or measur\* or diagnos\*) and (electric\* electro\* or magnet\* or infrared or IR or laser or ultraviolet or UV or ultraso\*) not (rat or monkey or dog or pig or cat or mouse or mice or rabbit or rodent\*)]. We excluded laboratory experiments on animals, studies that were not written in English, and reviews. We searched 728 articles obtained from the electronic databases, excluding 489 articles that included studies on animals, manual

acupuncture-only clinical trials, non-English-language articles, and review articles by screening the titles and abstracts. A total of 44 studies were excluded from the selected 239 articles because of duplication. Finally, 195 studies met the inclusion criteria and were evaluated in detail. The topics of these 195 articles were ES (133), LS (44), MS (16), and US (2), as shown in Fig. 1. Prior to describing the results of the detailed analysis, we introduce the features and research history of ES, LS, MS, and US in the following sections.

### 2.1. Electrical stimulation

Low electrical impedance and high conductance are recognized as typical electrical properties of acupoints and meridians.<sup>12-14</sup> In the Western hemisphere, the electrical properties of acupoints and meridians have been investigated since the 1950s. In 1958, Niboyet and Mery<sup>15</sup> reported the points with low skin impedance using the Wheatstone bridge, whereas in 1962, Kramar<sup>16</sup> showed that acupoints have high capacitance compared with neighboring points. Voll<sup>7</sup> devised an ES device to apply to acupoints and meridians, thereby establishing a method that was called "electroacupuncture according to Voll." This method of Voll<sup>7</sup> greatly stimulated clinical and research activities associated with ES at acupoints and meridians. In the East in 1956, Nakatani<sup>8</sup> reported that electrical pathways connected the points with low skin resistance and named them "Ryodoraku." Today, ES can be classified into five types: electroacupuncture (EA), transcutaneous electrical acupoint stimulation (TEAS), auricular electroacupuncture (AEA), transcutaneous electrical nerve stimulation (TENS), and electrical heat acupuncture (EHA). EA is an electrical, minimally invasive stimulation technique applied to acupoints. TEAS is an electrical, noninvasive stimulation technique applied to acupoints. AEA is a subtype of EA applied to acupoints of the ear. TENS is an electrical, noninvasive stimulation technique applied to the nervous system (nonacupoints). EHA is similar to EA with the exception that a needle heated by an electric current is used at acupoints. Of the 133 articles on ES, 54 pertained to EA, 69 to TEAS, six to AEA, three to TENS, and one to EHA. To simplify the discussion, we categorized ES into EAs and TEASs, where EAs represented




Fig. 1 – Flow diagram of literature survey.

all invasive techniques, such as EA, AEA, and EHA, and TEASs included all noninvasive techniques, such as TEAS, auricular TEAS, and TENS.

2.2. Laser stimulation

Studied since the 1970s, LS is used to expose acupoints of the human body to low-energy laser beams. A review article<sup>17</sup> noted that studies using LS were conducted between 1970 and 1972 in the USSR. Nevertheless, Friedrich Plog's<sup>18</sup> study published in 1976 is well known as the first report of implementation of LS at acupoints. Since the 1980s, LS has been recognized as an effective method for stimulating acupoints without needles. Applications of LS at acupoints were mostly described as noninvasive in the studies reviewed, with only a few being described as invasive. Here, we do not distinguish invasive techniques from noninvasive stimulation.

2.3. Magnetic and ultrasonic stimulation

MS is used to access the body's magnetic fields by stimulating acupoints, and MS of acupoints has been studied since the 1970s. Transcranial magnetic stimulation is one of the most frequently used MSs and was introduced by Barker<sup>19</sup> in 1985. In 1980, Inoue<sup>20</sup> applied for a patent for a device used for MS of body acupoints, and in 1982, Katayama<sup>21</sup> reported

the meridian magnetic analgesia of acupuncture stimulation (published in Japanese). The MS used in all 16 papers consisted of noninvasive stimulations at acupoints.

US is used to irritate acupoints using a narrow, cylindrical, high-frequency beam of sound. Characteristics of phonation and sound transmission in meridians were reported in the 1980s, and a study on US of acupoints was published by Jin<sup>22</sup> in 1984. Only two studies that we identified in the electronic databases were relevant.

3. Results

The aforementioned four types of ASDs were classified into the following 13 categories according to the stimulation purposes: (a) analgesic effect; (b) pain relief; (c) physiological change; (d) improvement of the alimentary system; (e) prevention of nausea and vomiting; (f) recovery of muscle fatigue or improvement of muscle strength; (g) reduction of body weight; (h) treatment of depression; (i) treatment of addiction, such as addiction to tobacco, narcotics, and alcohol; (j) treatment of stroke; (k) treatment of various diseases; (l) characteristics of stimulation; and (m) brain activity. Fig. 2 shows how the four types of ASDs were distributed between the 13 categories for research purposes. It also shows the ratio of randomized controlled trials (RCTs) to efficacies for the 13 categories. The




Fig. 2 – Distribution of the four ASDs with respect to the 13 research categories from (a) to (m) whereby the numbers of RCTs and the therapeutic effectiveness are shown for each category. Numbers reflect the article counts, with the numbers in parentheses for the four types indicating the number of cases of overlap between the stimulations, and the numbers in parentheses below the efficacy (%) are presented when the efficacy was unclear. In RCT (%) = A/(A + B) % and (O:X = A:B), A is the number of RCTs and B is the number of non-RCTs. The same formula was applied to the efficacy percentages. When the efficacy was unclear, indicated by the numbers in parentheses, we considered those studies as not effective in computing the percent values. For example, % value = A/(A + B + b) for efficacy [O:X = A:B(b)]. ASD, acupuncture-like stimulation device; ES, electrical stimulation; LS, laser stimulation; MS, magnetic stimulation; RCT, randomized controlled trial; US, ultrasound stimulation. O = yes, X = no. Example: RCT (O:X) = (RCT:non-RCT), Efficacy (O:X) = (efficacious:not efficacious).

**Table 1 – Summary of studies on analgesic effects with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Jiang et al <sup>23</sup>	TEAS	LI4, PC8	Healthy	46 individuals, TEAS/mock TEAS	Analgesic effect
Wang et al <sup>24</sup>	TEAS	LI4, PC6, ST36	Sinusotomy	60 patients, random TEAS/control: no stimulation	Analgesic effect
Zhang et al <sup>25</sup>	TEAS	T3 acupoints	Ambulatory surgery	72 women, random TEAS/sham	Recovery & decrease of anesthesia
Wu et al <sup>35</sup>	EA	GV1, BL57	Hemorrhoids	120 cases, random EA (40)/sham EA (40)/blank (40)	Effects of preemptive analgesia
Lan et al <sup>26</sup>	TEAS	Bi PC6, LI14, ipsilateral to surgery site	Total hip arthroplasty surgery	68 elderly patients, random TEAS/sham TEAS	Reduction of postoperative analgesic requirement
Zheng et al <sup>36</sup>	EA	ST36, GB31, GV24, EX-HN3	Orotracheally intubated patients	45 patients, random, no treatment/sham EA/EA	Sedation & analgesia
Cheing and Chan <sup>27</sup>	TEAS/TENS	Right elbow LI11, Nonacupoint (right superficial radial nerve)	Healthy	Randomized controlled trial, 45 individuals, random TEAS (15)/TENS (15)/control-no stim (15)	Hypoalgesic effects (acupuncture points & nerve points)
DeSantana et al <sup>31</sup>	TENS	Around the incision	Unilateral inguinal herniorrhaphy with epidural anesthetic technique	Prospective, randomized, double-blinded, placebo-controlled study, 40 patients, TENS (20)/placebo-TENS (20)	Hypoalgesic effect for postoperative pain
Barlas et al <sup>37</sup>	EA	Bi LI10, HT5/ipsilateral GB34, ST38	Healthy (acupuncture naïve)	Randomized, double-blinded, placebo-controlled study, 48 volunteers, control/placebo-EA no stim/high-intensity EA/low-intensity EA	Hypoalgesic response
Leung et al <sup>40</sup>	EA	Left SP1, LR1	Healthy	13 individuals, EA/before-EA/after-EA (time sequence)	Analgesic benefit
Litscher <sup>42</sup>	Acupressure/MA/LS	EX-HN3	Healthy	Randomized, controlled, blinded crossover trial, 20 volunteers, acupress/MA/LA; APs/non-APs	ECG similarities of acupressure-induced sedation & general anesthesia (all)
Zhang et al <sup>38</sup>	EA	Acupoints	Healthy	Eight8 individuals, EA/mock-EA	Analgesic effect
Attele et al <sup>28</sup>	TEAS	LI4, PC6	Healthy	22 individuals, TEAS/control	Analgesic effect
Chesterton et al <sup>29</sup>	TENS/TEAS	GB34, radial nerve or extrasegmental	Healthy	Randomized, double-blind, sham-controlled study, 240 participants, six6 TENS (180; 90 m, 90 f)/control (30; 15 m, 15 f)/sham TENS (30; 15 m, 15 f); 4/110 Hz, intensity, site	Hypoalgesic effect
Yuan et al <sup>30</sup>	TEAS	LI4, PC6	Healthy	20 individuals TEAS/morphine/TEAS + morphine/control	Analgesia effect
Morioka et al <sup>39</sup>	EA	ST36, GB34, BL60	Healthy	14 volunteers, EA/control	No difference in minimum alveolar anesthetic concentration
Lin et al <sup>41</sup>	EA	Bi ST36	Lower abdominal surgery	Randomly, 100 women, control (25)/sham-EA no stim (25)/LF-EA 2 Hz (25)/HF-EA 100 Hz (25)	Reduction of postoperative analgesic requirements & side effects (LF-EA, HF-EA)
Greif et al <sup>32</sup>	ATEAS	Auricular acupoints	Healthy	Randomized, double-blind, crossover trial, 20 volunteers (10 m, 10 f), ATEAS/no treatment	Reduction of anesthetic requirement
Chen et al <sup>33</sup>	TEAS/TENS	ST36/dermatomal level	Total abdominal hysterectomy or myomectomy procedures	Randomized controlled trial, 100 women, sham-TEAS no stim (25)/non-APs TEAS (25)/dermatomal-TENS (25)/TEAS (25)	TENS was as effective as TEAS, both were more effective than stim at non-APs



**Table 1 (Continued)**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Wang et al <sup>34</sup>	TEAS	LI4	Healthy women undergoing lower abdominal procedures	Random, 101 participants, PCA (26)/PCA + LP-TEAS (25)/PCA + HP-TEAS (25)/PCA + sham-TEAS no stim (25)	Decrease in PCA opioid requirement & opioid-related side effects (HP-TEAS)
Brokhaus and Elger <sup>43</sup>	LS/MA	Bi LI4, EX-UE	Healthy	Double-blind, 40 probationers, MA-LI4/LA-LI4, EX	Analgesic effect of MA on painful heat stim, no effect on pain (LA)

AP, acupuncture point; ASD, acupuncture-like stimulation device; ATEAS, auricular TEAS; EA, electroacupuncture; ECG, electrocardiogram; HF, high frequency; HP-TEAS, high power TEAS; LA, laser acupuncture; LF, low frequency; LP-TEAS, low power TEAS; LS, laser stimulation; MA, Manual acupuncture; PCA, patient-controlled analgesia; stim, stimulation; TEAS, transcutaneous electrical acupoint stimulation; TENS, transcutaneous electrical nerve stimulation.

numbers shown in the uppermost boxes in Fig. 2 signify the numbers of articles. The numbers of overlapping articles are shown in parentheses under the 13 categories of the four ASDs, and the numbers in parentheses below the efficacy (%) are presented when the efficacy was unclear.

To investigate the effectiveness of ASDs, we analyzed the efficacy of each stimulation type through the articles reporting effectiveness. The effectiveness of ES was stated in the fields of analgesic effect (94.7%), pain relief (90.9%), and reduction of nausea and vomiting (90.9%) based on the sample size of > 1000 trials. Based on the sample size of > 100 trials, ES was shown to be effective in improving the alimentary system (100%), improving muscle strength (100%), reducing body weight (100%), treating various addictions (60%), and treating stroke (100%), whereas LS was effective for pain relief (62.5%) and treating various addictions (100%). Based on a sample size of <100 trials, ES was shown to be a therapeutic possibility in various diseases such as orthostatic intolerance, autism spectrum disorders, supratentorial craniotomy, tinnitus, asthma, dyspnea, distress, and anxiety. LS presented potential in the treatment of nausea and vomiting, depression, menopausal symptoms, cholecystitis, renal failure, head injury, and interstitial cystitis. MS was a possible treatment for muscle and diving fatigue, whereas US demonstrated potential for relieving pain.

### 3.1. Analgesic effect

All the studies that reported an analgesic effect are shown in Table 1. Twelve articles reported an analgesic effect using TEASs,<sup>23-34</sup> seven articles reported an analgesic effect using EAs,<sup>35-41</sup> and two articles discussed LS.<sup>42,43</sup> A total of 834 individuals received TEASs to enable estimation of the analgesic effect of their clinical application using RCTs, and all articles reported that the TEASs had an analgesic or hypoalgesic effect or decreased opioid requirements. A total of 348 individuals received EAs, and an effect of the EAs on analgesia, sedation, hypoalgesia, pre-emptive analgesia, and reduction of analgesic requirements was found in 334 patients. In 20 individuals who received acupressure, manual acupuncture (MA), and LS, a sedative effect was observed, and an anesthetic effect was observed in 60 individuals who received LS. Two of the 21 papers<sup>39,43</sup> reported no significant analgesic effect of the EAs and LS.

### 3.2. Pain relief

As shown in Table 2, presenting the studies reporting the effect on pain relief, 15 articles reported using TEASs,<sup>44-58</sup> EAs were used in 18 studies,<sup>59-76</sup> other acupoint stimulations such as US were used in one study,<sup>77</sup> and LS was used in eight studies.<sup>78-85</sup> TEASs and EAs were compared in a total of 872 individuals to evaluate their effect on pain relief. Both had an effect on pain relief in two studies; however, the effect of EAs was reported to be superior to that of TEASs in one study. Of 1046 individuals who received TEASs, 926 experienced relief or a reduction in various types of pain. Of the 877 individuals who received EAs, 628 also experienced pain relief. Of the 435 individuals who received LS, 230 experienced relief of dysmenorrhea pain or carpal tunnel syndrome pain, whereas 50 individuals who received US experienced an effect on short-term segmental antinociception. Six of the 42 papers<sup>44,47,64,79,82,84</sup> reported no statistically significant effect on pain relief when TEASs, LS, EAs, LS combined with paracetamol and chlormezanone, and LS were applied to the acupoints of study participants.

### 3.3. Treatments of the alimentary system

As summarized in Table 3, ES (TEASs and EAs) was the primary ASD method for treating digestive disorders. Of these studies, seven that investigated TEAS<sup>86-92</sup> and three that evaluated EA<sup>93-95</sup> comprise this category. In total, 149 individuals who received TEAS experienced a beneficial effect on the alimentary system, as did 68 individuals who received EA. No study reported statistically insignificant results regarding stimulation of the alimentary system.

### 3.4. Prevention of nausea and vomiting

All the studies shown in Table 4 employed ASDs for the prevention of nausea and vomiting. ES was primarily applied for the prevention or treatment of nausea and vomiting, except for one study that used LS for this purpose. TEAS was the main method used for preventing nausea and vomiting: we retrieved nine articles on TEAS,<sup>96-104</sup> two on EA,<sup>105,106</sup> and one on LS.<sup>107</sup> A total of 830 individuals who received TEAS experienced an effect on prevention, reduction, or treatment of postoperative nausea and vomiting and nausea or vomiting. A total of 224 individuals who received EA also

**Table 2 – Summary of studies on pain relief with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Mucuk & Baser <sup>44</sup>	TEAS	LI4, SP6	Pregnant women	Bi LI4-TEAS (40)/SP6-TEAS (40)/control (40) no stim	Labor pain relief, not statistically significant
Sun et al <sup>45</sup>	TEAS	PC6	Driver fatigue	–	Able to withstand driver fatigue
Vassal et al <sup>46</sup>	TENS	Left common peroneal nerve	Healthy	20 individuals, TENS/sham TENS (left thigh)	Pain relief
Kim et al <sup>59</sup>	EA	Bi LI4, TE3, GV39, GV41, SP6, LR3, Ba Feng, Ba Xie	CIPN	Randomized, patient-assessor-blinded, controlled trial, 40 patients, EA (20)/sham EA (20)	Treatment for CIPN, trials
Lee et al <sup>60</sup>	EA	Bi ST36, GB39, SP9, PC6, LR3, GB41	PDN	3-armed, randomized, controlled pilot trial, EA (15)/sham EA (15)/usual care (15)	Treatment for PDN, trials
Mucuk et al <sup>47</sup>	TEAS	Bilateral LI4	Labor pain	Random TEAS/control no TEAS; all standard treatments	Pain relief, not statistically significant
Ni et al <sup>48</sup>	TEAS	Bilateral PC6	Children with congenital heart defects	70 eligible children, random, TEAS (34)/control-no stim (36)	Attenuation of myocardial injury in children undergoing cardiac surgery
Wu et al <sup>49</sup>	TEAS	LI4, SP6	Gynecology patients (primary dysmenorrhea)	Randomized controlled trial, 66 patients (f), TEAS (34)/control non-APs (32)	Mitigation of pain in dysmenorrhea
Yoshimizu et al <sup>72</sup>	EA/TEAS	For acupoints in trapezius muscle	Shoulder & neck pain	Randomized crossover trial, 90 patients, EA/TENS	Reduction in pain (EA > TEAS)
Musial et al <sup>73</sup>	EA	LI4, LI10	Healthy	Double-blind design, 125 individuals, EA (25)/tramadol (25)/ibuprofen (25)/placebo pill (25)/no treatment (25)	Reduction of experimentally induced ischemic pain
Choi et al <sup>61</sup>	EA/meditation	LI4, LI10	Vipassana meditators	Semirandomized trial, meditators(8)/nonmeditators (20)-EA/nonmeditators (20)-no EA	Reduction in the pain induced by SETT
Yeh et al <sup>50</sup>	TEAS	BL40, GB34, HT7, PC6	Spinal surgery receiving patients	Placebo- & sham-controlled study, random TEAS (30)/TEAS-sham point (30)/no TEAS (30)	Reduction in postoperative pain, analgesic usage
Montenegro et al <sup>51</sup>	TEAS	TE5, CV6	Healthy	32 volunteers, random TEAS/sham TEAS	Increase in the latency of pain threshold
Yeh et al <sup>53</sup>	TEAS	Acupoints	Lumbar spinal surgery	Randomized controlled repeated measures design, 99 patients, ES/sham-AP ES/no ES	Improvement of acute postoperative pain management without adversely affecting vital signs
Takamjani et al <sup>81</sup>	LS	Acupoints	Wrist pain	Randomized controlled trial, 70 women, LS (33)/control (37) no LS	Increase in mean value of pain threshold
Lee & Lee <sup>62</sup>	EA	Bi BL32, BL33, GB30	Chronic prostatitis/chronic pelvic pain syndrome	39 men, random 3 group exercise + EA/exercise + sham EA/exercise	Pain relief effect
Kempf et al <sup>78</sup>	LS	Bi SP6, LR3, LI4; right CV3, ST36	Minimum menstrual pain	Randomized controlled double blind pilot trial, 48 women, LA (18)/placebo-LA (30)	Dysmenorrhea treatment
Glazov et al <sup>82</sup>	LS	Acupoints	Chronic nonspecific low-back pain	Double blind, 2-group parallel randomized controlled trial, 100 participants, LA/sham-LA	Not showing a specific effect for chronic low-back pain

Table 2 (Continued)

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Chan et al <sup>74</sup>	EA	Acupoints on the wrist	Chronic neck pain	Single-blind, randomized, sham-controlled trial, 49 patients, EA (22)/sham-EA (27)	Significant improvements of chronic neck pain
Jubb et al <sup>63</sup>	EA	Acupoints	Osteoarthritic knee pain & disability	Blinded randomized trial, MA (34)/EA (34)/sham MA (34)	Symptomatic improvement
Srbely et al <sup>77</sup>	US	Right supraspinatus trigger point	Identifiable myofascial trigger points	Randomized controlled study, 50 individuals, random US/sham US (off)	Short-term segmental antinociceptive effects on TPs
Ye et al <sup>54</sup>	TEAS + PCA	LI4, PC8; Jiaogan, Shenmen, Shen, Waifei, Naogan, Pizhixia (ear acupoints)	Craniotomy & required pain relief following surgery	Randomized control, 40 patients, PCA + TEAS (20)/PCA (20)	Enhancement of the effect of pain relief & reduction of adverse reactions
Michalek-Sauberer et al <sup>64</sup>	AEA	Auricular shenmen, mouth, tooth	Molar tooth extraction	Prospective, randomized, double-blind, placebo-controlled study, 149 patients, AEA (76)/AMA (37)/sham AEA no stim no needle (36)	No reduction in either pain intensity or analgesic consumption in a molar tooth extraction model
Zhang et al <sup>65</sup>	EA	GB34, GB39	Healthy (right handedness)	12 volunteers, EA/sham-points EA/shallow EA subcutaneous needling	Pain relief
Yip et al <sup>55</sup>	TEAS + EMMW	–	Subacute neck or low-back pain	Randomly, 47 individuals, TEAS + EMMW (23)/control (24)	Reduction in pain intensity, stress, & stiffness level
Fang et al <sup>52</sup>	TEAS/EA	Acupoints	Periarthritis of shoulder at different stages	360 cases, TEAS (186)/EA (174)	Therapy for periarthritis of shoulder, no significant differences (TEAS/EA)
Aigner et al <sup>79</sup>	LS + paracetamol, chlormezanone	22 acupuncture points	Whiplash injuries	Prospective, randomized placebo-controlled trial, LA (23)/placebo-LA (22)	Ineffective in management of whiplash injuries
Sator-Katzenschlager et al <sup>66</sup>	AEA	Auricular 29, 55, 57	<i>In vitro</i> fertilization	94 women, random, AEA (32)/AMA (32)/pharm. (30)	Reduction of pain intensity
Wong et al <sup>75</sup>	EA	LI4, GB34, GB36, TE8	Operable non-small cell lung carcinoma patients who received thoracotomy	Random, 25 patients, EA (13)/sham-EA (12)	Management of post-thoracotomy wound pain
Weng et al <sup>56</sup>	TEAS	LI10, LI11	Tennis elbow pain for at least 3 mo	Randomly, 20 patients, 5 kHz modulated LF-TEAS 2 Hz (20)/5 kHz modulated HF-TEAS 100 Hz (20)/sham-TEAS, different time slots	Effective in the treatment of patients with tennis elbow pain (LF-TEAS, HF-TEAS)
Tsui & Cheing <sup>67</sup>	EA/EHA	6 acupuncture points	Chronic low-back pain	42 individuals, random EA/EHA/control; all exercise	Treatment of chronic low-back pain
Sator-Katzenschlager et al <sup>68</sup>	AEA	Auricular acupuncture points 29, 40, 55	Chronic low-back pain	Prospective, randomized, double-blind, controlled study, 61 patients, random AEA (31)/sham-AEA no stim (30)	Treatment of chronic low-back pain
Sator-Katzenschlager et al <sup>69</sup>	AEA	Cervical spine, shenmen, cushion	Chronic cervical pain patients without radicular symptoms with insufficient pain relief	Prospective, randomized, double-blinded, controlled study, 21 patients, EA (10)/control (11)	Treatment of chronic cervical pain

**Table 2 (Continued)**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Ng et al <sup>70</sup>	EA/TEAS	ST35/EX-LE4	OA-induced knee pain	Single-blinded randomized controlled trial, 24 individuals (1 m, 23 f), EA (8)/TEAS (8)/control standard therapy (8)	Reduction of OA-induced knee pain
Naeser et al <sup>80</sup>	LS/TENS	Shallow acupuncture points/wrist	CTS	Randomized, double-blind, placebo-controlled, crossover trial, 11 cases, red LS/IR LS/TEAS/sham (off)	Treating CTS pain
Tsui & Leung <sup>71</sup>	EA	GB34, ST38	Chronic tennis elbow	Single-blinded randomized controlled trial, 20 patients, MA/EA	Treating patients with tennis elbow
Zoghi & Jaberzadeh <sup>57</sup>	ATEAS/ATENS	4 auricular acupoints	Healthy	Double-blind within-subject design, randomly, 90 individuals, HV-ES (30)/HV-sham-ES non-APs (30)/no ES (30)	Increase in experimental pain threshold (HV-ES, sham)
Lorenazana <sup>58</sup>	TEAS	HT7, LI4	Episiotomy pain	Randomized, double-blind, controlled trial, 68 patients, TEAS (38)/control (30)	Relief of episiotomy pain (TEAS > lidocaine)
King et al <sup>83</sup>	ALS	Auricular acupoints	Healthy	80 individuals, ALS (41)/control (39) sham-ALS	Increase in mean pain threshold after treatment
Waylonis et al <sup>84</sup>	LS	Acupoints	Myofascial pain syndromes (fibrositis, fibromyalgia)	Crossover double-blind trials, 62 patients, LS/placebo	No statistical difference between the treatment and placebo groups
Kreczi & Klingler <sup>85</sup>	LS	Acupoints	Radicular and pseudoradicular pain syndromes	Prospective randomized single-blind crossover study, 21 patients, LS/mock LS	Mean pain levels (lower)
Ernst & Lee <sup>76</sup>	EA	LI4	Normal individuals	Crossover repeated-measure design, 5 individuals, control/EA/EA + naloxone/EA + placebo	Pain threshold increase

AEA, auricular electroacupuncture; ALS, auricular laser stimulation; AMA, auricular manual acupuncture; AP, acupuncture point; ASD, acupuncture-like stimulation device; ATEAS, auricular TEAS; ATENS, auricular TENS; CIPN, chemotherapy-induced peripheral neuropathy; CTS, carpal tunnel syndrome; EA, electroacupuncture; EHA, electrical heat acupuncture; EMMW, electromagnetic millimeter wave; ES, electrical stimulation; f, female; m, male; HF, high frequency; HV, high voltage; IR, infrared; LA, laser acupuncture; LF, low frequency; LS, laser stimulation; MA, manual acupuncture; OA, osteoarthritis; PCA, patient-controlled analgesia; PDN, painful diabetic neuropathy; pharm, pharmacological treatment; SETT, submaximum effort tourniquet technique; stim, stimulation; TEAS, transcutaneous electrical acupoint stimulation; TENS, transcutaneous electrical nerve stimulation; TP, trigger point; US, ultrasound stimulation.

experienced either the same effect or controlled emesis, whereas 40 individuals who received LS experienced a decrease in the incidence of vomiting. We observed that TEAS has been steadily applied in the prevention of nausea and vomiting, and exceeded EA in the number of clinical studies since 2003. This finding implies that the effectiveness of TEAS in preventing nausea and vomiting has been confirmed, and that TEAS was preferred to EA because of the infection risk and pain due to the use of needles with EA.

### 3.5. Improvement of the muscle system

Studies regarding ASDs that were related to the recovery of muscle fatigue or improvement of muscle strength are shown in Table 5. MS and ES were used to reduce muscle fatigue or improve muscle strength. This category included two studies on MS<sup>108,109</sup> and five studies<sup>110–114</sup> on ES. The two MS studies, which were conducted by the same research group, reported

the effective recovery of muscle fatigue. One study<sup>109</sup> reported better performance of MS than TEAS with respect to the therapeutic effect on muscle fatigue, and we expect more studies to validate this report.

### 3.6. Reduction in body weight

All the papers investigating the reduction in body weight were associated with ES, as shown in Table 6. EAs<sup>115,118,119</sup> and TEASs<sup>116,117</sup> were applied to facilitate the reduction in body weight. One study<sup>117</sup> stated that TEAS was as effective as EA in weight reduction. A total of 193 individuals who received ES experienced a reduction in body weight or fat, and an improvement in body mass index or body composition. All the studies reporting on the reduction in body weight claimed significant effects. More studies are required to substantiate the effectiveness of ES for body weight reduction.



**Table 3 – Summary of studies on the effects of the four ASDs on alimentary system**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
McNearney et al <sup>86</sup>	TEAS	PC6, ST36	SSc	17 patients, all TEAS	Enhancement of gastric myoelectrical functioning in SSc
Leung et al <sup>87</sup>	TEAS	LI4, PC6, ST36	Healthy	40 individuals, random TEAS/placebo TEAS	Reduction of rectal discomfort
Chen et al <sup>93</sup>	EA	ST36, ST37, ST25, ST28, CV4, CV6	Female constipation	Single-blind, randomized trial, 30 females, EA (14)/sham EA (16)	Improvement of constipation
Liu et al <sup>88</sup>	TEAS	PC6, ST36	Functional dyspepsia	Double-blind, crossover study, 27 patients, random acute-TEAS/chronic (2w) TEAS	Improvement of dyspepsia symptoms
Wang et al <sup>94</sup>	EA	ST36, LI4	Type 2 diabetes (symptoms of gastroparesis)	Single-blind, randomized pilot study, 19 patients, EA (9)/sham EA (10)	Reduction of the dyspeptic symptoms of diabetic gastroparesis
Sallam et al <sup>89</sup>	TEAS	Gastrointestinal (GI) acupoints	SSc	17 patients, TEAS/baseline	Treatment of upper GI symptoms
Xu et al <sup>95</sup>	EA	ST36, PC6	Functional dyspepsia	19 patients, acute-EA (10)/short-term (2w) EA (9)	Relief of dyspeptic symptoms
Zou et al <sup>92</sup>	TEAS	PC6	Healthy	Random, 26 volunteers, TEAS/sham	Inhibition of frequency of transient lower esophageal sphincter relaxations
Xing et al <sup>90</sup>	TEAS	ST36, PC6	Diarrhea-predominant IBS	7 patients, TEAS/sham-TEAS/control	Reduction of rectal sensitivity in IBS patients
Chang et al <sup>91</sup>	TEAS	ST36	Healthy (males)	15 volunteers (males) EA/TEAS	Enhancement of gastric myoelectrical regularity, bradygastria not significant

AP, acupuncture point; EA, electroacupuncture; GI, gastrointestinal; IBS, irritable bowel syndrome; SSc, scleroderma; TEAS, transcutaneous electrical acupoint stimulation.

### 3.7. Treatment of depression, addiction, and stroke

Two studies investigating the treatment of depression using LS,<sup>120,121</sup> five studies evaluating the treatment of various addictions (i.e., alcoholism and addictions to tobacco and narcotics) using ES<sup>122–124</sup> and LS,<sup>125,126</sup> and four studies examining the treatment of stroke using ES<sup>127–130</sup> are shown in Tables 7–9, respectively. LS was used by a research group to treat depression<sup>120,121</sup>, whereas two studies used ES devices<sup>123,124</sup>, one used LS<sup>125</sup> to treat tobacco dependence, one used an ES device in the treatment of drug abuse,<sup>122</sup> and one used LS to treat alcoholism.<sup>126</sup> Five studies showed that the use of ES and that of LS for treating various addictions were appropriate treatment adjuncts. ES was applied for treating stroke in four studies. All the studies in which stroke was treated, including treatment with a combination therapy consisting of TEAS and task-related training, reported treatment efficacy of TEAS or EA based on clinical trials involving 421 individuals. These results showed that ES is feasible for treating stroke. All the studies in these three categories claimed beneficial effects on the treatment of depression, various addictions, or stroke.

### 3.8. Physiological changes, diverse diseases, miscellaneous characteristics, and brain activities

All the papers regarding ASDs that induced physiological changes, treated various diseases, affected miscellaneous

characteristics, and induced brain activities are shown in Tables 10–13,<sup>131–174</sup> respectively. Most studies in these categories were focused on phenomenological observations or consisted of a small number of clinical trials. Many more case studies are required to demonstrate the effects of ASDs on diverse diseases. These various investigations may expand the application of modern ASDs. Due to the limited scope of this review, we did not further investigate the diverse aspects of these studies.

## 4. Discussion

EAs, which are invasive types of ES, were the first and most intensively studied modern applications of ASDs. Recently, the number of publications regarding the clinical effectiveness of noninvasive stimulations, such as TEAS, LS, MS, and US, has been increasing (Fig. 3). The increase is more substantial for noninvasive acupuncture-like techniques, most likely due to the growing demands for painless acupuncture or acupoint stimulations. Among the 195 articles analyzed, the studies involving ES (EAs and TEASs) predominated (133 articles, 68%), followed by LS studies (44 articles, 23%). Studies involving MS (16 articles, 8%) or US (2 articles, 1%) were less common. The publication of ES studies has steadily increased since the early 2000s, whereas LS and MS showed similar increment patterns with delayed start-up points (i.e., the increases began in 2009 and 2011, respectively). Despite its long history, ES had a steady

**Table 4 – Summary of studies on the effects of the four ASDs on nausea and vomiting**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Xu et al <sup>96</sup>	TEAS	PC6	PONV	Prospective, blind, & randomized study, 119 patients, TEAS/sham TEAS	Prevention of PONV after infratentorial craniotomy
Wang et al <sup>97</sup>	TEAS	Right PC6	Supratentorial craniotomy	Random TEAS (40)/control-nonacupoint (40), all standard general anesthesia	Prevalence of nausea, vomiting
Larson et al <sup>98</sup>	TEAS	Acupuncture points	Patients undergoing cosmetic surgery	Prospective, randomized, blinded, clinical trial, 122 patients, random standard pharm./pharm. + EA	Postoperative nausea & vomiting
Liu et al <sup>99</sup>	TEAS	Left-side PC6	Patients undergoing laparoscopic cholecystectomy	96 patients, random EA/placebo-EA no stim	Reduction of nausea & vomiting, pain relief
Habib et al <sup>102</sup>	TEAS	PC6/dorsum of wrist	Cesarean delivery with spinal anesthesia	Random, 91 patients, TEAS (47)/sham-APs TEAS (44)	No difference between the 2 groups (less PONV in 2 groups)
Kabalak et al <sup>100</sup>	TEAS	PC6, CV13	Tonsillectomy under general anesthesia	Randomized, controlled, prospective study, 90 children, TEAS (30)/pharm. dose (30)/no treatment (30)	Prophylaxis of postoperative retching & vomiting in pediatric tonsillectomy
Kramer et al <sup>101</sup>	TEAS	PC6	Patients receiving electroconvulsive therapy	11 patients, TEAS (9 good, 1 mixed, 1 no response)	Treating nausea & vomiting
Rusy et al <sup>105</sup>	EA	PC6	Tonsillectomy	120 patients, random EA (40)/sham-EA sham needle(40)/control no needle (40)	PONV prevention
Zárate et al <sup>103</sup>	TEAS	PC6	Laparoscopic cholecystectomy with standardized general anesthetic technique	Sham-controlled, double-blinded study, random, 221 outpatients, TEAS/placebo no stim	TEAS reduced postoperative nausea, but not vomiting
Shen et al <sup>106</sup>	EA	Antiemetic acupoints	High-risk breast cancer patients undergoing highly emetogenic chemotherapy regimen	3-arm, parallel-group, randomized controlled trial, LF-EA (37)/mock-EA (33)/no-EA (34)	Effective in controlling emesis (EA > pharm.)
Schlager et al <sup>107</sup>	LS	Bi PC6	Postoperative vomiting in children undergoing strabismus surgery	Double-blind, randomized, controlled study, 40 children, LS (20)/placebo (20)	Incidence of vomiting significantly lower
McMillan & Dundee <sup>104</sup>	TEAS	PC6	Cancer chemotherapy	-	Antiemetic action, useful adjunct to both the older antiemetics & the new antagonist ondansetron

ASD, acupuncture-like stimulation device; EA, electroacupuncture; LF, low frequency; PONV, postoperative nausea and vomiting; pharm, Pharmacological; stim, stimulation; TEAS, transcutaneous electrical acupoint stimulation.

but limited publication rate prior to 2000, whereas during the 1980s and 1990s, the number of publications on ES remained between zero article and two articles per year.

Fig. 4 shows the yearly publications of invasive (EAs) and noninvasive (TEASs) ES techniques. The total number of studies was similar between EAs (63 articles) and TEASs (70 articles). However, differences were observed in the number of publications per year; the publications associated with TEASs

showed a steady increase over time, which is in contrast to the stable annual publication pattern of EAs. Notably, the number of TEAS publications surpassed that of EAs in 2010. Specifically, TEASs were studied more than EAs over the past 5 years in the context of diseases with high therapeutic benefits, such as analgesic effect, pain relief, improvement of the alimentary system, and prevention of nausea and vomiting. The rising popularity of TEASs is presumably due to the increasing needs

**Table 5 – Summary of studies on the recovery of muscle fatigue or improvement of muscle strength with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Kim et al <sup>108</sup>	MS	LR9	Healthy (males)	20 participants (males), MS (10)/no MS (10)	Recovery of muscle fatigue
Kim et al <sup>109</sup>	TEAS/MS	An acupoint	Muscle fatigue	TEAS/MS/no stim	Therapeutic effect on muscle fatigue (MS better)
Zhou et al <sup>110</sup>	EA	ST36, ST39	Healthy (males)	randomized controlled trial, 43 young men, control/MA/EA-APs/EA-non-APs	Improvement of muscle strength in both limbs
Ngai et al <sup>111</sup>	TEAS	Bi LU7, EX-B1	Healthy (males)	11 individuals (males), TEAS/placebo-TEAS no stim	Higher postexercise FEV1, prolongation of submaximal exercise
Huang et al <sup>112</sup>	EA	ST36, ST39	Healthy (males)	30 men, random EA/control	Improvement of muscle strength of both limbs
Chiu et al <sup>113</sup>	TEAS + LS/exercise + LS/LS	Acupoints	Chronic neck pain	Randomized clinical trial, 218 patients, TEAS + IR/Exercise + IR (LS)/IR (LS)	Improvement in disability, isometric neck muscle strength, pain (TEAS, exercise)
Milne et al <sup>114</sup>	TEAS/EA	LI4, LI11	Healthy	TEAS/EA	Relief of muscle spasm & musculoskeletal pain, & restoration of mobility (TEAS)

AP, acupuncture point; ASD, acupuncture-like stimulation device; EA, electroacupuncture; FEV1, forced expiratory volume in 1 second; IR, infrared; stim, stimulation; LS, laser stimulation; MA, manual acupuncture; MS, magnetic stimulation; TEAS, transcutaneous electrical acupoint stimulation.

**Table 6 – Summary of studies on the reduction in body weight with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Schukro et al <sup>115</sup>	AEA	18, 87, 91 at ear	Obese females	Prospective, randomized, double-blinded study, 56 patients (females), AEA (28)/placebo dummy (28)	Reduction of body weight & BMI
Chien et al <sup>116</sup>	TEAS	ST36	Postmenopausal obese women	Prospective study, 49 women, random TEAS (24)/control no-TEAS (25)	Reduction in percentage body fat
Rerksuppaphol & Rerksuppaphol <sup>117</sup>	TEAS/EA	10 acupoints	Obese women	Prospective randomized open-label study, 45 women, TEAS/EA	Effective method for weight reduction as seen with EA
Lin et al <sup>118</sup>	EA	ST36, SP6	Postmenopausal women with obesity	Randomized controlled trial, 41 women, EA (20)/control (21)	Improvement of body composition
Jeong & Lee <sup>119</sup>	EA	Acupoints	Factual panniculitis	2 cases (females), EA	Weight reduction

AEA, auricular electroacupuncture; ASD, acupuncture-like stimulation device; EA, electroacupuncture; TEAS, transcutaneous electrical acupoint stimulation.

**Table 7 – Summary of studies on the treatment of depression with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Quah-Smith et al <sup>120</sup>	LS	LR14, LR8, CV14, HT7	Depressed participants	Random block on-off design, 10 nondepressed participants, 10 depressed participants	Antidepressant effect
Quah-Smith et al <sup>121</sup>	LS	LR14, CV14, LR8, HT7, KI3	major depressive disorder	Randomized, double blinded, placebo controlled trial, 47 participants, LA/placebo LA	Reduction of symptoms of depression

ASD, acupuncture-like stimulation device; LS, laser stimulation.

**Table 8 – Summary of studies on the treatment of smoking and addiction of drug and alcohol with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Penetar et al <sup>122</sup>	TEAS	PC6, TH5; LI4, PC8	Cocaine dependent or cannabis dependent	Single-blind, sham-controlled, crossover design, 20 volunteers (11 m, 9 f) TEAS/sham-TEAS/baseline ST	Modulation of mood & anxiety, no significant reduction in drug use or drug cravings
Lambert et al <sup>123</sup>	TEAS	LI4, PC8, PC6, TE5	Smoking	2 double-blind studies, 98 smokers, random TEAS-10 mA (20)/TEAS-5 mA (20)/placebo TEAS-0 mA (16)/TEAS-5 mA (19); intermittent 5 mA	Antagonizing the urge to smoke in dependent smokers
Kerr et al <sup>125</sup>	LS	4 acupoints	Smoking	Double-blind, randomized controlled trial, 387 volunteers, 3-LS 1-sham LS/4-LS/4-sham LS no stim	Assisting in smoking cessation by reducing the physical symptoms of withdrawal
Zalewska-Kaszubska & Obzejta <sup>126</sup>	ALS	Neck; 10 auricular acupoints	Alcoholics	53 patients, He-Ne LS (neck) + argon ALS	Adjunct treatment for alcoholism
Georgiou et al <sup>124</sup>	TEAS	SJ18, SJ17	Smoking cessation	Randomized controlled trial, 216 smokers, TEAS/control TEAS no stim	Insufficient power to detect real but small differences between treatment conditions

ALS, auricular laser stimulation; ASD, acupuncture-like stimulation device; f, female; LS, laser stimulation; m, male; stim, stimulation; ST, standard treatment; TEAS, transcutaneous electrical acupoint stimulation.

**Table 9 – Summary of studies on the treatment of stroke with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Ng & Hui-Chan <sup>127</sup>	TEAS + TRT	ST36, LV3, GB34, UB60	Poststroke	Case study, 1 man (age 61 y), TEAS + TRT	Decreased impairment & improved function in an individual with long-term chronic stroke
Gong et al <sup>128</sup>	EA	ST36	First-time cerebral infarction or hemorrhage, or a stroke history	Randomized, controlled, clinical study, 240 patients, EA (124)/control (116)	Effects on lower extremity motor function in stroke patients
Kim et al <sup>129</sup>	TEAS	Acupoints	Ischemic stroke with motor dysfunction	62 patients, 2 Hz-TEAS/120 Hz-TEAS	Helpful for motor recovery after ischemic stroke (LF-TEAS)
Wong et al <sup>130</sup>	TEAS	Acupoints	Patients with hemiplegia in stroke	Randomized, 118 patients, comprehensive rehabilitation + TEAS (59)/comprehensive rehabilitation (59)	Convenient & effective therapy for stroke

ASD, acupuncture-like stimulation device; EA, electroacupuncture; LF, low frequency; TEAS, transcutaneous electrical acupoint stimulation; TRT, task-related training.

for safety without needling, low infection risk, and relatively expedient utilization of clinical trials. The recent increase in studies of LS and MS, which are noninvasive, may be understood based on the same rationale.

According to a recent analysis, approximately 41% of clinical studies in acupuncture research from 1991 to 2009 addressed pain and analgesia.<sup>6</sup> Among the studies evaluating the four types of ASDs published through 2014, the percentage

of clinical studies addressing pain and analgesia was 33%. This reduction in the percentage of studies focused on pain and analgesia is directly related to the recently heightened interest in acupuncture research on brain activities. The percentage of publications focused on brain activities that have been published since 2010 constitutes 61% (19 articles) of all such publications since 2001 (31 articles). Excluding the emerging category of brain activity, approximately 38%

**Table 10 – Summary of studies on physiological changes with the four ASDs**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Cafaro et al <sup>131</sup>	LS	Bi LI2, ST5, ST6, ST7, SI19, BL 13	Sjögren's syndrome	26 female, patients, LA/sham	Salivary flow rate improvement
Kim et al <sup>175</sup>	MS	LI4	Healthy	–	Improvement of peripheral vascular system circulation
Li et al <sup>176</sup>	EA	LI4, TE5, BL63, LR3, ST36, BL40, BL10, BL20, BL2, EX-HN4	Supratentorial craniotomy	29 patients, control (10)/EA (9)/sham EA (10)	Prevention of decrease of immunoglobulin after surgery, no significant difference between EA & sham EA
Litscher et al <sup>132</sup>	LS	GV20, PC6	Healthy	Randomized crossover study, 11 volunteers (3 m, 8 f), MA (GV20;PC6)/red LA (GV20;PC6)/violet LA (GV20;PC6)	HR & HRV control
Tsuruoka et al <sup>177</sup>	US	LR3	Healthy	50 volunteers (40 m, 10 f), random US/MA	Increase of blood flow volume
Wang et al <sup>133</sup>	LS	Right LI4	Healthy	28 volunteers, random LA-LI4/LA-non LI4	Increase of left LI4 MBF, 40 min later after stimulation ceased, the MBF still increasing significantly
Raith et al <sup>134</sup>	LS	LI4	Premature neonates	10 neonates (7 m, 3 f), initial temp/5 min stim temp/10 min stim temp	Increase in the skin temperature
Lee et al <sup>178</sup>	MS	PC9	Healthy	1 individual	Parasympathetic activity of the autonomic nervous system
Jia et al <sup>179</sup>	EA	Bi ST36, ST37	Healthy	20 volunteers, EA/sham EA	Effect on autonomic function
Jones et al <sup>180</sup>	TEAS	Bilateral PC6	Healthy	16 volunteers, random TEAS/sham-TEAS non-APs/no TEAS no-stim	Change in artery
Lee et al <sup>181</sup>	EA	LI4, LI11	Healthy	Randomized crossover design, 14 participants, HF-EA 120 Hz/LF-EA 2 Hz	Increase in autonomic nervous activity (HF-EA), enhancing sympathovagal balance (both)
Chang et al <sup>182</sup>	EA	ST36, LI10	Healthy	15 volunteers, LF EA (low freq. 2 Hz)/HF EA (high freq. 100 Hz)	Not affecting cardiovagal activity in normal volunteers
Cunha et al <sup>140</sup>	LS/MA	10 acupoints	Circulatory deficiency	40 individuals, LS (20)/MA (20)	Significant increase in systolic pressure of lower limbs, consequent improvement of the revascularization index
Litscher et al <sup>135</sup>	LS	PC6	Healthy	Randomized, controlled study, 13 volunteers, LS/control-laser off	Decrease of HR
Kim et al <sup>183</sup> Lu et al <sup>184</sup>	EA MA, EA, TENS	PC5, PC6 Bi ST36, ST37, palm, dorsum	Healthy Healthy	EA (10)/sham-EA no stim (10) 20 volunteers, random sham-MA/MA/EA/TENS; before-A, during-A, after-A (time sequence)	EEG, ECG, HR change Cutaneous blood flow & temperature change
Zhang et al <sup>185</sup>	TEAS	LI4, LI11	Normal & elevated blood pressure	Randomly, 27 individuals, TEAS (13, 8 m, 5 f)/control (14, 9 m, 5 f)	Reduction of systolic blood pressure, but not diastolic blood pressure
Zhang et al <sup>136</sup>	LS	LI4, LI11	Healthy	Randomized controlled pilot study, 45 students + faculty, LA/sham-LA laser off	Reduction of blood pressure
Cakmak et al <sup>186</sup>	EA	ST29, ST25	Healthy (m)	Prospective, randomized study, 80 volunteers, MA/2 Hz-EA/10 Hz-EA	Increase in testicular blood flow, helpful in clinical treatment of infertile men (ST29, 10 Hz)



Table 10 (Continued)

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Arai et al <sup>187</sup>	TEAS	Bi PC5, PC6/shoulder	Parturients undergoing cesarean section under spinal anesthesia	Random, 36 singleton parturients, TEAS (12)/sham-APs TEAS (12)/no treatment (12)	Reduction of the severity & incidence of hypotension after spinal anesthesia in parturients
Cheung & Jones <sup>188</sup>	TEAS	Bilateral PC6	Healthy (m)	Single-blinded, randomized controlled trial, 28 individuals, treadmill, TEAS/pre-TEAS/placebo-TEAS	HR recovery after exercise
Banzer et al <sup>137</sup>	LS	Right forearm PC6	Healthy (nonsmoking males)	Randomized, double-blinded, placebo-controlled trial, 33 healthy (m), LA (18)/control no laser (15)	Improvement of blood flow
Szeles & Litscher <sup>189</sup>	AEA	Ear acupuncture	Healthy (f)	2 healthy (f), AEA	Modulation of blood flow
Litscher <sup>138</sup>	LS	Acupuncture points	Healthy	Randomized crossover study, 22 volunteers, LS	Changes in peripheral microcirculation & surface temperature of skin
Li et al <sup>190</sup>	MS (magnitopuncture)	GV14, PC6	Healthy (m)	Randomly, 40 individuals, MS/control MS non-APs	Modulating effect on sympathetic & parasympathetic nerve activities
Hsieh et al <sup>191</sup>	EA	ST36	Healthy	8 volunteers, before/during /after EA	Physiological mechanisms responsible
Litscher & Schikora <sup>139</sup>	LS	Vision-related acupuncture points	Healthy	Randomized crossover trial, 27 volunteers (13 m, 14 f), LA/MA	Increases of blood flow in ophthalmic artery
Cramp et al <sup>192</sup>	TENS/TEAS	Median nerve/LI4	Healthy	Randomly, 30 individuals (15 m, 15 f), control (10)/TENS (10)/TEAS (10)	Increase in cutaneous blood flow in the TENS median nerve
Litscher et al <sup>141</sup>	LS	Vision-related acupuncture points	Healthy	15 volunteers (10 m, 5 f), LS/MA	Increases in blood flow velocity in posterior cerebral artery
Balogun et al <sup>193</sup>	TEAS (HVG)	ST36, ST37	Healthy	11 individuals (5 m, 6 f), 2 Hz-TEAS/120 Hz-TEAS	No increase in peripheral hemodynamic functions in asymptomatic individuals
Williams et al <sup>194</sup>	TEAS	LR3, ST36, LI11	Diastolic hypertension	Random, 10 individuals, TEAS/sham-TEAS non-APs.	Reduction of diastolic blood pressure for TEAS
Dunn et al., <sup>195</sup>	TEAS	SP6, LR3	Pregnant women	Randomly, TEAS/control no stim	Increase in frequency & strength of uterine contractions

AEA, auricular electroacupuncture; AP, acupuncture point; ASD, acupuncture-like stimulation device; EA, electroacupuncture; ECG, electrocardiogram; EEG, electroencephalogram; f, female; HF, high frequency; HR, heart rate; HRV, heart rate variability; HVG, high voltage galvanic; LA, laser acupuncture; LF, low frequency; LS, laser stimulation; m, male; MA, manual acupuncture; MBF, meridian blood flow; MS, magnetic stimulation; stim, stimulation; TEAS, transcutaneous electrical acupoint stimulation; TENS, transcutaneous electrical nerve stimulation; US, ultrasound stimulation.

of the studies were focused on pain and analgesia, which is similar to the percentage of MA studies focused on pain and analgesia.

The effectiveness analysis showed that the effectiveness of ES with respect to the analgesic effect, pain relief, and reduction of nausea and vomiting was confirmed by clinical trials involving > 1000 individuals and many RCTs. Based on clinical trials involving > 100 individuals, ES was effective in improving the alimentary system, improving muscle strength, reducing body weight, and treating stroke. Likewise, LS was shown to be useful for providing pain relief and in treating various addictions. Interestingly, the addiction treatment effect was confirmed by LS studies but not by ES studies.

#### 4.1. Limitations

Our review is based on the four most influential databases, specifically Medline, PubMed, Cochrane Library, and Web of Science; moreover, we primarily analyzed Science Citation Index (SCI) or Science Citation Index Expanded (SCI-E) journal articles. The advantage of this approach is the inclusion of quality-guaranteed articles only. Laboratory experiments on animals, MA-only clinical trials, non-English-language articles, and review articles were excluded from the analysis. The details regarding device specifications or interventional designs, including stimulation strength, duration and interval, and patient and environmental conditions, were not analyzed due to space limitations.

**Table 11 – Summary of studies on the treatment of various diseases with the four ASDs**


Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Sun et al <sup>142</sup>	EA	Bilateral PC6	OI	Randomized, controlled, crossover design, EA (20)/no EA (10)	Treatment in attenuating OI
Zhang et al <sup>196</sup>	TEAS	LI4, PC6, ST36, SP6	Autistic children receiving rehabilitation training	76 children, TEAS (37)/no treatment (39)	Effective for treatment of autistic children with passive & aloof social interaction style
Yang et al <sup>197</sup>	TEAS	LI4, SJ5, ST36, BL63, LR3, GB40	Supratentorial craniotomy	Randomized controlled trial, EA/sham-EA	Significantly shortened speed of postoperative recovery
Sahmeddini et al <sup>198</sup>	EA	PC6, PC5	End-stage liver disease patients undergoing orthotropic deceased donor liver transplantation	Randomized, 40 patients, norepinephrine-vasoconstrictor/EA	Reduction of severity & incidence of hypotension during anesthesia for liver transplantation
Ng et al <sup>199</sup>	TEAS	Bi PC6	Open heart surgery	40 patients, random TEAS (20)/placebo-TEAS no stim (20)	Earlier return to preoperative BP, HR, & RPP values
Wang et al <sup>200</sup>	MA/EA	Bi GB8, TE17, GB2, GB20, GV20, TE3, ST36 (MA)/bi GB8, TE17 (EA)	Tinnitus	Randomized, single-blinded, placebo-controlled design, 50 patients (46 m, 4 f), MA/EA/placebo	Short-term general effects on tinnitus (EA)
O'Brien et al <sup>201</sup>	LS	10 acupoints	Active symptoms of menopause	Double-blind, randomized, placebo-controlled study, 40 women, LS/placebo LS (off)	Treatment of menopausal symptoms (no more efficacious than MA)
Ngai et al <sup>202</sup>	TEAS	Bi EX-B1, LU7	Patients with asthma	Randomized controlled trial, 30 individuals, random TEAS/TEAS + ST/sham-TEAS + ST	Reduction in the decline of forced expiratory volume in 1s FEV <sub>1</sub> (1) following exercise training
Burduli & Ranyuk <sup>203</sup>	LS + ST	Acupuncture points	Chronic noncalculous cholecystitis	73 patients, ST (35)/LA + ST (38)	Cholecystitis treatment
Su et al <sup>204</sup>	LS	Acupoints	Renal failure patients receiving regular hemodialysis	Randomized controlled trial, before/after LS	Decrease in both stress & fatigue levels
Lau & Jones <sup>205</sup>	TEAS	Bi Ex-B1	Chronic obstructive pulmonary disease	Randomized, placebo-controlled trial, 46 patients, TEAS/placebo-TEAS no stim	Management of dyspnea
Hsu et al <sup>206</sup>	EA	BL15	Healthy	10 volunteers, sham-EA/2 Hz-EA	Relaxation, calmness, & reduced feeling of tension or distress
Bray et al <sup>207</sup>	EA	Uni PC6, HT3, LR3/bi GB34, LI11, SI3	Healthy	80 individuals, EA-PC6, HT3, LR3/ EA-GB34, LI11, SI3/no stim; 5/60/100 Hz; uni/bilateral	Adjunct therapy for disorders of hypervigilance (to decrease arousal levels)
Litscher et al <sup>208</sup>	LS	ST7, TE22	Intensive care patient after severe head injury	34 volunteers (10 m, 24 f), 1 patient (head injury), acupressure/MA/LA	Reproducible functional changes in the brain
O'Reilly et al <sup>209</sup>	LS	SP6	Interstitial cystitis	Double-blind trial, random LS (29)/placebo (27)	Treatment & control cohorts experiencing similar improvements, no difference between active & sham
Li et al <sup>210</sup>	MS	GV14, PC6	Healthy	Randomly, 40 individuals, MS/control MS non-APs	Effects of driving fatigue

AP, acupuncture point; ASD, acupuncture-like stimulation device; BP, blood pressure; EA, electroacupuncture; f, female; FEV<sub>1</sub>, forced expiratory volume in 1 second; HR, heart rate; LA, laser acupuncture; LS, laser stimulation; m, male; MA, manual acupuncture; MS, magnetic stimulation; OI, orthostatic intolerance; RPP, rate pressure product; ST, standard treatment; stim, stimulation; TEAS, transcutaneous electrical acupoint stimulation.

**Table 12 – Clinical studies showing miscellaneous characteristics**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Chen et al <sup>143</sup>	LS	LU9, PC7, HT7, SI4, SJ4, LI5, SP3, LR3, KI4, BL65, GB40, ST42	Healthy	76 candidates	Complementary & interaction for current flow of meridians
Gopalan et al <sup>211</sup>	EA	Acupuncture points	Implanted with cardiac device	–	Safety in patients with a total artificial heart
Irnich et al <sup>212</sup>	LS/Seirin (sham-LS)	LI4, LU7, LR3	Healthy	Randomized, double-blinded, crossover design, 34 volunteers, LS (18)/sham-LS (16)	Valid placebo control in laser acupuncture studies (Seirin)
Litscher & Wang <sup>213</sup>	MA/LS	LU6	Healthy	1 person, MA/LA	Changes of electrical skin impedance
Thompson & Cummings <sup>214</sup>	EA	Acupuncture points in a limb	Healthy	–	No detectable currents in the chest (safety)
Leung et al <sup>215</sup>	TEAS/MA/EA	LI4	Healthy	15 individuals, TEAS/MA/EA	Difference in electrical conductance between APs & non-APs
Litscher et al <sup>216</sup>	LS	Acupuncture points	Healthy	29 volunteers (9 m, 20 f), LA/placebo-LA; before/after	Change in the median value of cold pain, no significant changes in parameters of thermal sensory & pain thresholds
Chang et al., <sup>217</sup>	EA/TEAS	Left LI4	Healthy	13 volunteers, 2 Hz-EA/2 Hz-TEAS/100 Hz-TEAS	Changes of cutaneous reflex

AP, acupuncture point; EA, electroacupuncture; f, female; LA, laser acupuncture; LS, laser stimulation; m, male; MA, manual acupuncture; TEAS, transcutaneous electrical acupoint stimulation.




**Fig. 3 – The number of published articles on the four ASDs per year.**

ASD, acupuncture-like stimulation device; ES, electrical stimulation; LS, laser stimulation; MS, magnetic stimulation; US, ultrasound stimulation.

**Table 13 – Summary of studies on the effect of the four ASDs on brain activity**

Reference	Stimulation type	Stimulation site	Symptom	Control	Effect
Guo et al., <sup>144</sup>	MS	PC6	Healthy	6 right-handed volunteers (3 m, 3 f)	Brain activity
Zhang et al <sup>145</sup>	MS	Left GB37	Healthy	GB37-MS/mock point-MS	Brain activity
Raith et al <sup>146</sup>	LS	Bi LI4	Term & preterm neonates	20 neonates (12 m, 8 f), LA period/postintervention period	Brain activity
Quah-Smith et al <sup>147</sup>	LS	LR8	Healthy	16 participants, random on-off block design, LA/MA	Brain activity
Zhang et al <sup>148</sup>	TEAS	LI4, PC8	Healthy	18 individuals (9 m, 9 f), all individuals TEAS	Brain activity
Yin et al <sup>149</sup>	MS	PC6	Healthy	—	Brain activity
Lee et al <sup>150</sup>	MS	PC9	Healthy	—	HRV & brain activity
Litscher <sup>151</sup>	LS	PC6	Healthy	40 volunteers, LA/MA	Brain activity
Wu et al <sup>152</sup>	LS	Palm	Healthy	single-blind randomized trial, 40 individuals (m), random LS (20)/sham LS (20)	Brain activity
Litscher et al <sup>153</sup>	LS	Bi PC6	Healthy (f)	1 volunteer (f), LA	Brain activity
Yu et al <sup>154</sup>	MS	PC6	Healthy	MS-PC6/ MS-mock point	Brain activity
Jiang et al., <sup>155</sup>	TEAS	LI4, PC8	Healthy	40 individuals, TEAS (40)	Brain activity
Hsieh et al <sup>156</sup>	LS	KI1	Healthy right handed	36 right-handed volunteers, random MW LA (12; 8 m, 4 f)/CW LA (12; 9 m, 4 f)/placebo LA(12)	Brain activity
Yu et al <sup>157</sup>	MS	PC6	Healthy	before MS/during MS/after MS	Brain activity
Kim et al <sup>158</sup>	MS	PC9	Healthy	—	Vascular & brain activity
Jo & Jo <sup>159</sup>	MS	HT4, HT6	Healthy	23 young adults (aged 19–22 y)	Brain activity (pole direction)
Zyloney et al <sup>160</sup>	EA	LI3, LI4 right hand	Healthy, right handed	48 individuals, random EA/sham EA	Brain activity
Quah-Smith et al <sup>161</sup>	LS	LR14, CV14, LR8, HT7	Healthy	10 individuals, random LA/LA-sham point	Brain activity
Xu et al <sup>162</sup>	MS	ST36, LI4	Healthy	MS/MS-mock point	Brain activity
Na et al <sup>163</sup>	EA	GB34	Healthy	12 individuals, EA/EA-sham points,	Brain activity
Xu et al <sup>164</sup>	MS	ST36	Healthy	Pre-MS/post-MS (0.5 Hz/1 Hz/3 Hz)	Brain activity
An et al <sup>165</sup>	EA	LI4, LI11	Healthy	Brain SPECT EA (20)/PET EA (13); before/during /after EA	Brain activity
Wang et al <sup>166</sup>	EA	Right LI4	Healthy	EA (9)/sham-point EA (5)	Brain activity
Zeng et al <sup>167</sup>	EA	LI4	Healthy (right handed)	EA	Brain activity
Litscher et al <sup>168</sup>	LS	Acupoints	Healthy	Randomized controlled crossover trial, 18 volunteers (7 m, 11 f), before/during-LA/after	Modulation of blood flow, brain activity
Zhang et al <sup>169</sup>	EA	Left leg ST36, SP6	Healthy (right handed)	48 individuals, 2 Hz-EA/100 Hz-EA	Analgesia effect/brain activity
Li et al <sup>170</sup>	EA	TE8, GV15	Healthy (Chinese males)	17 volunteers (m), EA-TE8 (11)/EA-GV15 (6)	Brain activity, typical language areas in the left inferior frontal cortex not activated
Kong et al <sup>171</sup>	EA	Left hand LI4	Healthy (right handed)	11 volunteers (6 m, 5 f), EA/MA	Brain activity
Siedentopf et al <sup>172</sup>	LS	Left foot BL67	Healthy (m)	10 volunteers (m), LA/dummy LA	Brain activity
Wu et al <sup>173</sup>	EA	GB34	Healthy	45 volunteers, EA (15)/mock-EA no stim (7)/minimal-EA superficial & light stim (8)/sham-EA non-Aps (15)	Modulation of hypothalamus limbic system
Chang et al <sup>174</sup>	MA/TEAS	LI4	Healthy	Randomly, control TEAS no stim/MA/2 Hz-TEAS/100 Hz-TEAS	Increases in amplitude of H-reflex (TEAS), 100 Hz TEAS has greater effect

AP, acupuncture point; ASD, acupuncture-like stimulation device; EA, electroacupuncture; f, female; HRV, heart rate variability; LA, laser acupuncture; LS, laser stimulation; m, male; MA, manual acupuncture; MS, magnetic stimulation; PET, positron emission tomography; SPECT, single-photon emission computed tomography; stim, stimulation; TEAS, transcutaneous electrical acupoint stimulation.



**Fig. 4** – The number of articles on ES methods with years, where EAs include the invasive techniques of EA, AEA, and EHA, and TEASs include the noninvasive techniques of TEAS and TENS.

**AEA, auricular electroacupuncture; EA, electroacupuncture; EHA, electrical heat acupuncture; ES, electrical stimulation; TEAS, transcutaneous electrical acupoint stimulation; TENS, transcutaneous electrical nerve stimulation.**

## 5. Conclusions

In the past decade, modern ASDs have been studied extensively for their clinical effectiveness and to test equivalence or noninferiority with traditional MA. Among the modern ASDs, ES was found to be most widely studied, and its popularity was sequentially followed by LS, MS, and US. Specifically, EAs, which are invasive types of ES, were the first and most intensively studied types of ASDs, whereas TEASs, which are noninvasive types of ES, have surpassed EAs in publication number since 2010. Very recently, noninvasive techniques, such as TEASs, LS, MS, and US have gained research attention, as evidenced by increasing annual publications.

The most extensively studied treatment effects were for analgesia and pain relief, whereas rapid growth has occurred in the research field of the effects of treatments on brain activities. The overall quality of the study designs was moderate, as 58% of the studies were based on RCTs and 96% of the RCT-based outcomes reported therapeutic benefits. ES was effective in providing an analgesic effect, pain relief, and a reduction of nausea and vomiting, based on clinical trials involving > 1000 individuals. Based on > 100 clinical trials, ES was shown to be effective in improving the alimentary system, improving muscle strength, reducing body weight, and treating stroke. LS was effective in pain relief and for treating various addictions. We anticipate more studies on the therapeutic effects of ASDs, particularly concerning noninvasive methods, to meet the growing needs of guaranteed safety, decreased risk of infection, decreased pain, and improved convenience.

## Conflicts of interest

No conflicts of interest are declared.

## Acknowledgments

This work was supported by a grant (K15012) from the Korea Institute of Oriental Medicine, Daejeon, Korea, funded by the Korean government.

## REFERENCES

1. World Health Organization. *Acupuncture: review and analysis of reports on controlled clinical trials*. Geneva, Switzerland: World Health Organization; 2002.
2. Choi SM, Park JE, Li SS, Jung H, Zi M, Kim TH, et al. A multicenter, randomized, controlled trial testing the effects of acupuncture on allergic rhinitis. *Allergy* 2013;68:365–74.
3. Paramore LC. Use of alternative therapies: estimates from the 1994 Robert Wood Johnson Foundation National Access to Care Survey. *J Pain Symptom Manage* 1997;13:83–9.
4. Helms JM. *Acupuncture energetics: a clinical approach for physicians*. Berkeley: Thieme Medical Acupuncture Publishers; 1995.
5. NIH Consensus Conference. Acupuncture. *JAMA* 1998;280:1518–24.
6. Han JS. Acupuncture analgesia: areas of consensus and controversy. *Pain* 2011;152(3 Suppl):S41–8.
7. Voll R. Twenty years of electroacupuncture diagnosis in Germany. A progress report. *Am J Acupunct* 1975;3:7–17.



8. Nakatani Y. Skin electric resistance and ryodoraku. *J Auton Nerv Syst* 1956;6:52.
9. Ahn AC, Colbert AP, Anderson BJ, Martinsen OG, Hammerschlag R, Cina S, et al. Electrical properties of acupuncture points and meridians: a systematic review. *Bioelectromagnetics* 2008;29:245–56.
10. Baxter GD, Bleakley C, McDonough S. Clinical effectiveness of laser acupuncture: a systematic review. *J Acupunct Meridian Stud* 2008;1:65–82.
11. Colbert AP, Cleaver J, Brown KA, Harling N, Hwang Y, Schiffke HC, et al. Magnets applied to acupuncture points as therapy—a literature review. *Acupunct Med* 2008;26:160–70.
12. Kaslow AL. A new technique for acupuncture point finding and stimulation 28. *Am J Acupunct* 1975;3:157–60.
13. Becker RO, Reichmanis M, Marino AA, Spadaro JA. Electrophysiological correlates of acupuncture points and meridians. *Psychoenergetic Syst* 1976;1:195–212.
14. Croley T. Electrical acupuncture point conductance in the living compared to that in the dead. *Am J Acupunct* 1986;14:57–60.
15. Niboyet J, Mery A. Experimentelle Studien über den Meridianverlauf. *Dt Ztschr f Akup* 2008;51:73–5 [In German].
16. Kramar F. Die biophysikalische Grundlagen der Akupunktur. *Deut Zeit fur Akupunktur* 1962;11:131–9 [In German, English abstract].
17. Hill S. Letter: acupuncture research in the USSR. *Am J Chin Med* 1976;4:204–5.
18. Baxter D. Laser acupuncture analgesia: an overview. *Acupunct Med* 1989;6:57–60.
19. Barker AT, Jalinous R, Freeston IL. Non-invasive magnetic stimulation of human motor cortex. *Lancet* 1985;325:1106–7.
20. Inoue K. *Magnetic treatment device*. Patent no. 5030196. USA: Inoue-Japax Research Incorporated; 1983.
21. Katayama I. Meridian magnetic analgesia under acupuncture stimulation. *Shikai Tenbo* 1982;59:1417–20.
22. Jin WC. A preliminary application and study on the ultrasonic acupuncture. *China Acupunct* 1984;4:1–3.
23. Jiang Y, Liu J, Liu J, Han J, Wang X, Cui C. Cerebral blood flow-based evidence for mechanisms of low-versus high-frequency transcutaneous electric acupoint stimulation analgesia: a perfusion fMRI study in humans. *Neuroscience* 2014;268:180–93.
24. Wang H, Xie Y, Zhang Q, Xu N, Zhong H, Dong H, et al. Transcutaneous electric acupoint stimulation reduces intra-operative remifentanyl consumption and alleviates postoperative side-effects in patients undergoing sinusotomy: a prospective, randomized, placebo-controlled trial. *Br J Anaesth* 2014;112:1075–82.
25. Zhang Q, Gao Z, Wang H, Ma L, Guo F, Zhong H, et al. The effect of pre-treatment with transcutaneous electrical acupoint stimulation on the quality of recovery after ambulatory breast surgery: a prospective, randomised controlled trial. *Anaesthesia* 2014;69:832–9.
26. Lan F, Ma YH, Xue JX, Wang TL, Ma DQ. Transcutaneous electrical nerve stimulation on acupoints reduces fentanyl requirement for postoperative pain relief after total hip arthroplasty in elderly patients. *Minerva Anestesiol* 2012;78:887–95.
27. Cheing GL, Chan WW. Influence of choice of electrical stimulation site on peripheral neurophysiological and hypoalgesic effects. *J Rehabil Med* 2009;41:412–7.
28. Attele AS, Mehendale S, Guan X, Dey L, Yuan C-S. Analgesic effects of different acupoint stimulation frequencies in humans. *Am J Chin Med* 2003;31:157–62.
29. Chesterton LS, Barlas P, Foster NE, Lundeborg T, Wright CC, Baxter GD. Sensory stimulation (TENS): effects of parameter manipulation on mechanical pain thresholds in healthy human subjects. *Pain* 2002;99:253–62.
30. Yuan CS, Attele AS, Dey L, Lynch JP, Guan XF. Transcutaneous electrical acupoint stimulation potentiates analgesic effect of morphine. *J Clin Pharmacol* 2002;42:899–903.
31. DeSantana JM, Santana-Filho VJ, Guerra DR, Sluka KA, Gurgel RQ, Silva WM. Hypoalgesic effect of the transcutaneous electrical nerve stimulation following inguinal herniorrhaphy: a randomized, controlled trial. *J Pain* 2008;9:623–9.
32. Greif R, Laciny S, Mokhtarani M, Doufas AG, Bakhshandeh M, Dorfer L, et al. Transcutaneous electrical stimulation of an auricular acupuncture point decreases anesthetic requirement. *Anesthesiology* 2002;96:306–12.
33. Chen L, Tang J, White PF, Sloninsky A, Wender RH, Naruse R, et al. The effect of location of transcutaneous electrical nerve stimulation on postoperative opioid analgesic requirement: acupoint versus nonacupoint stimulation. *Anesth Analg* 1998;87:1129–34.
34. Wang B, Tang J, White PF, Naruse R, Sloninsky A, Kariger R, et al. Effect of the intensity of transcutaneous acupoint electrical stimulation on the postoperative analgesic requirement. *Anesth Analg* 1997;85:406–13.
35. Wu J, Zhao Y, Yang C, Xue Q, Li N. Effects of electroacupuncture preemptive intervention on postoperative pain of mixed hemorrhoids. *Zhongguo Zhen Jiu (Chin Acupunct Moxibustion)* 2014;34:279–83.
36. Zheng X, Meng J-B, Fang Q. Electroacupuncture reduces the dose of midazolam monitored by the bispectral index in critically ill patients with mechanical ventilation: an exploratory study. *Acupunct Med* 2012;30:78–84.
37. Barlas P, Ting SLH, Chesterton LS, Jones PW, Sim J. Effects of intensity of electroacupuncture upon experimental pain in healthy human volunteers: a randomized, double-blind, placebo-controlled study. *Pain* 2006;122:81–9.
38. Zhang WT, Jin Z, Huang J, Zhang L, Zeng YW, Luo F, et al. Modulation of cold pain in human brain by electric acupoint stimulation: evidence from fMRI. *Neuroreport* 2003;14:1591–6.
39. Morioka N, Aka O, Doufas AG, Chernyak G, Sessler DI. Electro-acupuncture at the Zusanli, Yanglingquan, and Kunlun points does not reduce anesthetic requirement. *Anesth Analg* 2002;95:98–102.
40. Leung A, Khadivi B, Duann JR, Cho ZH, Yaksh T. The effect of Ting point (tendinomuscular meridians) electroacupuncture on thermal pain: A model for studying the neuronal mechanism of acupuncture analgesia. *J Altern Complement Med* 2005;11:653–61.
41. Lin JG, Lo MW, Wen YR, Hsieh CL, Tsai SK, Sun WZ. The effect of high and low frequency electroacupuncture in pain after lower abdominal surgery. *Pain* 2002;99:509–14.
42. Litscher G. Effects of acupressure, manual acupuncture and Laserneedle acupuncture on EEG bispectral index and spectral edge frequency in healthy volunteers. *Eur J Anaesth* 2004;21:13–9.
43. Brockhaus A, Elger CE. Hypoalgesic efficacy of acupuncture on experimental pain in man. Comparison of laser acupuncture and needle acupuncture. *Pain* 1990;43:181–5.
44. Mucuk S, Baser M. Effects of noninvasive electroacupuncture on labour pain and duration. *J Clin Nurs* 2014;23:1603–10.
45. Sun C, Hu C, Hao H, Niu C, Li L. Development of a uni-acupoint transcutaneous electric nerve stimulation device for electroacupuncture-like neuromodulation. In: *Conference proceedings: Annual International Conference of the IEEE Engineering in Medicine and Biology Society IEEE*

- Engineering in Medicine and Biology Society Conference*. 2013:6155–8.
46. Vassal F, Creac'h C, Convers P, Laurent B, Garcia-Larrea L, Peyron R. Modulation of laser-evoked potentials and pain perception by transcutaneous electrical nerve stimulation (TENS): a placebo-controlled study in healthy volunteers. *Clin Neurophysiol* 2013;124:1861–7.
  47. Mucuk S, Baser M, Ozkan T. Effects of noninvasive electroacupuncture on labor pain, adrenocorticotropic hormone, and cortisol. *Altern Ther Health Med* 2013;19:26–30.
  48. Ni X, Xie Y, Wang Q, Zhong H, Chen M, Wang F, et al. Cardioprotective effect of transcutaneous electric acupoint stimulation in the pediatric cardiac patients: a randomized controlled clinical trial. *Pediatr Anesth* 2012;22:805–11.
  49. Wu LL, Su CH, Liu CF. Effects of Noninvasive Electroacupuncture at Hegu (LI4) and Sanyinjiao (SP6) Acupoints on Dysmenorrhea: A Randomized Controlled Trial. *Journal of Alternative and Complementary Medicine* 2012;18:137–42.
  50. Yeh ML, Chung YC, Chen KM, Chen HH. Pain reduction of acupoint electrical stimulation for patients with spinal surgery: a placebo-controlled study. *Int J Nurs Stud* 2011;48:703–9.
  51. Montenegro EJN, Albuquerque NBd, Mariz LMRd, Costa RdCdS, Montarroyos CS, Motta MA. Ação da TENS acupuntural em acupontos na dor induzida pela hipotermia local (0-2° C). *Fisioter Mov* 2010;23:483–92 [In Portuguese, English abstract].
  52. Fang J, Zhang Y, Xuan L, Liu K, Chen L. Observation on clinical therapeutic effect of transcutaneous point electric stimulation on peri-arthritis of shoulder at different stages. *Zhongguo Zhen Jiu (Chin Acupunct Moxibustion)* 2006;26:11–4.
  53. Yeh ML, Chung YC, Chen KM, Tsou MY, Chen HH. Acupoint electrical stimulation reduces acute postoperative pain in surgical patients with patient-controlled analgesia: a randomized controlled study. *Altern Ther Health Med* 2010;16:10–8.
  54. Ye J, Zhu Z, Huang C, Wei J. Pain management using Han's acupoint nerve stimulator combined with patient-controlled analgesia following neurosurgery: a randomized case control study. *Neural Regen Res* 2008;3:809–12.
  55. Yip YB, Tse HM, Wu KK. An experimental study comparing the effects of combined transcutaneous acupoint electrical stimulation and electromagnetic millimeter waves for spinal pain in Hong Kong. *Complement Ther Clin Pract* 2007;13:4–14.
  56. Weng CS, Shu SH, Chen CC, Tsai YS, Hu WC, Chang YH. The evaluation of two modulated frequency modes of acupuncture-like tens on the treatment of tennis elbow pain. *Biomed Eng Appl Basis Commun* 2005;17:236–42.
  57. Zoghi M, Jaberzadeh S. Effects of high voltage electro-auriculotherapy on experimental pain threshold. *Physiotherapy* 2002;88:658–66.
  58. Lorenzana FD. A randomized controlled trial of the efficacy of transcutaneous electrical nerve stimulation (TENS) versus lidocaine in the relief of episiotomy pain. *Philippine J Obstet Gynecol* 1999;23:135–42.
  59. Kim JH, Kim EJ, Seo BK, Lee S, Jung SY, Lee MH, et al. Electroacupuncture for chemotherapy-induced peripheral neuropathy: Study protocol for a pilot multicentre randomized, patient-assessor-blinded, controlled trial. *Trials* 2013;14:254–60.
  60. Lee S, Kim JH, Shin KM, Kim JE, Kim TH, Kang KW, et al. Electroacupuncture to treat painful diabetic neuropathy: study protocol for a three-armed, randomized, controlled pilot trial. *Trials* 2013;14:225 [11p].
  61. Choi KE, Rampp T, Saha FJ, Dobos GJ, Musial F. Pain modulation by meditation and electroacupuncture in experimental submaximum effort tourniquet technique (SETT). *Explore J Sci Heal* 2011;7:239–45.
  62. Lee SH, Lee BC. Electroacupuncture relieves pain in men with chronic prostatitis/chronic pelvic pain syndrome: three-arm randomized trial. *Urology* 2009;73:1036–41.
  63. Jubb RW, Tukmachi ES, Jones PW, Dempsey E, Waterhouse L, Brailsford S. A blinded randomised trial of acupuncture (manual and electroacupuncture) compared with a non-penetrating sham for the symptoms of osteoarthritis of the knee. *Acupunct Med* 2008;26:69–78.
  64. Michalek-Sauberer A, Heinzl H, Sator-Katzenschlager SM, Monov G, Knolle E, Kress HG. Perioperative auricular electroacupuncture has no effect on pain and analgesic consumption after third molar tooth extraction. *Anesth Analg* 2007;104:542–7.
  65. Zhang JH, Cao XD, Li J, Tang WJ, Liu HQ, Feng XY. Neuronal specificity of needling acupoints at same meridian: a control functional magnetic resonance imaging study with electroacupuncture. *Acupunct Electrother Res* 2007;32:179–93.
  66. Sator-Katzenschlager SM, Woelfler MM, Kozek-Langenecker SA, Sator K, Sator PG, Li B, et al. Auricular electro-acupuncture as an additional perioperative analgesic method during oocyte aspiration in IVF treatment. *Hum Reprod* 2006;21:2114–20.
  67. Tsui MLK, Cheing GLY. The effectiveness of electroacupuncture versus electrical heat acupuncture in the management of chronic low-back pain. *J Altern Complement Med* 2004;10:803–9.
  68. Sator-Katzenschlager SM, Scharbert G, Kozek-Langenecker SA, Szeles JC, Finster G, Schiesser AW, et al. The short- and long-term benefit in chronic low back pain through adjuvant electrical versus manual auricular acupuncture. *Anesth Analg* 2004;98:1359–64.
  69. Sator-Katzenschlager SM, Szeles JC, Scharbert G, Michalek-Sauberer A, Kober A, Heinze G, et al. Electrical stimulation of auricular acupuncture points is more effective than conventional manual auricular acupuncture in chronic cervical pain: a pilot study. *Anesth Analg* 2003;97:1469–73.
  70. Ng MML, Leung MCP, Poon DMY. The effects of electro-acupuncture and transcutaneous electrical nerve stimulation on patients with painful osteoarthritic knees: a randomized controlled trial with follow-up evaluation. *J Altern Complement Med* 2003;9:641–9.
  71. Tsui P, Leung MCP. Comparison of the effectiveness between manual acupuncture and electro-acupuncture on patients with tennis elbow. *Acupunct Electrother Res* 2002;27:107–17.
  72. Yoshimizu M, Teo AR, Ando M, Kiyohara K, Kawamura T. Relief of chronic shoulder and neck pain by electro-acupuncture and transcutaneous electrical nervous stimulation: a randomized crossover trial. *Med Acupunct* 2012;24:97–103.
  73. Musial F, Choi KE, Gabriel T, Lüdtke R, Rampp T, Michalsen A, et al. The effect of electroacupuncture and tramadol on experimental tourniquet pain. *Acupunct Med* 2012;30:21–6.
  74. Chan DK, Johnson MI, Sun KO, Doble SJ, Jenkins S. Electrical acustimulation of the wrist for chronic neck pain: a randomized, sham-controlled trial using a wrist-ankle acustimulation device. *Clin J Pain* 2009;25:320–6.
  75. Wong RH, Lee TW, Sihoe AD, Wan IY, Ng CS, Chan SK, et al. Analgesic effect of electroacupuncture in postthoracotomy pain: a prospective randomized trial. *Ann Thorac Surg* 2006;81:2031–6.
  76. Ernst M, Lee M. Influence of naloxone on electro-acupuncture analgesia using an experimental

- dental pain test. Review of possible mechanisms of action. *Acupunct Electrother Res* 1986;12:5–22.
77. Srbely JZ, Dickey JP, Lowerison M, Edwards AM, Nolet PS, Wong LL. Stimulation of myofascial trigger points with ultrasound induces segmental antinociceptive effects: a randomized controlled study. *Pain* 2008;139:260–6.
  78. Kempf D, Berger D, Ausfeld-Hafter B. Laser needle acupuncture in women with dysmenorrhoea: a randomised controlled double blind pilot trial. *Forsch Komplementmed* 2009;16:6–12.
  79. Aigner N, Fialka C, Radda C, Vecsei V. Adjuvant laser acupuncture in the treatment of whiplash injuries: a prospective, randomized placebo-controlled trial. *Wien Klin Wochenschr* 2006;118:95–9.
  80. Naeser MA, Hahn KAK, Lieberman BE, Branco KF. Carpal tunnel syndrome pain treated with low-level laser and microamperes transcutaneous electric nerve stimulation: a controlled study. *Arch Phys Med Rehabil* 2002;83:978–88.
  81. Takamjani IE, Maroufi N, Amoli MJ, Nia SHS. The effect of low power LASER acupuncture on experimental pain threshold in normal subjects. *Caspian J Intern Med* 2010;1:145–8.
  82. Glazov G, Schattner P, Lopez D, Shandley K. Laser acupuncture for chronic non-specific low back pain: a controlled clinical trial. *Acupunct Med* 2009;27:94–100.
  83. King CE, Clelland JA, Knowles CJ, Jackson JR. Effect of helium–neon laser auriculotherapy on experimental pain threshold. *Phys Ther* 1990;70:24–30.
  84. Waylonis GW, Wilke S, O'Toole D, Waylonis DA, Waylonis DB. Chronic myofascial pain: management by low-output helium–neon laser therapy. *Arch Phys Med Rehabil* 1988;69:1017–20.
  85. Kreczi T, Klingler D. A comparison of laser acupuncture versus placebo in radicular and pseudoradicular pain syndromes as recorded by subjective responses of patients. *Acupunct Electrother Res* 1986;11:207–16.
  86. McNearney TA, Sallam HS, Hunnicutt SE, Doshi D, Chen JDZ. Prolonged treatment with transcutaneous electrical nerve stimulation (TENS) modulates neuro-gastric motility and plasma levels of vasoactive intestinal peptide (VIP), motilin and interleukin-6 (IL-6) in systemic sclerosis. *Clin Exp Rheumatol* 2013;31:S140–50.
  87. Leung WW, Jones AYM, Ng SSM, Wong CYN, Lee JFY. Acupuncture transcutaneous electrical nerve stimulation reduces discomfort associated with barostat-induced rectal distension: a randomized-controlled study. *World J Gastroenterol* 2013;19:381–8.
  88. Liu S, Peng S, Hou X, Ke M, Chen JDZ. Transcutaneous electroacupuncture improves dyspeptic symptoms and increases high frequency heart rate variability in patients with functional dyspepsia. *Neurogastroenterol Motil* 2008;20:1204–11.
  89. Sallam H, McNearney TA, Doshi D, Chen JDZ. Transcutaneous electrical nerve stimulation (TENS) improves upper GI symptoms and balances the sympathovagal activity in scleroderma patients. *Digest Dis Sci* 2007;52:1329–37.
  90. Xing JH, Larive B, Mekhail N, Soffer E. Transcutaneous electrical acustimulation can reduce visceral perception in patients with the irritable bowel syndrome: a pilot study. *Altern Ther Health Med* 2004;10:38–42.
  91. Chang CS, Chou JW, Ko CW, Wu CY, Chen GH. Cutaneous electrical stimulation of acupuncture points may enhance gastric myoelectrical regularity. *Digestion* 2002;66:106–11.
  92. Zou D, Chen WH, Iwakiri K, Rigda R, Tippett M, Holloway RH. Inhibition of transient lower esophageal sphincter relaxations by electrical acupoint stimulation. *Am J Physiol Gastrointest Liver Physiol* 2005;289:197–201.
  93. Chen CY, Ke MD, Kuo CD, Huang CH, Hsueh YH, Chen JR. The influence of electro-acupuncture stimulation to female constipation patients. *Am J Chin Med* 2013;41:301–13.
  94. Wang CP, Kao CH, Chen WK, Lo WY, Hsieh CL. A single-blinded, randomized pilot study evaluating effects of electroacupuncture in diabetic patients with symptoms suggestive of gastroparesis. *J Altern Complement Med* 2008;14:833–9.
  95. Xu S, Hou X, Zha H, Gao Z, Zhang Y, Chen JDZ. Electroacupuncture accelerates solid gastric emptying and improves dyspeptic symptoms in patients with functional dyspepsia. *Digest Dis Sci* 2006;51:2154–9.
  96. Xu M, Zhou SJ, Jiang CC, Wu Y, Shi WL, Gu HH, et al. The effects of P6 electrical acustimulation on postoperative nausea and vomiting in patients after infratentorial craniotomy. *J Neurosurg Anesthesiol* 2012;24:312–6.
  97. Wang XQ, Yu JL, Du ZY, Xu R, Jiang CC, Gao X. Electroacupoint stimulation for postoperative nausea and vomiting in patients undergoing supratentorial craniotomy. *J Neurosurg Anesthesiol* 2010;22:128–31.
  98. Larson JD, Gutowski KA, Marcus BC, Rao VK, Avery PG, Stacey DH, et al. The effect of electroacustimulation on postoperative nausea, vomiting, and pain in outpatient plastic surgery patients: a prospective, randomized, blinded, clinical trial. *Plastic Reconstr Surg* 2010;125:989–94.
  99. Liu YY, Duan SE, Cai MX, Zou P, Lai Y, Li YL. Evaluation of transcutaneous electroacupoint stimulation with the train-of-four mode for preventing nausea and vomiting after laparoscopic cholecystectomy. *Chin J Integr Med* 2008;14:94–7.
  100. Kabalak AA, Akcay M, Akcay F, Gogus N. Transcutaneous electrical acupoint stimulation versus ondansetron in the prevention of postoperative vomiting following pediatric tonsillectomy. *J Altern Complement Med* 2005;11:407–13.
  101. Kramer BA, Kadar AG, Clark K. Transcutaneous acupoint electrical stimulation in preventing and treating nausea and vomiting in patients receiving electroconvulsive therapy. *J ECT* 2003;19:194–6.
  102. Habib AS, Itchon-Ramos N, Phillips-Bute BG, Gan TJ. Transcutaneous acupoint electrical stimulation with the ReliefBand for the prevention of nausea and vomiting during and after cesarean delivery under spinal anesthesia. *Anesth Analg* 2006;102:581–4.
  103. Zárate E, Mingus M, White PF, Chiu JW, Scuderi P, Loskota W, et al. The use of transcutaneous acupoint electrical stimulation for preventing nausea and vomiting after laparoscopic surgery. *Anesth Analg* 2001;92:629–35.
  104. McMillan CM, Dundee JW. The role of transcutaneous electrical stimulation of Neiguan anti-emetic acupuncture point in controlling sickness after cancer chemotherapy. *Physiotherapy* 1991;77:499–502.
  105. Rusy LM, Hoffman GM, Weisman SJ. Electroacupuncture prophylaxis of postoperative nausea and vomiting following pediatric tonsillectomy with or without adenoidectomy. *Anesthesiology* 2002;96:300–5.
  106. Shen J, Wenger N, Glaspy J, Hays RD, Albert PS, Choi C, et al. Electroacupuncture for control of myeloablative chemotherapy-induced emesis: a randomized controlled trial. *JAMA* 2000;284:2755–61.
  107. Schlager A, Offer T, Baldissera I. Laser stimulation of acupuncture point P6 reduces postoperative vomiting in children undergoing strabismus surgery. *Br J Anaesth* 1998;81:529–32.
  108. Kim SB, Kim JY, Park SW, Lee NR, Kim YH, Lee KJ, et al. Effects of PEMFs (pulsed electromagnetic fields) stimulation on acupoint in quadriceps fatigue recovery. *Int J Precis Eng Manuf* 2012;13:1697–703.

109. Kim SB, Kim JY, Park SW, Lee NR, Lee SW, Kim YH, et al. Comparison of 2 methods of non-invasive treatment between transcutaneous electrical stimulation and pulsed electromagnetic field stimulation as replacement of invasive manual acupuncture. *Acupunct Electrother Res* 2012;37:247–61.
110. Zhou S, Huang LP, Liu J, Yu JH, Tian Q, Cao LJ. Bilateral effects of 6 weeks' unilateral acupuncture and electroacupuncture on ankle dorsiflexors muscle strength: a pilot study. *Arch Phys Med Rehabil* 2012;93:50–5.
111. Ngai SPC, Jones AYM, Hui-Chan CWY. Acu-TENS and postexercise expiratory flow volume in healthy subjects. *Evid Based Complement Altern Med* 2011;2011:1–7.
112. Huang LP, Zhou S, Lu Z, Tian Q, Li X, Cao LJ, et al. Bilateral effect of unilateral electroacupuncture on muscle strength. *J Altern Complement Med* 2007;13:539–46.
113. Chiu TTW, Hui-Chan CWY, Cheing G. A randomized clinical trial of TENS and exercise for patients with chronic neck pain. *Clin Rehabil* 2005;19:850–60.
114. Milne RJ, Dawson NJ, Butler MJ, Lippold OCJ. Intramuscular acupuncture-like electrical stimulation inhibits stretch reflexes in contralateral finger extensor muscles. *Exp Neurol* 1985;90:96–107.
115. Schukro RP, Heiserer C, Michalek-Sauberer A, Gleiss A, Sator-Katzenschlager S. The effects of auricular electroacupuncture on obesity in female patients—a prospective randomized placebo-controlled pilot study. *Complement Ther Med* 2014;22:21–5.
116. Chien LW, Lin MH, Chung HY, Liu CF. Transcutaneous electrical stimulation of acupoints changes body composition and heart rate variability in postmenopausal women with obesity. *Evid Based Complement Altern Med* 2011;2011:862121.
117. Rerksuppaphol L, Rerksuppaphol S. Efficacy of transcutaneous electrical acupoint stimulation compared to Electroacupuncture at the main acupoints for weight reduction in obese Thai women. *Int J Collab Res Intern Med Public Health* 2011;3:811–20.
118. Lin CH, Lin YM, Liu CF. Electrical acupoint stimulation changes body composition and the meridian systems in postmenopausal women with obesity. *Am J Chin Med* 2010;38:683–94.
119. Jeong KH, Lee MH. Two cases of factitial panniculitis induced by electroacupuncture. *Clin Exp Dermatol* 2009;34:E170–3.
120. Quah-Smith I, Suo C, Williams M, Sachdev P. The antidepressant effect of laser acupuncture: a comparison of the resting brain's default mode network in healthy and depressed subjects during functional magnetic resonance imaging. *Med Acupunct* 2013;25:124–33.
121. Quah-Smith I, Smith C, Crawford JD, Russell J. Laser acupuncture for depression: a randomised double blind controlled trial using low intensity laser intervention. *J Affect Disord* 2013;148:179–87.
122. Penetar DM, Burgos-Robles A, Trksak GH, MacLean RR, Dunlap S, Lee DW, et al. Effects of transcutaneous electric acupoint stimulation on drug use and responses to cue-induced craving: a pilot study. *Chin Med (UK)* 2012;7:1–10.
123. Lambert C, Berlin I, Lee TL, Hee SW, Tan ASL, Picard D, et al. A standardized transcutaneous electric acupoint stimulation for relieving tobacco urges in dependent smokers. *Evid Based Complement Altern Med* 2011;2011:195714.
124. Georgiou AJ, Spencer CP, Davies GK, Stamp J. Electrical stimulation therapy in the treatment of cigarette smoking. *J Subst Abuse* 1998;10:265–74.
125. Kerr CM, Lowe PB, Spielholz NI. Low level laser for the stimulation of acupoints for smoking cessation: a double blind, placebo controlled randomised trial and semi structured interviews. *J Chin Med* 2008;86:46–51.
126. Zalewska-Kaszubska J, Obzejta D. Use of low-energy laser as adjunct treatment of alcohol addiction. *Lasers Med Sci* 2004;19:100–4.
127. Ng SSM, Hui-Chan CWY. Transcutaneous electrical stimulation on acupoints combined with task-related training to improve motor function and walking performance in an individual 7 years poststroke: a case study. *J Neurol Phys Ther* 2010;34:208–13.
128. Gong W, Zhang T, Cui L, Yang Y, Sun X. Electro-acupuncture at Zusanli (ST 36) to improve lower extremity motor function in sensory disturbance patients with cerebral stroke: a randomized controlled study of 240 cases. *Neural Regen Res* 2009;4:935–40.
129. Kim YS, Hong JW, Na BJ, Park SU, Jung WS, Moon SK, et al. The effect of low versus high frequency electrical acupoint stimulation on motor recovery after ischemic stroke by motor evoked potentials study. *Am J Chin Med* 2008;36:45–54.
130. Wong AM, Su TY, Tang FT, Cheng PT, Liaw MY. Clinical trial of electrical acupuncture on hemiplegic stroke patients. *Am J Phys Med Rehab* 1999;78:117–22.
131. Cafaro A, Arduino PG, Gambino A, Romagnoli E, Broccoletti R. Effect of laser acupuncture on salivary flow rate in patients with Sjögren's syndrome. *Lasers Med Sci* 2015;30:1805–9.
132. Litscher G, Wang L, Wang X, Gaischek I. Laser acupuncture: two acupoints (Baihui, Neiguan) and two modalities of laser (658 nm, 405 nm) induce different effects in neurovegetative parameters. *Evid Based Complement Altern Med* 2013;2013:432764.
133. Wang G, Tian Y, Jia S, Litscher G, Zhang W. Evaluate laser needle effect on blood perfusion signals of contralateral Hegu acupoint with wavelet analysis. *Evid Based Complement Altern Med* 2012;2012:103729.
134. Raith W, Litscher G, Sapetschnig I, Bauchinger S, Ziehenberger E, Mueller W, et al. Thermographical measuring of the skin temperature using laser needle acupuncture in preterm neonates. *Evid Based Complement Altern Med* 2012;2012:614210.
135. Litscher G, Xie Z, Wang L, Gaischek I. Blue 405 nm laser light mediates heart rate—investigations at the acupoint Neiguan (Pe. 6) in Chinese adults. *North Am J Med Sci* 2009;1:226–31.
136. Zhang J, Marquina N, Oxinos G, Sau A, Ng D. Effect of laser acupoint treatment on blood pressure and body weight—a pilot study. *J Chiropr Med* 2008;7:134–9.
137. Banzer W, Huebscher M, Seib M, Vogt L. Short-time effects of laser needle stimulation on the peripheral microcirculation assessed by laser Doppler spectroscopy and near-infrared spectroscopy. *Photomed Laser Surg* 2006;24:575–80.
138. Litscher G. Cerebral and peripheral effects of Laserneedle®-stimulation. *Neurol Res* 2003;25:722–8.
139. Litscher G, Schikora D. Cerebral vascular effects of non-invasive laserneedles measured by transorbital and transtemporal Doppler sonography. *Lasers Med Sci* 2002;17:289–95.
140. Cunha RG, Rodrigues KC, Salvador M, Zangaro RA. Effectiveness of laser treatment at acupuncture sites compared to traditional acupuncture in the treatment of peripheral artery disease. In: *Engineering in Medicine and Biology Society (EMBC), 2010 Annual International Conference of the IEEE*. 2010:1262–5.



141. Litscher G, Wang L, Wiesner-Zechmeister M. Specific effects of laserpuncture on the cerebral circulation. *Lasers Med Sci* 2000;15:57–62.
142. Sun J, Sang H, Yang C, Dong H, Lei C, Lu Y, et al. Electroacupuncture improves orthostatic tolerance in healthy individuals via improving cardiac function and activating the sympathetic system. *Europace* 2013;15:127–34.
143. Chen CW, Tai CJ, Choy CS, Hsu CY, Lin SL, Chan WP, et al. Wave-induced flow in meridians demonstrated using photoluminescent bioceramic material on acupuncture points. *Evid Based Complement Altern Med* 2013;2013:739293.
144. Guo L, Wang Y, Yu H, Yin N, Li Y. Study of brain functional network based on sample entropy of EEG under magnetic stimulation at PC6 acupoint. *Biomed Mater Eng* 2014;24:1063–9.
145. Zhang X, Fu LD, Geng YH, Zhai X, Liu YH. Analysis of the effect of repeated-pulse transcranial magnetic stimulation at the Guangming point on electroencephalograms. *Neural Regen Res* 2014;9:549–54.
146. Raith W, Pichler G, Sapetschnig I, Avian A, Sommer C, Baik N, et al. Near-infrared spectroscopy for objectifying cerebral effects of laser acupuncture in term and preterm neonates. *Evid Based Complement Altern Med* 2013;2013:346852.
147. Quah-Smith I, Williams MA, Lundeborg T, Suo C, Sachdev P. Differential brain effects of laser and needle acupuncture at LR8 using functional MRI. *Acupunct Med* 2013;31:282–9.
148. Zhang Y, Jiang Y, Glielmi CB, Li L, Hu X, Wang X, et al. Long-duration transcutaneous electric acupoint stimulation alters small-world brain functional networks. *Magn Reson Imag* 2013;31:1105–11.
149. Yin N, Xu G-Z, Zhou Q. Construction and analysis of complex brain functional network under acupoint magnetic stimulation. *Acta Phys Sin* 2013;62:118704.
150. Lee HS, Hwang DG, Cha Y-Y. An analytical comparison in electroencephalography and electrocardiography under pulsed magnetic field and acupuncture stimulus on acupoint PC9. *J Magn* 2013;18:192–6.
151. Litscher G. Laser acupuncture-innovative basic research: visual and laser-induced evoked potentials. *Laser Ther* 2012;21:287–95.
152. Wu JH, Chang WD, Hsieh CW, Jiang JA, Fang W, Shan YC, et al. Effect of low-level laser stimulation on EEG. *Evid Based Complement Altern Med* 2012;2012:951272.
153. Litscher G, Bauernfeind G, Mueller-Putz G, Neuper C. Laser-induced evoked potentials in the brain after nonperceptible optical stimulation at the Neiguan acupoint: a preliminary report. *Evid Based Complement Altern Med* 2012;2012:292475.
154. Yu H, Xu G, Zhou Q, Yang S, Geng Y, Xie X. EEG characteristics under magnetic stimulation at acupuncture point and at mock point. *IEEE Trans Magn* 2012;48:2857–60.
155. Jiang Y, Hao Y, Zhang Y, Liu J, Wang X, Han J, et al. Thirty minute transcutaneous electric acupoint stimulation modulates resting state brain activities: a perfusion and BOLD fMRI study. *Brain Res* 2012;1457:13–25.
156. Hsieh CW, Wu JH, Hsieh CH, Wang QF, Chen JH. Different brain network activations induced by modulation and nonmodulation laser acupuncture. *Evid Based Complement Altern Med* 2011;2011:951258, <http://dx.doi.org/10.1155/2011/951258>.
157. Yu H, Xu G, Yang S, Zhou Q, Wan X, Li W, et al. Activation of cerebral cortex evoked by magnetic stimulation at acupuncture point. *IEEE Trans Magn* 2011;47:3052–5.
158. Kim S, Lee J, Lee HS. The effect of magnetic field direction on the EEG and PPG obtained from pulsed magnetic stimulus at acupoint PC9. *J Magn* 2011;16:259–62.
159. Jo HG, Jo GH. Electroencephalogram activity induced by magnetic stimulation on heart meridian. *Neurosci Lett* 2011;495:107–9.
160. Zyloney CE, Jensen K, Polich G, Loiotile RE, Cheetham A, LaViolette PS, et al. Imaging the functional connectivity of the Periaqueductal Gray during genuine and sham electroacupuncture treatment. *Mol Pain* 2010;6:80 [11p].
161. Quah-Smith I, Sachdev PS, Wen W, Chen X, Williams MA. The brain effects of laser acupuncture in healthy individuals: an fMRI investigation. *PLoS One* 2010;5:e12619.
162. Xu G, Zhang X, Yu H, Ho SL, Yang Q, Fu WN, et al. Complexity analysis of EEG under magnetic stimulation at acupoints. *IEEE Trans Appl Supercond* 2010;20:1029–32.
163. Na B-J, Jahng G-H, Park S-U, Jung W-S, Moon S-K, Park J-M, et al. An fMRI study of neuronal specificity of an acupoint: Electroacupuncture stimulation of Yanglingquan (GB34) and its sham point. *Neurosci Lett* 2009;464:1–5.
164. Xu G, Zhang X, Ho SL, Fu WN, Yan W, Wang Y. Complexity analysis of magnetic stimulation at the acupoint of Zusanli (St36) on EEG. *IEEE Trans Magn* 2009;45:4829–32.
165. An Y-S, Moon S-K, Min I-K, Kim D-Y. Changes in regional cerebral blood flow and glucose metabolism following electroacupuncture at LI 4 and LI 11 in normal volunteers. *J Altern Complement Med* 2009;15:1075–81.
166. Wang W, Liu L, Zhi X, Huang J-B, Liu D-X, Wang H, et al. Study on the regulatory effect of electro-acupuncture on Hegu point (L14) in cerebral response with functional magnetic resonance imaging. *Chin J Integr Med* 2007;13:10–6.
167. Zeng Y, Liang X-C, Dai J-P, Wang Y, Yang Z-L, Li M, et al. Electroacupuncture modulates cortical activities evoked by noxious somatosensory stimulations in human. *Brain Res* 2006;1097:90–100.
168. Litscher G, Rachbauer D, Ropele S, Wang L, Schikora D, Fazekas F, et al. Acupuncture using laser needles modulates brain function: first evidence from functional transcranial Doppler sonography and functional magnetic resonance imaging. *Lasers Med Sci* 2004;19:6–11.
169. Zhang WT, Jin Z, Cui GH, Zhang KL, Zhang L, Zeng YW, et al. Relations between brain network activation and analgesic effect induced by low vs. high frequency electrical acupoint stimulation in different subjects: a functional magnetic resonance imaging study. *Brain Res* 2003;982:168–78.
170. Li G, Liu HL, Cheung RTF, Hung YC, Wong KKK, Shen GGX, et al. An fMRI study comparing brain activation between word generation and electrical stimulation of language-implicated acupoints. *Hum Brain Map* 2003;18:233–8.
171. Kong JA, Ma L, Gollub RL, Wei JH, Yang XZ, Li DJ, et al. A pilot study of functional magnetic resonance imaging of the brain during manual and electroacupuncture stimulation of acupuncture point (LI-4 Hegu) in normal subjects reveals differential brain activation between methods. *J Altern Complement Med* 2002;8:411–9.
172. Siedentopf CM, Golaszewski SM, Mottaghy FM, Ruff CC, Felber S, Schlager A. Functional magnetic resonance imaging detects activation of the visual association cortex during laser acupuncture of the foot in humans. *Neurosci Lett* 2002;327:53–6.
173. Wu MT, Sheen JM, Chuang KH, Yang P, Chin SL, Tsai CY, et al. Neuronal specificity of acupuncture response: a fMRI study with electroacupuncture. *Neuroimage* 2002;16:1028–37.
174. Chang QY, Lin JG, Hsieh CL. Effect of manual acupuncture and transcutaneous electrical nerve stimulation on the H-reflex. *Acupunct Electrother Res* 2001;26:239–51.
175. Kim YJ, Yoo JS, Hwang DG, Lee HS. Comparative analysis of photoplethysmography under pulsed magnetic field and



- low level laser stimulus: motivation for blood flow increase using stimulus on acupoint LI4 (Hegu). *J Magn* 2014;19:32–6.
176. Li G, Li S, Wang B, An L. The effect of electroacupuncture on postoperative immunoinflammatory response in patients undergoing supratentorial craniotomy. *Exp Ther Med* 2013;6:699–702.
  177. Tsuruoka N, Watanabe M, Takayama S, Seki T, Matsunaga T, Haga Y. Brief effect of acupoint stimulation using focused ultrasound. *J Altern Complement Med* 2013;19:416–9.
  178. Lee J, Hwang DG, Yoo JS, Lee HS. Analysis of electroencephalogram and electrocardiogram at an acupoint PC9 during pulsed magnetic field stimulus. *J Magn* 2012;17:1133–7.
  179. Jia B-A, Cheng C-Y, Lin Y-W, Li T-C, Liu H-J, Hsieh C-L. The 2 Hz and 15 Hz electroacupuncture induced reverse effect on autonomic function in healthy adult using a heart rate variability analysis. *J Tradit and Complement Med* 2011;1:51–6.
  180. Jones AYM, Kwan YL, Leung NTF, Yu RPW, Wu CMY, Warburton DER. Electrical stimulation of acupuncture points and blood pressure responses to postural changes: a pilot study. *Am J Crit Care* 2011;20:E67–74.
  181. Lee JH, Kim KH, Hong JW, Lee WC, Koo S. Comparison of electroacupuncture frequency-related effects on heart rate variability in healthy volunteers: a randomized clinical trial. *J Acupunct Meridian Stud* 2011;4:107–15.
  182. Chang C-S, Ko C-W, Lien H-C, Chou M-C. Effect of electroacupuncture on St. 36 (Zusanli) and LI. 10 (Shousanli) acupuncture points on heart rate variability. *Am J Chin Med* 2010;38:231–9.
  183. Kim MS, Cho YC, Moon JH, Pak SC. A characteristic estimation of bio-signals for electro-acupuncture stimulations in human subjects. *Am J Chin Med* 2009;37:505–17.
  184. Lu C-C, Jan Y-M, Li T-C, Hsieh C-L. Electroacupuncture induces differential effects between Yin and Yang: a study using cutaneous blood flow and temperature recordings of the hand's dorsum and palm. *Am J Chin Med* 2009;37:639–45.
  185. Zhang J, Ng D, Sau A. Effects of electrical stimulation of acupuncture points on blood pressure. *J Chiropr Med* 2009;8:9–14.
  186. Cakmak YO, Akpınar IN, Ekinci G, Bekiroğlu N. Point- and frequency-specific response of the testicular artery to abdominal electroacupuncture in humans. *Fertil Steril* 2008;90:1732–8.
  187. Arai YC, Kato N, Matsura M, Ito H, Kandatsu N, Kurokawa S, et al. Transcutaneous electrical nerve stimulation at the PC-5 and PC-6 acupoints reduced the severity of hypotension after spinal anaesthesia in patients undergoing caesarean section. *Br J Anaesth* 2008;100:78–81.
  188. Cheung LC-T, Jones AY-M. Effect of Acu-TENS on recovery heart rate after treadmill running exercise in subjects with normal health. *Complement Ther Med* 2007;15:109–14.
  189. Szeles JC, Litscher G. Objectivation of cerebral effects with a new continuous electrical auricular stimulation technique for pain management. *Neurol Res* 2004;26:797–800.
  190. Li Z, Jiao K, Chen M, Wang C. Effect of magnitopuncture on sympathetic and parasympathetic nerve activities in healthy drivers—assessment by power spectrum analysis of heart rate variability. *Eur J Appl Physiol* 2003;88:404–10.
  191. Hsieh CL, Wu CHW, Lin JG, Chiu CC, Chen M, Hsieh CT. The physiological mechanisms of 2 Hz electroacupuncture: a study using blink and H reflex. *Am J Chin Med* 2002;30:369–78.
  192. Cramp AF, Noble JG, Lowe AS, Walsh DM. Transcutaneous electrical nerve stimulation (TENS): the effect of electrode placement upon cutaneous blood flow and skin temperature. *Acupunct Electrother Res* 2000;26:25–37.
  193. Balogun JA, Tang S, He Y, Hsieh JM, Katz JS. Effects of high-voltage galvanic stimulation of ST36 and ST37 acupuncture points on peripheral blood flow and skin temperature. *Disabil Rehabil* 1996;18:523–8.
  194. Williams T, Mueller K, Cornwall MW. Effect of acupuncture-point stimulation on diastolic blood pressure in hypertensive subjects: a preliminary study. *Phys Ther* 1991;71:523–9.
  195. Dunn PA, Rogers D, Halford K. Transcutaneous electrical nerve stimulation at acupuncture points in the induction of uterine contractions. *Obstet Gynecol* 1989;73:286–90.
  196. Zhang R, Jia MX, Zhang JS, Xu XJ, Shou XJ, Zhang XT, et al. Transcutaneous electrical acupoint stimulation in children with autism and its impact on plasma levels of arginine-vasopressin and oxytocin: a prospective single-blinded controlled study. *Res Dev Disabil* 2012;33:1136–46.
  197. Yang C, An L, Han R, Kang X, Wang B. Effects of combining electroacupuncture with general anesthesia induced by sevoflurane in patients undergoing supratentorial craniotomy and improvements in their clinical recovery profile & blood enkephalin. *Acupunct Electrother Res* 2012;37:125–38.
  198. Sahmeddini MA, Eghbal MH, Khosravi MB, Ghaffaripour S, Janatmakan F, Shokrizade S. Electro-acupuncture stimulation at acupoints reduced the severity of hypotension during anesthesia in patients undergoing liver transplantation. *J Acupunct Meridian Stud* 2012;5:11–4.
  199. Ng MCS, Jones AYM, Cheng LC. The role of Acu-TENS in hemodynamic recovery after open-heart surgery. *Evid Based Complement Altern Med* 2011;2011:301974.
  200. Wang K, Bugge J, Bugge S. A randomised, placebo-controlled trial of manual and electrical acupuncture for the treatment of tinnitus. *Complement Ther Med* 2010;18:249–55.
  201. O'Brien KA, Varigos E, Black C, Komesaroff PA. Laser acupuncture does not improve menopausal symptoms. *Menopause* 2010;17:636–41.
  202. Ngai SPC, Jones AYM, Hui-Chan CWY, Ko FWS, Hui DSC. Effect of Acu-TENS on post-exercise expiratory lung volume in subjects with asthma—a randomized controlled trial. *Respir Physiol Neurobiol* 2009;167:348–53.
  203. Burduli NM, Ranyuk LG. Effects of laser reflex therapy on a motor function of the gall bladder and physical properties of bile in patients with chronic acalculous cholecystitis. *Ter Arkh* 2009;81:57–61.
  204. Su LH, Wu KD, Lee LS, Wang H, Liu CF. Effects of far infrared acupoint stimulation on autonomic activity and quality of life in hemodialysis patients. *Am J Chin Med* 2009;37:215–26.
  205. Lau KSL, Jones AYM. A single session of Acu-TENS increases FEV1 and reduces dyspnoea in patients with chronic obstructive pulmonary disease: a randomised, placebo-controlled trial. *Aust J Physiother* 2008;54:179–84.
  206. Hsu CC, Weng CS, Liu TS, Tsai YS, Chang YH. Effects of electrical acupuncture on acupoint BL15 evaluated in terms of heart rate variability, pulse rate variability and skin conductance response. *Am J Chin Med* 2006;34:23–36.
  207. Bray PA, Mamiya N, Fann AV, Gellman H, Skinner RD, Garcia-Rill EE. Modulation of the sleep state-dependent P50 midlatency auditory-evoked potential by electric stimulation of acupuncture points. *Arch Phys Med Rehabil* 2005;86:2018–26.
  208. Litscher G, Wang L, Schwarz G, Schikora D. Increases of intracranial pressure and changes of blood flow velocity due to acupressure, needle and laserneedle acupuncture? *Forsch Komplementarmed Klass Naturheilkd* 2005;12:190–5.
  209. O'Reilly BA, Dwyer PL, Hawthorne G, Cleaver S, Thomas E, Rosamilia A, et al. Transdermal posterior tibial nerve laser

- therapy is not effective in women with interstitial cystitis. *J Urol* 2004;172:1880–3.
210. Li Z, Jiao K, Chen M, Wang C. Reducing the effects of driving fatigue with magnitopuncture stimulation. *Accid Anal Prev* 2004;36:501–5.
211. Gopalan R, Scott R, Arabia F, Chandrasekaran K. Electro-acupuncture therapy in a patient with a total artificial heart. *Acupunct Med* 2011;29:302–3.
212. Irnich D, Salih N, Offenbacher M, Fleckenstein J. Is sham laser a valid control for acupuncture trials? *Evid Based Complement Altern Med* 2011;2011:485945.
213. Litscher G, Wang L. Biomedical engineering meets acupuncture—development of a miniaturized 48-channel skin impedance measurement system for needle and laser acupuncture. *Biomed Eng Online* 2010;9:78.
214. Thompson JW, Cummings M. Investigating the safety of electroacupuncture with a Picoscope™. *Acupunct Med* 2008;26:133–9.
215. Leung AY, Park J, Schulteis G, Duann J-R, Yaksh T. The electrophysiology of de qi sensations. *J Altern Complement Med* 2006;12:743–50.
216. Litscher G, Wang L, Huber E, Schikora D, Schwarz G. Quantification of gender specific thermal sensory and pain threshold before and after laserneedle stimulation. *Biomed Tech* 2004;49:106–10.
217. Chang QY, Lin JG, Hsieh CL. Effect of electroacupuncture and transcutaneous electrical nerve stimulation at Hegu (LI.4) acupuncture point on the cutaneous reflex. *Acupunct Electrother Res* 2002;27:191–202.