

RESEARCH ARTICLE

Open Access

Identification of transcription factors and single nucleotide polymorphisms of *Lrh1* and its homologous genes in *Lrh1*-knockout pancreas of mice

Maochun Tang^{1†}, Li Cheng^{2†}, Rongrong Jia¹, Lei Qiu¹, Hua Liu¹, Shu Zhou¹, Xiuying Ma¹, Guoyong Hu², Xingpeng Wang² and Yan Zhao^{1*}

Abstract

Background: To identify transcription factors (TFs) and single nucleotide polymorphisms (SNPs) of *Lrh1* (also named *Nr5a2*) and its homologous genes in *Lrh1*-knockout pancreas of mice.

Methods: The RNA-Seq data GSE34030 were downloaded from Gene Expression Omnibus (GEO) database, including 2 *Lrh1* pancreas knockout samples and 2 wild type samples. All reads were processed through TopHat and Cufflinks package to calculate gene-expression level. Then, the differentially expressed genes (DEGs) were identified via non-parametric algorithm (NOISeq) methods in R package, of which the homology genes of *Lrh1* were identified via BLASTN analysis. Furthermore, the TFs of *Lrh1* and its homologous genes were selected based on TRANSFAC database. Additionally, the SNPs were analyzed via SAM tool to record the locations of mutant sites.

Results: Total 15683 DEGs were identified, of which 23 was *Lrh1* homology genes (3 up-regulated and 20 down-regulated). Fetoprotein TF (FTF) was the only TF of *Lrh1* identified and the promoter-binding factor of FTF was *CYP7A*. The SNP annotations of *Lrh1* homologous genes showed that 92% of the mutation sites were occurred in intron and upstream. Three SNPs of *Lrh1* were located in intron, while 1819 SNPs of *Phkb* were located in intron and 1343 SNPs were located in the upstream region.

Conclusion: FTF combined with *CYP7A* might play an important role in *Lrh1* regulated pancreas-specific transcriptional network. Furthermore, the SNPs analysis of *Lrh1* and its homology genes provided the candidate mutant sites that might affect the *Lrh1*-related production and secretion of pancreatic fluid.

Keyword: *Lrh1*-knockout pancreas, RNA-Seq, *Lrh1* homologous gene, Transcription factor, Single nucleotide polymorphisms

Background

The pancreas is an endocrine gland, producing insulin, glucagon, somatostatin, and pancreatic polypeptide, and also an exocrine gland, accounting for more than 98% of pancreatic gland and secreting pancreatic juice containing digestive enzymes [1]. These digestive enzymes help to further break down the carbohydrates, proteins and

lipids in the chime and thus support the absorption and digestion of nutrition in small intestine [2]. In the past decades, many research have focused on target genes and transcription factors (TFs) involved in the exocrine pancreas-specific transcriptional networks which are required for the production and secretion of pancreatic fluid that helps out the digestive system. Currently, many exocrine pancreas-specific genes and transcription factors have been identified, which may promote the understanding of the effect of exocrine pancreas on digestive system.

* Correspondence: 320zhaoyan@163.com

†Equal contributors

¹Department of Gastroenterology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, No.301, Yanchang Middle Road, Shanghai 200072, China

Full list of author information is available at the end of the article

Liver receptor homolog-1 (*Lrh1*; also called *Nr5a2*) is a nuclear receptor of ligand-activated transcription factors in liver by binding as a monomer to DNA sequence elements with the consensus sequence 5'-Py-CAAGGPyCPu-3' [3]. It has been suggested that *Lrh1* is progressively expressed in both the endocrine and exocrine pancreas [4]. Baquié M et al. [5] have found that *Lrh1* is expressed in human islets and protects β -cells against stress-induced apoptosis that may be mediated via the increased glucocorticoid production that blunts the pro-inflammatory response of islets. Meanwhile, Fayard E et al. [6] have demonstrated that both *Lrh1* and *CEL* (encoding carboxyl ester lipase) are co-expressed and confined to the exocrine pancreas. The identification of *CEL* as an *Lrh1*-target gene indicates that *Lrh1* plays an important role in enterohepatic cholesterol homeostasis associated with the absorption of cholesteryl esters and the assembly of lipoproteins by the intestine [7]. Besides, *Lrh1* is a downstream target in the *PDX-1* (lead to pancreas agenesis) regulatory cascade that is activated only during early stages of pancreas development and that governs pancreatic development, differentiation and function [8].

Recently, the rapid advent of next-generation sequencing has made this technology broadly available for researchers in various molecular and cellular biological fields. Holmstrom SR et al. [9] have determined the cistrome and transcriptome for the nuclear receptor LRH-1 in exocrine pancreas and revealed that *Lrh1* directly induces expression of genes encoding digestive enzymes and secretory and mitochondrial proteins based on Chromatin immunoprecipitation (ChIP)-seq and RNA-seq analyses. Besides, *Lrh1* cooperates with the pancreas transcription factor 1-L complex (PTF1-L) in regulation of exocrine pancreas-specific gene expression. However, many potential target genes and TFs of *Lrh1* based on RNA-seq analysis have not been revealed.

In the present study, we downloaded the raw RNA-seq data of Holmstrom SR et al. deposited in The National Center for Biotechnology Information (NCBI) database, which were analyzed using multiple bioinformatics tools in the purpose of finding specific TFs of *Lrh1* and its homology genes. Additionally, we also annotated the SNPs of *Lrh1* and its homology genes to predict their mutant sites. Our study might improve the understanding of the regulation network of *Lrh1*-related production and secretion of pancreatic fluid.

Methods

RNA-seq data acquisition

The RNA-seq data was downloaded from NCBI (<http://www.ncbi.nlm.nih.gov/>) Gene Expression Omnibus (GEO) database (GEO accession: GSE34030 [9]), including 2 *Lrh1* pancreas knockout samples and 2 wild type samples. RNA preparations were subjected to the Illumina RNA-seq protocol and the platform was GPL9185.

Data pre-processing, gene expression and homology gene of *Lrh1*

The raw data were downloaded from SRA (Sequence Read Archive) of NCBI and then converted to fastq reads using fastq-dump program of NCBI SRA Toolkit (-q 64) (<http://trace.ncbi.nlm.nih.gov/Traces/sra/sra.cgi?view=std>). Then, these reads were processed through TopHat [10] and Cufflinks [11] package to calculate gene-expression level. All parameters were set up according to the default settings of TopHat and Cufflinks. The DEGs were identified via non-parametric algorithm (NOISEq) methods in R package [12]. The thresholds value was False Discovery Rate (FDR) < 0.001. BLASTN analysis [13,14] of the selected DEGs was used to identify the homology genes of *Lrh1*. Homology genes here refer to the paralogous genes which share a high degree of sequence similarity (maximum expectation value was set to e^{-5}) with *Lrh1* in mice.

Function annotation of *Lrh1* homologous genes

For functional analysis of *Lrh1* homologous genes, DAVID (Database for Annotation, Visualization and Integrated Discovery) [15] was performed for Gene Ontology (GO) [16] function and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathway enrichment analysis.

Transcription factor (TF) of *Lrh1* homologous genes

Combined with TRANSFAC database [17], the TFs regulated the transcription of *Lrh1* and its homologous genes were identified. Then, the promoter-binding factors regulated via the selected TFs were analyzed based on the website (<http://www.nursa.org/molecule.cfm?molType=receptor&molId=5A2>).

Screening of SNPs

The fastq reads were mapped to marker sequences using bowtie [18]. And the aligned reads were called using the SAM tool [19]. In order to minimize the risk of false-positive SNP Callings, the threshold value was that ID was "*" with quality > 50, or ID was not "*" with quality > 20. These SNPs were annotated via SnpEff [20] to categorize the effects of variants in genome sequences. The identified SNPs were searched in the dbSNP database to identify diseased SNPs or de novo discovered SNPs.

Results

Identification and homology analysis of differentially expressed genes

After data processing, at FDR < 0.001, a total of 15683 DEGs were identified, including 10994 up-regulated and 4698 down-regulated genes. BLASTN analysis of DEGs showed 23 *Lrh1* homology genes. Among them, 3 were up-regulated and 20 were down-regulated (Table 1).

Table 1 *Lrh1* homology genes of differentially expressed genes

Regulation	<i>Lrh1</i> homology genes
Up-regulated	Fzd5, Klr1f, Phc1
Down-regulated	March8, Zfp282, Atrx, Sos2, Zfc3h1, Pofut1, Gm9079, Ykt6, Phkb, Galc, Setd1a, Fzd5, Kazn, Kcnmb1, Lamc2, Mylk4, Ache, Pbxip1, Lrh1, Phc1

Function and pathway annotation of *Lrh1* homologous genes

To determine the function of *Lrh1* homologous genes in pancreas, GO enrichment analysis and KEGG pathway enrichment analysis were used to analyze the up- and down-regulated *Lrh1* homologous genes. For function and pathway annotation, DEGs were enriched into hexose metabolic process and monosaccharide metabolic process, which were involved into glycometabolism (Figure 1). Meanwhile, KEGG pathway enrichment analysis identified insulin signaling pathway, indicating that the disorders of glycometabolism might be resulted from insulin resistance and/or insulin secretion (Figure 2). *PHKB*, an *Lrh1* homologous gene, participated in GO terms (hexose metabolic process and monosaccharide metabolic process) and KEGG pathway (insulin signaling pathway), was identified.

Potential TFs of *Lrh1* homologous genes

Fetoprotein transcription factor (FTF) (ID: T04754) of *Lrh1* was the only TF identified based on TRANSFAC database.

Meanwhile, the promoter-binding factor of *Lrh1* was *CYP7A* (Cholesterol 7 α -hydroxylase).


SNPs of *Lrh1* homologous genes

The annotation of SNPs of *Lrh1* homologous genes showed that the majority of SNPs were located in intron and upstream, accounting for nearly 92% of all SNPs (Tables 2 and 3). Three SNPs of *Lrh1* were distributed in intron. Meanwhile, total 1819 SNPs of *Phkb* were located in the intron and 1343 SNPs were located in the upstream region of *Phkb*.

Discussion

In the present study, combined with RNA-seq data of *Lrh1*-knockout pancreas samples, FTF was the only TF of *Lrh1* identified based on TRANSFAC database and may regulate cholesterol catabolism into bile acids by activation of the promoter-binding factor *CYP7A*. Many literatures have elucidated the function of *Lrh1/Nr5a2/FTF/CYP7A* via experimental studies [21-25].

FTF is highly expressed in the liver and intestine and is implicated in the regulation of cholesterol, bile acid and steroid hormone homeostasis [26]. Nearly 50% of the body cholesterol is catabolized to bile acids via bile acid biosynthetic pathway, of which cholic acid (hydroxylated at position 12) and chenodeoxycholic acid are the major primary bile acids and play an important






Figure 2 KEGG pathway annotations of *Lrh1* and its homology genes.

role in cholesterol homeostasis [19]. Chenodeoxycholic acid can repress *FTF* expression and is a more potent suppressor of HMG-CoA reductase and cholesterol 7 α -hydroxylase/*CYP7A1* (7 α -hydroxylase) than cholic acid [27]. It has been proposed that *Lrh1*, also known as *CYP7A* promoter-binding factor, *LRH1*, or *FTF*, is required for the transcription of the 7 α -hydroxylase gene [19,28]. The small heterodimer partner 1 (SHP) of the nuclear bile acid receptor, FXR (farnesoid X receptor) can dimerize with FTF and diminish its activity on the 7 α -hydroxylase promoter [29].

Although *Lrh1* has been demonstrated the function in feedback regulation of *CYP7A1* expression as part of the FXR-SHP-LRH-1 cascade, in which bile acids can inhibit their own synthesis, the mechanisms have not been well understood. Out C et al. [25] have suggested that *CYP7A1* expression is increased rather than decreased under chow-fed conditions in *Lrh1*-knockdown mice that is coincided with a significant reduction in expression of intestinal Fgf15, a suppressor of *CYP7A1*. Besides, Noshiro M et al. [30] have suggested that the circadian rhythm of

CYP7A is regulated by multiple transcription factors, including DBP, REV-ERB α/β , LXR α , HNF4 α DEC2, E4BP4, and PPAR α . Hepatocyte nuclear factor 4 α (HNF4 α) and FTF are two major TFs driving *CYP7A1* promoter activity in lipid homeostasis. Bochkis IM et al. [31] have shown that prospero-related homeobox (Prox1) directly interacts with both HNF4 α and FTF and potently co-represses *CYP7A1* transcription.

In the present study, we annotated the SNPs of *Lrh1* and its homologous genes, showing that the majority was located in intron and upstream. Quiles Romagosa M \acute{A} [32] has reported that a functional SNP located in *Lrh1* promoter is related to Body Mass Index (BMI) and these SNPs might play important roles in the obese phenotype. However, previous researches mostly focused on SNPs associated with pancreatic cancer cell growth

Table 2 The number of different type of SNPs of *Lrh1* and its homology genes

Type	Number	Proportion
Downstream	581	0.052234
Intron	8251	0.741796
Non-synonymous-coding	122	0.010968
Splice-site-donor	2	0.00018
Start-gained	4	0.00036
Stop-gained	5	0.00045
Synonymous-coding	97	0.008721
Upstream	1975	0.17756
UTR-3-prime	79	0.007102
UTR-5-prime	7	0.000629

Table 3 The annotation of SNPs of *Lrh1* and its homology genes

Gene name	Chromo	Number	Position
Zfc3h1	10	4	115385510
Galc	12	36	98232394-98232429
Mylk4	13	13	32711300-32711312
Fzd5	1	22	64737712-64737722
Lrh1	1	6	136849713-136870578
Lamc2	1	923	153145967-153191429
Pofut1	2	188	153239719-153275253
Ache	5	19	137288432-137288449
Zfp282	6	1016	47876551-47913485
8-Mar	6	4231	116342541-116360897
Setd1a	7	1834	127778726-127802950
Phkb	8	3197	85837045-86008353
Atrx	X	7	105863283-105906922

and proliferation. For example, a previous genome-wide association study has identified five SNPs on 1q32.1 associated with pancreatic cancer that mapped to *Lrh1* gene and its up-stream regulatory region [33].

Conclusions

In conclusion, FTF combined with *CYP7A* might play an important role in *Lrh1* regulated pancreas-specific transcriptional network. Furthermore, the SNPs analysis of *Lrh1* and its homology genes provided the candidate mutant sites that might affect the *Lrh1*-related production and secretion of pancreatic fluid. These common susceptibility loci for *Lrh1* and its homologous genes needed follow-up studies.

Highlights

1. Total 15683 DEGs were identified, of which 23 was *Lrh1* homology genes (3 up-regulated and 20 down-regulated).
2. Fetoprotein TF was the only TF of *Lrh1* identified based on TRANSFAC database and the promoter-binding factor of fetoprotein TF was *CYP7A*.
3. The SNP annotations of *Lrh1* homologous genes showed that 92% of mutation sites were occurred in intron and upstream. Three SNPs of *Lrh1* were located in intron, while 1819 SNPs of *Phkb* were located in intron and 1343 SNPs were located in upstream region.

Competing interest

The authors declare that they have no competing interests.

Authors' contributions

MT, XM and CL participated in the design of this study, and they both performed the statistical analysis. RJ, GYH and YZ carried out the study, together with LQ, collected important background information, and drafted the manuscript. HL, XPW and ZS conceived of this study, and participated in the design and helped to draft the manuscript. All authors read and approved the final manuscript.

Acknowledgements

This study was supported by grants from the National Natural Science Foundation of China (No. 81200320, 81300350), Shanghai Science and Technology Commission (No. 11JC1410000), Fund of Shanghai Health Bureau (No. 20114315) and Training Plan of Excellent Academic Researcher of Shanghai Tenth People's Hospital (No. 12XSGG105, No. 04.01.13037).

Author details

¹Department of Gastroenterology, Shanghai Tenth People's Hospital, Tongji University School of Medicine, No.301, Yanchang Middle Road, Shanghai 200072, China. ²Department of Gastroenterology, Shanghai First People's Hospital Affiliated Shanghai Jiaotong University, Shanghai 200080, China.

Received: 20 December 2013 Accepted: 27 March 2014
Published: 15 April 2014

References

1. Leung PS: Physiology of the pancreas. In *The Renin-Angiotensin System: Current Research Progress in The Pancreas: The RAS in the Pancreas*, Volume 690. Netherlands: Springer; 2010:13–27.

2. Whitcomb DC, Lowe ME: Human pancreatic digestive enzymes. *Dig Dis Sci* 2007, **52**:1–17.
3. Fernandez-Marcos PJ, Auwerx J, Schoonjans K: Emerging actions of the nuclear receptor LRH-1 in the gut. *Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease* 2011, **1812**:947–955.
4. Rausa FM, Galarneau L, Bélanger L, Costa RH: The nuclear receptor fetoprotein transcription factor is coexpressed with its target gene HNF-3 β in the developing murine liver intestine and pancreas. *Mech Dev* 1999, **89**:185–188.
5. Baquie M, St-Onge L, Kerr-Conte J, Cobo-Vuilleumier N, Lorenzo PI, Moreno CMJ, Cederroth CR, Nef S, Borot S, Bosco D: The liver receptor homolog-1 (LRH-1) is expressed in human islets and protects β -cells against stress-induced apoptosis. *Hum Mol Genet* 2011, **20**:2823–2833.
6. Fayard E, Schoonjans K, Annicotte J-S, Auwerx J: Liver receptor homolog 1 controls the expression of carboxyl ester lipase. *J Biol Chem* 2003, **278**:35725–35731.
7. Hui DY, Howles PN: Carboxyl ester lipase structure-function relationship and physiological role in lipoprotein metabolism and atherosclerosis. *J Lipid Res* 2002, **43**:2017–2030.
8. Annicotte J-S, Fayard E, Swift GH, Selander L, Edlund H, Tanaka T, Kodama T, Schoonjans K, Auwerx J: Pancreatic-duodenal homeobox 1 regulates expression of liver receptor homolog 1 during pancreas development. *Mol Cell Biol* 2003, **23**:6713–6724.
9. Holmstrom SR, Deering T, Swift GH, Poelwijk FJ, Mangelsdorf DJ, Kiewer SA, MacDonald RJ: LRH-1 and PTF1-L coregulate an exocrine pancreas-specific transcriptional network for digestive function. *Genes Dev* 2011, **25**:1674–1679.
10. Trapnell C, Pachter L, Salzberg SL: TopHat: discovering splice junctions with RNA-Seq. *Bioinformatics* 2009, **25**:1105–1111.
11. Trapnell C, Roberts A, Goff L, Pertea G, Kim D, Kelley DR, Pimentel H, Salzberg SL, Rinn JL, Pachter L: Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat Protoc* 2012, **7**:562–578.
12. Robles JA, Qureshi SE, Stephen SJ, Wilson SR, Burden CJ, Taylor JM: Efficient experimental design and analysis strategies for the detection of differential expression using RNA-Sequencing. *BMC Genomics* 2012, **13**:484.
13. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ: Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res* 1997, **25**:3389–3402.
14. Crawford JE, Guelbeogo WM, Sanou A, Traoré A, Vernick KD, Sagnon NF, Lazzaro BP: De novo transcriptome sequencing in *Anopheles funestus* using Illumina RNA-seq technology. *PLoS One* 2010, **5**:e14202.
15. Da Wei Huang BTS, Lempicki RA: Systematic and integrative analysis of large gene lists using DAVID bioinformatics resources. *Nat Protoc* 2008, **4**:44–57.
16. Hulsege I, Kommadath A, Smits MA: Globaltest and GOEAST: Two different approaches for Gene Ontology Analysis. *BMC Proceedings* 2009, **4**:S10.
17. Matys V, Kel-Margoulis OV, Fricke E, Liebich I, Land S, Barre-Dirrie A, Reuter I, Chekmenev D, Krull M, Hornischer K: TRANSFAC® and its module TRANSCOMP®: transcriptional gene regulation in eukaryotes. *Nucleic Acids Res* 2006, **34**:D108–D110.
18. Langmead B, Trapnell C, Pop M, Salzberg SL: Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* 2009, **10**:R25.
19. Gerbod-Giannone M-C, del Castillo-Olivares A, Janciauskiene S, Gil G, Hylemon PB: Suppression of cholesterol 7 α -hydroxylase transcription and bile acid synthesis by an α 1-antitrypsin peptide via interaction with α 1-fetoprotein transcription factor. *J Biol Chem* 2002, **277**:42973–42980.
20. Cingolani P, Platts A, Coon M, Nguyen T, Wang L, Land SJ, Lu X, Ruden DM: A program for annotating and predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila melanogaster* strain w1118; iso-2; iso-3. *Fly* 2012, **6**:80–92.
21. Mouzat K, Baron S, Marceau G, Caira F, Sapin V, Volle DH, Lumbroso S, Lobaccaro JM: Emerging roles for LXRs and LRH-1 in female reproduction. *Molecular and cellular endocrinology* 2013, **368**:47–58.
22. Falender AE, Lanz R, Malenfant D, Belanger L, Richards JS: Differential expression of steroidogenic factor-1 and FTF/LRH-1 in the rodent ovary. *Endocrinology* 2003, **144**:3598–3610.
23. Xu Z, Ouyang L, Castillo-Olivares AD, Pandak WM, Gil G: Alpha(1)-Fetoprotein Transcription Factor (FTF)/Liver Receptor Homolog-1 (LRH-1) Is an Essential Lipogenic Regulator. *Biochimica et biophysica acta* 1801, **2010**:473–479.

24. del Castillo-Olivares A, Gil G: Alpha 1-fetoprotein transcription factor is required for the expression of sterol 12alpha -hydroxylase, the specific enzyme for cholic acid synthesis. Potential role in the bile acid-mediated regulation of gene transcription. *J Biol Chem* 2000, **275**:17793–17799.
25. Out C, Hageman J, Bloks WW, Gerrits H, Sollewijn Gelpke MD, Bos T, Havinga R, Smit MJ, Kuipers F, Groen AK: Liver receptor homolog-1 is critical for adequate up-regulation of Cyp7a1 gene transcription and bile salt synthesis during bile salt sequestration. *Hepatology* 2011, **53**:2075–2085.
26. Xu Z, Ouyang L, del Castillo-Olivares A, Pandak WM, Gil G: α -Fetoprotein transcription factor (FTF)/liver receptor homolog-1 (LRH-1) is an essential lipogenic regulator. *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids* 2010, **1801**:473–479.
27. Fakheri RJ, Javitt NB: Autoregulation of cholesterol synthesis: Physiologic and pathophysiologic consequences. *Steroids* 2011, **76**:211–215.
28. del Castillo-Olivares A, Gil G: Role of FXR and FTF in bile acid-mediated suppression of cholesterol 7 α -hydroxylase transcription. *Nucleic Acids Res* 2000, **28**:3587–3593.
29. Del Castillo-Olivares A, Campos JA, Pandak WM, Gil G: Role of FTF/LRH-1 on bile acid biosynthesis. A known nuclear receptor activator that can act as a suppressor of bile acid biosynthesis. *J Biol Chem* 2004, **279**(16):16813–16821.
30. Noshiro M, Usui E, Kawamoto T, Kubo H, Fujimoto K, Furukawa M, Honma S, Makishima M, Honma K-i, Kato Y: Multiple mechanisms regulate circadian expression of the gene for cholesterol 7 α -hydroxylase (Cyp7a), a key enzyme in hepatic bile acid biosynthesis. *J Biol Rhythms* 2007, **22**:299–311.
31. Bochkis IM, Schug J, Diana ZY, Kurinna S, Stratton SA, Barton MC, Kaestner KH: Genome-wide location analysis reveals distinct transcriptional circuitry by paralogous regulators Foxa1 and Foxa2. *PLoS genetics* 2012, **8**:e1002770.
32. Quiles Romagosa M \acute{A} : NR5A2: a regulator of glucose metabolism. 2011, <http://repositorio.unican.es/xmlui/bitstream/handle/10902/555/%5B2%5D%20Quiles%20Romagosa%20MA.pdf?sequence=1>.
33. Petersen GM, Amundadottir L, Fuchs CS, Kraft P, Stolzenberg-Solomon RZ, Jacobs KB, Arslan AA, Bueno-de-Mesquita HB, Gallinger S, Gross M: A genome-wide association study identifies pancreatic cancer susceptibility loci on chromosomes 13q22. 1, 1q32. 1 and 5p15. 33. *Nat Genet* 2010, **42**:224–228.

doi:10.1186/1471-2350-15-43

Cite this article as: Tang et al.: Identification of transcription factors and single nucleotide polymorphisms of *Lrh1* and its homologous genes in *Lrh1*-knockout pancreas of mice. *BMC Medical Genetics* 2014 **15**:43.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

