



Published in final edited form as:

Allergy. 2020 January ; 75(1): 217–220. doi:10.1111/all.13978.

Allergenomics of the tick *Ixodes ricinus* reveal important α -Gal-carrying IgE-binding proteins in red meat allergy

Danijela Apostolovic, PhD¹, Jelena Mihailovic, MSc², Scott P. Commins, MD, PhD, Ing.³, Michiel Wijneveld⁴, Maria Kazimirova, PhD⁵, Maria Starkhammar, MD⁶, Hannes Stockinger, PhD⁴, Thomas A. E. Platts-Mills, MD PhD⁷, Tanja Cirkovic Velickovic, PhD^{2,8,9}, Carl Hamsten, PhD^{1,*}, Marianne van Hage, MD, PhD^{1,*}

¹Division of Immunology and Allergy, Department of Medicine Solna, Karolinska Institutet and University Hospital, Stockholm, Sweden;

²University of Belgrade - Faculty of Chemistry, Department of Biochemistry, Center of Excellence for Molecular Food Sciences, Belgrade, Serbia;

³Department of Medicine, University of North Carolina School of Medicine, Chapel Hill, NC, USA;

⁴Institute for Hygiene and Applied Immunology, Center for Pathophysiology, Infectiology and Immunology, Medical University of Vienna, Vienna, Austria;

⁵Institute of Zoology, Slovak Academy of Sciences, Bratislava, Slovakia;

⁶Department of Internal Medicine, Södersjukhuset, Stockholm, Sweden;

⁷Asthma and Allergic Diseases Center, University of Virginia Health System, Charlottesville, VA, USA;

⁸Ghent University Global Campus, Yeonsugu, Incheon, South Korea;

⁹Ghent University, Faculty of Bioscience Engineering, Ghent, Belgium.

Keywords

allergenomics; red meat allergy; saliva; ticks; α -Gal

To the Editor

Red meat allergy known as mammalian meat allergy, caused by IgE antibodies against galactose- α -1,3-galactose (α -Gal), is nowadays recognized worldwide and strong associations with tick bites have been identified for different tick species and geographic

This work is licensed under a [Creative Commons - Attribution-Noncommercial-No Derivative Works 3.0 Serbia](https://creativecommons.org/licenses/by-nc-nd/3.0/deed.en)

Correspondence to: Marianne van Hage, MD, PhD, Karolinska Institutet, Department of Medicine Solna, Division of Immunology and Allergy, Karolinska University Hospital Solna L2:04, SE - 171 76 Stockholm, Sweden, Tel +46-8-5177 5942, Fax +46-8-33 57 24, marianne.van.hage@ki.se.

*Shared authorship
Author's contribution

DA participated in all stages of the project, performed the experiments and interpreted the data. JM and TCV were involved in the proteomics analysis. MS, SC, TPM provided the patient material. MW, MK and HS provided tick saliva material. DA, CH and MvH wrote the manuscript. All authors provided critical review of the manuscript.

locations.¹ Time relationship between tick exposure and increased IgE levels to α -Gal has further supported the strong evidence that tick bites are the primary cause of the IgE antibodies.² All developmental stages of ticks can bite humans and in the US, high IgE levels to α -Gal following bites from larvae have been reported.³ While we have previously demonstrated the presence of α -Gal in the gut of the European tick *Ixodes ricinus*⁴, α -Gal-containing proteins in tick saliva from the South American, Japanese and European ticks, *A. sculptum*⁵, *Haemaphysalis longicornis*⁶ and *Hyalomma marginatum*⁷, have recently been reported. However, the α -Gal-content of the *I. ricinus* proteome has not been investigated yet. Here we used allergenomics⁸ and shotgun proteomics approaches to identify IgE-binding α -Gal carrying proteins in adult and larval stages as well as in saliva of *I. ricinus* ticks. Allergen-specific IgE antibody responses were assessed by ImmunoCAP (Thermo Fisher Scientific, Uppsala, Sweden) among 32 Swedish and 18 US mammalian meat allergic patients to reveal differences in IgE reactivity between two geographical population groups. IgE- and α -Gal binding capacity, as well allergenicity of *I. ricinus* tick proteins were evaluated (details on methods and patients characteristics are presented in the Supporting information).

Our data revealed that nearly all Swedish and American meat allergic patients had IgE responses against both the larval and adult stages of *I. ricinus* and we noted that the four individuals IgE negative to *I. ricinus* had comparably low IgE levels to α -Gal (Table S1). Similarly to previous reports, moderate to high correlations between IgE to α -Gal and adult ticks were seen in the two patient groups (Figure S1).⁴ Furthermore, the strong correlations between IgE reactivity to adult and larvae in both patient groups suggest that growth stages of *I. ricinus* ticks seems to be of less importance for the IgE recognition (Figure S1). In western-blot, IgE-reactive protein bands in the molecular weight range of 25–150 kDa in *I. ricinus* ticks were noted with comparable results for both tick stages using Swedish and American serum pools (Figure 1A and B respectively). To investigate whether the observed IgE reactivities were α -Gal related, the pools were pre-incubated with the α -Gal carrying glycoprotein bovine thyroglobulin prior to immunoblot analyses. The IgE binding to proteins from both adult and larval ticks was strongly diminished in Swedish as well as in US serum pools, revealing α -Gal as the main IgE target (Fig 1A and B lanes with α -Gal+) in both populations. Interestingly, proteins at the similar size were identified in *I. ricinus* saliva that bound red meat allergic patients' IgE and were recognized by the anti- α -Gal antibody (Figure 1C). Similar results have been reported for the analysis of saliva from the Japanese tick *H. longicornis*.⁶ In addition, we investigated the IgE binding capacity with IgE inhibition ELISA where protein extract from *I. ricinus* was able to inhibit IgE binding to HSA- α -Gal by up to 77% (Fig S2). We also evaluated the allergenic potential of *I. ricinus* using blood from 14 mammalian meat allergic patients and noted that adult ticks induced basophil activation in 13 patients (Figure 1D and Figure S3). The allergenic activity towards ticks was higher compared with HSA- α -Gal, however HSA- α -Gal showed to be more sensitive. This shows that *I. ricinus* protein epitopes also are of importance in basophil activation, which is in line with the IgE-binding capacity results (immunoblots and ELISA). When blood samples from four patients were stimulated with extract from *I. ricinus* larvae (Fig 1E), the allergenic activity was found to be similar to adult *I. ricinus* protein extract in three patients. None of the antigens activated basophils in two non-allergic individuals

(Figure S3), indicating that the observed reactions were IgE-dependent. Basophil activation with adult *I. ricinus* protein extract was dose-dependent reaching 72.7% of CD63-positive cells (median, 34.4%; range, 7.5% to 72.7%, at concentration 50µg/ml) giving a sensitivity of 93%. Furthermore, a strong correlation between %CD63-positive basophils for adult *I. ricinus* protein extract and HSA-α-Gal (Fig S4) was noted, pointing out the dominant role of the α-Gal epitope in activating red meat allergic patients' basophils.

We used an allergenomics approach with 2D PAGE and 2D immunoblots together with mass spectrometry to identify α-Gal-carrying IgE binding proteins in adults and larvae ticks (Fig S5 and Fig S6, for details please see supporting information). Analysis of the obtained MS/MS spectra gave high identification scores to 43 protein accession numbers for adult and 37 for larvae from the *Ixodida* order (Table S2), grouped into six protein groups: vitellogenins, SERPIN, actin, α-2-macroglobulin, chitinase like-lectins and transport or channel forming proteins (Table 1). Comparing data from the 2D immunoblots with IgE binding (Fig S5A–C), and the anti-α-Gal antibody (Fig S5D) protein spots in the range of 75–100 kDa were shown to contain α-Gal carrying proteins. These proteins belonged to the vitellogenins and α-2-macroglobulin protein groups (Table 1). The other protein groups (actin, SERPIN, chitinase-like lectins and transport forming proteins) seem not to carry α-Gal. However, since these proteins possess IgE-binding properties and possibly allergenic activity, as shown in the basophil activation test with the higher response to *I. ricinus* extract compared to HSA-α-Gal, point to the fact that they most likely play a role in immune responses against ticks. Vitellogenins (isoforms 1–4), which was the major α-Gal-carrying IgE binding protein group, are produced in mid gut cells, fat bodies and salivary glands of ticks and are present in glycosylated and non-glycosylated forms. These carbohydrate-binding proteins have major function in tick reproduction.⁹ The recognition of similar proteins in the 1D immunoblotting of saliva and identification of vitellogenins by mass spectrometry with high score (Table S3) indicated that they are abundantly present in tick saliva as well. Recently, Cabezas-Cruz and colleagues reported the expression of galactosyltransferase in *Ixodes scapularis*, an enzyme needed for the production of α-Gal.¹⁰ Thus, taken together these data support that ticks produce the α-Gal epitope. The α-Gal-containing vitellogenin and α-2-macroglobulin are probably delivered into the host's skin by tick bites and could presumably be involved in the induction of an anti-α-Gal immune response.

In conclusion, our results give new insight into mammalian meat allergy. Firstly, proteins from adults and larvae of the European tick *I. ricinus* are recognized by IgE from meat allergic patients and have allergenic activity, which are α-Gal and tick protein specific. In addition, larvae were not fed on the host which supports that α-Gal carrying proteins originate from ticks. Moreover, *I. ricinus* tick saliva contains IgE binding α-Gal carrying proteins which by allergenomics revealed to be vitellogenins. The results support the strong relationship with tick bites for the development of mammalian meat allergy. Thus, red meat allergic patients should be advised to avoid further tick bites.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgment

We thank Neda Bigdeli for excellent technical assistance. Funding: The Swedish Research Council, The Stockholm County Council (ALF project), The Swedish Asthma and Allergy Association's Research Foundation, The King Gustaf V 80th Birthday Foundation, The Swedish Heart-Lung Foundation, The Hesselman Foundation, The Konsul Th C Bergh Foundation, The Swedish Cancer and Allergy Foundation, The Magnus Bergvall Foundation, the Swedish Association for Allergology, the European Union's Horizon 2020 Research and Innovation Program - ID-LYME GA No.720480, the European Union's Horizon 2020 FoodEnTwin project, GA No.810752, and Ministry of Education Science and Technological Development Republic of Serbia GA No. 172024. The EC does not share responsibility for the content of the article.

Conflict of interest statements:

S.P. Commins has received support from Genentech as speaker's bureau and from UptoDate honorarium for topic author. The rest of the authors declare that they have no relevant conflicts of interest.

References

1. Hilger C, Fischer J, Wolbing F, Biedermann T. Role and Mechanism of Galactose-Alpha-1,3-Galactose in the Elicitation of Delayed Anaphylactic Reactions to Red Meat. *Curr Allergy Asthm R*. 2019;19(1).
2. Commins SP, James HR, Kelly LA, et al. The relevance of tick bites to the production of IgE antibodies to the mammalian oligosaccharide galactose-alpha-1,3-galactose. *J Allergy Clin Immun*. 2011;127(5):1286–U1310. [PubMed: 21453959]
3. Platts-Mills TA, Schuyler AJ, Tripathi A, Commins SP. Anaphylaxis to the carbohydrate side chain alpha-gal. *Immunol Allergy Clin North Am*. 2015;35(2):247–260. [PubMed: 25841549]
4. Hamsten C, Starkhammar M, Tran TA, et al. Identification of galactose-alpha-1,3-galactose in the gastrointestinal tract of the tick *Ixodes ricinus*; possible relationship with red meat allergy. *Allergy*. 2013;68(4):549–552. [PubMed: 23414348]
5. Araujo RN, Franco PF, Rodrigues H, et al. Amblyomma sculptum tick saliva: alpha-Gal identification, antibody response and possible association with red meat allergy in Brazil. *Int J Parasitol*. 2016;46(3):213–220. [PubMed: 26812026]
6. Chinuki Y, Ishiwata K, Yamaji K, Takahashi H, Morita E. Haemaphysalis longicornis tick bites are a possible cause of red meat allergy in Japan. *Allergy*. 2016;71(3):421–425. [PubMed: 26551325]
7. Mateos-Hernandez L, Villar M, Moral A, et al. Tick-host conflict: immunoglobulin E antibodies to tick proteins in patients with anaphylaxis to tick bite. *Oncotarget*. 2017;8(13):20630–20644. [PubMed: 28423486]
8. Di Girolamo F, Muraca M, Mazzina O, Lante I, Dahdah L. Proteomic applications in food allergy: food allergenomics. *Curr Opin Allergy Clin Immunol*. 2015;15(3):259–266. [PubMed: 25899690]
9. Donohue KV, Khalil SMS, Mitchell RD, Sonenshine DE, Roe RM. Molecular characterization of the major hemelipoglycoprotein in ixodid ticks. *Insect Mol Biol*. 2008;17(3):197–208. [PubMed: 18477238]
10. Cabezas-Cruz A, Espinosa PJ, Alberdi P, et al. Tick galactosyltransferases are involved in alpha-Gal synthesis and play a role during Anaplasma phagocytophilum infection and Ixodes scapularis tick vector development. *Sci Rep-Uk*. 2018;8.




Fig. 1. IgE-binding activity and allergenic activity of *I. ricinus* proteins from adult and larvae in mammalian meat allergic patients. A) IgE immunoblot with the Swedish serum pool with (α-Gal+) and without inhibition with bovine thyroglobulin (α-Gal-) and B) IgE immunoblot with the US serum pool with (α-Gal+) and without inhibition with bovine thyroglobulin (α-Gal-); Lane 1- protein extract from adult *I. ricinus* ticks; Lane 2 - protein extract from larvae *I. ricinus* ticks; C) IgE binding to tick saliva proteins Lane i) with the Swedish serum pool Lane ii) and after inhibition with bovine thyroglobulin; Lane iii) with monoclonal anti-α-Gal binding; M-Molecular markers. D) Allergenic activity of 14 Swedish meat allergic patients on HSA-α-Gal and *I. ricinus* adult E) Allergenic activity of four Swedish meat allergic patient on *I. ricinus* adult and larvae

Table 1 -Identified IgE and α -Gal binding proteins in *Ixodes ricinus* ticks

Protein group	IgE and α -Gal binding on 2D PAGE			Spots from 2D
	<i>I. ricinus</i> adults	<i>I. ricinus</i> larvae	α -Gal	
Vitellogenins	+	+	+	1-4,6-7,9-15
α -2-macroglobulin	-	+	+	13
SERPIN	+	+	-	8,16
Actin	+	+	-	8,16
Transport or channel forming proteins	+	-	-	5,8
Chitinase-like lectins	+	-	-	1,2,7

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript