

## EDGE ARTICLE

Cite this: *Chem. Sci.*, 2022, 13, 13851

All publication charges for this article have been paid for by the Royal Society of Chemistry

Received 29th September 2022

Accepted 4th November 2022

DOI: 10.1039/d2sc05423j

rsc.li/chemical-science

## An electrochemical multicomponent reaction toward C–H tetrazolization of alkyl arenes and vicinal azidotetrazolization of alkenes†

Yi Yu, Xiao-Bin Zhu, Yaofeng Yuan and Ke-Yin Ye \*

The widespread use of tetrazoles in medicine, biology, and materials science continuously promotes the development of their efficient and selective syntheses. Despite the prosperous development of multicomponent reactions, the use of the most abundant and inexpensive chemical feedstocks, *i.e.*, alkanes and alkenes, toward the preparation of diverse tetrazoles remains elusive. Herein, we developed an electrochemical multicomponent reaction (e-MCR) for highly efficient and selective C–H tetrazolization of alkyl arenes. When applied to alkenes, the corresponding vicinal azidotetrazoles were readily obtained, which were further demonstrated to be versatile building blocks and potential high-energy materials.

Tetrazoles constitute an important pharmacophore in medicinal chemistry thanks to their bioisosterism to carboxylic acid, furan, and amide moieties, as well as other beneficial physicochemical properties.<sup>1</sup> In addition, their nitrogen-rich character also makes tetrazoles one of the most popular explosives in high-energy materials.<sup>2</sup> Therefore, the development of efficient, straightforward, and ideally modular synthetic approaches toward diverse novel tetrazole scaffolds would significantly accelerate the relevant research in pharmaceutical, biology, and materials science.


The last few decades have witnessed tremendous advances in the highly efficient and selective C(sp<sup>3</sup>)-H functionalization of saturated hydrocarbons. Specifically, oxidative cross-dehydrogenative coupling reactions<sup>3</sup> between alkyl arenes and tetrazoles have been intensively investigated.<sup>4</sup> We reasoned that a multicomponent reaction (MCR), when combined with state-of-the-art C(sp<sup>3</sup>)-H functionalization, is an appealing approach to quickly construct complex tetrazoles in a highly modular manner ideally starting from the readily available chemical feedstock.<sup>5</sup> However, direct use of the most abundant and inexpensive unsaturated hydrocarbons in the MCR reaction for the synthesis of tetrazoles remains elusive.

Electrosynthesis provides new opportunities for the functionalization of C(sp<sup>3</sup>)-H bonds<sup>6</sup> owing to its unique capability to tune reactivity and selectivity *via* the precise control of applied potentials.<sup>7</sup> Therefore, we envisaged that an electrochemical multicomponent reaction (e-MCR) of saturated


hydrocarbons, nitriles, and azides, may constitute a highly straightforward and modular approach for the preparation of diverse tetrazoles (Scheme 1A, left).

However, the realization of such a hypothesis needs to surpass several competing reaction pathways of alkyl arenes, *i.e.*, electrochemical C–H acetamidation<sup>8</sup> with acetonitrile (Scheme 1B), and C–H azidation<sup>9</sup> with azide (Scheme 1C), respectively. In addition, we aim to develop an operationally simple and practical e-MCR, which would mitigate the sophisticated reaction setup as well as the requisite presence of irradiation and a transition-metal catalyst or organocatalyst in the realm of electrophotocatalysis.<sup>10</sup>


Herein, we report our development of a highly efficient e-MCR for the facile and modular synthesis of tetrazoles,

(A) Electrochemical MCR toward tetrazoles of alkanes and alkenes (*this work*)

(B) Known competing reactivity (I): C–H acetamidation (Lambert)



(C) Known competing reactivity (II): C–H azidation (Lei, Ackermann)

Scheme 1 Tetrazoles *via* electrochemical multicomponent reactions.

Key Laboratory of Molecule Synthesis and Function Discovery (Fujian Province University), College of Chemistry, Fuzhou University, Fuzhou 350108, China. E-mail: kyye@fzu.edu.cn

† Electronic supplementary information (ESI) available. See DOI: <https://doi.org/10.1039/d2sc05423j>




directly using readily available alkanes, acetonitrile, and azidotrimethylsilane. In addition, the same strategy could be further applied to electrochemical difunctionalization of alkenes,<sup>11</sup> from which the thus obtained vicinal azidotetrazoles (Scheme 1A, right) were further demonstrated to be versatile building blocks as well as potential high-energy materials.


After extensive optimization of reaction conditions (see the ESI†), we found that the constant-current electrolysis ( $I = 75$  mA) of ethylbenzene (**1**) and azidotrimethylsilane (**2**) in an undivided cell, with a graphite anode and a platinum cathode, readily afforded the C–H tetrazolization product (**3**) in 68% yield (Scheme 2). The use of excess amounts of  $\text{TMSN}_3$  was to compromise its unproductive consumption of  $\text{N}_2$  during electrolysis.<sup>9a</sup> Note that while a relatively low current electrolysis ( $I = 20$  mA) was also amenable to achieving this reactivity, the high current density further rendered this reaction complete within 20 min, providing a fast and straightforward approach to access the tetrazole moiety directly from alkyl arenes. Particularly, the addition of HFIP (hexafluoroisopropanol) was pivotal to achieving the anticipated reactivity.<sup>12</sup>

Regarding the possible mechanism, ethylbenzene (**1**) is likely to first undergo single-electron oxidation and then deprotonation, *via* a well-established phenyl radical cation intermediate, to generate a benzyl radical (**A**), which is followed by second electrochemical oxidation to generate the benzyl carbocation (**B**).<sup>13</sup> The nucleophilic interception by MeCN then forms a Ritter-type nitrilium ion intermediate (**C**),<sup>8</sup> which finally reacts with azide to afford the C–H tetrazolization product (**3**).<sup>14</sup> Another possible route for the formation of the benzyl radical (**A**), *i.e.*, *via* the hydrogen-atom transfer (HAT) of the benzylic C–H bond by the electrochemically generated azido radical, is less likely according to our mechanistic investigations (*vide infra*).

The substrate scope of this electrochemical C–H tetrazolization of alkyl arenes was then investigated (Scheme 3). This reaction tolerated a multitude of substituted ethylbenzenes (**4–6**), toluene (**7**), and other alkyl-substituted benzenes (**8, 9**). Diverse functionalities, such as ketones (**10**), sulfonyl benzamide (**11**), trifluoroacetamide (**12**), and succinimide (**13**) well survived under the current reaction settings. Despite their high propensity toward nucleophilic substitution by  $\text{TMSN}_3$ , the aliphatic chloride (**14**) and protected alcohols (**15, 16**) remained intact. In addition, functionalized cyclic alkyl arenes (**17, 18**), and thiophene derivatives (**19**) also proceeded with the




Scheme 2 Electrochemical C–H tetrazolization of ethylbenzene.



Scheme 3 Substrate scope of the electrochemical C–H tetrazolization.<sup>a</sup> <sup>a</sup>Conditions: undivided cell, graphite rod anode ( $\phi$  6 mm, about  $20 \pm 1$  mm immersion depth in solution), Pt cathode, alkyl arenes (0.3 mmol),  $\text{TMSN}_3$  (0.51 mmol, 1.7 equiv.),  $n\text{Bu}_4\text{NClO}_4$  (0.3 mmol), MeCN/HFIP = 10 : 1, 75 mA, 20 min, ( $F = 3.11$  F  $\text{mol}^{-1}$ ) at room temperature under an  $\text{N}_2$  atmosphere, isolated yield; <sup>b</sup> $\text{LiClO}_4$  (0.3 mmol), 75 mA, 23 min ( $F = 3.58$  F  $\text{mol}^{-1}$ ); <sup>c</sup>20 mA, 90 min ( $F = 3.73$  F  $\text{mol}^{-1}$ ).

anticipated electrochemical C–H tetrazolization smoothly. The current electrochemical C–H tetrazolization could be also extended to the late-stage functionalization of a variety of alkyl arenes derived from biologically relevant compounds and pharmaceuticals, such as FKGK 11 analogue (**20**), ibuprofen (**21**), celestolide (**22**), and retinoic acid receptor agonist analogue (**23**). Unfortunately, acetonitrile is currently the only amenable organonitrile in this electrochemical C–H tetrazolization.


In addition, the e-MCR strategy is readily applicable to the vicinal difunctionalization of alkenes. While the constant-current electrolysis only afforded a moderate yield of the desired product, we found the constant-cell potential electrolysis ( $E_{\text{cell}} = 2.8$  V) of styrene (**24**) and  $\text{TMSN}_3$  (**2**) in a MeCN/HFIP mixed solvent resulted in the formation of a vicinal azidotetrazole in a higher yield (**25**, 68%, Table 1, entry 1). In this case, the graphite plate anode afforded a slightly higher yield than that of the graphite rod as previously established.<sup>15</sup> Again, the choice of HFIP as the co-solvent was crucial since the concomitant formation of the competing azidoacetamide (**26**) occurred when HOAc or  $\text{H}_2\text{O}$  was used instead (entries 2 and 3). The electrolyte is another important parameter to tune product selectivity. For instance, though the use of  $n\text{Bu}_4\text{NBF}_4$  afforded both azidotetrazole (**25**) and azidoacetamide (**26**) almost unselectively (entry 4),  $n\text{Bu}_4\text{NH}_2\text{SO}_4$  completely suppressed the azidotetrazolization but promoted the azidoacetamidation albeit in a low yield (entry 5).

Table 1 Reaction optimization of the vicinal azidotetrazoliation<sup>a</sup>

Entry	Electrolyte	Co-solvent	25 (%)	26 (%)
1	<sup>n</sup> Bu <sub>4</sub> NClO <sub>4</sub>	HFIP	68 <sup>b</sup>	0
2	<sup>n</sup> Bu <sub>4</sub> NClO <sub>4</sub>	HOAc	28	13
3	<sup>n</sup> Bu <sub>4</sub> NClO <sub>4</sub>	H <sub>2</sub> O	21	12
4	<sup>n</sup> Bu <sub>4</sub> NBF <sub>4</sub>	H <sub>2</sub> O	15	18
5	<sup>n</sup> Bu <sub>4</sub> NHSO <sub>4</sub>	H <sub>2</sub> O	0	31

<sup>a</sup> Reaction conditions: undivided cell, graphite plate anode, Pt cathode, 24 (0.3 mmol), TMSN<sub>3</sub> (0.75 mmol, 2.5 equiv.), electrolyte (0.3 mmol), MeCN/co-solvent = 12.5:1, at room temperature under an N<sub>2</sub> atmosphere, the yield was determined by <sup>1</sup>H NMR with 1,3,5-trimethoxybenzene as the internal standard. <sup>b</sup> Isolated yield.

We then investigated the substrate scope of the electrochemical vicinal azidotetrazoliation. As illustrated in Scheme 4, a wide array of substituted styrenes were well tolerated to afford the anticipated vicinal azidotetrazoles in moderate to good yields (27–31). In addition,  $\alpha$ -alkyl-substituted styrenes generally served as good substrates for the preparation of vicinal azidotetrazoles (32–37). The 1,2-disubstituted linear alkenes



**Scheme 4** Substrate scope of the electrochemical vicinal azidotetrazoliation. <sup>a</sup> Reaction conditions: undivided cell, graphite plate anode, Pt cathode, alkene (0.3 mmol), 2 (0.75 mmol, 2.5 equiv.), <sup>n</sup>Bu<sub>4</sub>NClO<sub>4</sub> (0.3 mmol), MeCN/HFIP = 12.5:1, at room temperature under an N<sub>2</sub> atmosphere, isolated yield; <sup>b</sup> 2.5 V, 4.5 h; <sup>c</sup> MeCN/DCM/HFIP = (8.75:3.75:1) as the co-solvent.

(38, 39) proceeded with the anticipated vicinal azidotetrazoliation smoothly, too. Notably, the sterically hindered trisubstituted (40, 41) and tetrasubstituted alkenes (42), as well as heteroaryl alkenes (43, 44) were all amenable substrates. Besides acetonitrile, other aliphatic nitriles, such as isobutyronitrile (45) and butyronitrile (46), were also positively engaged in the electrochemical syntheses of vicinal azidotetrazoles. Unfortunately, the current protocol does not tolerate benzonitrile, probably owing to its relatively weak nucleophilicity. Again, this protocol can be successfully applied to late-stage vicinal azidotetrazoliation of complex alkenes derived from drug derivatives, including probenecid (47), fenofibric acid (48), and febuxostat (49).

Without electrical input, this e-MCR reactivity could not be replicated by using other common strong oxidants (Table 2). For the C–H tetrazoliation (4), low conversions were generally observed and a substantial amount of starting materials were recovered. Similarly, no desired vicinal azidotetrazoliation product (27) could be detected even though high conversions were realized. Instead, only the formation of a trace amount of aldehyde was detected. The direct comparison between electro-synthesis and chemical synthesis in the formation of targeted tetrazoles further substantiates the unique potential of electrochemistry in the expansion of novel chemical space and reactivity.

The nitrogen-rich character of the obtained vicinal azidotetrazoles raises safety concerns about their thermal stability since both the azido and tetrazole moieties are frequently present as explosives in high-energy materials.<sup>16</sup> Differential scanning calorimetry (DSC) and thermogravimetric (TG) analyses were then performed to elucidate their thermal sensitivity (Fig. 1).<sup>17</sup> Indeed, the azidotetrazole (25) bears a high value of decomposition enthalpy ( $-1270$  J g<sup>-1</sup> or  $-291$  kJ mol<sup>-1</sup>), implying that special attention should be paid to its

Table 2 Attempts to replace electricity with chemical oxidants<sup>a</sup>

Reaction scheme showing the attempted synthesis of vicinal azidotetrazoles (4 and 27) using chemical oxidants instead of electricity. Conditions: MeCN, TMSN<sub>3</sub>.

Entry	Oxidant	4, yield <sup>a</sup> (conv., %)	27, yield <sup>a</sup> (conv., %)
1	Anode	68 (>95)	70 (>95)
2	NFSI	n.d. (11)	n.d. (51)
3	TBHP	n.d. (20)	n.d. (73)
4	CAN	n.d. (36)	n.d. (35)
5	<i>m</i> -CPBA	n.d. (14)	n.d. (22)
6	DDQ	n.d. (15)	n.d. (91)
7	PhI(OAc) <sub>2</sub>	n.d. (30)	n.d. (>95)
8	Mn(OAc) <sub>3</sub>	n.d. (69)	n.d. (>95)
9	AgNO <sub>3</sub>	n.d. (>95)	n.d. (>95)

<sup>a</sup> Yield and conversion were determined by NMR of the crude reaction mixture with PhOCF<sub>3</sub> or CH<sub>2</sub>Br<sub>2</sub> as the internal standard; n.d. = not detected.





Fig. 1 DSC (left) and TG (right) curves of the azidotetrazole **25**. (a) Maximum temperature in the DSC curve; (b) temperature at which the time of maximum rate is 24 h.

scaled preparation and usage. However, the high onset temperature value (233 °C) also suggests that this azidotetrazole is relatively less sensitive to heat and would release its energy only at high temperatures. This is further substantiated by its  $TD_{24}$  data, which typically would serve as a consecutive prediction of a reasonable safety margin.<sup>18</sup> In addition, the TG analyses indicated that the major decomposition mechanism of this azidotetrazole is the thermal extrusion of the tetrazole moiety at  $T_m$ , probably initiated *via* the loss of molecular nitrogen (Fig. 1, right).<sup>19</sup>

The azido group of these vicinal azidotetrazoles provides a good platform for their further synthetic elaborations (Scheme 5). For instance, compounds vicinally functionalized both by a tetrazole and a triazole were conveniently prepared *via* click reactions between the azidotetrazoles and alkynes, such as phenylacetylene (**50**), and 5-ethynyl-2'-deoxyuridine (**51**). When treated with trimethyl phosphite, the azido moiety was readily transformed into its corresponding phosphoramidate (**52**). Finally, a base-promoted dehydroazidation of the vicinal azidotetrazole afforded the vinyltetrazole (**53**) in a quantitative yield. Note that the tetrazole moiety remained intact during all these derivatizations.

To gain some more insights into the reaction mechanism, several mechanistic experiments were performed. Cyclic voltammetry studies indicated that while the oxidation potential of ethyl benzene was rather high ( $E_{p/2} = 2.43$  V vs. Ag/AgCl), the addition of HFIP dramatically facilitated its oxidation ( $E_{p/2} = 2.01$  V vs. Ag/AgCl). This positive effect of HFIP was attributed to its unique capability to stabilize radical cation intermediates in



Scheme 5 Derivatization of azidotetrazoles. Reaction conditions: (a) phenylacetylene (3 equiv.), CuI (30 mol%), THF, r.t., 4 h; (b) 5-ethynyl-2'-deoxyuridine (3 equiv.), CuI (30 mol%), THF, 60 °C, 5 h; (c) P(OMe)<sub>3</sub> (1.3 equiv.), toluene, 80 °C, 3 h; (d) <sup>t</sup>BuOK (2 equiv.), THF, r.t., 1 min.

the anodic oxidation of the substrate (Fig. 2A).<sup>12</sup> As illustrated in Fig. 2B, the relatively low oxidation potential of TMSN<sub>3</sub> ( $E_{p/2} = 0.92$  V vs. Ag/AgCl) should favor its prior oxidation to ethyl benzene or styrene ( $E_{p/2} = 2.21$  V vs. Ag/AgCl). Indeed, the involvement of an azido radical was supported by the radical clock experiment (Fig. 2C).

In the electrochemical C–H tetrazolation, the Hammett studies ( $\rho = -2.97$ ) of various *p*-substituted ethyl benzenes exhibiting a much more negative  $\rho$  than that of a typical HAT process ( $\rho > -1$ ),<sup>20a</sup> revealed that a positive charge was built up in the transition state of the rate-limiting step (Fig. 2D).<sup>20b</sup> The possible formation of the benzyl radical *via* HAT was further disapproved by the fact that benzylic C–H bonds of similar bond dissociation energies but different electronic properties were still well distinguished (**21'**, Fig. 2E).<sup>21</sup> Parallel to the observations in Xu's photoelectrochemical C–H cyanation,<sup>22</sup> benzyltrimethylsilane (**56**) proceeded *via* the cleavage of the C–Si bond to afford tetrazole (**7**) exclusively, which again disproved the one-step HAT process but favored a radical cation mechanism (Fig. 2F). Taken together, the formation of the benzyl radical from ethylbenzene is more likely *via* a stepwise electron transfer/proton transfer process but not HAT.<sup>22,23</sup> The





Fig. 2 Mechanistic experiments.



Scheme 6 Proposed mechanism.

involvement of a carbocation intermediate was further substantiated by the formation of the Ritter-type product in the absence of  $TMSN_3$  (Fig. 2G).

A proposed mechanistic rationale for the vicinal azidotetrazolization reaction is shown in Scheme 6. Similar to the C–H tetrazolization of alkyl arenes, the  $\beta$ -azido carbocation (II), generated *via* the sequential addition of the azido radical ( $N_3^\bullet$ ) and single-electron oxidation, is the key intermediate in the azidotetrazolization reaction (IV, path a).<sup>24</sup> Notably, the competing diazidation (III)<sup>25</sup> is disfavoured since a radical pathway requires the cross-coupling of two transient radicals (path b).<sup>26</sup> On the other hand, a polar pathway to the diazidation product (path c) is mitigated owing to the much excess of MeCN as the carbocation trapping agent on the anode surface, out-competing the low-concentrated  $TMSN_3$  because of its easy oxidation character (see the ESI†). This rationale of product selectivity is also applicable to the electrochemical C–H tetrazolization since both reactions share similar Ritter-type nitrilium ion intermediates toward the formation of the tetrazole moiety.

In summary, we have developed an electrochemical multi-component reaction for the modular assembly of diverse tetrazoles from readily available alkanes, acetonitrile, and azido-trimethylsilane. The fine-tuning of electrolytic conditions detoured other competing pathways but afforded the anticipated benzylic tetrazoles highly selectively. When applied to alkenes, the corresponding vicinal azidotetrazoles were readily obtained, which have been demonstrated to be versatile building blocks and potential high-energy materials.<sup>27</sup> The current work indicates that the e-MCR might be broadly applicable for the facile syntheses of diverse nitrogen-containing heterocycles.

## Data availability

Detailed synthetic procedures and complete characterization data for all new compounds can be found in the ESI.†

## Author contributions

K. Y. Y. conceived the idea and supervised the project. Y. Y. conducted the majority of the experimental work. K. Y. Y., Y. Y., X. B. Z., and Y. F. Y. analyzed the data and contributed to the preparation of this manuscript.

## Conflicts of interest

There are no conflicts to declare.

## Acknowledgements

Financial support from the National Natural Science Foundation of China (22171046, 21901041, and 22071025), State Key Laboratory of Physical Chemistry of Solid Surfaces, Xiamen University (202008), Hundred-Talent Project of Fujian (50012742), and Fuzhou University (510841) is gratefully acknowledged.

## Notes and references

- (a) L. V. Myznikov, A. Hrabalek and G. I. Koldobskii, *Chem. Heterocycl. Compd.*, 2007, **43**, 1–9; (b) N. Dhiman, K. Kaur and V. Jaitak, *Bioorg. Med. Chem.*, 2020, **28**, 115599; (c) P. Patowary, B. Deka and D. Bharali, *Malar. Control Elimin.*, 2021, **10**, 167.
- (a) A. A. Dippold, D. Izsák, T. M. Klapötke and C. Pflüger, *Chem.–Eur. J.*, 2016, **22**, 1768–1778; (b) T. Wang, H. Gao and J. n. M. Shreeve, *Z. Anorg. Allg. Chem.*, 2021, **647**, 157–191.
- (a) C.-J. Li, *Acc. Chem. Res.*, 2009, **42**, 335–344; (b) H. Wang, X. Gao, Z. Lv, T. Abdelilah and A. Lei, *Chem. Rev.*, 2019, **119**, 6769–6787.
- (a) J. Wu, Y. Zhou, Y. Zhou, C.-W. Chiang and A. Lei, *ACS Catal.*, 2017, **7**, 8320–8323; (b) Z. Ruan, Z. Huang, Z. Xu, S. Zeng, P. Feng and P.-H. Sun, *Sci. China: Chem.*, 2021, **64**, 800–807.
- C. G. Neochoritis, T. Zhao and A. Dömling, *Chem. Rev.*, 2019, **119**, 1970–2042.
- Y. Yuan and A. Lei, *Acc. Chem. Res.*, 2019, **52**, 3309–3324.
- (a) R. Francke and R. D. Little, *Chem. Soc. Rev.*, 2014, **43**, 2492–2521; (b) T. H. Meyer, I. Choi, C. Tian and L. Ackermann, *Chem*, 2020, **6**, 2484–2496; (c) K. D. Moeller, *Chem. Rev.*, 2018, **118**, 4817–4833; (d) A. Wiebe, T. Gieshoff, S. Möhle, E. Rodrigo, M. Zirbes and S. R. Waldvogel, *Angew. Chem., Int. Ed.*, 2018, **57**, 5594–5619; (e) M. Yan, Y. Kawamata and P. S. Baran, *Chem. Rev.*, 2017, **117**, 13230–13319.
- (a) T. Shen and T. H. Lambert, *Science*, 2021, **371**, 620–626; (b) T. Shen and T. H. Lambert, *J. Am. Chem. Soc.*, 2021, **143**, 8597–8602; (c) L.-H. Li, Z.-J. Niu, Y.-X. Li and Y.-M. Liang, *Chem. Commun.*, 2018, **54**, 11148–11151; (d) K. Ishihara, T. Shioiri and M. Matsugi, *Org. Lett.*, 2020, **22**, 6244–6247.
- (a) T. H. Meyer, R. C. Samanta, A. Del Vecchio and L. Ackermann, *Chem. Sci.*, 2021, **12**, 2890–2897; (b) L. Niu, C. Jiang, Y. Liang, D. Liu, F. Bu, R. Shi, H. Chen, A. D. Chowdhury and A. Lei, *J. Am. Chem. Soc.*, 2020, **142**, 17693–17702.
- (a) L. Capaldo, L. L. Quadri and D. Ravelli, *Angew. Chem., Int. Ed.*, 2019, **58**, 17508–17510; (b) J. P. Barham and B. König, *Angew. Chem., Int. Ed.*, 2020, **59**, 11732–11747; (c) Y. Yu, P. Guo, J.-S. Zhong, Y. Yuan and K.-Y. Ye, *Org. Chem.*

- Front.*, 2020, 7, 131–135; (d) J. Liu, L. Lu, D. Wood and S. Lin, *Sci. China: Chem.*, 2020, 6, 1317–1340.
- 11 (a) R. I. McDonald, G. Liu and S. S. Stahl, *Chem. Rev.*, 2011, **111**, 2981–3019; (b) J. C. Siu, N. Fu and S. Lin, *Acc. Chem. Res.*, 2020, 53, 547–560.
- 12 (a) L. Ebersson, M. P. Hartshorn, O. Persson and F. Radner, *Chem. Commun.*, 1996, 2105–2112; (b) I. Colomer, A. E. R. Chamberlain, M. B. Haughey and T. J. Donohoe, *Nat. Rev. Chem.*, 2017, 1, 0088; (c) J. L. Röckl, M. Dörr and S. R. Waldvogel, *ChemElectroChem*, 2020, 7, 3686–3694.
- 13 Z.-W. Hou, D.-J. Liu, P. Xiong, X.-L. Lai, J. Song and H.-C. Xu, *Angew. Chem., Int. Ed.*, 2021, 60, 2943–2947.
- 14 S. J. Wittenberger, *Org. Prep. Proced. Int.*, 1994, 26, 499–531.
- 15 D. M. Heard and A. J. J. Lennox, *Angew. Chem., Int. Ed.*, 2020, 59, 18866–18884.
- 16 (a) C. B. Storm, J. R. Stine and J. F. Kramer, in *Chemistry and Physics of Energetic Materials*, ed. S. N. Bulusu, Springer Netherlands, Dordrecht, 1990, pp. 605–639; (b) J. Rein, J. M. Meinhardt, J. L. Hofstra Wahlman, M. S. Sigman and S. Lin, *ChemRxiv*, 2021, preprint, DOI: [10.33774/chemrxiv-2021-16f6w](https://doi.org/10.33774/chemrxiv-2021-16f6w).
- 17 (a) S. P. Green, K. M. Wheelhouse, A. D. Payne, J. P. Hallett, P. W. Miller and J. A. Bull, *Org. Process Res. Dev.*, 2020, 24, 67–84; (b) T. M. Klapötke, C. Miró Sabaté and M. Rusan, *Z. Anorg. Allg. Chem.*, 2008, 634, 688–695.
- 18 F. Stoessel, Thermal Stability, in *Thermal Safety of Chemical Processes: Risk Assessment and Process Design*, Wiley, 2008, pp. 279–309.
- 19 (a) V. G. Kiselev, P. B. Cheblakov and N. P. Gritsan, *J. Phys. Chem. A*, 2011, 115, 1743–1753; (b) B. Yuan and E. R. Bernstein, *J. Chem. Phys.*, 2016, 144, 234302; (c) A. I. Lesnikovich, S. V. Levchik, A. I. Balabanovich, O. A. Ivashkevich and P. N. Gaponik, *Thermochim. Acta*, 1992, 200, 427–441; (d) V. G. Prokudin, V. S. Poplavsky and V. A. Ostrovskii, *Russ. Chem. Bull.*, 1996, 45, 2094–2100.
- 20 (a) W. D. Thierow and G. J. Gleicher, *J. Am. Chem. Soc.*, 1969, 91, 7150–7154; (b) C. Hansch, A. Leo and R. W. Taft, *Chem. Rev.*, 1991, 91, 165–195.
- 21 The data of BDEs were obtained using web-based computational tool ALFABET, <https://bde.ml.nrel.gov/>.
- 22 C.-Y. Cai, X.-L. Lai, Y. Wang, H.-H. Hu, J. Song, Y. Yang, C. Wang and H.-C. Xu, *Nat. Catal.*, 2022, 5, 943–951.
- 23 (a) M. Freccero, A. Pratt, A. Albinì and C. Long, *J. Am. Chem. Soc.*, 1998, 120, 284–297; (b) A. J. Fry, J. M. Porter and P. F. Fry, *J. Org. Chem.*, 1996, 61, 3191–3194; (c) E. Baciocchi, T. Del Giacco, C. Rol and G. V. Sebastiani, *Tetrahedron Lett.*, 1989, 30, 3573–3576.
- 24 S. Hajra, D. Sinha and M. Bhowmick, *J. Org. Chem.*, 2007, 72, 1852–1855.
- 25 (a) N. Fu, G. S. Sauer, A. Saha, A. Loo and S. Lin, *Science*, 2017, 357, 575–579; (b) S. Lin, J. Parry and N. Fu, *Synlett*, 2018, 29, 257–265; (c) J. C. Siu, G. S. Sauer, A. Saha, R. L. Macey, N. Fu, T. Chauviré, K. M. Lancaster and S. Lin, *J. Am. Chem. Soc.*, 2018, 140, 12511–12520; (d) J. C. Siu, J. B. Parry and S. Lin, *J. Am. Chem. Soc.*, 2019, 141, 2825–2831; (e) H. M. Nelson, J. C. Siu, A. Saha, D. Cascio, S. N. MacMillan, S.-B. Wu, C. Lu, J. A. Rodríguez, K. N. Houk and S. Lin, *Org. Lett.*, 2021, 23, 454–458.
- 26 (a) A. Studer, *Chem.–Eur. J.*, 2001, 7, 1159–1164; (b) H. Fischer, *Chem. Rev.*, 2001, 101, 3581–3610; (c) A. Studer, *Chem. Soc. Rev.*, 2004, 33, 267–273; (d) M. Yan, J. C. Lo, J. T. Edwards and P. S. Baran, *J. Am. Chem. Soc.*, 2016, 138, 12692–12714.
- 27 During the submission of this manuscript, Weng and co-workers reported a conceptually similar electrochemical azidation/heterocyclization of tryptophans. Y. Weng, X. Xu, H. Chen, Y. Zhang and X. Zhuo, *Angew. Chem., Int. Ed.*, 2022, e202206308.