

RESEARCH ARTICLE

# Expression of the eight GABA<sub>A</sub> receptor $\alpha$ subunits in the developing zebrafish central nervous system

Bryan Monesson-Olson<sup>1,2</sup>, Jon J. McClain<sup>2</sup>, Abigail E. Case<sup>2</sup>, Hanna E. Dorman<sup>2</sup>, Daniel R. Turkewitz<sup>3</sup>, Aaron B. Steiner<sup>3</sup>, Gerald B. Downes<sup>1,2\*</sup>

**1** Neuroscience and Behavior Graduate Program, University of Massachusetts, Amherst, MA, United States of America, **2** Biology Department, University of Massachusetts, Amherst, MA, United States of America, **3** Department of Biology and Health Sciences, Pace University, Pleasantville, NY, United States of America

\* [gbdowndes@bio.umass.edu](mailto:gbdowndes@bio.umass.edu)



## Abstract

GABA is a robust regulator of both developing and mature neural networks. It exerts many of its effects through GABA<sub>A</sub> receptors, which are heteropentamers assembled from a large array of subunits encoded by distinct genes. In mammals, there are 19 different GABA<sub>A</sub> subunit types, which are divided into the  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\epsilon$ ,  $\pi$ ,  $\theta$  and  $\rho$  subfamilies. The immense diversity of GABA<sub>A</sub> receptors is not fully understood. However, it is known that specific isoforms, with their distinct biophysical properties and expression profiles, tune responses to GABA. Although larval zebrafish are well-established as a model system for neural circuit analysis, little is known about GABA<sub>A</sub> receptors diversity and expression in this system. Here, using database analysis, we show that the zebrafish genome contains at least 23 subunits. All but the mammalian  $\theta$  and  $\epsilon$  subunits have at least one zebrafish ortholog, while five mammalian GABA<sub>A</sub> receptor subunits have two zebrafish orthologs. Zebrafish contain one subunit,  $\beta_4$ , which does not have a clear mammalian ortholog. Similar to mammalian GABA<sub>A</sub> receptors, the zebrafish  $\alpha$  subfamily is the largest and most diverse of the subfamilies. In zebrafish there are eight  $\alpha$  subunits, and RNA *in situ* hybridization across early zebrafish development revealed that they demonstrate distinct patterns of expression in the brain, spinal cord, and retina. Some subunits were very broadly distributed, whereas others were restricted to small populations of cells. Subunit-specific expression patterns in zebrafish resembled were those found in frogs and rodents, which suggests that the roles of different GABA<sub>A</sub> receptor isoforms are largely conserved among vertebrates. This study provides a platform to examine isoform specific roles of GABA<sub>A</sub> receptors within zebrafish neural circuits and it highlights the potential of this system to better understand the remarkable heterogeneity of GABA<sub>A</sub> receptors.

## OPEN ACCESS

**Citation:** Monesson-Olson B, McClain JJ, Case AE, Dorman HE, Turkewitz DR, Steiner AB, et al. (2018) Expression of the eight GABA<sub>A</sub> receptor  $\alpha$  subunits in the developing zebrafish central nervous system. PLoS ONE 13(4): e0196083. <https://doi.org/10.1371/journal.pone.0196083>

**Editor:** Harold A. Burgess, National Institutes of Health, UNITED STATES

**Received:** January 11, 2018

**Accepted:** April 5, 2018

**Published:** April 27, 2018

**Copyright:** © 2018 Monesson-Olson et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was supported by a University of Massachusetts Faculty Research/Healey Endowment Grant and National Science Foundation Grant 1456866. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

## Introduction

Neural networks throughout the central nervous system rely upon a diversity of neurotransmitter systems for both their initial formation and mature function. GABA is the major inhibitory neurotransmitter throughout most of the mature nervous system [1]. It exerts its effects through its receptors, which are divided into two classes, GABA<sub>A</sub> and GABA<sub>B</sub>. GABA<sub>A</sub> receptors are ligand-gated chloride channels responsible for most of the rapid effects of GABA, while GABA<sub>B</sub> receptors are heterotrimeric G-protein coupled receptors. In mammalian systems, GABA<sub>A</sub> receptors demonstrate incredible subtype diversity and are targets for classes of clinically important drugs, such as benzodiazepines and barbituates [2, 3]. GABA<sub>A</sub> receptors are heteropentamers composed of various combinations of 19 different subunits:  $\alpha$ 1–6,  $\beta$ 1–3,  $\gamma$ 1–3,  $\delta$ ,  $\epsilon$ ,  $\pi$ ,  $\theta$  and  $\rho$ 1–3 (previously referred to as GABA<sub>C</sub> receptors) [4, 5]. Each subunit is encoded by a discrete gene that is spatially and developmentally regulated to generate distinct, but often overlapping, expression patterns [6–9]. Alternative splicing and RNA editing of some subunits further increases the number of subtypes available [10]. Although this extensive receptor heterogeneity is not fully understood, some subunits confer distinct biophysical and pharmacological properties, interact with specific cytoplasmic proteins, and localize to specific subcellular domains [11–13]. Ultimately, this receptor diversity provides a capacity to tailor responses to GABA within neural circuits.

The  $\alpha$  subunits play a prominent role in GABA<sub>A</sub> receptor function. They are thought to be essential components of ‘classic’ GABA<sub>A</sub> receptors, which exclude the pharmacologically distinct receptors composed from  $\rho$  subunits [3]. Exogenous expression studies and analysis of native receptors indicate that each receptor contains two  $\alpha$  subunits which, along with  $\beta$  subunits, form the GABA binding site.  $\alpha$  subunits can also dictate isoform-selective pharmacology. For example, receptors containing  $\alpha$ 1,  $\alpha$ 2,  $\alpha$ 3 and  $\alpha$ 5 subunits confer benzodiazepine-sensitivity, whereas receptors that contain  $\alpha$ 4 and  $\alpha$ 6 subunits do not bind to clinically used benzodiazepines [2]. In addition,  $\alpha$  subunits regulate subcellular localization. GABA receptors that contain  $\alpha$ 1,  $\alpha$ 2, or  $\alpha$ 3 demonstrate a synaptic localization and mediate transient or phasic inhibition, while receptors that contain  $\alpha$ 4,  $\alpha$ 5 or  $\alpha$ 6 are principally extrasynaptic and mediate tonic inhibition [14, 15]. Through both transient and tonic inhibition, these receptors mediate the majority of inhibition in neural circuits throughout the vertebrate brain.

Developing zebrafish have several features that make them well-suited to study the formation and function of neural circuits [16, 17]. First, zebrafish embryos and larvae develop external to the mother and are optically transparent. These features make the central nervous system easily accessible throughout development and particularly amenable to optical physiological approaches, such as optogenetics or calcium imaging. Second, many cell types and mechanisms are conserved among vertebrates, yet the central nervous system contains fewer cells compared to mammalian preparations. Lastly, larval zebrafish develop rapidly, therefore many sensory and motor systems are present and functional within five days post-fertilization. Combined, these features have helped establish developing zebrafish as an important system to examine neural circuit formation and function.

GABA<sub>A</sub> receptors are expressed in larval zebrafish and essential for normal nervous system function. For example, bath application of pentylenetetrazole, a convulsant and noncompetitive GABA<sub>A</sub> receptor antagonist, induces hyperactive behavior that can serve as a model of seizures [18]. Similarly, insecticides known to target GABA<sub>A</sub> receptors generate hyperactive behavior in larvae [19]. Pharmacological blockade of GABA<sub>A</sub> receptors has also been shown to regulate the activity of specific cells, including retinal bipolar cells [20], optic tectum neurons [21], mitral cells in the olfactory bulb [22], spinal cord CoPA interneurons and motoneurons [23], and the Mauthner cells, a pair of well-studied reticulospinal neurons found in the

amphibian and teleost hindbrain [24]. In these studies, the GABA<sub>A</sub> receptor isoforms were not identified. However, patch-clamp recordings of the Mauthner cells showed three distinct GABA<sub>A</sub> kinetic profiles, which were proposed to be caused by different receptor isoforms [25].

There is limited information about the heterogeneity of GABA<sub>A</sub> receptors in zebrafish. The most extensive study to date identified 23 GABA<sub>A</sub> receptor subunits and examined their expression broadly in adult zebrafish brain via RT-qPCR [26]. Despite playing major roles in developing animals, little is known about the expression of GABA<sub>A</sub> receptor subunits in zebrafish embryos and larvae.  $\alpha 1$  has been shown to be expressed at 50 and 72 hours post-fertilization (hpf),  $\gamma 2$  at 50 hpf, and  $\alpha 2a$  (then named  $\alpha 2$ ) from ~14–96 hpf. [27–29]. Unpublished  $\alpha 6a$  expression data has also been deposited in the ZFIN database [30]. The expression patterns of other GABA<sub>A</sub> receptor subunits has not been reported.

In this study we performed phylogenetic analysis, which indicates that zebrafish contain at least 23 different GABA<sub>A</sub> receptor subunits. Although we observed some differences between the zebrafish and mammalian GABA<sub>A</sub> receptor subunit gene families, zebrafish contain orthologs for most of the GABA<sub>A</sub> receptor subunits found in mammals. To examine when and where these GABA<sub>A</sub> receptors are expressed in the zebrafish nervous system, we performed whole-mount RNA *in situ* hybridization. We present the embryonic and larval expression of the eight identified  $\alpha$  subunit-encoding genes: *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4*, *gabra5*, *gabra6a* and *gabra6b*. These data show that each subunit has a distinct expression pattern, broadly similar to the reported expression of amphibian and rodent orthologs. Combined, these results suggest that GABA<sub>A</sub> isoform-specific roles are conserved among vertebrates, and they establish a foundation to use the zebrafish system to better understand how GABA<sub>A</sub> receptor heterogeneity mediates neural circuit formation and function.

## Materials and methods

### Phylogenetic analysis

Homologous gene queries were performed using the National Center for Biotechnology Information tBLASTn search tool. Protein sequences for each of the 19 mouse GABA<sub>A</sub> receptor subunits were used to search the zebrafish nucleotide collection. Matching zebrafish sequences were evaluated to determine if they were splice variants from the same gene or generated from different genes. When splice variants were identified, only the longest variant was used for subsequent analysis. 19 mouse and 23 zebrafish protein sequences were used to assemble a phylogenetic tree. The genes and Genbank accession numbers for the mouse sequences are: *Gabra1*- NP\_034380, *Gabra2*- NP\_032092, *Gabra3*- NP\_032093, *Gabra4*- NP\_034381, *Gabra5*- NP\_795916, *Gabra6*- NP\_00109311, *Gabrb1*- NP\_032095, *Gabrb2*- NP\_032096, *Gabrb3*- NP\_032097, *Gabrd*- NP\_032098, *Gabre*- NP\_059065, *Gabrg1*- NP\_034382, *Gabrg2*- NP\_032099, *Gabrg3*- NP\_032100, *Gabrp*- NP\_666129, *Gabrq*- NP\_065234, *Gabrr1*- NP\_032101, *Gabrr2*- NP\_032102, and *Gabrr3*- NP\_001074659. The Genbank accession numbers for the zebrafish sequences are: *gabra1*- NM\_001077326, *gabra2a*- XM\_009307207, *gabra2b*- XM\_017359049, *gabra3*- XM\_009295708, *gabra4*- NM\_001017822, *gabra5*- XM\_001339475, *gabra6a*- XM\_005173112, *gabra6b*- XM\_002667357, *gabrb1*- XM\_002664133, *gabrb2*- NM\_001024387, *gabrb3*- XM\_005166079, *gabrb4*- XM\_005173874, *gabrd*- XM\_695007, *gabrg1*- XM009307208, *gabrg2*- NM\_001256250, *gabrg3*- XM\_009302568, *gabrp*- XM\_005173293, *gabrr1*- NM\_001025553, *gabrr2a*- NM\_001045376, *gabrr2b*- XM\_009294512, *gabrr3a*- NM\_001128760, *gabrr3b*- XM\_009297450, *gabrz*- XM\_005156247. To construct the tree, a ClustalW alignment of amino acid sequences was performed using the Geneious software package [31]. An unrooted PHYML consensus tree was then generated using 100 bootstrap replicates [32].

## Fish maintenance and breeding

Zebrafish were raised and maintained using established husbandry procedures. Embryos were kept at 28.5 °C in E3 media and staged according to morphological criteria [33]. All experiments were performed using Tuebingen (Tu) or Tupfel long fin (TLF) wild type embryos. From 0–24 hpf, embryos were grown in 0.01% Methylene Blue in E3 medium. At 24 hpf the solution was changed to 0.0045% Phenylthiourea (PTU) in E3 to inhibit pigment formation. This solution was changed every 24 hours until the fish were sacrificed by anesthetizing them in 0.4% MS-222 until the cessation of movement then transferring the fish into 4% paraformaldehyde. All animal protocols were approved by the University of Massachusetts Institutional Animal Care and Use Committees (IACUC).

## Whole-mount *in situ* hybridization

Antisense digoxigenin probes were generated against *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4*, *gabra5*, *gabra6a* and *gabra6b* (S1 Table). *In situ* probes were synthesized using a digoxigenin RNA labeling kit with SP6 or T7 RNA polymerases (Roche Diagnostics, Mannheim, Germany). Whole-mount, colorimetric *in situ* hybridization was performed using established protocols [34] and examined using a compound microscope (Zeiss, Thornwood, NY) attached to a digital camera (Zeiss, Thornwood, NY). Cross sections were generated by imbedding *in situ* hybridization stained embryos in 1.5% agar, 5% sucrose. The blocks were submerged in a 30% sucrose solution overnight then cut into 20  $\mu$ m thick sections using a cryostat (Leica, Buffalo Grove, IL). Representative larvae at 24, 48, and 96 hpf were mounted in either 100% glycerol or dehydrated through a methanol series, equilibrated in a 2:1 benzylbenzoate/benzylalcohol solution and mounted in a 10:1 Canada balsam/methyl salicylate mixture. Images were captured using a Zeiss Discovery v12 stereomicroscope with a 1.5x objective or a Zeiss Axioskop Microscope with a 10x objective and an AxioCam MRc camera. All images were processed using Adobe Photoshop, in which multiple focal planes were merged to produce single representative images. The identify of neuroanatomical structures was determined using zebrafish brain atlases [35–37].

## Reverse-Transcriptase PCR


RT-PCR was used to analyze whether the *gabra3* and *gabra3*-like sequence were portions of the same transcript. Primers designed against *gabra3*-like (Primer 1, 5' -GGACGGCGGATGATGAGAAA-3') and *gabra3* (Primer 2, 5' -CACGACCGTCCTGACTA-3', Primer 3, 5' -GTG GAGTAGATGTGGTGGGC-3') were used to amplify cDNA from wild-type zebrafish larvae. RNA was extracted from 48 hpf zebrafish larvae using the RNAeasy kit (Qiagen, Venlo, Netherlands) and reverse transcribed using the Accuscript RT-PCR system (Stratagene). The PCR products were sequenced to confirm that they are portions from the same transcript.

## Results

### The zebrafish GABA<sub>A</sub> receptor subunit gene family is similar in size and diversity to the mammalian GABA<sub>A</sub> receptor gene family

19 GABA<sub>A</sub> receptor subunit genes have been identified in humans, mice, and rats [4, 5]. To establish the number and organization of zebrafish GABA<sub>A</sub> receptor subunits, we used mouse GABA<sub>A</sub> receptor subunit amino acid sequences to query zebrafish genome databases. We identified 23 zebrafish GABA<sub>A</sub> receptor subunits, each encoded by a distinct gene. Splice variants were frequently observed. In these cases, the principle splice isoform, as indicated by the databases, was selected for analysis. 12 of the 19 mouse GABA<sub>A</sub> receptor subunits were found





**Fig 1. Phylogenetic analysis shows that the zebrafish GABA<sub>A</sub> subunit gene family is similar in size, diversity, and organization to the mouse GABA<sub>A</sub> subunit gene family.** Amino acid sequence alignments were used to generate a consensus tree using 100 bootstrap replicates. The genes that encode the GABA<sub>A</sub> subunits are shown at the tip of each branch and bootstrap proportions are shown at the branch points. The zebrafish and mouse GABA<sub>A</sub> subunit sequences showed high amino acid identity and grouped into the  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$ ,  $\pi$ , and  $\rho$  subfamilies. Most mouse GABA<sub>A</sub> subunits have a single zebrafish ortholog, while six mouse GABA<sub>A</sub> subunits have two zebrafish orthologs, likely due to gene duplication.

<https://doi.org/10.1371/journal.pone.0196083.g001>

to have a single ortholog in zebrafish (Fig 1). Amino acid identities between mouse and zebrafish orthologs ranged from 53.1 to 86.3%. Five additional mouse subunits,  $\alpha 2$ ,  $\alpha 6$ ,  $\rho 2$ ,  $\rho 3$ , and  $\pi$ , each exhibit similarity with two subunits in zebrafish: *gabrac2a* and *gabrac2b*, *gabrac6a* and *gabrac6b*, *gabrr2a* and *gabrr2b*, *gabrr3a* and *gabrr3b*, *gabrp* and *gabrz*, respectively. The relatively high percentage of amino acid identity of these zebrafish subunits with each other suggests that the zebrafish paralogs are duplicated genes. Duplicated genes are often observed in zebrafish, and they are thought to be due to a whole genome duplication within the teleost lineage [38, 39]. It was reported previously that there is a *gabrac3* and an  $\alpha 3$ -like gene. While

the  $\alpha 3$ -like gene has been localized to chromosome 21, the genomic location of the  $\alpha 3$ -like gene is not known. Our database analysis suggested that these sequences could be non-overlapping portions of the same gene, therefore RT-PCR was performed using primers targeting  $\alpha 3$ -like and *gabra3* sequences (S1 Fig). Transcripts were amplified that spanned the region between *gabra3* and  $\alpha 3$ -like sequences, indicating that these are different portions of the same gene and that there is only one  $\alpha 3$  encoding gene, *gabra3*.

There are two mouse GABA<sub>A</sub> receptor subunits with no clear orthologs, the  $\theta$  and  $\epsilon$  subunits. The  $\theta$  subunit, encoded by the gene *Gabrq*, exhibits only 31.3% amino acid identity with zebrafish  $\beta 4$ , encoded by *gabrb4*, which is the greatest percentage identity of this subunit with any zebrafish sequence. There are three  $\beta$  subunits in mice compared to the four  $\beta$  subunits in zebrafish so it is possible that, despite the low amino acid sequence identity, zebrafish  $\beta 4$  is orthologous to the mouse  $\theta$  subunit. No clear zebrafish ortholog has been identified for the mouse  $\epsilon$  subunit, which is encoded by the *Gabre* gene.


### $\alpha$ subunits demonstrate distinct expression patterns across early zebrafish development

Determining the temporal and spatial expression of GABA<sub>A</sub> subunits is essential to identify isoform-specific roles in zebrafish neural circuit development and function. Given the large size of the  $\alpha$  subunit subfamily and their leading role in GABA<sub>A</sub> receptor function, we examined the expression of the eight  $\alpha$  subunit-encoding genes: *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4*, *gabra6a* and *gabra6b*. Using whole-mount RNA *in situ* hybridization, we determined their expression patterns at 24, 48, and 96 hpf. These time points span key stages of locomotor network development in developing zebrafish [40].

Expression of *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4* and *gabra5* was detected at 24 hpf (Fig 2). Neither *gabra6a* or *gabra6b* were detected at this developmental stage. *gabra2a* and *gabra2b* were detected broadly and did not appear to be spatially restricted (Fig 2C–2F). To distinguish whether these gene transcripts were widespread or background due to the probe, a second probe was generated for each gene (S1 Table). These probes yielded widespread staining, very similar to the initial probes used, which suggests that *gabra2a* and *gabra2b* are widely distributed. Consistent with these findings, a previous study reported that *gabra2a* exhibits a diffuse, broad pattern of expression in embryonic and larval zebrafish [28].

*gabra1*, *gabra3*, *gabra4*, and *gabra5* were each expressed in discrete cells in the brain or spinal cord at 24 hpf. *gabra1* was detected in the olfactory placodes, the ventral rostral cluster, the ventral caudal cluster and small clusters of cells within each rhombomere of the hindbrain (Fig 2A and 2B). Although expressed in discrete cells, *gabra3* was widely expressed, and found within the tegmentum, hindbrain, and ventral and intermediate domains in the spinal cord (Fig 2G and 2H). *gabra4* was expressed in a diffuse manner throughout the brain but selectively within a population of ventral spinal cord cells (Fig 2I and 2J). These *gabra4* expressing cells are cuboidal in appearance and along the central canal, which suggests they may be Kolmer-Agduhr (KA) neurons [41]. Lastly, *gabra5* is expressed in small groups of cells in each rhombomere of the hindbrain and in a dorsal domain of the spinal cord (Fig 2K and 2L). The large soma size of these spinal cord cells, their dorsal position, and the number of labeled cells per somite (~1) suggests that they are CoPA neurons, which have been previously reported to receive GABAergic input [23, 42].

The expression of all GABA<sub>A</sub> receptor subunit genes, except for *gabra6a*, was observed at 48 hpf (Fig 3). Similar to the pattern observed at 24 hpf, the *gabra2* paralogs were again detected broadly, with little spatial restriction (Fig 3C–3F). Similar to its expression at 24 hpf, *gabra1* is expressed in the olfactory bulbs, the pallium and discrete clusters of cells in medial



**Fig 2. Expression of GABA<sub>A</sub>  $\alpha$  subunits at 24 hpf.** *gabra1* (A, B), *gabra2a* (C, D), *gabra2b* (E, F), *gabra3* (G, H), *gabra4* (I, J), and *gabra5* (K, L) were detected at this time point. Whole-mount lateral (A, C, E, G, I, K) views are shown along with dorsal views of the head (B, D, F, H, J, L). The scale bar (A) is 0.1 mm. The brackets in G, I, and K indicate the regions shown at higher magnification in the corresponding insets. The arrowheads within the insets indicate the dorsal and ventral boundaries of the spinal cord. Abbreviations: drc, dorsal rostral cluster; hbc, hindbrain cluster; op, olfactory placode; SC, spinal cord; t, tectum; vcc, ventral caudal cluster; vrc, ventral rostral cluster.

<https://doi.org/10.1371/journal.pone.0196083.g002>


portions of the medulla (Fig 3A and 3B). *gabra3* was observed in relatively small clusters of cells in the pallium, thalamus, and the medulla (Fig 3G and 3H). *gabra4* transcripts were identified in distinct cells in the subpallium and lateral portions of the medulla. *gabra5* transcripts were detected most prominently in the medulla. This staining suggests  $\alpha 5$  is expressed in the Mauthner cells, and that  $\alpha 5$  containing receptors are likely some of the GABA<sub>A</sub> isoforms identified via electrophysiology [25]. *gabra6b* is expressed prominently in the olfactory bulbs.

Although *gabra1*, *gabra3*, *gabra4*, and *gabra5* transcripts were all detected in the medulla, they do not appear to be in the same cells. Instead, an intriguing medial to lateral organization was observed. *gabra1* is expressed strongly in the most medial cells, *gabra3* and *gabra5* transcripts were found in an intermediate domain, and *gabra4* was observed laterally (compare insets in Fig A, G, I, K).

Transcripts for all  $\alpha$  subunits are detected in larval zebrafish at 96 hpf (Fig 4). As with the earlier developmental time points, *gabra2a* and *gabra2b* transcripts were detected broadly, although *gabra2b* appears less diffuse and more discrete compared to at 24 and 48 hpf (Fig 4C–4F). *gabra1*, *gabra3*, and *gabra6b* are also expressed more broadly compared to earlier stages. In contrast, *gabra4*, *gabra5* and *gabra6a* transcripts were detected in smaller groups of cells. *gabra4* is expressed in the retina, the posterior tuberculum area, which is a portion of the diencephalon, and the tectum, cerebellum, and medulla (Fig I, J). *gabra5* is expressed in discrete cells in the pallium, hypothalamus, cerebellum and medulla (Fig K, L). Within the medulla, *gabra5* expression in the Mauthner Cells is robust, as it is at earlier developmental stages.

Transcripts for several  $\alpha$  subunits are expressed in the retina at 96 hpf, however most are restricted to one or two cell layers (Fig 4). *gabra1*, *gabra2a*, *gabra2b*, *gabra3*, *gabra4*, and





**Fig 3. Expression of the GABA<sub>A</sub> α subunits in the brain at 48hpf.** *gabra1* (A, B), *gabra2a* (C, D), *gabra2b* (E, F) *gabra3* (G, H) *gabra4* (I, J) *gabra5* (K, L) and *gabra6b* (M, N) were detected at this time point. Lateral (A, C, E, G, I, K, M) and dorsal (B, D, F, H, J, L, N) views are shown. The scale bar (A) is 0.1 mm. Brackets in A, G, I, and K indicate the region shown in cross sections within the insets. Abbreviations: cb, cerebellum; di, diencephalon; hth, hypothalamus; M, Mauthner cell; mo, medulla oblongata; ob, olfactory bulb; p, pallium; sp, subpallium; t, tegmentum; th, thalamus.


<https://doi.org/10.1371/journal.pone.0196083.g003>

*gabra6b* transcripts are all expressed in distinct cell layers in the retina. Although most of these genes are also expressed in other areas of the nervous system, *gabra6a* does not demonstrate robust expression outside of the retina. *gabra6a* is expressed prominently in the photoreceptor cell layer.

## Discussion

In this study, we showed that the GABA<sub>A</sub> receptor gene family exhibits a number and diversity very similar to those found in mammals. We identified sequences for 23 GABA<sub>A</sub> receptor encoding genes, and phylogenetic analysis indicates that most isoforms are conserved across





**Fig 4. Expression of the GABA<sub>A</sub> α subunits in the brain at 96hpf.** Transcripts encoding all α subunits were detected at this time point. Lateral (A, C, E, G, I, K, M, O) and dorsal (B, D, F, H, J, L, N, P) are shown. The scale bar (A) is 0.1 mm. Abbreviations: cb, cerebellum; gcl, hypothalamus; M, Mauthner cell; mo, medulla oblongata; ob, olfactory bulb; opn, optic nerve; p, pallium; pcl, photoreceptive cell layer; po, preoptic region; poc, post optic commissure; pta, posterior tubercular area; sp, subpallium; t, tegmentum; te, telencephalon; th, thalamus; to, tectum opticum.

<https://doi.org/10.1371/journal.pone.0196083.g004>

vertebrates. We determined the expression of the eight zebrafish α subunit encoding genes, the largest and most diverse gene family, and observed that they are expressed in distinct, often overlapping expression patterns. Taken together, these data argue that larval zebrafish can serve as a useful model to investigate the functional roles of GABA<sub>A</sub> receptor subtypes within developing neural circuits.

A previous study identified 23 GABA<sub>A</sub> receptor genes in zebrafish [26]. Our data largely confirm their results, in addition, we identified the previously unreported subunit p3b, which is encoded by *gabrr3b*. An α3 and an α3-like gene were also described, but our analysis indicates that zebrafish likely contain only one α3 gene. Given the amount of zebrafish genome



data that is available, it seems unlikely that additional GABA<sub>A</sub> receptor subunit encoding genes exist. However, similar to mammalian systems, sequence data indicate that several zebrafish GABA<sub>A</sub> receptor encoding genes are alternatively spliced, which further enhances the already extensive diversity of receptor isoforms.

The evolution of the GABA<sub>A</sub> receptor gene family has been examined by using sequence data from a wide variety of species, including humans, rodents, canary, chicken, frogs, pufferfish, tunicates, *C. elegans* and *Drosophila* [43, 44]. Our results are in line with these studies, finding that mammalian  $\theta$  and  $\epsilon$  subunits are unusual when compared to other subunits. Similar to other non-mammalian vertebrates, zebrafish do not have clear orthologs for the  $\theta$  and  $\epsilon$  subunits; their existence outside of the mammalian lineage is unclear [45]. Even among mammalian species, for example comparing humans to rodents, these two subunits are very diverse, which suggests they have evolved at a much faster rate than other subunits [46]. The roles of  $\theta$  and  $\epsilon$  subunits may be unique to mammals or, alternatively, fulfilled in zebrafish and other species using other subtypes. Experiments to investigate the functional ability of zebrafish subtypes to substitute for  $\theta$  and  $\epsilon$  in rodent systems could distinguish between these possibilities.

The  $\alpha$  subunits showed a wide range of expression patterns across early zebrafish development (Table 1). On one end of the spectrum are *gabra2a* and *gabra2b*, which appeared to be expressed very broadly at each of the three time-points examined. At the other end of the spectrum is *gabra6a*, which we detected only at 96 hpf and mostly in the photoreceptor cell layer of the retina. The other subunits fall between these extremes, with expression in discrete cells in various brain and spinal cord regions. When GABA<sub>A</sub> receptor  $\alpha$  subtypes are expressed in the same region, they often occupy different domains. For example, *gabra1*, *gabra3*, *gabra4*, and *gabra5* are all expressed in the medulla, but are organized in medial to lateral stripes. The zebrafish medulla has a structural and functional organization in which neurons of shared neurotransmitter, phenotype, age, morphology and functional properties are arranged into stripes [47, 48]. Another example of different GABA<sub>A</sub>  $\alpha$  subtypes occupying different domains within the same region is found in the spinal cord. *gabra4* transcripts were found in ventral cells, likely KA neurons, *gabra5*-expressing cells were observed more dorsally, likely CoPA neurons, and *gabra3* transcripts were detected more broadly in both ventral and more dorsal domains. It is not yet clear how the expression of different GABA<sub>A</sub> subtypes correspond to the organizational stripes of the hindbrain or cell types in the spinal cord, but it seems likely that some subtypes demonstrate cell-type specific expression and confer distinct responses to GABA. Conclusive cell-type identification will require co-labeling experiments that examine GABA<sub>A</sub> subunit expression along with specific markers, techniques that determine neuronal identity by revealing cell morphology, and/or precise registration in one of the zebrafish digital brain atlases [49–52].

The expression patterns of GABA<sub>A</sub>  $\alpha$  subunits are well-conserved across species. The expression of five  $\alpha$  subunits has been described in *Xenopus laevis* during development [53]. Consistent with our observations in zebrafish,  $\alpha 2$  was found to be expressed broadly and early in development. In fact,  $\alpha 2$  was found to be deposited in eggs as a maternal message.  $\alpha 6$  showed the latest onset of expression during development and demonstrated robust photoreceptor expression. Likewise, the expression patterns of  $\alpha 1$ ,  $\alpha 3$ , and  $\alpha 5$  are similar between frogs and zebrafish. These similarities in expression extend to mammalian systems. For example, in rat, similar to zebrafish  $\alpha 2$ , transcripts were detected relatively early in development and were widespread, and  $\alpha 6$  is expressed much later in development and exhibits restricted expression [7]. One notable difference is that in mammals  $\alpha 6$  transcripts were detected in the cerebellum and retinal expression has not been reported, whereas in zebrafish  $\alpha 6a$  exhibits robust retinal expression and  $\alpha 6b$  is more widespread. It is unclear how much these differences in expression between zebrafish and mammals can be attributed to the various development

**Table 1. Expression of GABA<sub>A</sub> receptor  $\alpha$  subunits at 24, 48, and 96 hours post fertilization.**

		$\alpha 1$	$\alpha 2a$	$\alpha 2b$	$\alpha 3$	$\alpha 4$	$\alpha 5$	$\alpha 6a$	$\alpha 6b$
24 hpf	dorsal rostral cluster		•	•					
	hindbrain cluster	•	•	•	•	•	•		
	olfactory placode	•							
	spinal cord				•	•	•		
	tegmentum				•				
	ventral caudal cluster	•	•	•	•	•			
	ventral rostral cluster	•	•	•					
48 hpf	cerebellum		•	•					
	diencephalon	•							
	hypothalamus						•		
	Mauthner cell						•		
	medulla oblongata	•			•	•			
	olfactory bulb	•							•
	pallium				•	•	•		
	subpallium				•				
	tegmentum				•				
	thalamus				•				
96 hpf	cerebellum	•	•	•		•	•	•	•
	hypothalamus	•		•			•		
	Mauthner cell						•		
	medulla oblongata	•		•	•	•	•		
	olfactory bulb								•
	optic nerve	•							
	pallium			•	•	•	•		
	post-optic commissure			•					
	pre-optic region			•					
	tectum opticum	•					•		
	tegmentum	•		•	•				
		thalamus				•			
	retina-	•		•	•	•		•	•
	photoreceptors						•		

Shaded boxes indicate broad, diffuse expression. Dots indicate punctate expression.

<https://doi.org/10.1371/journal.pone.0196083.t001>

stages and tissues selected for investigation versus authentic differences in patterns of expression.

Developing zebrafish are a powerful system for neural circuit analysis and, more recently, they have been cultivated as a model of epilepsy [17, 54–58]. Although GABA<sub>A</sub> receptors are robust regulators of many neural circuits and play a central role in epilepsy syndromes, GABA<sub>A</sub> receptors have not been well characterized in zebrafish. By reporting the diversity of zebrafish GABA<sub>A</sub> receptor subunits and describing the expression patterns of the  $\alpha$  subfamily during early development, we have laid a foundation to leverage the strengths of the zebrafish system to investigate isoform specific roles of GABA<sub>A</sub> receptors and further develop zebrafish as a model of epilepsy. In mammalian systems, genetic inactivation of individual  $\alpha$  subunits has revealed surprisingly subtle defects compared to what would be predicted from pharmacological blockade. Compensatory upregulation of other subunits is thought to mask some of the effects of inactivating a single subunit [59, 60]. The ability easily mutate multiple genes in

zebrafish may help overcome such compensatory changes in expression to shed new light into subtype specific roles of GABA<sub>A</sub> receptors.

## Supporting information

**S1 Fig. RT-PCR results suggest that zebrafish contain only one  $\alpha 3$  gene.** (A) Schematic of *gabra3* cDNA. The portions that are identical to *gabra3*-like and *gabra3* sequences are shaded green. The white region that connects the two was identified through RT-PCR. The location of the three primers used for PCR are shown as numbered arrows. (B) RT-PCR results. Primers 2 and 3 served as a positive control since they amplify sequence from the known *gabra3* region. Primers 1 and 3 amplify a previously unknown region that links *gabra 3*-like and *gabra3* sequence, showing that they are contained within same transcript and likely portions of the same gene.

(PDF)

**S1 Table. GABA<sub>A</sub>  $\alpha$  subunit antisense RNA probe information.** Note- sizes are shown in base pairs. The probe start site is defined here as the start codon of the open reading frame. Negative values indicate that the probe contains 5' untranslated sequence. Some probes were generated via RT-PCR with a primer that contains an RNA polymerase promoter, so subcloning into vectors wasn't performed.

(PDF)

## Acknowledgments

The authors thank members of the Downes lab, Kelly Anne McKeown at Westfield State University, and Josef Trapani at Amherst College for helpful advice and discussion.

## Author Contributions

**Conceptualization:** Gerald B. Downes.

**Formal analysis:** Bryan Monesson-Olson, Jon J. McClain, Gerald B. Downes.

**Funding acquisition:** Gerald B. Downes.

**Investigation:** Bryan Monesson-Olson, Jon J. McClain, Abigail E. Case, Gerald B. Downes.

**Methodology:** Abigail E. Case, Hanna E. Dorman, Gerald B. Downes.

**Project administration:** Gerald B. Downes.

**Resources:** Bryan Monesson-Olson, Jon J. McClain, Hanna E. Dorman, Daniel R. Turkewitz, Aaron B. Steiner.

**Supervision:** Gerald B. Downes.

**Visualization:** Bryan Monesson-Olson, Jon J. McClain, Abigail E. Case, Gerald B. Downes.

**Writing – original draft:** Gerald B. Downes.

**Writing – review & editing:** Bryan Monesson-Olson, Jon J. McClain, Abigail E. Case, Hanna E. Dorman, Daniel R. Turkewitz, Aaron B. Steiner, Gerald B. Downes.

## References

1. Bowery NG, Smart TG. GABA and glycine as neurotransmitters: a brief history. *Br J Pharmacol*. 2006; 147 Suppl 1:S109–19. Epub 2006/01/13. <https://doi.org/10.1038/sj.bjp.0706443> PMID: 16402094; PubMed Central PMCID: PMC1760744.

2. Mohler H. GABA(A) receptor diversity and pharmacology. *Cell Tissue Res.* 2006; 326(2):505–16. Epub 2006/08/29. <https://doi.org/10.1007/s00441-006-0284-3> PMID: 16937111.
3. Sigel E, Steinmann ME. Structure, function, and modulation of GABA(A) receptors. *J Biol Chem.* 2012; 287(48):40224–31. Epub 2012/10/06. <https://doi.org/10.1074/jbc.R112.386664> PMID: 23038269; PubMed Central PMCID: PMC3504738.
4. Chua HC, Chebib M. GABAA Receptors and the Diversity in their Structure and Pharmacology. *Adv Pharmacol.* 2017; 79:1–34. Epub 2017/05/23. <https://doi.org/10.1016/bs.apha.2017.03.003> PMID: 28528665.
5. Olsen RW, Sieghart W. International Union of Pharmacology. LXX. Subtypes of gamma-aminobutyric acid(A) receptors: classification on the basis of subunit composition, pharmacology, and function. Update. *Pharmacol Rev.* 2008; 60(3):243–60. Epub 2008/09/16. <https://doi.org/10.1124/pr.108.00505> PMID: 18790874; PubMed Central PMCID: PMC2847512.
6. Laurie DJ, Seeburg PH, Wisden W. The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. II. Olfactory bulb and cerebellum. *J Neurosci.* 1992; 12(3):1063–76. Epub 1992/03/01. PMID: 1312132.
7. Laurie DJ, Wisden W, Seeburg PH. The distribution of thirteen GABAA receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci.* 1992; 12(11):4151–72. PMID: 1331359.
8. Wisden W, Laurie DJ, Monyer H, Seeburg PH. The distribution of 13 GABAA receptor subunit mRNAs in the rat brain. I. Telencephalon, diencephalon, mesencephalon. *J Neurosci.* 1992; 12(3):1040–62. Epub 1992/03/01. PMID: 1312131.
9. Delgado-Lezama R, Loeza-Alcocer E, Andres C, Aguilar J, Guertin PA, Felix R. Extrasynaptic GABA (A) receptors in the brainstem and spinal cord: structure and function. *Curr Pharm Des.* 2013; 19(24):4485–97. Epub 2013/01/31. PMID: 23360278.
10. Simon J, Wakimoto H, Fujita N, Lalande M, Barnard EA. Analysis of the set of GABA(A) receptor genes in the human genome. *J Biol Chem.* 2004; 279(40):41422–35. Epub 2004/07/20. <https://doi.org/10.1074/jbc.M401354200> PMID: 15258161.
11. Olsen RW, Sieghart W. GABA A receptors: subtypes provide diversity of function and pharmacology. *Neuropharmacology.* 2009; 56(1):141–8. Epub 2008/09/02. <https://doi.org/10.1016/j.neuropharm.2008.07.045> PMID: 18760291; PubMed Central PMCID: PMC3525320.
12. Tretter V, Moss SJ. GABA(A) Receptor Dynamics and Constructing GABAergic Synapses. *Front Mol Neurosci.* 2008; 1:7. Epub 2008/10/24. <https://doi.org/10.3389/neuro.02.007.2008> PMID: 18946540; PubMed Central PMCID: PMC35256003.
13. Fritschy JM, Panzanelli P. GABAA receptors and plasticity of inhibitory neurotransmission in the central nervous system. *Eur J Neurosci.* 2014; 39(11):1845–65. Epub 2014/03/19. <https://doi.org/10.1111/ejn.12534> PMID: 24628861.
14. Farrant M, Nusser Z. Variations on an inhibitory theme: phasic and tonic activation of GABA(A) receptors. *Nat Rev Neurosci.* 2005; 6(3):215–29. Epub 2005/03/02. <https://doi.org/10.1038/nrn1625> PMID: 15738957.
15. Egawa K, Fukuda A. Pathophysiological power of improper tonic GABA(A) conductances in mature and immature models. *Front Neural Circuits.* 2013; 7:170. Epub 2013/10/30. <https://doi.org/10.3389/fncir.2013.00170> PMID: 24167475; PubMed Central PMCID: PMC3807051.
16. Wolman M, Granato M. Behavioral genetics in larval zebrafish: learning from the young. *Dev Neurobiol.* 2012; 72(3):366–72. Epub 2012/02/14. <https://doi.org/10.1002/dneu.20872> PMID: 22328273.
17. McLean DL, Fetcho JR. Movement, technology and discovery in the zebrafish. *Curr Opin Neurobiol.* 2011; 21(1):110–5. Epub 2010/10/26. <https://doi.org/10.1016/j.conb.2010.09.011> PMID: 20970321; PubMed Central PMCID: PMC3057273.
18. Baraban SC, Taylor MR, Castro PA, Baier H. Pentylentetrazole induced changes in zebrafish behavior, neural activity and c-fos expression. *Neuroscience.* 2005; 131(3):759–68. Epub 2005/02/26. <https://doi.org/10.1016/j.neuroscience.2004.11.031> PMID: 15730879.
19. Stehr CM, Linbo TL, Incardona JP, Scholz NL. The developmental neurotoxicity of fipronil: notochord degeneration and locomotor defects in zebrafish embryos and larvae. *Toxicol Sci.* 2006; 92(1):270–8. Epub 2006/04/14. <https://doi.org/10.1093/toxsci/kfj185> PMID: 16611622.
20. Connaughton VP, Nelson R, Bender AM. Electrophysiological evidence of GABAA and GABAC receptors on zebrafish retinal bipolar cells. *Vis Neurosci.* 2008; 25(2):139–53. Epub 2008/04/30. <https://doi.org/10.1017/S0952523808080322> PMID: 18442437; PubMed Central PMCID: PMC35256003.
21. Del Bene F, Wyart C, Robles E, Tran A, Looger L, Scott EK, et al. Filtering of visual information in the tectum by an identified neural circuit. *Science.* 2010; 330(6004):669–73. Epub 2010/10/30. <https://doi.org/10.1126/science.1192949> PMID: 21030657; PubMed Central PMCID: PMC3243732.

22. Mack-Bucher JA, Li J, Friedrich RW. Early functional development of interneurons in the zebrafish olfactory bulb. *Eur J Neurosci*. 2007; 25(2):460–70. Epub 2007/02/08. <https://doi.org/10.1111/j.1460-9568.2006.05290.x> PMID: 17284187.
23. Hubbard JM, Böhm UL, Prendergast A, Tseng PB, Newman M, Stokes C, et al. Intraspinal Sensory Neurons Provide Powerful Inhibition to Motor Circuits Ensuring Postural Control during Locomotion. *Curr Biol*. 2016; 26(21):2841–53. Epub 2016/10/11. <https://doi.org/10.1016/j.cub.2016.08.026> PMID: 27720623.
24. Triller A, Rostaing P, Korn H, Legendre P. Morphofunctional evidence for mature synaptic contacts on the Mauthner cell of 52-hour-old zebrafish larvae. *Neuroscience*. 1997; 80(1):133–45. Epub 1997/09/01. PMID: 9252227.
25. Roy B, Ali DW. Multiple types of GABA<sub>A</sub> responses identified from zebrafish Mauthner cells. *Neuroreport*. 2014; 25(15):1232–6. Epub 2014/08/28. <https://doi.org/10.1097/WNR.000000000000258> PMID: 25162782.
26. Cocco A, Ronnberg AM, Jin Z, Andre GI, Vossen LE, Bhandage AK, et al. Characterization of the gamma-aminobutyric acid signaling system in the zebrafish (*Danio rerio* Hamilton) central nervous system by reverse transcription-quantitative polymerase chain reaction. *Neuroscience*. 2017; 343:300–21. Epub 2016/07/28. <https://doi.org/10.1016/j.neuroscience.2016.07.018> PMID: 27453477.
27. Hortopan GA, Dinday MT, Baraban SC. Spontaneous seizures and altered gene expression in GABA signaling pathways in a mind bomb mutant zebrafish. *J Neurosci*. 2010; 30(41):13718–28. Epub 2010/10/15. <https://doi.org/10.1523/JNEUROSCI.1887-10.2010> PMID: 20943912; PubMed Central PMCID: PMC2962868.
28. Gonzalez-Nunez V. Role of *gabra2*, GABA<sub>A</sub> receptor alpha-2 subunit, in CNS development. *Biochem Biophys Reports*. 2015; 3:11.
29. Baxendale S, Holdsworth CJ, Meza Santoscoy PL, Harrison MR, Fox J, Parkin CA, et al. Identification of compounds with anti-convulsant properties in a zebrafish model of epileptic seizures. *Dis Model Mech*. 2012; 5(6):773–84. Epub 2012/06/26. <https://doi.org/10.1242/dmm.010090> PMID: 22730455; PubMed Central PMCID: PMC3484860.
30. Thisse B, Thisse C. Fast Release Clones: A High Throughput Expression Analysis. *ZFIN Direct Data Submission*. 2004.
31. Thompson JD, Higgins DG, Gibson TJ. CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res*. 1994; 22(22):4673–80. Epub 1994/11/11. PMID: 7984417; PubMed Central PMCID: PMC308517.
32. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst Biol*. 2010; 59(3):307–21. Epub 2010/06/09. <https://doi.org/10.1093/sysbio/syq010> PMID: 20525638.
33. Kimmel CB, Ballard WW, Kimmel SR, Ullmann B, Schilling TF. Stages of embryonic development of the zebrafish. *Dev Dyn*. 1995; 203(3):253–310. <https://doi.org/10.1002/aja.1002030302> PMID: 8589427.
34. Thisse C, Thisse B. High-resolution in situ hybridization to whole-mount zebrafish embryos. *Nat Protoc*. 2008; 3(1):59–69. <https://doi.org/10.1038/nprot.2007.514> PMID: 18193022.
35. Mueller T, Wullmann MF. Atlas of early zebrafish brain development: a tool for molecular neurogenetics. Second edition. ed. xiii, 244 pages p.
36. Figueira M, Bayley P, Navratilova P, Becker TS, Wilson SW, Clarke JD. Morphogenesis underlying the development of the everted teleost telencephalon. *Neural Dev*. 2012; 7:32. Epub 2012/09/20. <https://doi.org/10.1186/1749-8104-7-32> PMID: 22989074; PubMed Central PMCID: PMC3520737.
37. Sabaliauskas NA, Foutz CA, Mest JR, Budgeon LR, Sidor AT, Gershenson JA, et al. High-throughput zebrafish histology. *Methods*. 2006; 39(3):246–54. Epub 2006/07/28. <https://doi.org/10.1016/j.ymeth.2006.03.001> PMID: 16870470.
38. Amores A, Force A, Yan YL, Joly L, Amemiya C, Fritz A, et al. Zebrafish hox clusters and vertebrate genome evolution. *Science*. 1998; 282(5394):1711–4. Epub 1998/11/30. PMID: 9831563.
39. Postlethwait JH, Yan YL, Gates MA, Horne S, Amores A, Brownlie A, et al. Vertebrate genome evolution and the zebrafish gene map. *Nat Genet*. 1998; 18(4):345–9. Epub 1998/04/16. <https://doi.org/10.1038/ng0498-345> PMID: 9537416.
40. Saint-Amant L, Drapeau P. Time course of the development of motor behaviors in the zebrafish embryo. *J Neurobiol*. 1998; 37(4):622–32. PMID: 9858263.
41. Bernhardt RR, Patel CK, Wilson SW, Kuwada JY. Axonal trajectories and distribution of GABAergic spinal neurons in wildtype and mutant zebrafish lacking floor plate cells. *J Comp Neurol*. 1992; 326(2):263–72. <https://doi.org/10.1002/cne.903260208> PMID: 1479075.



42. Bernhardt RR, Chitnis AB, Lindamer L, Kuwada JY. Identification of spinal neurons in the embryonic and larval zebrafish. *J Comp Neurol*. 1990; 302(3):603–16. <https://doi.org/10.1002/cne.903020315> PMID: 1702120.
43. Tsang SY, Ng SK, Xu Z, Xue H. The evolution of GABAA receptor-like genes. *Mol Biol Evol*. 2007; 24(2):599–610. Epub 2006/12/01. <https://doi.org/10.1093/molbev/msl188> PMID: 17135332.
44. Darlison MG, Pahal I, Thode C. Consequences of the evolution of the GABA(A) receptor gene family. *Cell Mol Neurobiol*. 2005; 25(3–4):607–24. Epub 2005/08/03. <https://doi.org/10.1007/s10571-005-4004-4> PMID: 16075381.
45. Martyniuk CJ, Aris-Brosou S, Drouin G, Cahn J, Trudeau VL. Early evolution of ionotropic GABA receptors and selective regimes acting on the mammalian-specific theta and epsilon subunits. *PLoS One*. 2007; 2(9):e894. Epub 2007/09/20. <https://doi.org/10.1371/journal.pone.0000894> PMID: 17878929; PubMed Central PMCID: PMC1975676.
46. Sinkkonen ST, Hanna MC, Kirkness EF, Korpi ER. GABA(A) receptor epsilon and theta subunits display unusual structural variation between species and are enriched in the rat locus ceruleus. *J Neurosci*. 2000; 20(10):3588–95. Epub 2000/05/11. PMID: 10804200.
47. Higashijima S, Mandel G, Fetcho JR. Distribution of prospective glutamatergic, glycinergic, and GABAergic neurons in embryonic and larval zebrafish. *J Comp Neurol*. 2004; 480(1):1–18. Epub 2004/10/30. <https://doi.org/10.1002/cne.20278> PMID: 15515020.
48. Kinkhabwala A, Riley M, Koyama M, Monen J, Satou C, Kimura Y, et al. A structural and functional ground plan for neurons in the hindbrain of zebrafish. *Proc Natl Acad Sci U S A*. 2011; 108(3):1164–9. Epub 2011/01/05. <https://doi.org/10.1073/pnas.1012185108> PMID: 21199947; PubMed Central PMCID: PMC3024665.
49. Downes GB, Waterbury JA, Granato M. Rapid in vivo labeling of identified zebrafish neurons. *Genesis*. 2002; 34(3):196–202. <https://doi.org/10.1002/gene.10120> PMID: 12395384.
50. Randlett O, Wee CL, Naumann EA, Nnaemeka O, Schoppik D, Fitzgerald JE, et al. Whole-brain activity mapping onto a zebrafish brain atlas. *Nat Methods*. 2015; 12(11):1039–46. Epub 2016/01/19. <https://doi.org/10.1038/nmeth.3581> PMID: 26778924; PubMed Central PMCID: PMC4710481.
51. Ronneberger O, Liu K, Rath M, Ruebeta D, Mueller T, Skibbe H, et al. ViBE-Z: a framework for 3D virtual colocalization analysis in zebrafish larval brains. *Nat Methods*. 2012; 9(7):735–42. Epub 2012/06/19. <https://doi.org/10.1038/nmeth.2076> PMID: 22706672.
52. Marquart GD, Tabor KM, Brown M, Strykowski JL, Varshney GK, LaFave MC, et al. A 3D Searchable Database of Transgenic Zebrafish Gal4 and Cre Lines for Functional Neuroanatomy Studies. *Front Neural Circuits*. 2015; 9:78. Epub 2015/12/05. <https://doi.org/10.3389/fncir.2015.00078> PMID: 26635538; PubMed Central PMCID: PMC4656851.
53. Kaeser GE, Rabe BA, Saha MS. Cloning and characterization of GABAA alpha subunits and GABAB subunits in *Xenopus laevis* during development. *Dev Dyn*. 2011; 240(4):862–73. Epub 2011/03/09. <https://doi.org/10.1002/dvdy.22580> PMID: 21384470; PubMed Central PMCID: PMC3071254.
54. Fetcho JR, Higashijima S, McLean DL. Zebrafish and motor control over the last decade. *Brain Res Rev*. 2008; 57(1):86–93. Epub 2007/09/11. <https://doi.org/10.1016/j.brainresrev.2007.06.018> PMID: 17825423; PubMed Central PMCID: PMC2237884.
55. Friedrich RW, Genoud C, Wanner AA. Analyzing the structure and function of neuronal circuits in zebrafish. *Front Neural Circuits*. 2013; 7:71. Epub 2013/05/01. <https://doi.org/10.3389/fncir.2013.00071> PMID: 23630467; PubMed Central PMCID: PMC3632777.
56. Hale ME, Katz HR, Peek MY, Fremont RT. Neural circuits that drive startle behavior, with a focus on the Mauthner cells and spiral fiber neurons of fishes. *J Neurogenet*. 2016; 30(2):89–100. Epub 2016/06/16. <https://doi.org/10.1080/01677063.2016.1182526> PMID: 27302612.
57. Griffin A, Krasniak C, Baraban SC. Advancing epilepsy treatment through personalized genetic zebrafish models. *Prog Brain Res*. 2016; 226:195–207. Epub 2016/06/22. <https://doi.org/10.1016/bs.pbr.2016.03.012> PMID: 27323944.
58. Cunliffe VT. Building a zebrafish toolkit for investigating the pathobiology of epilepsy and identifying new treatments for epileptic seizures. *J Neurosci Methods*. 2016; 260:91–5. Epub 2015/07/30. <https://doi.org/10.1016/j.jneumeth.2015.07.015> PMID: 26219659.
59. Kralic JE, Sidler C, Parpan F, Homanics GE, Morrow AL, Fritschy JM. Compensatory alteration of inhibitory synaptic circuits in cerebellum and thalamus of gamma-aminobutyric acid type A receptor alpha1 subunit knockout mice. *J Comp Neurol*. 2006; 495(4):408–21. Epub 2006/02/18. <https://doi.org/10.1002/cne.20866> PMID: 16485284.
60. Peng Z, Zhang N, Chandra D, Homanics GE, Olsen RW, Houser CR. Altered localization of the delta subunit of the GABAA receptor in the thalamus of alpha4 subunit knockout mice. *Neurochem Res*. 2014; 39(6):1104–17. Epub 2013/12/20. <https://doi.org/10.1007/s11064-013-1202-1> PMID: 24352815; PubMed Central PMCID: PMC4024081.