

Leptin and hypertension in obesity

Paco E Bravo¹
 Stephen Morse¹
 David M Borne²
 Erwin A Aguilar¹
 Efrain Reisin¹

¹Sections of Nephrology and

²Comprehensive Medicine,
 Department of Internal Medicine,
 Louisiana State University Health
 Sciences Center, New Orleans, LA,
 USA

Abstract: Leptin, a peptide discovered more than 10 years ago, decreases food intake and increases sympathetic nerve activity to both thermogenic and nonthermogenic tissue. Leptin was initially believed to be an anti-obesity hormone, owing to its metabolic effects. However, obese individuals, for unknown reasons, become resistant to the satiety and weight-reducing effect of the hormone, but preserve leptin-mediated sympathetic activation to nonthermogenic tissue such as kidney, heart, and adrenal glands. Leptin has been shown to influence nitric oxide production and natriuresis, and along with chronic sympathetic activation, especially to the kidney, it may lead to sodium retention, systemic vasoconstriction, and blood pressure elevation. Consequently, leptin is currently considered to play an important role in the development of hypertension in obesity.

Keywords: leptin, renal sympathetic nerve activity, blood pressure, obesity, selective leptin resistance, nitric oxide, natriuresis

Introduction

Obesity is considered a world health problem especially in western industrialized countries, where its incidence and prevalence are rising steadily, along with several comorbidities associated with it, such as hypertension, diabetes, dyslipidemia, atherosclerosis, and chronic renal failure. Among these, hypertension – a very important cardiovascular risk factor – has been observed in roughly 50% of obese individuals, which has led researchers to consider obesity as one of the most common causes of hypertension. Hence, hypertension in obesity has become a topic of extensive ongoing research.

Now, several mechanisms have been implicated in the association between obesity and hypertension, including activation of sympathetic nervous system, abnormal renal sodium handling, insulin resistance, and physical compression of the kidney (Haynes et al 1998).

In this respect, sympathetic activation appears to mediate at least part of the obesity-induced hypertension, and leptin, the adipocyte-derived hormone, has recently been postulated as one of the possible causes of this sympathetic activation in obesity.

Leptin is a 167 amino acid hormone discovered in 1994 that is almost exclusively produced by adipose tissue and possibly secreted by a constitutive mechanism. The effects of this peptide are mediated by receptors (Ob-R), most of them located in the hypothalamus, belonging to the class I cytokine receptor family. As of yet, 6 leptin receptor isoforms are known (Ahima and Flier 2000).

Leptin is considered a homeostatic hormone regulating food intake and body weight. Acting on the hypothalamic nuclei, leptin decreases appetite, and increases energy expenditure through sympathetic activation, which consequently decreases adipose tissue mass and body weight. The hormone levels are decreased during fasting and increased after several days of overfeeding as an effort to help regulate energy balance in humans. Due to latter homeostatic control mechanism, leptin is an anti-

Correspondence: Efrain Reisin
 Chief of the Section of Nephrology,
 Department of Internal Medicine,
 Louisiana State University Health
 Sciences Center 1542 Tulane Avenue,
 New Orleans, LA 70112, USA
 Tel +1 504 568 8655
 Fax +1 504 454 8218
 Email ereisi@lsuhsc.edu

obesity hormone, based on the hypothetical fact that high leptin levels would prevent the occurrence of obesity. Unfortunately, this is not the case, and so the strong correlation between serum leptin levels and body fat mass found in obese individuals now suggests the existence of an endogenous leptin-resistant mechanism in obesity (Considine et al 1996).

In addition to regulating food intake and adipose tissue mass, leptin has been found to be involved in cardiovascular physiological processes like sympathetic nerve system activation, renal hemodynamics, blood vessel tone, and blood pressure. This review will focus on the leptin-mediated sympathetic activation and its relevance to the mechanism of obesity-related hypertension.

Concept of selective leptin resistance

As stated above, obesity is associated with high leptin levels, reflecting the increased amount of adipose tissue in obese individuals. As a consequence, it has been postulated that the apparent loss of the anorexic and weight-reducing effects of leptin in obesity, is a result of a leptin-resistance mechanism. Nevertheless, obesity is associated with increased sympathetic nerve activity, and leptin has been proven to participate in autonomic nervous system control, in part, by increasing renal sympathetic nerve activity (RSNA). Therefore, it is contradictory that, in the presence of a leptin resistance state, the hormone can contribute to the sympathetic activation seen in obesity. This has led to the novel concept of selective leptin resistance, in which resistance appears to be primarily limited to the metabolic (satiety and weight-reducing) actions of leptin, sparing the renal sympathetic activation effects. This concept has emerged from observations made first in agouti yellow obese (Ay) mice (Correia et al 2002; Rahmouni et al 2002), and corroborated recently in a diet-induced obesity model (Rahmouni et al 2005). In these studies, leptin administered peripherally produced increase in RSNA in both obese and lean mice, but failed to decrease food intake and body weight in the obese mice to a similar degree as it did in their lean littermates. Interestingly, this same effect was observed when leptin was injected into the lateral ventricle of the brain (Rahmouni et al 2002, 2005). Here, leptin increased RSNA again, without changing significantly food intake and body weight in Ay and diet-induced obese mice, as compared with their respective lean littermates. These latter studies also served to demonstrate that is unlikely that leptin

resistance is due to an impaired leptin transport mechanism across the blood–brain barrier as intracerebroventricular administration of the hormone also failed to enhance food intake and weight loss. Moreover, the increased RSNA achieved by central leptin administration was not accompanied by changes in plasma leptin levels, suggesting that leptin-induced RSNA was centrally-mediated.

In agreement with these observations, Munzberg et al (2004) recently described that leptin resistance is not global within the hypothalamus, and found that the arcuate nuclei, which contains leptin-mediated anorexigenic neurons, becomes resistant to leptin in very early stages of diet-induced obesity in mice. On the contrary, other hypothalamic and extrahypothalamic sites appear to continue relatively sensitive to the actions of leptin, including the ventromedial hypothalamic nuclei, which has been previously shown to mediate leptin-induced sympathetic nerve activation (Sato et al 1999).

This hypothesis, however, must be proved in human studies as well, and if true, it would help explain how leptin could contribute to sympathetic activation and hypertension despite the fact that there is resistance to the satiety and weight-reducing effects of leptin in obesity.

Renal effects of leptin: the role of the sympathetic nervous system and nitric oxide

In the kidney, leptin may affect blood pressure mainly by two opposing ways: the first is through renal sympathetic activation (Haynes, Sivitz, et al 1997), and the second is related to nitric oxide (NO) synthesis (Vecchione et al 2002). The renal effect of leptin also depends on the exposure time to the hormone. As it has been observed, acutely administered leptin usually does not increase blood pressure in animal or human studies, but when given for longer periods of time, leptin does elevate blood pressure in animal models.

Acute effects

Traditionally, obesity-induced renal sympathetic activation has been thought to contribute to the development of hypertension by enhancing sodium retention, since renal denervation attenuates the antinatriuretic and hypertensive effect of obesity in dogs (Kassab et al 1995). Although systemic administration of leptin stimulates sympathetic nerve activity to the kidney (Haynes, Morgan, et al 1997), if infused for short

periods of time, leptin produces increased sodium excretion and urine output with no changes in blood pressure in lean animals (Jackson and Li 1997; Villarreal et al 1998; Beltowski, Wojcicka, Gorny, et al 2002; Beltowski, Jochem, et al 2004). The increased natriuresis and fractional sodium excretion are not accompanied by substantial changes in glomerular filtration rate or the renin–aldosterone axis suggesting a decreased sodium tubular transport. Beltowski, Wojcicka, et al (2002) described a decreased sodium–potassium exchanger (Na/K-ATPase) activity in the renal medulla following acute leptin administration. New evidence suggests that leptin stimulates systemic NO release which opposes the pressor and antinatriuretic effect of leptin-induced sympathetic activation (Beltowski, Wojcicka, Borkowska, et al 2002; Beltowski, Jochem, et al 2004). In this regard, Vecchione et al (2002) demonstrated that leptin stimulates NO production in endothelial cells and blood vessels. In other studies, leptin evoked a hypotensive effect when the sympathetic nervous system output was blocked pharmacologically, implying a likely leptin-mediated vasorelaxant effect (Fruhbeck 1999; Lembo et al 2000).

It has been demonstrated that endogenously-produced NO inhibits renal sodium reabsorption in most of the tubule segments apparently through a decrease in the activity of both apical transporters and basolateral Na/K-ATPases (Ortiz and Garvin 2002). Likewise, Villarreal et al (2004) found that chronic inhibition of NO synthesis impairs the acute leptin-mediated natriuretic effect in normotensive lean rats. The kidney has been shown to express high amounts of leptin receptors (Serradeil-Le Gal et al 1997), yet it is not clear whether leptin evokes a direct effect on leptin receptors or indirectly via induction of NO release in renal tubules.

Interestingly, the short-term leptin-induced natriuretic effect seen in lean animals was attenuated in obese rats and was even blunted in spontaneously hypertensive rats (SHR) (Villarreal et al 1998; Beltowski, Wojcicka, Gorny, et al 2002). This suggests a certain level of peripheral leptin resistance, probably as a result of several factors including renal receptor down-regulation induced by long-standing hyperleptinemia (Coatmellec-Taglioni et al 2003) and/or the presence of chronic antinatriuretic mechanisms, typical of these states, such as an enhanced RSNA and renin–angiotensin system (Villarreal et al 1998; Beltowski, Wojcicka, Gorny, et al 2002), which may have overcome leptin-mediated natriuresis. In agreement with this statement, Villarreal et al (2000) found that SHR after renal denervation

recovered the sodium excretory capacity in response to acute leptin administration.

Chronic effects

Conversely, long-term leptin administration increases both RSNA and blood pressure. Recent data indicate that chronic hyperleptinemia decreases natriuresis and urinary excretion of NO metabolites (NOx) (Beltowski, Jamroz-Wisniewska, et al 2004). In humans, a negative correlation between leptin and NOx was observed recently in obese hypertensive individuals (Golan et al 2002). Apparently, long-term hyperleptinemic states like obesity increase the level of systemic and intrarenal oxidative stress, leading to NO deficiency (Beltowski, Wojcicka, et al 2004). Leptin, chronically, would reduce natriuresis by up-regulation of the Na/K-ATPase pump, especially in the renal medulla, thereby promoting sodium retention (Beltowski, Jamroz-Wisniewska, et al 2004). This antinatriuretic response to chronic hyperleptinemia, could be either secondary to renal sympathetic activation (Echtenkamp and Dandridge 1989; Kirchheim et al 1989; Beach 1992) and/or deficiency of systemic and intrarenal NO production (Beltowski, Wojcicka, et al 2004). Chronic inhibition of NO synthesis in the renal medulla may reduce medullary blood flow by 30%, which is associated with sodium retention and high blood pressure (Mattson et al 1994).

Supporting these observations, Bickel et al (2001) reported an increased number of different sodium tubular transporters in the kidney of obese rats. These animals exhibited a decreased natriuretic effect after a saline load when compared with their lean age mates. In addition, thiazide-sensitive Na/Cl cotransporters' abundance has been shown to be associated with inhibition of NO synthesis.

Clinical relevance

Chronic hyperleptinemia, as mentioned, augments the blood pressure by means of different mechanisms in animal models. In humans, there have been several studies attempting to link leptin to hypertension (Figure 1). Serum leptin levels are elevated in obesity due to increase amount of adipose tissue which is the main source of the hormone and possibly secondary to some degree of central resistance to its action. A number of studies have found leptin to be positively correlated with systolic and diastolic blood pressure in both obese (Kunz et al 2000; Golan et al 2002; Itoh et al 2002; Al-Hazimi and Syiamic 2004; Canatan et al 2004a; Schutte et al 2005) and non-obese individuals (Uckaya et al 1999; Adamczak et al 2000; Takizawa et al

Renal and Blood Pressure Effect of Leptin




Figure 1 Renal and blood pressure effect of leptin. Adapted from Hall et al 2001. Copyright © 2001. Reproduced with permission from Hall JE, Hildebrandt DA, Kuo J. 2001. Obesity hypertension: role of leptin and sympathetic nervous system. *Am J Hypertension*, 14(6 Pt 2):103S-115S.

2001; Barba et al 2003; Canatan et al 2004b). Al-Hazimi and Syiamic (2004), for example, found that serum leptin and angiotensin II levels were strong predictors of elevated blood pressure in obese women. Likewise, other investigators (Kunz et al 2000; Golan et al 2002; Canatan et al 2004b) reported higher leptin levels in obese hypertensives compared with obese normotensive individuals, even after controlling for confounders such as age and body mass index (BMI). In another study leptin correlated stronger with systolic and diastolic blood pressure than did with BMI (or waist circumference) in untreated male adults (Barba et al 2003). Moreover, one study found that after a 3-month weight reduction program, leptin

continued to correlate positively with mean arterial blood pressure in the hypertensive, but not in the normotensive obese group (Itoh et al 2002). Interestingly, leptin has been found to be elevated in several hypertensive states of pregnancy as well (Anato et al 2000; Vitoratos et al 2001). For instance, Vitoratos et al (2001) reported increased leptin levels in pre-eclamptics compared with normotensive pregnant women of the same gestational age. Of note is also the observation that a positive correlation between leptin and platelet count levels exist in pre-eclamptic women (Anato et al 2000).

In disagreement to some of these data, one study found that leptin levels were higher in hypertensives than in

normotensive African Americans, but once these individuals were adjusted for obesity, no significant relationship was further observed between leptin and blood pressure (El-Gharbawy et al 2002).

In summary, it seems evident that serum leptin levels are significantly elevated in obese hypertensive individuals when compared with obese normotensives. However this relationship between leptin and blood pressure may not always be as evident after controlling for BMI in some subjects, making the assumption of leptin as a one of the potential causes of hypertension in obesity, less consistent.

Possible treatment considerations

The elevated leptin concentrations seen in obese hypertensive humans are mainly due to abdominal adipose tissue secretion. Current studies have elucidated that some antihypertensives might be more relevant than others in terms of reducing leptin levels in obesity. For example, angiotensin II has recently been found to stimulate leptin production in human adipose tissue (Skurk et al 2005), possibly by means of activation of the angiotensin II type 1 receptor subtype. This effect was completely abolished when the angiotensin receptor blocker (ARB) candesartan was employed prior to angiotensin II administration in human fat cells. In agreement with this observation, valsartan, another ARB, aside from lowering blood pressure, decreased leptin levels and BMI in obese individuals, when compared with the calcium channel blocker (CCB) felodipine (Fogari et al 2005). Similarly, the angiotensin-converting enzyme (ACE) inhibitor, enalapril, in combination with a weight reduction program, evidenced the greatest benefits in terms of weight loss and diminution of plasma norepinephrine, insulin, and leptin levels in comparison with control groups treated with weight reduction program alone or combined with the CCB amlodipine (Masuo et al 2001). In another study, the beta-blocker (BB) pindolol showed a marked suppressive effect on serum leptin levels, not seen in hypertensive individuals on perindopril, or felodipine (Ficek et al 2002).

In contrast to these findings, Sonmez et al (2001) in an earlier study, found no clear effect of the ARB Losartan on leptin levels in young hypertensive individuals despite its hypotensive action. Likewise, both enalapril and clonidine reduced heart sympathetic activity and blood pressure in another clinical trial, but failed to decrease serum leptin levels in normotensive obese and non-obese subjects after 7 days of treatment (Amador et al 2004).

It is worth mentioning that obesity is associated with sodium retention and increased extracellular volume inducing hypervolemia and increased cardiac output, leading to the activation of the renin–angiotensin and sympathetic nervous system; all of these factors contributing ultimately to the development of hypertension and eccentric-concentric left ventricular hypertrophy: changes that may induce congestive heart failure, arrhythmia, and sudden death (Figure 2) (Zhang and Reisin 2000).

Although the studies are not yet consistent, it seems that pharmacologic suppression of the sympathetic nervous system, the renin–angiotensin system and, lately hyperleptinemia with ACE inhibitors, ARB and BB, may resemble the blood pressure reduction effect induced by weight loss. Diuretics, on the other hand, by eliminating urinary sodium and extracellular fluid, might also play a significant role in the treatment of hypertension in obesity, as these individuals are known for being sodium retainers and hypervolemics. Nevertheless, BBs and diuretics may affect insulin sensitivity and cause hyperglycemia and so their use should be carried out with close monitoring of glycemia in this patient population.

Summary

In summary, leptin the adipose tissue-derived hormone by way of distinct neurochemical pathways stimulates sympathetic nerve activity in thermogenic and nonthermogenic tissue, affecting the metabolic and cardiovascular system respectively. Leptin, acutely, could have a dual influence on blood pressure control, in which the net effect would depend on the balance between the pressor action through activation of the sympathetic nervous system and a possible natriuretic and peripheral vasorelaxant effect of the hormone on the renal tubules and endothelium. In contrast, chronic hyperleptinemia may lead to abnormal renal sodium retention and vasoconstriction associated with renal sympathetic activation and NO deficiency, both contributing to pressure elevation in obese individuals, who may develop resistance to the satiety effect of leptin with preservation of the cardiovascular effect. Treatment of these individuals might be focused in overcoming the hemodynamic alterations seen in obesity, such as antinatriuresis and overactivity of the sympathetic and renin–angiotensin systems.




Figure 2 Potential pathways in which obesity can cause cardiovascular dysfunction. Effects of obesity-hypertension on the heart. Adapted from Zhang et al 2000.

Copyright © 2004. Reproduced with permission from Zhang R, Reisin E. 2000. Obesity-hypertension: the effects on cardiovascular and renal systems. *Am J Hypertens*, 13:1308-14.

Abbreviations: BP, blood pressure; CHF, congestive heart failure; CO, cardiac output; LV, left ventricular; NP, natriuretic peptides; RAS, renin-angiotensin system; SNS, sympathetic nervous system.

References

- Adamczak M, Kokot F, Wiecek AW. 2000. Relationship between plasma renin profile and leptinaemia in patients with essential hypertension. *J Hum Hypertens*, 14:503-9.
- Ahima RS, Flier JS. 2000. Leptin. *Ann Rev Physiol*, 62:413-37.
- Al-Hazimi AM, Syiamic AY. 2004. Relationship between plasma angiotensinII, leptin and arterial blood pressure. *Saudi Med J*, 25:1193-8.
- Amador N, Perez-Luque E, Malacara JM, et al. 2004. Leptin and heart sympathetic activity in normotensive obese and non-obese subjects. *Ital Heart J*, 5:29-35.
- Anato V, Garmendia JV, Bianco NE, et al. 2000. Serum leptin levels in different types of hypertension during pregnancy. *Res Commun Mol Pathol Pharmacol*, 108:147-53.
- Barba G, Russo O, Siani A, et al. 2003. Plasma leptin and blood pressure in men: graded association independent of body mass and fat pattern. *Obes Res*, 11:160-6.
- Beach RE. 1992. Renal nerve-mediated proximal tubule solute reabsorption contributes to hypertension in spontaneously hypertensive rats. *Clin Exp Hypertens A*, 14:685-97.
- Beltowski J, Wojcicka G, Gorny D, et al. 2002. Human leptin administered intraperitoneally stimulates natriuresis and decreases renal medullary Na⁺, K⁺-ATPase activity in the rat –impaired effect in dietary-induced obesity. *Med Sci Monit*, 8:BR221-9.
- Beltowski J, Jamroz-Wisniewska A, Borkowska E, et al. 2004. Up-regulation of renal Na⁽⁺⁾, K⁽⁺⁾-ATPase: the possible novel mechanism of leptin-induced hypertension. *Pol J Pharmacol*, 56:213-22.
- Beltowski J, Jochem J, Wojcicka G, et al. 2004. Influence of intravenously administered leptin on nitric oxide production, renal hemodynamics and renal function in the rat. *Regul Pept*, 120:59-67.
- Beltowski J, Wojcicka G, Borkowska E. 2002. Human leptin stimulates systemic nitric oxide production in the rat. *Obes Res*, 10:939-46.
- Beltowski J, Wojcicka G, Marciniak A, et al. 2004. Oxidative stress, nitric oxide production, and renal sodium handling in leptin-induced hypertension. *Life Sci*, 74:2987-3000.
- Bickel CA, Verbalis JG, Knepper MA, et al. 2001. Increased renal Na-K-ATPase, NCC, and beta-ENaC abundance in obese Zucker rats. *Am J Physiol Renal Physiol*, 281:F639-48.
- Canatan H, Bakan I, Akbulut M, et al. 2004a. Comparative analysis of plasma leptin levels in both genders of patients with essential hypertension and healthy subjects. *Endocr Res*, 30:95-105.
- Canatan H, Bakan I, Akbulut M, et al. 2004b. Relationship among levels of leptin and zinc, copper, and zinc/copper ratio in plasma of patients with essential hypertension and healthy normotensive subjects. *Biol Trace Elem Res*, 100:117-23.
- Coatmellec-Taglioni G, Dausse JP, Giudicelli Y, et al. 2003. Sexual dimorphism in cafeteria diet-induced hypertension is associated with gender-related difference in renal leptin receptor down-regulation. *J Pharmacol Exp Ther*, 305:362-7.

- Considine RV, Sinha MK, Heiman ML, et al. 1996. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med*, 334:292-5.
- Correia ML, Haynes WG, Rahmouni K, et al. 2002. The concept of selective leptin resistance: evidence from agouti yellow obese mice. *Diabetes*, 51:439-42.
- Echtenkamp SF, Dandridge PF. 1989. Influence of renal sympathetic nerve stimulation on renal function in the primate. *Am J Physiol*, 257:F204-9.
- El-Gharbawy AH, Kotchen JM, Grim CE, et al. 2002. Gender-specific correlates of leptin with hypertension-related phenotypes in African Americans. *Am J Hypertens*, 15:989-93.
- Ficek J, Kokot F, Chudek J, et al. 2002. Influence of antihypertensive treatment with perindopril, pindolol or felodipin on plasma leptin concentration in patients with essential hypertension. *Horm Metab Res*, 34:703-8.
- Fogari R, Derosa G, Zoppi A, et al. 2005. Comparison of the effects of valsartan and felodipine on plasma leptin and insulin sensitivity in hypertensive obese patients. *Hypertens Res*, 28:209-14.
- Fruhbeck G. 1999. Pivotal role of nitric oxide in the control of blood pressure after leptin administration. *Diabetes*, 48:903-8.
- Golan E, Tal B, Dror Y, et al. 2002. Reduction in resting metabolic rate and ratio of plasma leptin to urinary nitric oxide: influence on obesity-related hypertension. *Isr Med Assoc J*, 4:426-30.
- Hall JE, Hildebrandt DA, Kuo J. 2001. Obesity hypertension: role of leptin and sympathetic nervous system. *Am J Hypertension*, 14(6 Pt 2):103S-115S.
- Haynes WG, Morgan DA, Walsh SA, et al. 1997. Receptor-mediated regional sympathetic nerve activation by leptin. *J Clin Invest*, 100:270-8.
- Haynes WG, Morgan DA, Walsh SA, et al. 1998. Cardiovascular consequences of obesity: role of leptin. *Clin Exp Pharmacol Physiol*, 25:65-9.
- Haynes WG, Sivitz WI, Morgan DA, et al. 1997. Sympathetic and cardiorenal actions of leptin. *Hypertension*, 30:619-23.
- Itoh K, Imai K, Masuda T, et al. 2002. Relationship between changes in serum leptin levels and blood pressure after weight loss. *Hypertens Res*, 25:881-6.
- Jackson EK, Li P. 1997. Human leptin has natriuretic activity in the rat. *Am J Physiol*, 272:F333-8.
- Kassab S, Kato T, Wilkins FC, et al. 1995. Renal denervation attenuates the sodium retention and hypertension associated with obesity. *Hypertension*, 25:893-7.
- Kirchheim H, Ehmke H, Persson P. 1989. Sympathetic modulation of renal hemodynamics, renin release and sodium excretion. *Klin Wochenschr*, 67:858-64.
- Kunz I, Schorr U, Klaus S, et al. 2000. Resting metabolic rate and substrate use in obesity hypertension. *Hypertension*, 36:26-32.
- Lembo G, Vecchione C, Fratta L, et al. 2000. Leptin induces direct vasodilation through distinct endothelial mechanisms. *Diabetes*, 49:293-7.
- Masuo K, Mikami H, Ogihara T, et al. 2001. Weight reduction and pharmacologic treatment in obese hypertensives. *Am J Hypertens*, 14:530-8.
- Mattson DL, Lu S, Nakanishi K, et al. 1994. Effect of chronic renal medullary nitric oxide inhibition on blood pressure. *Am J Physiol*, 266:H1918-26.
- Munzberg H, Flier JS, Bjorbaek C. 2004. Region-specific leptin resistance within the hypothalamus of diet-induced obese mice. *Endocrinology*, 145:4880-9.
- Ortiz PA, Garvin JL. 2002. Role of nitric oxide in the regulation of nephron transport. *Am J Physiol Renal Physiol*, 282:F777-84.
- Rahmouni K, Haynes WG, Morgan DA, et al. 2002. Selective resistance to central neural administration of leptin in agouti obese mice. *Hypertension*, 39:486-90.
- Rahmouni K, Morgan DA, Morgan GM, et al. 2005. Role of selective leptin resistance in diet-induced obesity hypertension. *Diabetes*, 54:2012-18.
- Satoh N, Ogawa Y, Katsuura G, et al. 1999. Sympathetic activation of leptin via the ventromedial hypothalamus: leptin-induced increase in catecholamine secretion. *Diabetes*, 48:1787-93.
- Schutte R, Huisman HW, Schutte AE, et al. 2005. Leptin is independently associated with systolic blood pressure, pulse pressure and arterial compliance in hypertensive African women with increased adiposity: the POWIRS study. *J Hum Hypertens*, 19:535-41.
- Serradeil-Le Gal C, Raufaste D, Brossard G, et al. 1997. Characterization and localization of leptin receptors in the rat kidney. *FEBS Lett*, 404:185-91.
- Skurk T, van Harmelen V, Blum WF, et al. 2005. Angiotensin II promotes leptin production in cultured human fat cells by an ERK1/2-dependent pathway. *Obes Res*, 13:969-73.
- Sonmez A, Kisa U, Uckaya G, et al. 2001. Effects of losartan treatment on T-cell activities and plasma leptin concentrations in primary hypertension. *J Renin Angiotensin Aldosterone Syst*, 2:112-16.
- Takizawa H, Ura N, Saitoh S, et al. 2001. Gender difference in the relationships among hyperleptinemia, hyperinsulinemia, and hypertension. *Clin Exp Hypertens*, 23:357-68.
- Uckaya G, Ozata M, Sonmez A, et al. 1999. Plasma leptin levels strongly correlate with plasma renin activity in patients with essential hypertension. *Horm Metab Res*, 31:435-8.
- Vecchione C, Maffei A, Colella S, et al. 2002. Leptin effect on endothelial nitric oxide is mediated through Akt-endothelial nitric oxide synthase phosphorylation pathway. *Diabetes*, 51:168-73.
- Villarreal D, Reams G, Freeman RH. 2000. Effects of renal denervation on the sodium excretory actions of leptin in hypertensive rats. *Kidney Int*, 58:989-94.
- Villarreal D, Reams G, Freeman RH, et al. 1998. Renal effects of leptin in normotensive, hypertensive, and obese rats. *Am J Physiol*, 275:R2056-60.
- Villarreal D, Reams G, Samar H, et al. 2004. Effects of chronic nitric oxide inhibition on the renal excretory response to leptin. *Obes Res*, 12:1006-10.
- Vitoratos N, Chrystodoulacos G, Kouskouni E, et al. 2001. Alterations of maternal and fetal leptin concentrations in hypertensive disorders of pregnancy. *Eur J Obstet Gynecol Reprod Biol*, 96:59-62.
- Zhang R, Reisin E. 2000. Obesity-hypertension: the effects on cardiovascular and renal systems. *Am J Hypertens*, 13:1308-14.

