

Ventilation time and prognosis after stroke thrombectomy: the shorter, the better!

S. Fandler-Höfler^a , S. Heschl^b, M. Kneihsl^a, P. Argüelles-Delgado^b, K. Niederkorn^a, A. Pichler^a, H. Deutschmann^c, F. Fazekas^a, A. Berghold^d, C. Enzinger^{a,c} and T. Gatttringer^{a,c} 

^aDepartment of Neurology, Medical University of Graz, Graz; ^bDepartment of Anaesthesiology and Intensive Care Medicine, Medical University of Graz, Graz; ^cDepartment of Radiology, Division of Neuroradiology, Vascular and Interventional Radiology, Medical University of Graz, Graz; and ^dInstitute for Medical Informatics, Statistics and Documentation, Medical University of Graz, Graz, Austria

Keywords:

endovascular procedures, intensive care, mechanical ventilation, neurocritical care, pneumonia, prognosis, stroke, thrombectomy

Received 20 December 2019
Accepted 15 February 2020

European Journal of Neurology 2020, **27**: 849–855

doi:10.1111/ene.14178

Background and purpose: The aim was to investigate the clinical impact of the duration of artificial ventilation in stroke patients receiving mechanical thrombectomy (MT) under general anaesthesia.

Methods: All consecutive ischaemic stroke patients who had been treated at our centre with MT for anterior circulation large vessel occlusion under general anaesthesia were identified over an 8-year period. Ventilation time was analysed as a continuous variable and patients were grouped into extubation within 6 h ('early'), 6–24 h ('delayed') and > 24 h ('late'). Favourable outcome was defined as modified Rankin Scale scores of 0–2 at 3 months post-stroke. Pneumonia rate and reasons for prolonged ventilation were also assessed.

Results: Amongst 447 MT patients (mean age 69.1 ± 13.3 years, 50.1% female), the median ventilation time was 3 h. 188 (42.6%) patients had a favourable 3-month outcome, which correlated with shorter ventilation time (Spearman's rho 0.39, $P < 0.001$). In patients extubated within 24 h, early compared to delayed extubation was associated with improved outcome (odds ratio 2.40, 95% confidence interval 1.53–3.76, $P < 0.001$). This was confirmed in multivariable analysis ($P = 0.01$). A longer ventilation time was associated with a higher rate of pneumonia during neurointensive care unit/stroke unit stay (early/delayed/late extubation: 9.6%/20.6%/27.7%, $P < 0.01$). Whilst stroke-associated complications represented the most common reasons for late extubation (> 24 h), delayed extubation (6–24 h) was associated with admission outside of core working hours ($P < 0.001$).

Conclusions: Prolonged ventilation time after stroke thrombectomy independently predicts unfavourable outcome at 3 months and is associated with increased pneumonia rates. Therefore, extubation should be performed as early as safely possible.

Background

Mechanical thrombectomy (MT) has become the recommended treatment for acute ischaemic stroke caused by anterior circulation large vessel occlusion [1]. Currently, there is no consensus whether this

procedure should be performed under a specific form of anaesthesia, as previous studies have provided conflicting evidence whether general anaesthesia (GA), conscious sedation or local anaesthesia lead to better functional neurological outcome [2–4]. Therefore, all three methods are frequently used in clinical routine [5].

General anaesthesia with intubation offers the advantage of full patient immobilization, which might be associated with improved angiographic image

Correspondence: T. Gatttringer, Department of Neurology, Medical University of Graz, Auenbruggerplatz 22, Graz A-8036, Austria (tel.: +43/316/385 80231; fax: +43/316/385 16808; e-mail: thomas.gatttringer@medunigraz.at).

quality and shorter procedure time. In contrast, GA and intubation could delay time to groin puncture and consecutively vessel recanalization. GA might also increase the risk of relevant blood pressure drops, thereby impeding cerebral blood flow due to vasodilatory side effects of most anaesthetic drugs, as well as (ventilator-associated) pneumonia [6].

In general, longer ventilation time augments the risk for both of the latter complications in neurocritical care patients [7,8]. However, rash extubation may cause complications such as blood pressure surges, aspiration and respiratory difficulties. To date, there is no guideline regarding the optimal timing of extubation after MT. Only one retrospective study addressed this topic and found worse outcome in patients who were extubated after more than 24 h. Interestingly, the authors did not identify outcome differences between patients who were extubated at different time points within 24 h, but this might be explained by the rather small sample size ($n = 103$) of their study cohort [9].

In the present study, the clinical impact of ventilation time was investigated in a large cohort of consecutive patients undergoing MT for anterior circulation large vessel occlusion. It was hypothesized that a longer ventilation time (particularly >24 h) would be associated with a higher pneumonia rate and unfavourable functional outcome at 3 months. Furthermore, it was specifically analysed whether 'early' (<6 h) compared to 'delayed' extubation (6–24 h) had an effect on the clinical prognosis, and what reasons caused delays in extubation after MT.

Methods

For this cohort study, all consecutive patients ≥ 18 years who had received MT because of anterior circulation large vessel occlusion stroke (occlusion of the internal carotid and/or middle cerebral artery in the M1 or M2 segments) between January 2011 and April 2019 were identified. At our centre, MT is routinely performed by interventional radiologists using stent retrievers under GA provided by neuroanaesthesiologists. Patients were excluded from this study if MT was done under conscious sedation ($n = 15$) or if a subsequent decompressive hemicraniectomy was performed ($n = 4$). A study flowchart is shown in Fig. 1.

Clinical data including the duration of ventilation, the exact time point of extubation and early stroke complications within the neurointensive care unit or stroke unit stay such as pneumonia were prospectively and electronically recorded as part of an ongoing thrombectomy cohort registry at our centre [10]. Pneumonia was defined according to established criteria: the presence of an infiltrate on chest X-ray, plus

at least two of fever $>38^{\circ}\text{C}$, leucocytosis or leukopenia, purulent secretions [11,12].

The time from intubation to extubation was analysed as a continuous variable. Furthermore, patients were trichotomized according to three predefined subgroups in order to depict common clinical scenarios: (i) 'early' extubation within 6 h, in order to encompass all patients extubated directly after the procedure in the angio suite or within the first hours of admission at the neurointensive care unit; (ii) 'delayed' extubation (6–24 h); and (iii) 'late' extubation (>24 h).

The primary outcome variable was defined as functional neurological outcome at 3 months according to the modified Rankin Scale (mRS), dichotomized into favourable (scores of 0–2) versus unfavourable (scores 3–6) outcome. Three-month mRS scores were rated by experienced stroke physicians during an in-person visit at the stroke outpatient department or by telephone interview if the former was not possible.

The study was approved by the ethics committee of the Medical University of Graz. Because this was a retrospective registry study, individual consent was waived.

Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics for Windows, Version 25.0 (IBM Corp., Armonk, NY, USA). Normally distributed continuous variables were compared by the unpaired Student's *t* test; for other distributions the Mann–Whitney *U* test was used. For categorical variables, the chi-squared test was used. Furthermore, a binary multivariable regression model for predictors of favourable outcome at 3 months was calculated entering all baseline variables with $P < 0.1$ in the univariable analysis; then stepwise backward elimination was performed using a threshold of $P < 0.1$. *P* values equal to or less than 0.05 were considered statistically significant.

Results

Patient characteristics and outcome

In all, 447 patients (mean age 69.1 ± 13.3 years, 50.1% female) were identified and in 441 3-month follow-up data were available. The majority of patients had an M1-segment occlusion of the middle cerebral artery (65.3%); 58.6% had been treated with intravenous thrombolysis before MT. Median National Institutes of Health Stroke Scale (NIHSS) at admission was 15 (interquartile range 11–18) and successful recanalization (thrombolysis in cerebral infarction grades 2b–3) was achieved in 88.5% of the patients (Table 1).

At 3 months post-stroke 188 patients (42.6%) had a favourable outcome (mRS 0–2). In univariable analysis,




Figure 1 Study flowchart. [Colour figure can be viewed at wileyonlinelibrary.com]

higher age, higher NIHSS at admission, the presence of arterial hypertension, chronic heart disease, diabetes mellitus, atrial fibrillation and unsuccessful recanalization were associated with unfavourable functional outcome at 3 months (Table 1).

Ventilation time

Median ventilation time was 3 h (range 1–530); seven patients (1.6%) were admitted to our hospital already intubated by prehospital emergency physicians or in the referring hospital. The majority of patients (85.5%) were extubated within 24 h; only 65 patients (14.5%) had late extubation (>24 h). 258 patients (57.7%) were extubated early (within 6 h of intubation), whilst 124 patients (27.7%) had delayed extubation (6–24 h, Table 2).

Time of mechanical ventilation correlated with higher mRS values at 3 months (Spearman's rho 0.39, $P < 0.001$). Similarly, patients with favourable outcome (mRS 0–2) at 3 months had lower median ventilation times (2 h, interquartile range 1–6, vs. 11 h, interquartile range 2–24, in patients with mRS 3–6; $P < 0.001$).

Differences between early, delayed and late extubation

Patients with late extubation (>24 h) more frequently had concomitant heart disease (30.8% vs. 17.5%, $P = 0.01$), more severe strokes (median NIHSS 17 vs.

14, $P < 0.001$), more frequently received intravenous thrombolysis (70.8% vs. 56.2%, $P = 0.03$) and had lower rates of successful recanalization after MT (76.6% vs. 90.5%, $P = 0.001$). Favourable functional 3-month outcome was rather rare in this subgroup (7.8%, Table 2).

Analysing only those patients who had been extubated within 24 h, patients with early extubation (within 6 h) had a higher frequency of stroke of unknown onset (24.8% vs. 15.3%, $P = 0.04$) and slightly lower NIHSS scores at admission (median 14 vs. 15, $P < 0.001$) compared to patients with delayed extubation (6–24 h, Table 2). Early extubation was associated with favourable outcome [odds ratio (OR) 2.40, 95% confidence interval (CI) 1.53–3.76, $P < 0.001$].

In a multivariable model investigating the effect of early versus delayed extubation on functional neurological outcome and entering all outcome-relevant baseline variables with a P value of <0.1 in univariable analysis (Table 1), hypertension, atrial fibrillation, chronic heart disease, occlusion location and periprocedural dissection were removed using stepwise backward elimination.

Early extubation remained a significant predictor of favourable functional outcome at 3 months (OR 1.93, 95% CI 1.15–3.24, $P = 0.01$), together with successful recanalization (OR 6.97, 95% CI 2.38–20.4). Age (OR

Table 1 Clinical data and timing of extubation in the study cohort regarding 3-month outcome

	Study cohort <i>n</i> = 441	mRS 0–2 <i>n</i> = 188 (42.6%)	mRS 3–6 <i>n</i> = 253 (57.4%)	<i>P</i> value
Clinical data				
Age (years, mean ± SD)	69.1 ± 13.3	64.3 ± 13.4	72.4 ± 12.2	<0.001
Male sex	220 (49.9%)	96 (51.0%)	124 (49.0%)	0.67
Arterial hypertension	307 (69.6%)	106 (56.4%)	201 (79.4%)	<0.001
Dyslipidaemia	98 (20.7 %)	39 (20.7%)	59 (23.3%)	0.52
Chronic heart disease ^a	84 (19.0 %)	26 (13.8%)	58 (22.9%)	0.02
Diabetes mellitus	73 (16.6%)	19 (10.1%)	54 (21.3%)	0.002
Atrial fibrillation	182 (41.3%)	60 (31.9%)	122 (48.2%)	0.001
Stroke of unknown symptom onset	90 (20.4%)	35 (18.6%)	55 (21.7%)	0.42
Pre-stroke mRS (median, IQR)	0 (0–0)	0 (0–0)	0 (0–0)	<0.001
NIHSS at admission (median, IQR)	15 (11–18)	13 (9–16)	16 (13–19)	<0.001
MCA/M1 occlusion	288 (65.3%)	122 (64.9%)	166 (65.6%)	0.88
MCA/M2 occlusion	59 (13.4%)	32 (17.0%)	27 (10.7%)	0.06
Intracranial ICA occlusion	93 (21.1%)	31 (16.5%)	62 (24.5%)	0.05
Acute stroke treatment				
Intravenous thrombolysis	258 (58.6%)	118 (62.8%)	140 (55.6%)	0.13
Symptom onset to groin puncture (min, median, IQR)	200 (160–247)	200 (158–244)	200 (160–252)	0.61
Symptom onset to recanalization (min, median, IQR)	260 (215–310)	250 (204–303)	266 (225–315)	0.42
Successful recanalization (TICI 2b–3)	385 (88.5%)	181 (97.3%)	204 (81.9%)	<0.001
Peri-interventional dissection	12 (2.7%)	8 (4.3%)	4 (1.6%)	0.09
Timing of extubation				
Time to extubation (h, median, range)	3 (1–530)	2 (1–115)	11 (1–530)	<0.001
Early extubation (within 6 h)	256 (58.0%)	142 (75.5%)	114 (45.1%)	<0.001
Delayed extubation (6–24 h)	121 (27.4%)	41 (21.8%)	80 (31.6%)	<0.001
Late extubation (>24 h)	64 (14.5%)	5 (2.7%)	59 (23.3%)	<0.001

ICA, internal carotid artery; IQR, interquartile range; MCA, middle cerebral artery; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; TICI, thrombolysis in cerebral infarction; *P*-values <0.05 are shown in bold font. ^aCoronary artery disease, heart failure, cardiomyopathy or valve disease.

0.95 per year, 95% CI 0.93–0.97), NIHSS at admission (OR 0.90 per point, 95% CI 0.86–0.95) and pre-stroke mRS (OR 0.66 per point, 95% CI 0.47–0.93) were significant predictors of unfavourable outcome (Table 3).

Reasons for delayed and late extubation

Of the 65 patients with late extubation (>24 h), reasons for prolonged ventilation were brain oedema (44.6%), insufficient alertness for other reasons (26.1%), respiratory insufficiency (15.9%), status epilepticus (4.6%) and oronasopharyngeal bleeding (3.1%, multiple reasons possible). However, 16.9% of patients with late extubation did not have an obvious medical reason documented for late extubation – all of those patients were extubated within 24–48 h. Symptomatic intracranial haemorrhage was more frequent in patients with late extubation (9.2%); there was no difference between patients with early (2.7%) and delayed extubation (1.6%; Table 2).

In the group of 121 patients with delayed extubation (6–24 h), none of the patients had failed an earlier wake-up trial. Delayed extubation was associated with patient admission outside of core working hours

(within core working hours 86.8% of patients were extubated early, whilst outside core working hours only 52.1% of patients were extubated early; OR 6.06, 95% CI 3.59–10.22, *P* < 0.001, Table 2).

Pneumonia

Overall, early pneumonia rate within the neurointensive care unit/stroke unit stay was 15.8%. Time to extubation was strongly associated with early pneumonia (median ventilation time 15 h in patients with pneumonia versus 3 h in those without, *P* < 0.001). Patients with delayed extubation (6–24 h) more frequently had pneumonia compared to patients with early extubation (20.6% vs. 9.6%, OR 2.43, 95% CI 1.26–4.69, *P* = 0.007). Likewise, patients with late extubation (>24 h) showed even higher rates of pneumonia (27.7% vs. 13.3% in all patients who were extubated within 24 h, OR 2.49, 95% CI 1.76–4.69, *P* = 0.004, Table 2).

Discussion

In this large cohort of consecutive ‘real world’ patients undergoing MT for anterior circulation large

Table 2 Clinical differences between patients with early, delayed or late extubation after stroke thrombectomy

	Early extubation (≤6 h) <i>n</i> = 258 (57.7%)	Delayed extubation (6–24 h) <i>n</i> = 124 (27.7%)	<i>P</i> value	Late extubation (>24 h) <i>n</i> = 65 (14.5%)	<i>P</i> value ^a
Clinical data					
Age (years, mean ± SD)	68.6 ± 13.3	70.8 ± 13.2	0.15	67.6 ± 13.4	0.34
Male sex	132 (51.2%)	67 (54.0%)	0.34	35 (53.8%)	0.52
Arterial hypertension	172 (66.7%)	92 (74.2%)	0.14	49 (75.4%)	0.31
Dyslipidaemia	59 (22.9%)	32 (25.8%)	0.53	8 (12.3%)	0.04
Chronic heart disease ^b	42 (16.3%)	25 (20.2%)	0.35	20 (30.8%)	0.01
Diabetes mellitus	37 (14.3%)	21 (16.9%)	0.51	16 (24.6%)	0.06
Atrial fibrillation	99 (38.4%)	59 (47.6%)	0.09	27 (41.5%)	0.98
Stroke of unknown symptom onset	64 (24.8%)	19 (15.3%)	0.04	8 (12.3%)	0.08
Pre-stroke mRS (median, IQR)	0 (0–0)	0 (0–0)	0.12	0 (0–1)	0.02
NIHSS at admission (median, IQR)	14 (10–17)	15 (13–18)	<0.001	17 (14–19)	<0.001
MCA/M1 occlusion	172 (66.7%)	80 (64.5%)	0.68	41 (63.1%)	0.65
Intracranial ICA occlusion	49 (19.0%)	25 (20.2%)	0.79	20 (30.8%)	0.04
Admission outside core working hours	111 (43.0%)	102 (82.3%)	<0.001	43 (66.2%)	0.12
Intervention					
Intravenous thrombolysis	147 (57.2%)	67 (54.0%)	0.56	46 (70.8%)	0.03
Time to groin puncture (min, median, IQR)	195 (154–245)	210 (169–250)	0.16	204 (160–254)	0.78
Time to recanalization (min, median, IQR)	250 (204–306)	279 (233–324)	0.06	277 (222–302)	0.94
Successful recanalization (TICI 2b–3)	235 (92.5%)	106 (86.2%)	0.05	49 (76.6%)	0.001
Peri-interventional dissection	10 (3.9%)	2 (1.6%)	0.24	0 (0%)	0.15
Outcome parameters					
Pneumonia rate within NICU/stroke unit	20 (9.6%)	22 (20.6%)	0.007	18 (27.7%)	0.004
Symptomatic intracranial haemorrhage	7 (2.7%)	2 (1.6%)	0.51	6 (9.2%)	0.004
Duration of stay in the NICU/stroke unit (days, median, IQR)	3 (2–6)	4 (2–5)	0.86	6 (2–11)	<0.001
NIHSS at discharge from NICU/stroke unit (median, IQR)	5 (2–12)	10 (5–18)	<0.001	18 (10–22)	<0.001
mRS 3 months post-stroke ^c (median, IQR)	2 (1–4)	4 (2–5)	<0.001	6 (4–6)	<0.001
Favourable outcome at 3 months (mRS 0–2)	142 (55.5%)	41 (33.9%)	<0.001	5 (7.8%)	<0.001

ICA, internal carotid artery; IQR, interquartile range; MCA, middle cerebral artery; mRS, modified Rankin Scale; NICU, neurointensive care unit; NIHSS, National Institutes of Health Stroke Scale; TICI, thrombolysis in cerebral infarction; *P*-values <0.05 are shown in bold font.

^aCalculated between patients with late extubation versus early or delayed extubation; ^bCoronary artery disease, heart failure, cardiomyopathy or valve disease; ^cSix patients were lost to follow-up.

vessel occlusion under GA, prolonged ventilation time was associated with a higher pneumonia rate and, besides well-known prognostic factors such as age, NIHSS and unsuccessful recanalization, independently predicted unfavourable functional outcome at 3 months post-stroke. Whilst it was not surprising that patients with late extubation (>24 h) had worse outcome, the most interesting and novel finding of our work was the independent prognostic effect of early extubation within the first 24 h.

To the best of our knowledge, there exists only one previously published report on this topic. In that German study investigating the clinical impact of the duration of mechanical ventilation in 103 stroke patients after MT, patients who were ventilated longer than 24 h had worse 3-month outcome and a higher frequency of pneumonia, but no differences were observed within the group of patients with ventilation times ≤24 h [9]. The weaknesses of that study include

the rather small sample size, the retrospective data collection and the relatively long median overall ventilation time of 18.5 h compared to 3 h in our cohort. Furthermore, reasons for late extubation were not specifically investigated. Conversely to that work, a strong indication for a dose-dependent effect of ventilation time on functional neurological outcome at 3 months was found. In particular, early extubation (within 6 h) after stroke thrombectomy was strongly associated with a favourable 3-month prognosis. This finding is novel and clinically relevant for the post-interventional management of stroke patients treated with MT under GA. Based on our results, MT patients with anterior circulation large vessel occlusion undergoing the procedure under GA should be extubated early after the intervention, unless there are medical reasons against it. However, it is important to note that our results are most probably not transferable to patients who are treated with thrombectomy

Table 3 Binary multivariable logistic regression model for the target variable 'favourable outcome' (modified Rankin Scale 0–2) in patients extubated within 24 h

Test variable	Odds ratio	95% confidence interval	<i>P</i> value
Clinical data and intervention			
Age (per year)	0.95	0.93–0.97	<0.001
Pre-stroke mRS (per point)	0.66	0.47–0.93	0.02
NIHSS at admission (per point)	0.90	0.86–0.95	<0.001
Peri-interventional management			
Successful recanalization (TICI 2b–3 vs. 0–2a)	6.97	2.38–20.4	<0.001
Early extubation (within 6 h vs. 6–24 h)	1.93	1.15–3.24	0.01

mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale; TICI, thrombolysis in cerebral infarction. *P*-values <0.05 are shown in bold font.

for large vessel occlusion of the vertebrobasilar circulation, as brainstem infarction often affects functional pathways responsible for respiration, swallowing and protective reflexes, which may cause additional difficulties regarding extubation.

When interpreting our work, important potential confounding factors have to be considered. In clinical practice, physicians might be more reluctant to swiftly extubate patients if they have severe strokes, more comorbidities or if a critical disease course is expected (e.g. a failed thrombectomy in a younger patient). This is probably the main explanation for the unfavourable outcome of patients with late extubation >24 h. In most of those patients, it was possible to identify specific medical reasons leading to later extubation (most frequently neurological and/or respiratory stroke complications). However, those factors mostly did not apply to extubations within 24 h in our study cohort. On the contrary, it was found that admission outside core working hours was an important predictor of delayed extubation. This means that non-medical reasons played an important role in the decision to extubate early or delayed within the first 24 h after MT.

It was also found that patients with delayed extubation had slightly higher NIHSS scores at admission. Nonetheless, multivariable analysis including all assessed outcome-relevant factors still identified early extubation as an independent prognostic factor at 3 months post-stroke. There are multiple possible reasons for unfavourable outcome resulting from delayed extubation. First, it is well known that pneumonia leads to worse outcome in stroke patients [13]. Ventilator-associated pneumonia is one of the most important complications of mechanical ventilation and its

occurrence is strongly associated with the duration of ventilation [14]. This was also seen in our study, where patients with late extubation had the highest pneumonia rate, whilst patients who were extubated early had pneumonia much less frequently. Importantly, pneumonia occurrence was only assessed within the first days when patients were admitted on the neurointensive care unit/stroke unit. This could be seen as a limitation of this study; however, our primary aim was to analyse pneumonia as a complication of mechanical ventilation rather than stroke-related pneumonia, which occurs at variable time points during the hospital stay and depends on different circumstances. Respiratory insufficiency was a reason for prolonged ventilation >24 h in 11 patients and may have been caused by pneumonia in some of those cases.

Delayed extubation also requires sedative drugs to be given for longer periods. The majority of sedative drugs can cause hypotension, which might negatively affect cerebral blood flow in the critical time period after recanalization, possibly influencing final infarct size. Additionally, hyperoxaemia may be present for a longer time in patients with delayed extubation, which might contribute to secondary brain injury and worse outcomes [15,16]. Furthermore, early diagnostic stroke workup may be impeded in patients who remain intubated for a longer time, potentially leading to delays in specific treatment of stroke causes. Moreover, early rehabilitative measures including mobilization are only possible to a limited extent in intubated patients.

Extubation outside core working hours and in particular during night-time are often considered more prone to complications with an increased mortality rate, as recently described in a large cohort study across different US intensive care unit settings [17]. In contrast, several smaller studies have shown that in specific settings night-time extubation may even be associated with better patient outcomes such as decreased reintubation rates, length of ventilation and hospital stay [18,19].

Whilst our study has several strengths such as the large cohort of consecutive MT patients, the high follow-up rate and the exact documentation of reasons for delayed extubation, its main limitation is the retrospective design. Even though data were collected prospectively and our study showed a strong association of delayed extubation and unfavourable 3-month outcome, the study design cannot prove causality. Furthermore, uniform pre- and post-interventional imaging to assess infarct size and blood pressure data were not available. Therefore, our findings should be seen as hypothesis-generating and need to be confirmed in prospective studies.

Conclusion

Prolonged ventilation time in patients undergoing endovascular stroke treatment under GA is independently related to unfavourable neurological outcome at 3 months post-stroke. This finding is clinically relevant for the post-interventional management of patients with ischaemic stroke of the anterior cerebral circulation, and indicates that extubation after stroke thrombectomy should be performed as early as safely possible.

Disclosure of conflicts of interest

The authors declare no financial or other conflicts of interest.

Data availability statement

The datasets generated during this study are available from the corresponding author upon reasonable request.

References

- Goyal M, Menon BK, van Zwam WH, *et al.* Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet* 2016; **387**: 1723–1731.
- Campbell BCV, van Zwam WH, Goyal M, *et al.* Effect of general anaesthesia on functional outcome in patients with anterior circulation ischaemic stroke having endovascular thrombectomy versus standard care: a meta-analysis of individual patient data. *Lancet Neurol* 2018; **17**: 47–53.
- Ilyas A, Chen C-J, Ding D, *et al.* Endovascular mechanical thrombectomy for acute ischemic stroke under general anesthesia versus conscious sedation: a systematic review and meta-analysis. *World Neurosurg* 2018; **112**: e355–e367.
- Zhang Y, Jia L, Fang F, Ma L, Cai B, Faramand A. General anesthesia versus conscious sedation for intracranial mechanical thrombectomy: a systematic review and meta-analysis of randomized clinical trials. *J Am Heart Assoc* 2019; **8**: e011754.
- Fargen KM, Arthur AS, Spiotta AM, *et al.* A survey of neurointerventionalists on thrombectomy practices for emergent large vessel occlusions. *J Neurointerv Surg* 2017; **9**: 142–146.
- Hindman BJ, Dexter F. Anesthetic management of emergency endovascular thrombectomy for acute ischemic stroke, part 2: Integrating and applying observational reports and randomized clinical trials. *Anesth Analg* 2019; **128**: 706–717.
- Coplin WM, Pierson DJ, Cooley KD, Newell DW, Rubinfeld GD. Implications of extubation delay in brain-injured patients meeting standard weaning criteria. *Am J Respir Crit Care Med* 2000; **161**: 1530–1536.
- Asehnoune K, Roquilly A, Cinotti R. Respiratory management in patients with severe brain injury. *Crit Care* 2018; **22**: 76.
- Nikoubashman O, Schürmann K, Probst T, *et al.* Clinical impact of ventilation duration in patients with stroke undergoing interventional treatment under general anesthesia: the shorter the better? *AJNR Am J Neuroradiol* 2016; **37**: 1074–1079.
- Kneihsl M, Niederkorn K, Deutschmann H, *et al.* Abnormal blood flow on transcranial duplex sonography predicts poor outcome after stroke thrombectomy. *Stroke* 2018; **49**: 2780–2782.
- Johanson WG, Pierce AK, Sanford JP, Thomas GD. Nosocomial respiratory infections with Gram-negative bacilli. The significance of colonization of the respiratory tract. *Ann Intern Med* 1972; **77**: 701–706.
- Rea-Neto A, Youssef NCM, Tuche F, *et al.* Diagnosis of ventilator-associated pneumonia: a systematic review of the literature. *Crit Care* 2008; **12**: R56.
- Finlayson O, Kapral M, Hall R, *et al.* Risk factors, inpatient care, and outcomes of pneumonia after ischemic stroke. *Neurology* 2011; **77**: 1338–1345.
- Kalanuria AA, Ziai W, Zai W, Mirski M. Ventilator-associated pneumonia in the ICU. *Crit Care* 2014; **18**: 208.
- Taccone FS, Crippa IA, Vincent J-L. Normobaric hyperoxia after stroke: a word of caution. *Expert Rev Neurother* 2018; **18**: 91–93.
- Robba C, Bonatti G, Battaglini D, Rocco PRM, Pelosi P. Mechanical ventilation in patients with acute ischaemic stroke: from pathophysiology to clinical practice. *Crit Care* 2019; **23**: 388.
- Gershengorn HB, Scales DC, Kramer A, Wunsch H. Association between overnight extubations and outcomes in the intensive care unit. *JAMA Intern Med* 2016; **176**: 1651–1660.
- Everhart KK, Khorsand S, Khandelwal N, Michaelsen KE, Spiekerman CF, Joffe AM. Nighttime extubation does not increase risk of reintubation, length of stay, or mortality: experience of a large, urban, teaching hospital. *Anesth Analg* 2019; **128**: 918–923.
- Tischenkel BR, Gong MN, Shiloh AL, *et al.* Daytime versus nighttime extubations: a comparison of reintubation, length of stay, and mortality. *J Intensive Care Med* 2016; **31**: 118–126.