LETTERS

Taylor & Francis

OPEN ACCESS Check for updates

Could mandatory vaccination increase coverage in health-care Workers? The experience of Bari Policlinico General Hospital

Antonio Di Lorenzo^a, Silvio Tafuri^a, Andrea Martinelli^a, Giusy Diella^a, Luigi Vimercati^b, and Pasquale Stefanizzi D^a

^aDepartment of Biomedical Science and Human Oncology, Aldo Moro University, Bari, Italy; ^bDepartment of Interdisciplinary Medicine, Occupational Health Division, Aldo Moro University, Bari, Italy

ABSTRACT

Influenza is a major issue for health-care systems all over the world and health-care workers (HCWs) are a subgroup at increased risk of exposure to respiratory pathogens, including influenza, with potential threat for both their own and their patients' health. Despite this and despite the anti-influenza vaccination being offered to all HCWs, vaccination coverage among them is still low in Europe, generally less than 30%. During 2020s influenza season, influenza vaccination begun mandatory in Puglia (Italy) and Bari Policlinico General Hospital designed an active call system in order to increase its employees' compliance to vaccination. This system was able to determine an increase of vaccination coverage from 23.1% to 77.6%.

ARTICLE HISTORY

Received 8 September 2021 Revised 4 October 2021 Accepted 26 October 2021

KEYWORDS Influenza; vaccination; health-care workers; Italy

Dear Editor,

Despite important efforts of Public Health institutions, influenza remains a major issue for the health of the population. This is also due to suboptimal vaccination coverage among the target population, even though active vaccination offer has been recommended for many years. Health-care workers (HCWs) are a subgroup at increased risk of exposure to respiratory pathogens, including influenza, with potential threat for their health and for the patients' safety.¹In fact, infected HCWs could be the source of contagion of both other colleagues and patients.

Influenza epidemics cause a significant increase in sickness absenteeism from work among HCWs. A 2019 Italian study estimated the overall cost of this phenomenon at \in 1,76M.¹

Vaccination coverage among HCWs is still low in Europe, generally less than 30%, despite several recommendations from international authorities, such as the European Center for Disease Prevention and Control (ECDC). In particular, the ECDC states that a vaccination coverage above 75% of the atrisk population, including HCWs, should be obtained in order to grant an adequate control of the disease.²

In Italy, each of the 20 Regions is responsible for the management of Health-care services. During 2020s influenza season, Puglia adopted the Regional Law 19 June 2018 n. 27 and the Regional Regulation n. 10/2020,³ which provided influenza vaccination as mandatory for HCWs operating in high-risk settings. Bari Policlinico General Hospital is the most important Hospital in the South of Italy, with 1000 beds and >6000 HCWs. Since 2010, influenza vaccination is offered actively and free of charge for Policlinico HCWs. In compliance with Regional Law, Policlinico made anti- influenza vaccination mandatory for HCWs in order for them to be allowed to operate in the following Operative Units (OUs), considered at high infective risk (Intensive Care Unit, Organ Transplant Center, Hematology and Transplant, Gynecology and Obstretics, Respiratory System Diseases, Infectious Diseases, Physical and Rehabilitative Medicine, General Medicine, Nephrology, Neonatology, Neurology, Oncology, Emergency Room, Rheumatology).

In order to achieve the highest possible compliance to antiinfluenza vaccination, both a vaccination clinic and on-site daily vaccination activities were implemented.

Thanks to this effort, by 15 December 2020, 1634 HCWs operating in Policlinico's high-risk OUs out of 2383 (68.6%) had been vaccinated, while 749 (31.4%) of them had not.

Following this first phase of vaccination campaign, an active call system was activated in order to collect more adhesions among the non-compliant HCWs. A letter, signed by the Head of Vaccination Clinic, the Occupation Physician and the Director of Risk Management Unit, was sent to the non-vaccinated subjects operating in high-risk facilities, inviting them to undergo the vaccination and reminding them of the law requirements for HCWs regarding influenza vaccination. Personnel was also presented with the possibility of being suspended or transferred in other units.

Following this formal reminder, an additional 215 HCWs underwent the anti-influenza vaccination, thus providing a grand total of 1849 (77.6%) vaccinated HCWs against 534 (22.4%) who kept refusing the vaccine. The data were last collected on 25 March 2021.

This coverage was higher than the figure of previous season: in fact, in the same Units, coverage was 23.8% (324/1360) in 2019 and 28.1% (378/1347) in 2018.

© 2021 The Author(s). Published with license by Taylor & Francis Group, LLC.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (http://creativecommons.org/licenses/by-nc-nd/4.0/), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

CONTACT Silvio Tafuri 🖾 silvio.tafuri@uniba.it 😰 Department of Biomedical Science and Human Oncology, Aldo Moro University, Bari, Italy

In conclusion, mandatory vaccination seems to be a tool consistent with the purpose of increasing the vaccination coverage in HCWs. Our active re-call system adopted the quota of vaccinated HCWs in high-risk OUs by 9%, meaning that approximately one-third of HCWs who had previously refused the vaccination changed their mind after the written reminder. The repetition of the same formula during the next influenza season may achieve even better results, especially with more iterations of the active call to non-compliant HCWs, even implementing other interventions such as declination statements and educational audits, which proved to be highly effective in various studies.⁴

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

The authors reported that there is no funding associated with the work featured in this article.

ORCID

Pasquale Stefanizzi (D) http://orcid.org/0000-0002-3279-0196

References

- Dini G, Toletone A, Sticchi L, Orsi A, Bragazzi NL, Durando P. Influenza vaccination in healthcare workers: a comprehensive critical appraisal of the literature. Hum Vaccines Immunother. 2018 Mar 4;14(3):772–89. Epub 2017 Oct 20. PMID: 28787234; PMCID: PMC5861785. doi:10.1080/21645515.2017.1348442.
- European Centre for Disease Prevention and Control. Implementation of the council recommendation on seasonal influenza vaccination (2009/1019/EU). Stockholm, Sweden: ECDC; 2014.
- Regolamento Regionale 25 giugno 2020, n. 10, Disposizioni per l'esecuzione degli obblighi di vaccinazione degli operatori sanitari. Legge regionale 19 giugno 2018, n. 27, Bollettino Ufficiale della Regione Puglia - n. 94 *suppl.* del 26-6-2020.
- Lytras T, Kopsachilis F, Mouratidou E, Papamichail D, Bonobas S. Interventions to increase seasonal influenza vaccine coverage in healthcare workers: a systematic review and meta-regression analysis. Hum Vaccines Immunother. 2016 Mar 3;12(3):671–81. PMID: 26619125; PMCID: PMC4964628. doi:10.1080/ 21645515.2015.1106656.