

ORIGINAL ARTICLE

CFTR founder mutation causes protein trafficking defects in Chinese patients with cystic fibrosis

Gordon K. C. Leung¹, Dingge Ying², Christopher C. Y. Mak¹, Xin-Ying Chen³, Weiyi Xu³, Kit-San Yeung¹, Wai-Lap Wong¹, Yoyo W. Y. Chu¹, Gary T. K. Mok¹, Christy S. K. Chau⁴, Jenna McLuskey⁵, Winnie P. T. Ong⁶, Huey-Yin Leong⁶, Kelvin Y. K. Chan⁷, Wanling Yang¹, Jeng-Haur Chen³, Albert M. Li⁸, Pak C. Sham², Yu-Lung Lau^{1,4}, Brian H. Y. Chung^{1,4,9,*} & So-Lun Lee^{1,4,9,*}

¹Department of Paediatrics and Adolescent Medicine, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong, Hong Kong

²Department of Psychiatry, LKS Faculty of Medicine, The University of Hong Kong, Hong Kong, Hong Kong

³HKU Shenzhen Institute of Research and Innovation, School of Biomedical Sciences, The University of Hong Kong, Hong Kong, Hong Kong

⁴Department of Paediatrics and Adolescent Medicine, Queen Mary Hospital, Hong Kong, Hong Kong

⁵NHS Lothian, Lothian NHS Board, Edinburgh, UK

⁶Department of Genetics, Kuala Lumpur Hospital, Kuala Lumpur, Malaysia

⁷Department of Obstetrics and Gynaecology, Tsan Yuk Hospital, Hong Kong, Hong Kong

⁸Department of Paediatrics, Faculty of Medicine, The Chinese University of Hong Kong, Hong Kong, Hong Kong

⁹Department of Paediatrics and Adolescent Medicine, The Duchess of Kent Children's Hospital at Sandy Bay, Hong Kong

Keywords

Bronchiectasis, Chinese, cystic fibrosis, founder mutation, trafficking defects

Correspondence

So-Lun Lee, Department of Paediatrics and Adolescent Medicine, The Duchess of Kent Children's Hospital at Sandy Bay, 12 Sandy Bay Road, Pokfulam, Hong Kong. Tel: +852 2974 0392; E-mail: slleem@hku.hk and

Brian H. Y. Chung, Department of Paediatrics and Adolescent Medicine, LKS Faculty of Medicine, The University of Hong Kong, HKSAR. Rm 103, 1/F, New Clinical Building, Queen Mary Hospital, Hong Kong, Hong Kong. Tel: +852 2255 4482; Fax: +852 2855 1523; E-mail: bhychung@hku.hk

Funding Information

This work was supported by the SK Yee Medical Foundation [grant number: 200940007] and The Society for the Relief of Disabled Children [grant number: 260007609].

Received: 17 July 2016; Revised: 16 October 2016; Accepted: 19 October 2016

Molecular Genetics & Genomic Medicine 2017; 5(1): 40–49

doi: 10.1002/mgg3.258

*These authors contributed equally to this work.

Abstract**Background**

Cystic fibrosis (CF) is a rare condition in Asians. Since 1985, only about 30 Chinese patients have been reported with molecular confirmation.

Method

Using our in-house next-generation sequencing (NGS) pipeline for childhood bronchiectasis, we identified disease-causing *CFTR* mutations in CF patients in Hong Kong. After identifying p.I1023R in multiple patients, haplotype analysis was performed with genome-wide microarray to ascertain the likelihood of this being a founder mutation. We also assessed the processing and gating activity of the mutant protein by Western hybridization and patch-clamp test.

Results

Molecular diagnoses were confirmed in four patients, three of whom shared a missense mutation: *CFTR*:c.3068T>G:p.I1023R. The results suggested that p.I1023R is a founder mutation in southern Han Chinese. In addition, the processing and gating activity of the mutant protein was assessed by gel electrophoresis and a patch-clamp test. The mutant protein exhibited trafficking defects, suggesting that the dysfunction is caused by reduced cell surface expression of the fully glycosylated proteins.

Conclusion

Together with other previously reported mutations, the specific founder mutation presented herein suggests a unique *CFTR* mutation spectrum in the southern Chinese populations, and this finding has vital implications for improving molecular testing and mutation-specific treatments for Chinese patients with CF.

Introduction

Cystic fibrosis (CF) (OMIM 219700) is a common autosomal recessive genetic disorder in Caucasian populations with an incidence of approximately 1 in 2500–3500 newborns (Southern et al. 2007; Salvatore et al. 2011). Compared with Western countries, CF is a considerably rarer condition in Asia. For example, the incidence of CF is approximately 1 in 350,000 in the Japanese population (Yamashiro et al. 1997). The condition is caused by mutations in the coding gene of the membrane glycoprotein, cystic fibrosis transmembrane conductance regulator (CFTR), which is responsible for regulating the flow of chloride ions across the epithelial membrane. To date, the number of documented disease-causing mutations in *CFTR* (RefSeq NM_000492.3) exceeds 2000 (Cystic Fibrosis Mutation Database 2016). However, molecular confirmation of mutation has only been reported for 32 Chinese patients from 1985 to 2016 (see Fig. 1 and Table S1). One explanation for the relatively high incidence of CF in the West is the founder effect of certain mutations (Castellani et al. 2008). The most common mutation, p.Phe508del ($\Delta F508$), accounts for greater than two-thirds of CF alleles worldwide, though it is not a predominant mutation in Asian CF patients (Wong et al. 2003). Although founder mutations have not yet been identified in the Chinese population, among the *CFTR* mutations found in the Chinese population, the novel variant c.3068T>G:p.Ile1023Arg (p.I1023R) was identified in a pair of siblings with CF from Taiwan (Ho et al. 2007). Here, we describe the identification of four additional patients with the p.I1023R variant in southern China, and our results suggest that p.I1023R is a Chinese-specific founder mutation that leads to trafficking defects and reduced expression of functionally mature CFTR protein.

Materials and Methods

Patient recruitment

This study was approved by the HKWC-HKU Institutional Review Board. Informed consent was obtained from patients (or parents) after clinical consultation. Twenty-one patients diagnosed with early-onset bronchiectasis were referred to our unit for clinical evaluation and a sweat test. Four patients with available genomic DNA presented a strong clinical suspicion of CF and were sweat-test positive (chloride ion concentration >60 mmol/L). Molecular diagnoses were performed for all patients.

Experimental workflow

For a detailed description of the workflow, please refer to the supplementary data (Appendix S1).

Results

Detection of disease-causing mutations in *CFTR*

Four unrelated Chinese patients from Hong Kong were positive for *CFTR* mutations, of whom three harbored a p.I1023R mutation. These patients were all compound heterozygotes with p.I1023R and one other disease-causing mutation. Through personal communication, we identified another nonconsanguineous southern Chinese CF patient from Malaysia with a homozygous p.I1023R mutation confirmed by Sanger sequencing in our laboratory. Clinical and genetic data for the patients are summarized in Table 1.




Figure 1. *CFTR* mutation spectrum in Chinese CF patients (findings from this study and previous studies reported in both English and Chinese medical literatures from 1985 to 2016). Only variants observed in patients with complete molecular diagnosis were included. The upper panel refers to *CFTR* mutations identified from Northern China (i.e., Beijing), while the lower panel refers to *CFTR* mutations identified from Southern China (i.e., Sichuan, Taiwan and Hong Kong). The geographical origins of mutations were determined according to either the information provided from the manuscript on patient recruitment, or the institute address of the corresponding authors. *CFTR* mutations from two reports (Zielenski et al. 1995; Wagner et al. 1999) were not included as the corresponding institute was located outside China.

Table 1. Clinical presentation of Chinese patients with molecularly confirmed CF. Organisms in **bold** indicate bacteria isolated from first-time culture. y.o., year old; m.o., month old; Age at Dx, age at diagnosis¹; PFT, pulmonary functional test; MI, meconium ileus; PI, pancreatic insufficiency²; HRCT, high-resolution computed tomography; Ac, *Acinetobacter* sp.; BC, *Burkholderia cepacia*; HI, *Haemophilus influenzae*; KP, *Klebsiella pneumoniae*; MC, *Moraxella catarrhalis*; NTM, non-tuberculosis mycobacterium; PA, *Pseudomonas aeruginosa*; PM, *Pseudomonas pseudomallei*; SA, *Stenotrophomonas aureus*; SM, *Stenotrophomonas maltophilia*; USG, ultrasound.

Patients	Gender	Age at onset of symptom	Age at Dx	Age at time of report	Sweat test results (mmol/L)	Respiratory symptoms	Organism	Latest PFT	MI	PI	Other symptoms	Current status	Mutation 1	Mutation 2
I	Male	2 y.o.	17 y.o.	Succumbed at 23 y.o.	121,126	Chronic productive cough since 2 y.o.; suspected pulmonary TB and treated at 15 y.o.; persistent cough and sputum with a severe superimposed chest infection; HRCT chest at 17 y.o. showed significant bronchiectatic changes at right upper lobe.	PA , SA, HI, PM, BC	FEV ₁ : 16% pred. FVC: 29% pred. FEV ₁ /FVC: 47%	No	No	Soft drug abuser; possibility of lung transplant was rejected as he had BC infection.	Developed progressive respiratory failure, palliative non-invasive ventilation implemented at 23 y.o. and succumbed 3 months later.	c.1766+5G>T	c.3068T>G; p.I1023R
II	Male	4 m.o.	6 m.o.	14 y.o.	100,112	Two episodes of pneumonia due to SA at 4 and 6 m.o.; chest radiography showed haziness in both lungs; HRCT chest was not done as there has not been any further episodes of significant chest infections.	SA , MC, HI	FEV ₁ : 78% pred. FVC: 79% pred. FEV ₁ /FVC: 91%	No	No	Hyponatraemia, hypokalaemia and metabolic alkalosis at initial presentation; negative screening for adrenal insufficiency; initial growth along 10 th percentile.	Height and weight reached 25th percentile; active in various sport activities	c.1766+5G>T	c.3140-26A>G

(Continued)

Table 1. Continued.

Patients	Gender	Age at onset of symptom	Age at Dx	Age at report of symptom	Sweat test results (mmol/L)	Respiratory symptoms	Organism	Latest PFT	MI	PI	Other symptoms	Current status	Mutation 1	Mutation 2
III	Male	At birth	2 m.o.	9 y.o.	108, 112	Recurrent pneumonia in first 2 years; NTM infection at 5 y.o. and was eradicated; HRCT chest at 5 y.o. showed bronchiectasis, segmental left lingular collapse and patchy parenchymal opacities in both lungs; infrequent respiratory symptoms.	SA , Ac, PA, KP, SM, NTM	FEV ₁ : 110% pred. FVC: 98% pred. FEV ₁ /FVC: 83%	Yes	Yes	Peanut and drug allergy; eczema.	Height and weight at around 10th percentile; pulmonary exacerbations requiring hospitalization about once per year; on antibiotics prophylaxis and airway clearance management.	c.868C>T: p.Q290X	c.3068T>G: p.I1023R
IV	Female	At birth	9 y.o.	12 y.o.	123	Recurrent pneumonia; HRCT chest at 9 y.o. showed significant bronchiectatic changes; persistent cough and sputum.	PA , SA	FEV ₁ : 68% pred. FVC: 74% pred. FEV ₁ /FVC: 84%	Yes	Yes	Abdominal USG showed fatty change in liver; body height at 50th percentile and body weight at 3rd percentile at diagnosis	Height caught up to 90th percentile and weight reached 50th percentile; pulmonary exacerbations requiring hospitalization about once per year; on antibiotics and airway clearance management.	c.1657C> T: p.R533X	c.3068T>G: p.I1023R

(Continued)

Table 1. Continued.

Patients	Gender	Age at onset of symptom	Age at Dx	Age at time of report	Sweat test results (mmol/L)	Respiratory symptoms	Organism	Latest PFT	MI	PI	Other symptoms	Current status	Mutation 1	Mutation 2
V	Female	4 m.o.	13 m.o.	5 y.o.	122,124	Persistent cough since 4 m.o.; recurrent pneumonia and rhinosinusitis; HRCT chest at 13 m.o. showed bronchiectatic changes.	PA	Not available as the patient is too young for PFT	No	Yes	Ventricular septal defect which closed spontaneously at 6 m.o.; hyponatraemia, hypokalaemia and hypochloraemia at initial presentation; abdominal USG at 3 y.o. suggested possible liver cirrhosis and early portal hypertension but not confirmed by liver biopsy; liver function remained normal; height was between 10th to 25th percentile and weight was less than 3rd percentile at diagnosis; required supplemental feeding via gastrostomy.	Height caught up to 25th percentile and weight reached 10th percentile since 4 y.o.; oxygen dependent until 4.5 y.o.	c.3068T>G: p.I1023R	c.3068T>G: p.I1023R

¹Diagnosis was made based on clinical phenotype and abnormal sweat test results after referral to our unit.

²Patients with pancreatic insufficient were given pancreatic enzyme supplement.




Figure 3. Functional analysis of p.I1023R-CFTR. **A.** Protein processing of wild-type CFTR, p.F508del-CFTR and p.I1023R-CFTR. The same immunoblotting images are presented. Each lane contains 50 μ g total protein. Band C (or C) represents fully glycosylated and mature CFTR, whereas Band B (or B) represents partial glycosylated and immature CFTR. **B.** The percentage ratio of Band C intensity is normalized to the sum of band B and band C. Data are the mean + SE ($N = 6$). The asterisk indicates that the values between two groups are significantly different ($p < 0.001$, one-way ANOVA, Holm-Sidak method). **C.** Representative 10 second recordings show the activities of a single wild-type CFTR and p.I1023R-CFTR in the excised inside-out membrane patches at 1 mM ATP and room temperature. The dotted lines indicate closed channels. **D** and **E.** The single-channel current amplitude (i) and open probability (P_o) are presented. Data are the mean + SE ($N = 3$).

of evidence suggest that all four p.I1023R carrier families are southern Han Chinese. Therefore, the above evidence as well as sibling data from the first report on p.I1023R from Taipei, which is located in the northern region of Taiwan (Ho *et al.* 2007; Liu *et al.* 2014), indicate that the mutation is a founder mutation confined to southern regions of China.

Functional analysis demonstrates that p.I1023R leads to a trafficking defect during CFTR maturation without affecting the gating function of the protein

To better understand the molecular defects of the p.I1023R mutant CFTR protein, we investigated protein

processing properties by resolving CFTR protein profile, as previously described (Ostedgaard et al. 2007). In this analysis, wild-type CFTR was characterized by the presence of fully glycosylated proteins was normally transported to the cell membrane, and presented increased expression compared with mutated p.I1023R-CFTR (Fig. 3A,B), for which a significant proportion of less-glycosylated protein was found. This finding suggests that the p.I1023R mutation may lead to a trafficking defect, with reduced protein processing from the endoplasmic reticulum to the Golgi apparatus. However, a single-channel study of p.I1023R-CFTR using the patch-clamp method (Chen et al. 2009) demonstrated that the single-channel current amplitude (i) and open probability (P_o) of the CFTR chloride channel were not altered by the p.I1023R mutation (Fig. 3C–E). Hence, the dysfunction caused by p.I1023R appears to be largely caused by the reduced expression of fully glycosylated and mature protein.

Discussion

CF has seldom been reported in Chinese individuals. Here, a molecular diagnosis was performed for four local Chinese patients. Our unit is the only center that offers sweat testing in Hong Kong, a city located in southern China with a population of approximately 7.3 million. We estimated that the incidence of CF is approximately 1 in 300,000 live births in Hong Kong, which is similar to that in Japan (Yamashiro et al. 1997).

We identified the missense mutation p.I1023R in three of four CF patients from Hong Kong as well as in an additional Chinese patient (homozygous) from Malaysia. All four patients with the p.I1023R mutation exhibited recurrent pneumonia and bronchiectasis. *Pseudomonas aeruginosa* was isolated from all patients, and a more diverse respiratory tract flora was noted in two patients. Three patients had pancreatic insufficiency, and two had meconium ileus. In the limited number of cases presenting with the classical clinical features of CF, we did not observe an association of p.I1023R with a unique clinical phenotype.

The identification of multiple patients with p.I1023R led us to hypothesize that p.I1023R is a potential founder mutation in southern Han Chinese. We defined the following criteria and examined the likelihood of our hypothesis: (1) all patients and p.I1023R carriers share a haplotype associated with the p.I1023R mutation; (2) the p.I1023R-associated haplotype was shared among affected families with a genetic distance greater than 1 cM; (3) the mutant allele was rare and specific to the population (Fachal et al. 2012); (4) the p.I1023R-associated haplotype was unique in the normal population; and (5) all carriers were delineated to the same

geographic region. Our experimental data were well fit to these criteria well, and we concluded that p.I1023R is indeed a founder mutation in southern Han Chinese. The additional patient from Malaysia who was homozygous for p.I1023R further supported our conclusion. Because the founder haplotype of this mutation was well defined, the time of emergence of p.I1023R can be estimated. Assuming all five p.I1023R lineages from four families to be independent, all of the patients from this study inherited the mutation from a common ancestor approximately 15.6 generations (i.e., meiotic events) ago.

Although p.I1023R is the most common CFTR mutation in our cohort, it was not observed in the cohort of 20 CF patients recently reported by investigators from Beijing (Liu et al. 2015; Tian et al. 2016). Another Chinese-specific mutation, p.G970D, appears to be more commonly observed in northern Chinese with CF. We summarize our findings along with previously reported CFTR mutations. Interestingly, an obvious difference is observed in the spectrum of CFTR mutations between the northern and southern regions of China (Fig. 1). Although further studies must be performed to elucidate the complete picture of CFTR molecular epidemiology in China, current evidence suggests that compared with Caucasians, Chinese CF patients have a different CFTR mutation spectrum. As translating CF molecular tests into clinical practice, such as through the use of a 23-variant ACMG CF panel (Watson et al. 2004), will result in a low diagnostic yield when applied to Chinese patients, a customized Chinese-specific CF panel may be required to investigate the incidence of CF in Chinese patients. This Chinese-specific CF panel is not only relevant for Asian countries but also for other parts of the world that present significant population of Chinese immigrants. For example, the 2011 US Census reported that the Chinese American population included approximately 3.8 million individuals and accounted for 1.2% for the overall population.

Here, we illustrate that p.I1023R causes a disruption in CFTR intracellular processing, though the conductance (i) and channel open probability (P_o) remained similar to that of the wild-type protein. Our calculation of normal CFTR expression (i.e., 25%) is likely an overestimate (Moran and Zegarra-Moran 2008). We tested expression of the mutant protein via the overexpression of foreign plasmids in a cell line and analyzing the total protein content from the cell extract, including the CFTR protein trapped in the Golgi apparatus and cell membrane. However, the band intensity ratio did not reflect the actual proportion of functional CFTR in the cell membrane compared with wild-type CFTR. In addition, we did not define an association between the physiological values of p.I1023R-associated CFTR expression and a clinical

phenotype of CF. Moreover, our data clearly indicate that maturation of the p.I1023R-CFTR protein maturation was inhibited.

In general, genetic mutations primarily disrupt CFTR functions by reducing protein expression or channel activity. We functionally characterized p.I1023R as a class II *CFTR* mutation (reducing protein expression) based on the functional properties of the gene product as originally proposed by Tsui (1992). Similar to p.I1023R, several other mutations, such as p.A561E, p.S1235R, and p.N1303K, are also to class II mutations according to their pathological mechanism (Welsh and Smith 1993; Mendes *et al.* 2003). Understanding the pathological mechanisms of each class of mutation can shed light on the development of mutation-specific therapies. For instance, the protein maturation process affected by p.A561E was enhanced by lumacaftor treatment (Awatade *et al.* 2014), whereas human cell lines with p.S1235R and p.N1303K responded positively to treatment with small molecular correctors (Rapino *et al.* 2015). Thus, our findings can be used as the basis for future clinical trials in drug development.

To conclude, we provide the first report of a founder mutation, p.I1023R, in *CFTR* in southern Chinese individuals as well as a functional characterization of this mutation. Because this mutation and other known mutations found in Chinese patients are not currently included in the conventional *CFTR* gene panel for the molecular diagnosis of CF, we suggest a different *CFTR* gene panel or mutation screening strategy for CF patients of Chinese origin.

Acknowledgments

We are grateful to the families who participated in this study and thank SK Yee Medical Foundation and The Society for the Relief of Disabled Children for their financial support. We also thank the Centre for Genomic Sciences at The University of Hong Kong for technical support.

Conflict of Interest

The authors declare that they have no actual or potential conflicts of interest, including any financial, personal or other relationships with people or organizations regarding the submitted work.

References

- Awatade, N. T., I. Uliyakina, C. M. Farinha, L. A. Clarke, K. Mendes, A. Solé, *et al.* 2014. Measurements of functional responses in human primary lung cells as a basis for personalized therapy for cystic fibrosis. *EBioMedicine* 22:147.

- Castellani, C., H. Cuppens, M. Macek, J. J. Cassinian, E. Kerern, P. Durie, *et al.* 2008. Consensus on the use and interpretation of cystic fibrosis mutation analysis in clinical practice. *J. Cyst. Fibros.* 7:179–196.
- Chen, J. H., Z. Cai, and D. N. Sheppard. 2009. Direct sensing of intracellular pH by the cystic fibrosis transmembrane conductance regulator (CFTR) Cl⁻ channel. *J. Biol. Chem.* 284:35495–35506.
- Cystic Fibrosis Mutation Database. 2016. Available at: <http://www.genet.sickkids.on.ca/app> [Online]. (accessed 30th May 2016).
- Exome Aggregation Consortium. 2015. Available at: <http://exac.broadinstitute.org/> [Online]. Cambridge, MA. (accessed 26th January 2015).
- Fachal, L., L. Rodriguez-Pazos, M. Ginarte, J. Toribio, A. Salas, and A. Vega. 2012. Multiple local and recent founder effects of TGM1 in Spanish families. *PLoS ONE* 7:e33580.
- Ho, Y., W. Wu, K. Wong, C. Huang, C. Niu, S. Shyur, *et al.* 2007. Novel mutation (I1023R) in two Taiwanese siblings of cystic fibrosis. *J. Pediatr. Respirat. Dis.*, 5: 152–158.
- Liu, L., S. Shyur, S. Chu, L. Huang, Y. Kao, W. Lei, *et al.* 2014. Cystic fibrosis: experience in one institution. *J. Microbiol. Immunol. Infect.*, 47:358–361.
- Liu, Y., L. Wang, X. Tian, K. F. Xu, W. Xu, X. Li, *et al.* 2015. Characterization of gene mutations and phenotypes of cystic fibrosis in Chinese patients. *Respirology* 20:312–318.
- Mendes, F., M. Roxo Rosa, A. Dragomir, C. M. Farinha, G. M. Roomans, M. D. Amaral, *et al.* 2003. Unusually common cystic fibrosis mutation in Portugal encodes a misprocessed protein. *Biochem. Biophys. Res. Commun.* 311:665–671.
- Moran, O., and O. Zegarra-Moran. 2008. On the measurement of the functional properties of the CFTR. *J. Cyst. Fibros.* 7:483–494.
- Ostedgaard, L. S., C. S. Rogers, Q. Dong, C. O. Randak, D. W. Vermeer, T. Rokhlina, *et al.* 2007. Processing and function of CFTR-DeltaF508 are species-dependent. *Proc. Natl Acad. Sci. USA* 104:15370–15375.
- Rapino, D., I. Sabirzhanova, M. Lopes-Pacheco, R. Grover, W. B. Guggino, and L. Cebotaru. 2015. Rescue of NBD2 mutants N1303K and S1235R of CFTR by small-molecule correctors and transcomplementation. *PLoS ONE* 10: e0119796.
- Salvatore, D., R. Buzzetti, E. Baldo, M. P. Forneris, V. Lucidi, D. Manunza, *et al.* 2011. An overview of international literature from cystic fibrosis registries. Part 3. Disease incidence, genotype/phenotype correlation, microbiology, pregnancy, clinical complications, lung transplantation, and miscellanea. *J. Cyst. Fibros.* 10:71–85.
- Southern, K. W., A. Munck, R. Pollitt, G. Travert, L. Zanolla, J. Dankert-Roelse, *et al.* 2007. A survey of newborn screening for cystic fibrosis in Europe. *J. Cyst. Fibros.*, 6:57–65.

- Tian, X., Y. Liu, J. Yang, H. Wang, T. Liu, W. Xu, *et al.* 2016. p. G970D is the most frequent CFTR mutation in Chinese patients with cystic fibrosis. *Hum. Genome Var.* 3:15063.
- Tsui, L. C. 1992. The spectrum of cystic fibrosis mutations. *Trends Genet.* 8:392–398.
- Wagner, J. A., A. Vassilakis, K. Yee, M. Li, G. Hurlock, M. E. Krouse, *et al.* 1999. Two novel mutations in a cystic fibrosis patient of Chinese origin. *Hum. Genet.* 104:511–515.
- Watson, M. S., G. R. Cutting, R. J. Desnick, D. A. Driscoll, K. Klinger, M. Mennuti, *et al.* 2004. Cystic fibrosis population carrier screening: 2004 revision of American College of Medical Genetics mutation panel. *Genet. Med.* 6:387–391.
- Welsh, M. J., and A. E. Smith. 1993. Molecular mechanisms of CFTR chloride channel dysfunction in cystic fibrosis. *Cell* 73:1251–1254.
- Wong, L. J. C., Ö. M. Alper, B. T. Wang, M. H. Lee, and S. Y. Lo. 2003. Two novel null mutations in a Taiwanese cystic fibrosis patient and a survey of East Asian CFTR mutations. *Am. J. Med. Genet. A* 120:296–298.
- Yamashiro, Y., T. Shimizu, S. Oguchi, T. Shioya, S. Nagata, and Y. Ohtsuka. 1997. The estimated incidence of cystic fibrosis in Japan. *J. Pediatr. Gastroenterol. Nutr.* 24:544–547.
- Yang, W., N. Shen, D. Q. Ye, Q. Liu, Y. Zhang, X. X. Qian, *et al.* 2010. Genome-wide association study in Asian populations identifies variants in ETS1 and WDFY4 associated with systemic lupus erythematosus. *PLoS Genet.* 6:e1000841.
- Zielenski, J., D. Markiewicz, S. P. Lin, F. Huang, T. L. Yangfeng, and L. C. Tsui. 1995. Skipping of exon 12 as a consequence of a point mutation (1898+5G→T) in the cystic fibrosis transmembrane conductance regulator gene found in a consanguineous Chinese family. *Clin. Genet.* 47:125–132.

Supporting Information

Additional Supporting Information may be found online in the supporting information tab for this article:

Figure S1. Birthplace of the individuals with p.I1023R mutation involved.

Figure S2. Principal components analysis (PC1 vs. PC2) of 612 Hans Chinese (gray dots) from Hong Kong and 12 individuals from the affected family with p.I1023R (black dots).

Figure S3. Target coverage of the childhood bronchiectasis panel. The panel includes 16 selected candidate genes for primary ciliary dyskinesia (PCD) and cystic fibrosis (CF).

Table S1. A list of reported CF Chinese patients with CFTR mutations identified.

Appendix S1. Experimental workflow.