

Effect of mean serum potassium level on in-hospital and long-term outcomes in ST segment elevation myocardial infarction

To the Editor,

Serum potassium level plays a substantial role in cardiovascular disorders. Based on previous studies, current guidelines for serum potassium in cardiovascular disorders recommend maintaining the serum potassium level of 4–4.5 mEq/L in patients with AMI (1, 2). Recent studies have examined the recommendations of the guidelines, and the lowest mortality was observed with serum potassium level of 3.5–4 mEq/L in patients with AMI (3, 4). Because of important improvements in revascularization and drug therapies, it is important to re-evaluate the impact of serum potassium level with respect to mortality and ventricular arrhythmias in patients with AMI. To address this critical diversity, further investigations are needed for confirmation. We report a study evaluating the impact of mean serum potassium level on in-hospital and long-term outcomes in a large patient population with STEMI (5). In total, 3,760 consecutive patients diagnosed with STEMI were retrospectively analyzed. At least two serum potassium measurements were taken from each patient, and the mean serum potassium levels were categorized accordingly: <3.0, 3.0–<3.5, 3.5–<4.0, 4.0–<4.5, 4.5–<5.0, 5.0–<5.5, and ≥ 5.5 mEq/L. Hierarchical logistic regression and Cox-proportional regression analysis were used to establish the relationship between mean serum potassium levels and clinical outcomes. The lowest in-hospital and long-term mortality was determined in patients with serum potassium level of 4–<4.5 mEq/L, whereas mortality was higher in patients with serum potassium levels of ≥ 5.0 and <3.5 mEq/L. There was a U-shaped association between mean serum potassium level and mortality. In multivariable hierarchical logistic regression analysis, in-hospital mortality risks were higher for patients with serum potassium level of ≥ 5 mEq/L [odds ratio (OR), 2.60; 95% confidence interval (CI), 1.30–4.2 and OR, 3.22; 95% CI, 1.14–9.07 for patients with serum potassium levels of 5–<5.5 mEq/L and ≥ 5.5 mEq/L, respectively]. In a multivariable Cox-proportional regression analysis, the mortality risk was higher for patients with serum potassium levels of ≥ 5 mEq/L [hazard ratio (HR), 2.11; 95% CI, 1.23–4.74 and HR, 4.20; 95% CI, 1.08–8.23, for patients with potassium levels of 5–<5.5 mEq/L and ≥ 5.5 mEq/L, respectively]. In-hospital and long-term mortality risks were also higher for patients with serum potassium levels of ≤ 3.5 mEq/L. Conversely, ventricular arrhythmias were higher only for patients with serum potassium levels of ≤ 3.5 mEq/L. Therefore, serum potassium level plays a substantial role in patients with AMI in terms of mortality and ventricular arrhythmias;

however, there is no current consensus on optimal serum potassium level in patients with AMI. The current study challenges the current guidelines in clinical practice, which recommend maintaining serum potassium level at 4.0–5.0 mEq/L in patients with AMI. In addition, our findings were in line with recent studies with respect to maintaining serum potassium level at 3.5–4.5 mEq/L. Although the lowest mortality and ventricular arrhythmia range in the current study was 4.0–4.5 mEq/L, it was 3.5–4.0 mEq/L in the recent studies.

Muhammed Keskin, Adnan Kaya¹, Mustafa Adem Tatlısu², Mert Ilker Hayroğlu

Department of Cardiology, Siyami Ersek Thoracic and Cardiovascular Surgery Center, Training and Research Hospital; İstanbul-Turkey

¹Department of Cardiology, Suruç State Hospital; Şanlıurfa-Turkey

²Department of Cardiology, Texas A&M University; Texas-USA

References

1. Macdonald JE, Struthers AD. What is the optimal serum potassium level in cardiovascular patients? *J Am Coll Cardiol* 2004; 43: 155-61.
2. Cohn JN, Kowey PR, Whelton PK, Prisant LM. New guidelines for potassium replacement in clinical practice: a contemporary review by the National Council on Potassium in Clinical Practice. *Arch Intern Med* 2000; 160: 2429-36.
3. Choi JS, Kim YA, Kim HY, Oak CY, Kang YU, Kim CS, et al. Relation of serum potassium level to long-term outcomes in patients with acute myocardial infarction. *Am J Cardiol* 2014; 113: 1285-90.
4. Uluganyan M, Ekmekçi A, Murat A, Avşar Ş, Ulutaş TK, Uyarel H, et al. Admission serum potassium level is associated with in-hospital and long-term mortality in ST-elevation myocardial infarction. *Anatol J Cardiol* 2016; 16: 10-5.
5. Keskin M, Kaya A, Tatlısu MA, Hayroğlu Mİ, Uzman O, Börklü EB, et al. The effect of serum potassium level on in-hospital and long-term mortality in ST elevation myocardial infarction. *Int J Cardiol* 2016; 221: 505-10.

Address for Correspondence: Dr. Muhammed Keskin

Dr. Siyami Ersek Hastanesi, Kardiyoloji Bölümü

Tıbbiye Cad., No: 25, Üsküdar, İstanbul-Türkiye

Phone: +90 216 542 44 44 Fax: +90 216 337 97 19

E-mail: drmuhammedkeskin@gmail.com

©Copyright 2016 by Turkish Society of Cardiology - Available online

at www.anatoljcardiol.com

DOI:10.14744/AnatolJCardiol.2016.7450

