

Research article

Open Access

Transient change in GABA_A receptor subunit mRNA expression in *Lurcher* cerebellar nuclei during Purkinje cell degeneration

C Linnemann*, I Schmech, P Thier and C Schwarz

Address: Department of Cognitive Neurology, Hertie-Institute for Clinical Brain Research, University of Tübingen, Otfried-Müller-Strasse 27, 72076 Tübingen, Germany

Email: C Linnemann* - Christoph.Linnemann@uni-tuebingen.de; I Schmech - IsabellaSchmech@hotmail.com; P Thier - Thier@uni-tuebingen.de; C Schwarz - Cornelius.Schwarz@uni-tuebingen.de

* Corresponding author

Published: 27 July 2006

Received: 10 March 2006

BMC Neuroscience 2006, 7:59 doi:10.1186/1471-2202-7-59

Accepted: 27 July 2006

This article is available from: <http://www.biomedcentral.com/1471-2202/7/59>

© 2006 Linnemann et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: *Lurcher* mice suffer from a complete Purkinje cell (PC) loss in the first four postnatal weeks. Parallel to this degeneration, GABAergic synapses in the deep cerebellar nuclei (DCN), the major recipient of the inhibitory PC projection, increase synaptic conductance. Here, we further investigated this phenomenon, using real-time RT-PCR to assess GABA_A receptor subunit gene expression during PC degeneration.

Results: We observed a specific reduction in $\gamma 2$ subunit gene expression, while $\alpha 1-5$, $\beta 1-2$, $\gamma 1,3$ and δ subunits were unaffected. We made two further specific findings. First, the difference in gene expression was shown in tissue from DCN only. Neither the hippocampus nor coronal sections through the forebrain showed such effects. Furthermore, the involvement of different levels of corticosterone, a possible humeral trigger for differences in gene expression, could be excluded. Second, like the known potentiation of GABAergic synapses, the $\gamma 2$ down-regulation was present only after the onset of degeneration at p14. The difference in $\gamma 2$ mRNA expression, however, appeared transient, since it was no longer detectable in adult *Lurcher* mice.

Conclusion: In conclusion, the down-regulation of $\gamma 2$ subunits may be related to differences in synaptic efficacy and, as such, may reflect the initial phase of adaptive responses of DCN tissue to massive GABAergic deafferentation. Its transient course, however, does not support the idea that modulations in GABAergic transmission are at the basis of the well-known DCN-based functional benefit of *Lurcher* mice present throughout their life.

Background

Inhibitory projections are sometimes placed at decisive locations within the motor system and are therefore important for programming and controlling the execution of movement. One of these projections is the GABAergic Purkinje cell (PC) projection onto the deep cerebellar nuclei (DCN) neurons. This projection system undergoes a degeneration in the spontaneous, autosomal semidom-

inant mouse mutation *Lurcher*, yielding nonviable homozygous mice, while heterozygous individuals (*Lc/+*) show ataxia expressed by a tendency to fall [1]. The syndrome is based on a complete loss of Purkinje cells during postnatal development followed by a secondary degeneration of intracerebellar and precerebellar cells [2-4]. The phenotype arises from a mutation in the orphan receptor subunit glutamate $\delta 2$ [5] which is expressed almost exclu-

sively in PCs and thus determines the specificity of the primary effect [6]. The first signs of PC degeneration are observed at postnatal day 8 (p8; [7]) and by p26, approximately 90% of PCs have disappeared [2]. The majority of DCN neurons survive, albeit deprived of their inhibitory PC input [8,9].

Behaviorally, *Lc/+* show deficits in standard tests of motor behavior like the rotarod and other tests of equilibrium [10-15]. However, the symptoms are mild, since differences to wild type animals were seen only in test situations that confront the motor system with higher demands such as, for instance, higher rotation frequencies in the rotarod test [12]. Experiments using lesions of the DCN in *Lc/+* clearly showed that the mild symptoms reflect a residual functionality of the cerebellar system because lesioned animals showed a grossly deteriorated motor performance in comparison with non-lesioned mutants. Scores measured with the rotarod test dropped virtually to zero without recovery during repeated sessions [16]. Furthermore, spatial orientation, which is mildly impaired in *Lc/+* [17,18], deteriorated drastically after DCN lesions [19,20]. These behavioral data suggest that the DCN play a beneficial role for the motor performance of *Lc/+*. Electrophysiological recordings in DCN neurons of *Lc/+* strongly support this hypothesis. *Lc/+* exhibited a potentiation of GABA_A-receptor-mediated average synaptic conductance (g_{syn}) in close temporal relationship to PC death [21]. This potentiation of g_{syn} in the DCN might have an adaptive value, compensating for the loss of the extrinsic inhibition. A possible basis for the modulation of GABAergic synaptic transmission are changes in the subunit composition of GABA_A receptors [22]. We asked in the present study whether the DCN change their GABA_A receptor subunit composition in temporal relationship to the degeneration of the inhibitory afferents. Guided by the occurrence of conductance changes between p11 and p14 [21], we used real-time RT-PCR to investigate the gene expression of GABA_A receptor subunits α 1-5, β 1-2, γ 1-3 and δ in DCN tissue of *Lc/+* compared with wild type mice at these critical ages and after the completion of PC degeneration at the age of 8 weeks. Furthermore, we addressed the question as to whether the observed changes are specific to the DCN. In addition, we performed whole cell voltage-clamp recordings of CA1 pyramidal neurons in mutants and controls to determine whether the potentiation in g_{syn} previously observed in DCN is a generalized and ubiquitous phenomenon or specific to the denervated DCN.

Results

Using real-time RT-PCR, mRNA expression levels of GABA_A receptor subunits α 1-5, β 1-2, γ 1-3 and δ were investigated in tissue of DCN obtained from *Lurcher* mutants aged p11 and p14 and compared to wild type

controls of the respective age. Since gene expression of GABA_A receptor subunits [23] and of internal controls such as hypoxanthine-phosphoribosyltransferase (HPRT) [24,25] are known to be subject to changes during postnatal development, we restricted our comparison of gene expression to age-matched mutants and controls.

Animals aged p11 showed no clear signs of the characteristic *Lc/+* phenotype in terms of an ataxic gait or tendency to fall. This time point was also before the onset of changes of g_{syn} [21]. In accordance with the lack of differences in behavior and electrophysiological recordings, at p11 we did not find any difference in GABA_A receptor subunit gene expression between *Lc/+* and wild type mice either. Figure 1 shows the ratios GABA_A receptor subunit/HPRT [$n = 5$ animals each except for wild type α 2 ($n = 4$) and γ 3 ($n = 3$)]. We next investigated the gene expression in the DCN tissue of animals aged p14. At p14, the animals were clearly distinguishable by their mild ataxia, due to the progressive PC degeneration. At this stage, *Lc/+* exhibit a potentiation of g_{syn} in DCN [21]. We observed a statistically significant decrement of the γ 2 subunit mRNA expression in *Lc/+* [1.6-fold down-regulation, ratio GABA_A receptor subunit/HPRT 1.9390 ± 0.9230 (*Lc/+*, $n = 9$) versus 3.0410 ± 1.1530 (wild type, $n = 10$), $P = 0.036$]. With regard to the other subunits, no differences [$n = 5$ animals each except for *Lc/+* α 5, β 1, γ 1 ($n = 9$) and wild type α 5, β 1, γ 1 ($n = 10$)] were found. Figure 2 summarizes these results. The decrement of γ 2 subunit mRNA expression in *Lc/+* DCN tissue was of transient nature, since it could not be detected in DCN tissue of eight week old mutants ($n = 8$) compared to age-matched controls ($n = 8$, $P = 1.000$; Figure 3).

Such decrement of γ 2 mRNA at p14 was absent in Lurcher hippocampus (CA1/2) ($n = 5$ animals each) and in coronal slices cut through the forebrain at the optic chiasm ($n = 5$ animals each; Figure 4) – suggesting that the difference in GABA_A receptor subunit gene expression occurred specifically in the DCN. Furthermore, differences in GABAergic transmission, as found in DCN during PC degeneration [21], were not observed in hippocampal slices taken from mice aged p15-18. Voltage-clamp recordings from hippocampal CA1 pyramidal neurons ($n = 6$ for mutant neurons, $n = 5$ for wild type neurons) revealed dense patterns of mIPSCs (Figure 5A, top traces). These were GABA_A-receptor-mediated events, as judged from the equilibrium potential under chloride symmetrical conditions and pharmacological interference (Figure 5). The equilibrium potentials were close to the expected 0 mV in both mutant and control neurons [$+1.8 \pm 6.4$ mV (*Lc/+*, $n = 6$); $+2.8 \pm 3.9$ mV (wild type, $n = 5$), $P = 0.758$]. The mIPSCs could be blocked regularly in both *Lc/+* and wild type mice ($n = 6$ recordings) by application of 10 μ M bicuculline, indicating that they were GABAergic (Figure




Figure 1

mRNA expression [GABA_A receptor subunit/hypoxanthine-phosphoribosyltransferase (HPRT)] in DCN tissue of *Lc/+* (black bars) and wild type mice (WT; grey bars; Mean ± SD) aged postnatal day 11 (p11). No statistically significant differences were found for any of the investigated subunits. The data shown in the inset are from [21] and give the average synaptic conductance (g_{syn}) values (with SD as error bars) obtained from mIPSCs recorded from DCN of animals <math><p14</math>. No statistically significant differences were detected between *Lc/+* [black bars, 0.50 ± 0.39 nS ($n = 3$)] and WT [grey bars; 0.39 ± 0.16 nS ($n = 4$), $P = 0.62$].

5C). The average synaptic conductance values were statistically not different between the two groups (Figure 6A). The parameters reflecting the kinetic properties of the mIPSCs (rise 10–90 as well as decay times) did not differ significantly either (Figures 6B and 6C). An analysis of the frequencies of the mIPSCs revealed no differences between mutants and controls (Figure 6D).

To rule out the possibility that different corticosterone levels at the time of animal preparation might affect our electrophysiological and/or gene expression results, serum corticosterone levels were measured in mutant and control mice aged p14. To avoid variations due to sex differences, we analyzed the data obtained from male and female mice separately. We found no differences in corticosterone serum level between mutants and controls (Figure 7).

Discussion and conclusion

The residual motor functionality of *Lurcher* mice is based on the intactness of the DCN and outlasts the period in which PCs degenerate [16]. To elucidate the role of candidate cellular mechanisms for this compensation, their

specificity for the structure (DCN) and time of occurrence (degeneration of PCs) must be investigated. Mechanisms showing no clear specificity and relationship to the DCN (or connected structures) or to the occurrence of the PC degeneration can be excluded as a basis for this functional compensation. In a previous study from our laboratory, *Lurcher* mice displayed a potentiation of miniature IPSCs in the DCN in close temporal relationship to PC degeneration [21]. Our present data complement these earlier results by showing that the synaptic conductance of GABAergic IPSCs is unaffected in another brain structure, the hippocampus.

The responsiveness of the hypothalamo-pituitary-adrenal system is known to be enhanced in *Lurcher* mice. Under stress, this mutant shows increased levels of serum corticosterone compared to wild type controls [26,27]. Corticosterone has a known modulatory role for GABA_A-receptor-mediated conductance [28] as well as GABA_A receptor subunit gene expression [29,30]. The absence of significant differences of serum corticosterone between *Lurcher* and wild type mice (at the ages and under the experimental conditions investigated here), thus supports




Figure 2

mRNA expression [GABA_A receptor subunit/hypoxanthine-phosphoribosyltransferase (HPRT)] in DCN tissue of *Lc/+* (black bars) and wild type mice (WT; grey bars; Mean ± SD) aged postnatal day 14 (p14). *Statistically significant difference for $\gamma 2$ (n = 9 for mutants/10 for WT; P = 0.036); no statistically significant differences were observed in any of the other investigated subunits. The data shown in the inset are from [21] and give the average synaptic conductance (g_{syn}) values (with SD as error bars) obtained from mIPSCs recorded from DCN of animals \geq p14. *Statistically significant difference between *Lc/+* [black bars; 1.23 ± 0.51 nS (n = 10)] and WT [grey bars; 0.57 ± 0.21 nS (n = 19), $P \leq 0.001$].

the idea that the changes observed in DCN of *Lurcher* mice are not unspecific phenomena that occur throughout the brain. The results instead suggest that these changes are specific to the DCN. The down-regulation of $\gamma 2$ mRNA matches the spatiotemporal specificity of the electrophysiological changes in GABAergic transmission. This was detected in the DCN, but not in material taken from the hippocampus or from a section through the forebrain. Furthermore, its de novo occurrence at p14 indicates a temporal relationship to the course of PC degeneration. However, since it was not detectable in adulthood (eight weeks of age), the reduction of subunit gene expression cannot be the direct structural cause of the permanent mild expression of ataxia in *Lurcher* mice. If future work is to show a functional link between subunit composition and synaptic potentiation, the latter must be subjected to the same reasoning and must be excluded from the candidates of mechanisms directly responsible for functional adaptation in the DCN. The main question to be discussed below is therefore, to what extent can a changed subunit expression relate to changes in synaptic transmission in the *Lurcher* mouse.

First and foremost, the differences in gene expression were calculated with reference to the ubiquitously expressed HPRT. Therefore, the measurements should be insensitive to known differences of volume and cell densities in *Lurcher* DCN [8,9]. This may help to explain apparent conflicting results obtained in adult *Lurcher* mice using in situ hybridization, where increased $\gamma 2$ mRNA levels in *Lurcher* mice aged p108 based on the assessment of autoradiographic grain density were reported [31]. The different results may be partly explained by the different ages of animals and the different methods used, but the shrinkage of the DCN (confer [8,9]) may well have contributed to an increase in grain density (as discussed in [31]). The insensitivity of the real-time RT-PCR to shrinkage of the DCN is further supported by the exclusive reduction of the $\gamma 2$ subunit mRNA expression. If general differences in cell densities, volume and presence of GABAergic synapses in DCNs of *Lurcher* and wild type mice had accounted for differences in mRNA expression, the consequences should have affected all, or at least many, subunits. Our findings imply that the relative composition of postsynaptic GABA_A receptors during the time of PC degeneration was different in *Lurcher* mice. This conclusion is in line with the probable postsynaptic location of the mechanism,




Figure 3
mRNA expression [GABA_A receptor subunit γ 2/hypoxanthine-phosphoribosyltransferase (HPRT)] in DCN tissue of *Lc/+* relative to wild type mice (WT; Mean \pm SD) aged 11 days [n = 5 (*Lc/+*), n = 5(WT)] 14 days [n = 9 (*Lc/+*), n = 10 (WT)], 8 weeks [n = 8 (*Lc/+*), n = 8 (WT)]. *Statistically significant difference between mutants and controls aged 14 days; no statistically significant differences were observed between mutants and controls aged 11 days as well as 8 weeks.

leading to the increment in miniature synaptic conductance [21]. There are two possible explanations as to how the present findings may be related to changes in synaptic conductance. First, it is well established that the compositions of subunits determine single channel conductances [32-35]; see for review [22]. Second, subunit composition specifically affects the dynamic control of GABA_A receptor expression at the cell surface and may thus effectively determine its presence at the postsynaptic membrane ([36-39]; see for review also [40]).

If modulation of single channel conductances were at the basis of the synaptic potentiation, then our present data, together with earlier results, [21] would imply that a reduced γ 2 content increases channel conductance and hence synaptic conductance. This prediction, however, would be in conflict with previously published data of several investigators. Verdoorn et al. [32] analyzed GABA-gated chloride channels expressed in human embryonic kidney cells. When the γ 2 subunit was present, the resulting GABA_A receptors had a larger conductance. Moreover, Lorez et al. [35] investigated the channel properties of neuronal GABA_A receptors from brains of embryonic and




Figure 4
mRNA expression [GABA_A receptor subunit γ 2/hypoxanthine-phosphoribosyltransferase (HPRT)] in hippocampus (CA1/2) and in coronal sections cut through the forebrain at the optic chiasm of *Lc/+* (black bars) and wild type mice (WT, grey bars, Mean \pm SD) aged postnatal day 14. No statistically significant differences were found for hippocampus or for forebrain (coronal slice at the optic chiasm).

postnatal mice lacking the γ 2 subunit and found smaller single channel conductances. To rescue this scenario, one would need to assume that the decrement of the γ 2 subunit mRNA expression found here is accompanied by the replacement of γ 2 by another subunit that has the potential to increase the g_{syn} , but that has not been detected so far. Replacements for GABA_A receptor subunit composition during postnatal development under physiological conditions are well known. For instance, α 2 subunits are partially replaced by α 1 subunits in DCN during the first two postnatal weeks [23]. Such switches have indeed the potential to change the synaptic strength of GABA_A receptors [41,42].

The second possibility is that subunit composition is altered to dynamically regulate the receptor availability at the cell surface. DCN neurons in rodents express mainly




Figure 5
A: Recordings of mIPSCs in hippocampal CA1 pyramidal neurons of *Lc/+* [postnatal day (p) 15] and wild type mice (WT; p17) under chloride symmetrical conditions at the indicated holding potentials (V_h). **B:** The mean peak amplitudes (I_{mean}) of the mIPSCs were plotted versus the V_h values and a regression line ($r^2 = 0.97$ for *Lc/+*, $r^2 = 0.98$ for WT) was generated to determine the equilibrium potential that was close to the expected 0 mV (+2.2 mV for *Lc/+*; -1.6 mV for WT). **C:** Application of bicuculline 10 μM blocked the mIPSCs, indicating that they were GABA_A-receptor-mediated events.




Figure 6
 Mean values (with SD as error bars) for the average synaptic conductance (g_{syn} , **A**), the rise 10–90 (**B**), the decay times (**C**) and the frequency (**D**) obtained from recordings of mIPSCs in hippocampal CA1 neurons of *Lc/+* (black bars) and wild type mice (WT, grey bars). Mice were aged p15-18. No statistically significant differences were found for any of the parameters.

$\alpha 1$, $\beta 2$ and $\gamma 2$ subunits [31,43]. The presence of the $\alpha 1$ subunit appears to be required for the surface expression of $\beta 2$ (as investigated in Madin-Darby canine kidney cells [37] and in COS7 cells [36]). Our study shows that the content of $\alpha 1$ and $\beta 2$ mRNA remained unchanged in *Lurcher* mice compared to the age-matched controls. None of these facts suggest that the amount of generated GABA_A receptors and their subsequent transport to the cell membrane is impaired in *Lurcher* mice. On the other hand, it may also be possible to regulate receptor presence by interfering with the removal of receptors from the membrane. GABA_A receptors undergo a constitutive endocytosis after association with the adaptin AP2 complex, a process dependent on the presence of β and γ subunits [39]. Moreover, blockade of endocytosis enhances the amplitude of GABA_A-mediated mIPSCs in hippocampal neurons [39]. More specifically, Connolly et al. [38] ascer-




Figure 7
 No statistically significant differences were found in corticosterone serum levels in *Lc/+* and wild type mice (WT) aged postnatal day 14. Corticosterone serum levels were measured as described in the Methods section. n indicates the number of animals.

tained that cell surface stability of GABA_A receptors depends on the presence of $\gamma 2$ subunits. They observed that receptors containing $\gamma 2$ are selectively removed from the cell surface upon activation of protein kinase C with concomitant reduction of GABA-induced currents. In summary, a tentative hypothesis – to be tested by further experiments – can be formulated from these considerations as follows: The specific $\gamma 2$ reduction of subunit expression during PC degeneration in *Lurcher* mice might modulate endocytosis such as to allow the *Lurcher* DCN neurons to specifically increase the GABA_A receptor cell surface levels at the postsynaptic membrane. This speculation receives some experimental support from the results of a non-stationary fluctuation analysis on mIPSCs in wild type and *Lurcher* mice. This analysis gave no indication of changes of single channel conductances, while the peak number of open receptors underlying *Lurcher* mIPSCs was estimated to be of a factor two higher than in wild type mice [21].

Methods

Animals and molecular genotyping

Heterozygous B6CBACa-A^w-1/A-*Lc* (Jackson Laboratories, Bar Harbor, ME, USA) were mated with B6CBA (Harlan-

Table 1: Primers for GABA_A receptor subunits and the internal control

	forward	reverse
GABA _A alpha1	5'-gccccgagctgtgcaa-3'	5'-ggtccagcccaagatagtc-3'
GABA _A alpha2	5'-acagtcgaagcgaatgtcc-3'	5'-aacggagtcagaagcattgtaagt-3'
GABA _A alpha3	5'-aacaaatctgtggaagtagcacagg-3'	5'-tctctgtcccaacaacatggcc-3'
GABA _A alpha4	5'-ataagtgcggagtccccatg-3'	5'-tcccaaatccaaagggcag-3'
GABA _A alpha5	5'-gcagcttgaggactttccaatg-3'	5'-ggtaagcataactgccaatttca-3'
GABA _A beta1	5'-accagagtccaatgaaaagaac-3'	5'-gctgaggagaatattgccgtg-3'
GABA _A beta2	5'-ccacatccgaagcagtaatggg-3'	5'-tgctggaggcatcatagggc-3'
GABA _A gamma1	5'-cactctgtactcactgtgtgt-3'	5'-tccttcctcactagctcagtc-3'
GABA _A gamma2	5'-caaatacatggagcattggaagc-3'	5'-tcatagtcacatctgactttggc-3'
GABA _A gamma3	5'-ttcagttgaggcagctgatca-3'	5'-ctgaggccatgaagcacaac-3'
GABA _A delta	5'-gtggccagcattgacatc-3'	5'-tctgatgcaggaaacagctcag-3'
HPRT	5'-gcctaagatgagcgaagttg-3'	5'-tactaggcagatggccacagg-3'

Winkelmann, Borchon, Germany) mice. The animals were kept and used in experiments according to the institutional and national animal care guidelines (also conforming with NIH guidelines on animal care). Animals from the *Lurcher* progeny not showing the characteristic *Lc/+* phenotype (ataxic gait and tendency to fall) were genotyped as described previously [21].

Real-time RT-PCR

Following deep anesthesia by application of ketamine (150 mg/kg i.p.), the brain was removed and prepared as follows under RNase-free conditions. DCN tissue was dissected from parasagittal cerebellar slices (400–500 µm thickness). Hippocampal CA1/2 tissue was isolated from horizontal cerebral slices (400–500 µm thickness). Fore-brain slices were obtained from 1000–1200 µm thick coronal slices, cut through the forebrain at the optic chiasm. The tissue samples were immediately transferred into RNA later RNA Stabilization Reagent (Qiagen, Hilden, Germany) to stabilize and protect RNA in tissue. Total RNA was isolated using RNeasy Mini Kit (Qiagen) including a rotor-stator homogenization (Ultra Turrax T8, IKA-Werke, Staufen, Germany) of the samples. All samples were treated with DNase (*DNA-free*, Ambion, Huntingdon, United Kingdom) and tested for contaminating genomic DNA by PCR using a non-intron spanning primer pair for β-actin (forward 5'-caccgccaccagttcgcca-3', reverse 5'-caggtcccggccagcaggt-3'; MWG-Biotech, Ebersberg, Germany). First-strand cDNA was prepared from total RNA by reverse transcription using the Superscript II RNase H⁻ Reverse Transcriptase (Invitrogen, Karlsruhe, Germany) and random hexamers. Gene expression was measured in the ABI Prism 7700 Sequence Detection System (Applied Biosystems, Foster City, California, USA). Primers (MWG-Biotech, Ebersberg, Germany) were selected to result in amplicons less than 150 bp to increase the efficiency of PCR amplification. The sequences of the primers are listed in Table 1. Product specificity of the PCR products was confirmed by

dissociation curve analysis and/or agarose gel electrophoresis. Amplification of specimens and serial dilutions of standards (PAGE-purified oligonucleotides; Thermo Electron, Ulm, Germany) was carried out in a total volume of 15 µl including Sybrgreen Master Mix (Applied Biosystems) and primers (final concentration 300 nM) under the following conditions: 50 °C for 2 min; 95 °C for 10 min; 40 cycles with 95 °C for 15 s and 60 °C for 1 min. Standard curve extrapolations were calculated for the genes of interest to obtain the quantity of starting amount for each sample. All results were obtained from duplicates and normalized with respect to the internal control HPRT to correct for sample to sample variations.

Electrophysiology

Mice aged p15-18 (without statistically significant age differences between mutants and controls) were deeply anesthetized by application of ketamine (150 mg/kg i.p.). The brains were removed and placed in ice-cold modified artificial cerebrospinal fluid [ACSF, containing (in mM) 126 sucrose, 2.5 KCl, 1.3 NaH₂PO₄, 3 MgCl₂, 26 NaHCO₃, 0.1 CaCl₂ and 20 D-glucose] that was bubbled with 95%O₂-5%CO₂. Horizontal slices of 275 µm thickness that included the hippocampus were obtained using a vibratome (VT1000s, Leica, Nussloch, Germany) and transferred into modified ACSF at room temperature. The modified ACSF was replaced within the next 90 min with ACSF containing (in mM) 125 NaCl, 2.5 KCl, 1.3 NaH₂PO₄, 2 MgCl₂, 26 NaHCO₃, 2 CaCl₂ and 20 D-glucose, oxygenated with 95%O₂-5%CO₂ at room temperature. Experiments were performed with this ACSF to which the selective non-NMDA receptor antagonist 6,7-dinitroquinoxaline-2,3-dione (25 µM; Tocris, Bristol, UK), tetrodotoxine (1 µM, Tocris) and, in some recordings, also the GABA_A antagonist bicuculline (10 µM; Sigma, Taufkirchen, Germany) were added.

Whole-cell voltage-clamp recordings of hippocampal CA1 pyramidal neurons were conducted with patch pipettes

that had a resistance of 2.5–3.5 M Ω when filled with a solution containing (in mM) 124 CsCl, 10 K⁺-HEPES, 5 EGTA, 4.6 MgCl₂, 4 K⁺-ATP, 0.4 Na⁺-GTP, 0.1 CaCl₂, and 5 QX-314 (Tocris), adjusted to pH 7.3 with CsOH, thus obtaining chloride symmetrical experimental conditions. CA1 pyramidal neurons were identified and the patch procedure was visualized using a motorized (Luigs&Neumann, Ratingen, Germany) microscope (Axioscope, Zeiss, Göttingen, Germany) with water immersion objective ($\times 40$, Zeiss numerical aperture 0.75), infrared illumination, Normaski optics and infrared sensitive CCD camera (Newvicon C2400-07-C Hamamatsu, Japan). The recordings were performed with an EPC-7 amplifier (Heka, Lambrecht, Germany). The data was sampled at a rate of 25 kHz (Spike2, CED, Cambridge, UK). Neurons were voltage-clamped at -70.6 ± 0.7 mV. Data was not quantitatively analyzed if, under these conditions, the serial resistance (17.9 ± 7.3 M Ω , $n = 11$) had changed by more than 20% or if the membrane resistance (221.8 ± 66.8 M Ω , $n = 11$) of the recorded neuron was below 150 M Ω . Neither the membrane nor the serial resistances differed significantly between genotypes.

Analysis of the miniature inhibitory postsynaptic currents (mIPSCs) was performed with the MiniAnalysis program (Synaptosoft, Decatur, GA, USA). For analysis of the synaptic waveforms, only those synaptic events that had not been interrupted by other synaptic events were evaluated. g_{syn} values were calculated as $g_{syn} = I_{mean}/(V_h - E)$, where I_{mean} is the mean peak amplitude of the mIPSCs, V_h is the holding potential and E is the measured equilibrium potential for the mPSCs of the respective recording. The decay time mentioned in this paper is the time it takes the current values to decay to $1/e$ of the peak values of the mIPSC amplitudes.

¹²⁵I radioimmunoassay

Following deep anesthesia by application of ketamine (150 mg/kg i.p.), blood was collected by intracardial puncture and immediately transferred into a plasma separator tube filled with lithium heparin (BD Microtainer PST LH, Becton Dickinson, Heidelberg, Germany) on ice. After centrifugation, the sera were stored at -20°C until the ¹²⁵I radioimmunoassay was performed. The Coat-A-Count Rat Corticosterone (TRKC 1; DPC Biermann, Bad Nauheim, Germany) procedure is a solid-phase ¹²⁵I radioimmunoassay. ¹²⁵I-labelled rat corticosterone competed for 2 h at room temperature with corticosterone in the serum specimens for antibody sites immobilized to the wall of the tubes. The counts per minute measured in the tubes with a gamma counter (Multi-Crystal Counter LB 2104, Berthold, Bad Wildbad, Germany), converted by way of a calibration curve yielded to a measure of the corticosterone serum level. The minimal detectable dose is approximately 5.7 ng/ml. The blood collections were

always performed in the same time window in the afternoon, to avoid variations due to circadian rhythm.

Statistical analysis

Statistically significant differences and correlations were tested with the Student's t-test for normally distributed data, and otherwise with the Mann-Whitney-Test (italicized P values throughout the paper). Statistical significance was assumed if $P < 0.05$.

Authors' contributions

CL, PT and CS designed and analyzed the experiments and drafted the manuscript. CL and IS carried out the experiments. All authors read and approved the final manuscript.

Acknowledgements

The sequencing procedures were performed by the sequencing service facility of the Center for Interdisciplinary Clinical Research, Tübingen. We thank Ute Großhennig and Britta Baumann for technical assistance. We are also grateful to Dr. Martin Möck of the Department of Anatomy, University of Tübingen, and Dr. Bernd Wissinger of the Molecular Genetics Laboratory, University Eye Hospital Tübingen for discussions, advice and support.

This work was supported by the German Science Foundation (SFB 430-C6, SCHW 577/5-1), the Center for Interdisciplinary Clinical Research, Tübingen (IZKF IC1) and the Hermann and Lilly Schilling Foundation.

References

1. Phillips RJS: "**Lurcher**", new gene in the linkage group XI of the house mouse. *J Genet* 1960, **57**:35-42.
2. Caddy KW, Biscoe TJ: **Structural and quantitative studies on the normal C3H and Lurcher mutant mouse**. *Phil Trans R Soc Lond B* 1979, **287**:167-201.
3. Wetts R, Herrup K: **Interaction of granule, Purkinje and inferior olivary neurons in Lurcher chimaeric mice. I. Qualitative studies**. *J Embryol Exp Morphol* 1982, **68**:87-98.
4. Wetts R, Herrup K: **Interaction of granule, Purkinje and inferior olivary neurons in lurcher chimeric mice. II. Granule cell death**. *Brain Res* 1982, **250**:358-362.
5. Zuo J, De Jager PL, Takahashi KA, Jiang W, Linden DJ, Heintz N: **Neurodegeneration in Lurcher mice caused by mutation in $\delta 2$ glutamate receptor gene**. *Nature* 1997, **388**:769-773.
6. Yang N, Horn R: **Evidence for voltage-dependent S4 movement in sodium channels**. *Neuron* 1995, **15**:213-218.
7. Dumesnil-Bousez N, Sotelo C: **Early development of the Lurcher cerebellum: Purkinje cell alterations and impairment of synaptogenesis**. *J Neurocytol* 1992, **21**:506-529.
8. Heckroth JA: **Quantitative morphological analysis of the cerebellar nuclei in normal and lurcher mutant mice. I. Morphology and cell number**. *J Comp Neurol* 1994, **343**:173-182.
9. Sultan F, König T, Möck M, Thier P: **Quantitative organization of neurotransmitters in the deep cerebellar nuclei of the Lurcher mutant**. *J Comp Neurol* 2002, **452**:311-323.
10. Lalonde R, Botez MI, Joyal CC, Caumartin M: **Motor abnormalities in lurcher mutant mice**. *Physiol Behav* 1992, **51**:523-525.
11. Lalonde R: **Motor learning in lurcher mutant mice**. *Brain Res* 1994, **639**:351-353.
12. Le Marec N, Caston J, Lalonde R: **Impaired motor skills on static and mobile beams in lurcher mutant mice**. *Exp Brain Res* 1997, **116**:131-138.
13. Le Marec N, Lalonde R: **Sensorimotor learning and retention during equilibrium tests in Purkinje cell degeneration mutant mice**. *Brain Res* 1997, **768**:310-316.
14. Le Marec N, Lalonde R: **Treadmill performance of mice with cerebellar lesions: I. Purkinje cell degeneration mutant mice**. *Behav Neurosci* 1998, **112**:225-232.

15. Porrás-García E, Cendelin J, Dominguez-del-Toro E, Vozech F, Delgado-García JM: **Purkinje cell loss affects differentially the execution, acquisition and prepulse inhibition of skeletal and facial motor responses in Lurcher mice.** *Eur J Neurosci* 2005, **21**:979-988.
16. Caston J, Vasseur F, Stelz T, Chianale C, Delhaye-Bouchaud N, Mariani J: **Differential roles of cerebellar cortex and deep cerebellar nuclei in the learning of the equilibrium behavior: studies in intact and cerebellectomized lurcher mutant mice.** *Brain Res Dev Brain Res* 1995, **86**:311-316.
17. Lalonde R, Joyal CC, Guastavino J-M, Botez MI: **Hole poking and motor coordination in lurcher mutant mice.** *Physiol Behav* 1993, **54**:41-44.
18. Lalonde R, Joyal CC, Côte C, Botez MI: **Simultaneous visual discrimination learning in lurcher mutant mice.** *Brain Res* 1993, **618**:19-22.
19. Hilber P, Jouen F, Delhaye-Bouchaud N, Mariani J, Caston J: **Differential roles of cerebellar cortex and deep cerebellar nuclei in learning and retention of a spatial task: studies in intact and cerebellectomized lurcher mutant mice.** *Behav Genet* 1998, **28**:299-308.
20. Caston J, Vasseur F, Delhaye-Bouchaud N, Mariani J: **Delayed spontaneous alternation in intact and cerebellectomized control and lurcher mutant mice: differential role of cerebellar cortex and deep cerebellar nuclei.** *Behav Neurosci* 1997, **111**:214-218.
21. Linnemann C, Sultan F, Pedroarena CM, Schwarz C, Thier P: **Lurcher mice exhibit potentiation of GABA_A-receptor-mediated conductance in cerebellar nuclei neurons in close temporal relationship to Purkinje cell death.** *J Neurophysiol* 2004, **91**:1102-1107.
22. Hevers W, Lüddens H: **The diversity of GABA_A receptors. Pharmacological and electrophysiological properties of GABA_A channel subtypes.** *Mol Neurobiol* 1998, **18**:35-86.
23. Fritschy JM, Paysan J, Enna A, Möhler H: **Switch in the expression of GABA_A-receptor subtypes during postnatal development: An immunohistochemical study.** *J Neurosci* 1994, **14**:5302-5324.
24. Gutensohn W, Guroff G: **Hypoxanthine-guanine-phosphoribosyltransferase from rat brain (purification, kinetic properties, development and distribution).** *J Neurochem* 1972, **19**:2139-2150.
25. Brosh S, Sperling O, Bromberg Y, Sidi Y: **Developmental changes in the activity of enzymes of purine metabolism in rat neuronal cells in culture and in whole brain.** *J Neurochem* 1990, **54**:1776-1781.
26. Frederic F, Chautard T, Brochard R, Chianale C, Wollman E, Oliver C, Delhaye-Bouchaud N, Mariani J: **Enhanced endocrine response to novel environment stress and endotoxin in Lurcher mutant mice.** *Neuroendocrinology* 1997, **66**:341-347.
27. Hilber P, Lorivel T, Delarue C, Caston J: **Stress and anxious-related behaviors in Lurcher mutant mice.** *Brain Res* 2004, **1003**:108-112.
28. Teschemacher A, Zeise ML, Zieglgänsberger W: **Corticosterone-induced decrease of inhibitory postsynaptic potentials in rat hippocampal pyramidal neurons in vitro depends on cytosolic factors.** *Neurosci Lett* 1996, **215**:83-86.
29. Orchinik M, Carroll SS, Li Y-H, McEwen BS, Weiland NG: **Heterogeneity of Hippocampal GABA_A Receptors: Regulation by Corticosterone.** *J Neurosci* 2001, **21**:330-339.
30. Stone DJ, Walsh JP, Sebro R, Stevens R, Pantazopolous H, Benes FM: **Effects of pre- and postnatal corticosterone exposure on the rat hippocampal GABA system.** *Hippocampus* 2001, **11**:492-507.
31. Luntz-Leybman V, Frosthalm A, Fernando L, DeBlas A, Rotter A: **GABA_A/benzodiazepine receptor gamma 2 subunit gene expression in developing normal and mutant mouse cerebellum.** *Brain Res Mol Brain Res* 1993, **19**:9-21.
32. Verdoorn TA, Draguhn A, Ymer S, Seeburg PH, Sakman B: **Functional properties of recombinant rat GABA_A receptors depend upon subunit composition.** *Neuron* 1990, **4**:919-928.
33. Fisher JL, Macdonald RL: **Single channel properties of recombinant GABA_A receptors containing gamma 2 or delta subtypes expressed with alpha 1 and beta 3 subtypes in mouse L929 cells.** *J Physiol* 1997, **505**:283-297.
34. Neelands TR, Macdonald RL: **Incorporation of the pi subunit into functional gamma-aminobutyric Acid(A) receptors.** *Mol Pharmacol* 1999, **56**:598-610.
35. Lorez M, Benke D, Luscher B, Mohler H, Benson JA: **Single-channel properties of neuronal GABA_A receptors from mice lacking the gamma 2 subunit.** *J Physiol* 2000, **527**:111-131.
36. Bollan K, King D, Robertson LA, Brown K, Taylor PM, Moss SJ, Connolly CN: **GABA_A receptor composition is determined by distinct assembly signals within alpha and beta subunits.** *J Biol Chem* 2003, **278**:4747-4755.
37. Connolly CN, Wooltorton JRA, Smart TG, Moss SJ: **Subcellular localization of gamma-aminobutyric acid type A receptors is determined by receptor beta subunits.** *Proc Natl Acad Sci USA* 1996:9899-9904.
38. Connolly CN, Kittler JT, Thomas P, Uren JM, Brandon NJ, Smart TG, Moss SJ: **Cell surface stability of gamma-aminobutyric acid type A receptors. Dependence on protein kinase C activity and subunit composition.** *J Biol Chem* 1999, **274**:36565-36572.
39. Kittler JT, Delmas P, Jovanovic JN, Brown DA, Smart TG, Moss SJ: **Constitutive endocytosis of GABA_A receptors by an association with the adaptin AP2 complex modulates inhibitory synaptic currents in hippocampal neurons.** *J Neurosci* 2000, **20**:7972-7977.
40. Moss SJ, Smart TG: **Constructing inhibitory synapses.** *Nat Rev Neurosci* 2001, **2**:240-250.
41. Okada M, Onodera K, Van Renterghem C, Sieghart W, Takahashi T: **Functional correlation of GABA_A receptor subunits expression with the properties of IPSCs in the developing thalamus.** *J Neurosci* 2000, **20**:2202-2208.
42. Ortinski PI, Lu C, Takagaki K, Fu Z, Vicini S: **Expression of distinct subunits of GABA_A receptor regulates inhibitory synaptic strength.** *J Neurophysiol* 2004, **92**:1718-1727.
43. Persohn E, Malherbe P, Richards JG: **Comparative molecular neuroanatomy of cloned GABA_A receptor subunits in the rat CNS.** *J Comp Neurol* 1992, **326**:193-216.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

