

Prostaglandin E₂ (PGE₂) Exerts Biphasic Effects on Human Tendon Stem Cells

Jiaying Zhang, James H-C. Wang*

MechanoBiology Laboratory, Departments of Orthopaedic Surgery, Bioengineering, Mechanical Engineering and Materials Science, and Physical Medicine and Rehabilitation, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of America

Abstract

Prostaglandin E₂ (PGE₂) has been reported to exert different effects on tissues at low and high levels. In the present study, cell culture experiments were performed to determine the potential biphasic effects of PGE₂ on human tendon stem/progenitor cells (hTSCs). After treatment with PGE₂, hTSC proliferation, stemness, and differentiation were analyzed. We found that high concentrations of PGE₂ (>1 ng/ml) decreased cell proliferation and induced non-tenocyte differentiation. However, at lower concentrations (<1 ng/ml), PGE₂ markedly enhanced hTSC proliferation. The expression levels of stem cell marker genes, specifically SSEA-4 and Stro-1, were more extensive in hTSCs treated with low concentrations of PGE₂ than in cells treated with high levels of PGE₂. Moreover, high levels of PGE₂ induced hTSCs to differentiate aberrantly into non-tenocytes, which was evident by the high levels of PPAR γ , collagen type II, and osteocalcin expression in hTSCs treated with PGE₂ at concentrations >1 ng/ml. The findings of this study reveal that PGE₂ can exhibit biphasic effects on hTSCs, indicating that while high PGE₂ concentrations may be detrimental to tendons, low levels of PGE₂ may play a vital role in the maintenance of tendon homeostasis *in vivo*.

Citation: Zhang J, Wang JH-C (2014) Prostaglandin E₂ (PGE₂) Exerts Biphasic Effects on Human Tendon Stem Cells. PLoS ONE 9(2): e87706. doi:10.1371/journal.pone.0087706

Editor: Giovanni Camussi, University of Torino, Italy

Received: August 28, 2013; **Accepted:** January 2, 2014; **Published:** February 4, 2014

Copyright: © 2014 Zhang, Wang. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: This work was supported in part by National Institutes of Health under award numbers AR061395, AR060920, and AR049921. No additional external funding received for this study. The funding agency had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: wanghc@pitt.edu

Introduction

Tendons transmit muscular forces to bone and, as a result, they are subjected to large mechanical loads *in vivo*. Consequently, tendons are frequently injured, especially during intense sport activities. Tendon injuries are generally difficult to treat; tendinopathy, a chronic tendon disorder involving tendon inflammation and/or degeneration, is a particularly significant challenge in orthopaedics and sports medicine. Thus far, strategies that stimulate the complete regeneration of tendons after injury have not been developed. To this end, a better understanding of tendon cell biology is essential to devise improved treatment options for tendon injuries such as tendinopathy [1].

One of the major causative factors that contribute to the development of tendinopathy is excessive mechanical loading (or overuse and over-loading) placed on tendons [2,3]. Such excessive mechanical loading has been shown to increase the production of prostaglandin E₂ (PGE₂) in cultures of human tendon fibroblasts (tenocytes) *in vitro* [3,4]. In addition, PGE₂ production was shown to increase after exercise in the peritendinous space of Achilles tendons *in vivo* [5].

Although PGE₂ levels increase after mechanical loading, baseline levels of PGE₂ are present in the patellar and Achilles tendons of mice under normal conditions without mechanical loading such as treadmill running [6]. This suggests that PGE₂ could have an impact on the tendon stem/progenitor cells (TSCs) that reside in tendons [6–8] and could play an important physiological role in the maintenance of tendon homeostasis.

Therefore, PGE₂ may have biphasic effects depending on its concentration. A better understanding of the concentration-dependent effects of PGE₂ on tendon cells, particularly TSCs, may shed new light on tendon physiology and pathology. Thus, in this study we hypothesized that lower concentrations of PGE₂ increase TSC proliferation and decrease non-tenocyte differentiation of TSCs, while higher concentrations produce the opposite effects. To test this hypothesis, we carried out cell culture experiments by treating human TSCs (hTSCs) with low and high levels of PGE₂. We also performed *in vivo* implantation experiments to determine the differentiation fate of hTSCs after treatment with various concentrations of PGE₂ *in vitro*.

Materials and Methods

Ethics Statement

The Gift of Hope Organ and Tissue Donor Network (Elmhurst, IL) provided normal human knee tissues, after obtaining written consent from donors' families and approval from the local ethics committee (Gift of Hope Organ and Tissue Donor Network). The University of Pittsburgh IRB also approved the study protocol for using human tendon tissues in the cell culture and animal studies performed in this study. These specimens were used for investigation only and no human subjects were involved in this project. Data obtained for the study was not through intervention or interaction with individuals and does not have any identifiable private information. Further, the University of Pittsburgh IACUC

approved the protocol for the use of rats in the *in vivo* implantation experiments.

hTSC Culture

hTSCs were isolated from the patellar tendons of six human donors (20 to 44 years old) using our previously published protocol [8]. Briefly, after removing the paratenons, the core portions of the patellar tendons were cut into small pieces and digested with collagenase type I (3 mg/ml) and dispase (4 mg/ml) at 37°C for 1 hr. After centrifugation at 3,000 rpm for 15 min and removal of the enzyme-containing supernatant, a single-cell suspension was obtained, which was cultured in growth medium (DMEM plus 20% FBS) at 37°C with 5% CO₂. After 8 to 10 days in culture dishes, hTSCs formed colonies. The stem cell colonies were then isolated and cultured in DMEM with 20% FBS. These hTSCs at passage 1 were used in the following experiments.

Verification of the Stemness of hTSCs

The stemness of human tendon stem cells (hTSCs) from the patellar tendon used in this study was verified by immunocytochemical analysis of three stem cell markers, including octamer-binding transcription factor 4 (Oct-4), Nanog, and nucleostemin (NS). hTSCs were first seeded into 12-well plates at a density of 20,000 cells/well with 1.5 ml medium and cultured for 3 days. Then, the hTSCs were fixed in 4% paraformaldehyde in PBS for 20 min at room temperature and washed in 0.5% Triton-X-100 in PBS for 15 min. Subsequently, the fixed cells were incubated with either mouse anti-human Oct-4 (1:500), rabbit anti-human Nanog (1:500), or goat anti-human nucleostemin (1:500) overnight at 4°C. After washing three times with PBS, the cells were again incubated for 2 hrs at room temperature with either Cy3-conjugated goat anti-mouse IgG antibodies (1:1000) for Oct-4, Cy3-conjugated goat anti-rabbit IgG (1:500) for Nanog, or Cy3-conjugated donkey anti-goat IgG antibodies (1:500) for Nucleostemin. Nuclei were stained with Hoechst fluorochrome 33342 (1 µg/ml; Sigma, St. Louis, MO). Stained cells were then examined using fluorescence microscopy. All antibodies were obtained from Chemicon International (Temecula, CA), BD Biosciences (Franklin Lakes, NJ), Neuromics (Edina, MN), or Santa Cruz Biotechnology Inc. (Santa Cruz, CA).

Measurement of Proliferation of hTSCs Treated with PGE₂

hTSCs were seeded in 6-well plates (6×10^4 /well) and six different concentrations of PGE₂ (0, 0.01, 0.1, 1, 10, and 100 ng/ml) were added to the culture. Three replicates were maintained for each concentration. The medium was changed every day and PGE₂ was replenished. After 6 days, cell number was measured using a digital cellometer (Nexcelcom Bioscience, Lawrence, MA), and the population doubling time (PDT), which is a measure of cell proliferation, was calculated based on the formula: $\log_2[N_c/N_0]$, where N_c is the total number of cells at confluence, and N_0 is the initial number of cells seeded [8].

Determination of the Effect of PGE₂ Treatment on hTSC Stemness

Stemness of hTSCs was determined by immunocytochemistry and FACS analysis. For immunocytochemistry, hTSCs were seeded in 12-well plates (3×10^4 /well) and treated with six different PGE₂ concentrations ranging from 0 to 100 ng/ml for 5 days, with three replicates for each concentration. The effect of PGE₂ treatment on hTSC stemness was then determined by performing immunocytochemistry for stem cell markers SSEA-4 and Stro-1. Briefly, cells were fixed in 4% paraformaldehyde in PBS for

30 min at room temperature. After washing with PBS, the cells were incubated at room temperature with mouse anti-human SSEA-4 (1:350; Invitrogen, Cat. # 414000) for 3 hrs or mouse anti-human Stro-1 (1:200; Invitrogen, Cat. # 398401) for 4 hrs. The cells were then washed three times with PBS, followed by incubation with Cy3-conjugated goat anti-mouse IgG (1:500; Invitrogen, Cat. # A10521) secondary antibody at room temperature for 2 hrs. After a final wash with PBS, the nuclei were stained with Hoechst fluorochrome 33342, as described above. Stained cells were examined and images of cells were obtained using a fluorescence microscope (Nikon eclipse microscope, TE2000-U).

Semi-quantification of Positively-stained hTSCs

For the semi-quantification of stem cell markers *in vitro*, seven random images were captured from each well at a magnification of 20x under the Nikon eclipse microscope. The positively-stained cells in each picture were manually identified and analyzed using SPOTTM imaging software (Diagnostic Instruments, Inc., Sterling Heights, MI). The positive staining percentage was calculated by dividing the number of positively-stained cells by the total number of cells under the microscopic field. The average value of all seven images from each well represented the percentage of positive staining, which indicates the stemness of hTSCs in the respective PGE₂ concentrations.

Fluorescence Activated Cell Sorting (FACS) Analysis of hTSCs

To determine the effect of PGE₂ treatment on hTSC stemness by FACS analysis, hTSCs (1.5×10^6 in 50 µl PBS) were incubated with 20 µl of the appropriate serum in a centrifuge tube at 4°C for 30 min. Subsequently, 0.4 µg of mouse anti-human SSEA-4 (Cell Signaling, Cat. #4755S) or mouse anti-human Stro-1 (Millipore, Cat. #MAB4315) primary antibody was added and incubated at 4°C overnight. The cells were then washed three times with 2% FBS-PBS, followed by centrifugation at 500 g for 5 min/each time. Then the cells were treated with 1 µg Cy3 conjugated goat anti-mouse IgG secondary antibody at room temperature for 2 hrs. The cells treated with the second antibody only were used as a staining negative control. Finally, the cells were washed twice with PBS and fixed in 1% paraformaldehyde, followed by FACS analysis on a BD LSR II Flow Cytometer (BD Biosciences).

Determination of hTSC Differentiation *in vitro* by qRT-PCR

To determine the effect of PGE₂ treatment on the differentiation of hTSCs, we performed quantitative RT-PCR (qRT-PCR) to measure gene expression using a QIAGEN QuantiTect SYBR Green PCR Kit (QIAGEN). Briefly, total RNA was isolated from hTSCs using the RNeasy Mini Kit with an on-column DNase I digest (Qiagen, Valencia, CA). Then first-strand cDNA was reverse transcribed using SuperScript II (Invitrogen, Grand Island, NY) in a 20 µl reaction containing 1 µg total RNA. Conditions for the cDNA synthesis included 65°C for 5 min, 4°C for 1 min, 42°C for 50 min, and finally 72°C for 15 min. qRT-PCR was performed in a 25 µl PCR reaction mixture with 2 µl cDNA (~100 ng RNA) in a Chromo 4 Detector (MJ Research, Maltham, MA) by incubating at 94°C for 5 min, followed by 30 to 60 cycles of a three temperature program consisting of 1 min at 94°C, 40 sec at 57°C, and 40 sec at 72°C. The PCR reaction was terminated after a 10 min extension at 70°C and stored at 4°C until further analysis. Expression of stem cell markers (Oct-4 and Nanog), tenocyte markers (collagen type I and tenascin C),

adipocyte marker (PPAR γ), chondrocyte marker (Sox9), and osteocyte marker (Runx2) were measured using the primers listed in **Table 1**. GAPDH was used as an internal control. All primers were synthesized by Invitrogen. Expression of each target gene was normalized to GAPDH gene expression and the relative gene expression levels were calculated from the formula $2^{-\Delta\Delta CT}$. Details of the calculation are described in our previous study [9]. The mean and standard deviation (SD) of the CT values were determined from at least three replicates.

Determination of hTSC Differentiation *in vivo* by Implantation

To verify the effect of PGE₂ treatment on the differentiation of hTSCs *in vivo*, eight female nude rats (10 weeks old; 200–250 g) were used. hTSCs at passage 2 were seeded into 24-well plates (8×10^6 cells/well) and cultured in DMEM with or without various concentrations of PGE₂ for 6 days, with a change of medium every day. For implantation experiments, the cells were trypsinized from each well and mixed with 0.25 ml Matrigel (BD Scientific) to enable gel formation after implantation. These hTSC-Matrigel mixtures were placed subcutaneously in the back of anesthetized rats. Three pieces of hTSC-Matrigel were positioned in three distinct places on each side of each rat's back. Four weeks after implantation, tissue samples were harvested from the implanted area and placed in pre-labeled base molds filled with frozen section medium (Neg 50; Richard-Allan Scientific; Kalamazoo, MI). The tissue blocks were stored at -80°C until histological analysis.

Immunohistochemical and Histological Analyses

Each frozen tissue block was cut into 10 μm thick sections, placed on glass slides, and then allowed to dry overnight at room temperature. The tissue sections were fixed in 4% paraformaldehyde for 30 min and further washed three times with PBS. They were then incubated at room temperature with mouse anti-human PPAR γ antibody (Santa Cruz Biotechnology, Inc., Cat. #271392, Santa Cruz, CA) diluted to 1:350 for 2 hrs, mouse anti-collagen type II antibody (1:300; Millipore, Cat. #MAB8887, Temecula,

CA) for 2 hrs, or mouse anti-human osteocalcin antibody (1:300; Santa Cruz Biotechnology, Inc., Cat. #74495, Santa Cruz, CA) for 3 hrs. After washing with PBS, Cy3-conjugated goat anti-mouse IgG (1:500; Santa Cruz Biotechnology) was added as secondary antibody and incubated at room temperature for 1 hr, followed by staining the nuclei with Hoechst fluorochrome 33342 (1 $\mu\text{g}/\text{ml}$; Sigma, St. Louis, MO) at room temperature for 5 min. Additionally, cell morphology and distribution in those tissues that received hTSCs, which had been treated with various concentrations of PGE₂ in culture, were assessed by staining with hematoxylin and eosin (H&E). Finally, all tissue sections were examined under a fluorescence microscope.

Semi-quantification of Positively Stained Tissue Sections

Each tissue section was examined under a microscope (Nikon eclipse, TE2000-U) and five random images were taken for the semi-quantification of hTSC differentiation *in vivo*. SPOTTM imaging software (Diagnostic Instruments, Inc., Sterling Heights, MI) was used to process positively stained areas, which were manually identified by examining the images taken. The total area viewed under the microscope was divided by the positively stained area to calculate the proportion of positive staining. Five tissue sections were used for each group and five images were obtained per tissue section. These values were averaged to represent the percentage positive staining in all the groups treated with various PGE₂ concentrations, which indicated the extent of cell differentiation.

Statistical Analysis

Data are expressed as mean \pm standard deviation (mean \pm SD). Unless otherwise indicated, at least three replicates were used for each experimental condition. For statistical analysis of data, one-way ANOVA or a student *t*-test was used wherever appropriate. All comparisons were between each PGE₂-treated group and the respective control. A P-value less than 0.05 was considered to indicate statistically-significant differences between the groups compared.

Table 1. Primers used in qRT-PCR for gene expression analysis.

Gene	Primer Sequence	Accession numbers	Reference
Oct-4	Forward 5'-CGC AAG CCC TCA TTT CAC-3'	NM_002701	[32]
	Reverse 5'-CAT CAC CTC CAC CAC CTG-3'		
Nanog	Forward 5'-TCC TCC TCT TCC TCT ATA CTA AC-3'	NM_024865	[33]
	Reverse 5'-CCC ACA ATC ACA GGC ATA C-3'		
Tenascin C	Forward 5'- CGG GGC TAT AGA ACA CCA GT-3'	NM_002160.2	[34]
	Reverse 5'- AAC ATT TAA GTT TCC AAT TTC AGG TT-3'		
Collagen I	Forward 5'-AGG GTG AGA CAG GCG AAC AG-3'	NM_000088	[35]
	Reverse 5'-CTC TTG AGG TGG CTG GGG CA-3'		
PPAR γ	Forward 5'- GGC TTC ATG ACA AGG GAG TTT C-3'	NM_138711	[36]
	Reverse 5'- CTT TAT GGA GCC CAA GTT TGA GTT-3'		
Sox9	Forward 5'- CCC CAA CAG ATC GCC TAC AG-3'	NM_000346	[37]
	Reverse 5'- GAG TTC TGG TCG GTG TAG TC-3'		
Runx2	Forward 5'- ATG CTT CAT TCG CCT CAC AAA-3'	NM_001015051	[38]
	Reverse 5'- CCA AAA GAA GTT TTG CTG ACA TGG-3'		
GAPDH	Forward 5'-GCC AAA AGG GTC ATC ATC-3'	NM_002046	[32]
	Reverse 5'-ATG ACC TTG CCC ACA GCC TT-3'		

doi:10.1371/journal.pone.0087706.t001

Results

Verification of the Stemness of hTSCs

Prior to using hTSCs for cell culture experiments in this study, we first verified the stemness of these tendon cells. Microscopic examination of hTSCs revealed the typical cobblestone-shaped morphology of tendon stem cells under phase contrast microscopy (**Fig. 1A**). Further, cells in culture also showed robust expression of all three stem cells markers, Oct-4 (**Fig. 1B**), Nanog (**Fig. 1C**), and nucleostemin (**Fig. 1D**), in immunohistochemical analyses. These characteristics indicated that the cells derived from the human patellar tendons were indeed tendon-specific stem cells.

Effect of PGE₂ on the Proliferation of hTSCs

After establishing that the cells in culture were hTSCs, we investigated cell proliferation after PGE₂ treatment of hTSCs by determining their population doubling time (PDT). Treatment of hTSCs with a lower concentration (0.01 ng/ml) of PGE₂ significantly increased cell proliferation, as evidenced by decreased PDT when compared to the control (**Fig. 2**). PGE₂ treatment at a higher concentration (0.1 ng/ml) also induced similar proliferative effects, although to a smaller extent. At concentrations of 1 and 10 ng/ml, the proliferation of hTSCs was not significantly different from the control without PGE₂ treatment. At the highest concentration (100 ng/ml), TSC proliferation was significantly decreased.

Effect of PGE₂ Treatment on the Stemness of hTSCs

Immunofluorescence assays for stem cell markers revealed that hTSCs treated with a low concentration of PGE₂ (0.01 ng/ml) expressed SSEA-4 (**Fig. 3B**) and Stro-1 (**Fig. 4B**) more extensively




Figure 2. Population doubling time (PDT) of hTSCs treated with various concentrations of PGE₂. hTSCs were seeded in 6-well plates and cultured for six days on medium containing six different concentrations of PGE₂. PDT increased with increasing concentration of PGE₂, meaning that increased PGE₂ resulted in decreased cell proliferation (*p<0.05 when compared to control cells without PGE₂ treatment).

doi:10.1371/journal.pone.0087706.g002

than controls (without PGE₂ treatment) (**Fig. 3A, 4A**) and those treated with higher concentrations of PGE₂ (10 or 100 ng/ml) (**Fig. 3E, 3F, 4E, 4F**). Indeed, the expression levels of both stem cell markers were significantly inhibited by higher concentrations of PGE₂ (10 or 100 ng/ml) (**Fig. 3, 4**). However, semi-quantification of the staining results revealed that the levels of both SSEA-4 (**Fig. 3G**) and Stro-1 (**Fig. 4G**) were similar between the control hTSCs and hTSCs treated with 0.01 ng/ml PGE₂. Consistent with the microscopic observations, higher concentra-




Figure 1. Verification of the stemness of hTSCs. Cobblestone shaped morphology of hTSCs visualized under phase contrast microscopy (**A**). hTSCs also expressed Oct-4 (**B**), Nanog (**C**), and nucleostemin (**D**). Staining for all three stem cell markers was nearly 100% positive with the respective antibodies. Bar = 50 μ m.

doi:10.1371/journal.pone.0087706.g001

tions of PGE₂ significantly reduced staining for both stem cell markers. Particularly, the concentration-dependent effect of PGE₂ on Stro-1 was more profound than its effect on SSEA-4 (**Fig. 3G, 4G**), with 81% reduction at 100 ng/ml, 76% at 10 ng/ml, 52% at 1 ng/ml, and 38% at 0.1 ng/ml for Stro-1, and 61% at 100 ng/ml, 40% at 10 ng/ml, 17% at 1 ng/ml, and 12% at 0.1 ng/ml for SSEA-4.

Additionally, FACS analysis of the stem cell markers also corroborated the immunocytochemical findings. Specifically, as PGE₂ concentrations increased from 0 to 0.01 ng/ml, more cells positively stained with SSEA-4 and Stro-1 (**Fig. 5**, blue dots) were evident; however, when PGE₂ concentrations were further increased to 1 and 100 ng/ml, few positively-stained cells were detected. Quantification of the results from two independent FACS experiments also confirmed these observations (**Fig. 6**).

To further characterize the stemness of hTSCs after treatment with PGE₂, we examined the expression of stem cell genes using qRT-PCR. We found that the gene expression levels of Nanog and Oct-4 were significantly ($p < 0.05$) up-regulated in hTSCs treated with lower concentrations (0.01 and 0.1 ng/ml) of PGE₂ (**Fig. 7**). Notably, the expression level of Oct-4 was twice as high as that of Nanog at 0.01 ng/ml PGE₂ concentration. When treated with higher concentrations (1, 10, and 100 ng/ml) of PGE₂, expression levels of both Nanog and Oct-4 were down-regulated and almost reached the levels of controls without PGE₂ treatment.

Effect of PGE₂ on the Differentiation of hTSCs

We next examined the effects of PGE₂ on hTSC differentiation by determining the expression of tenocyte and non-tenocyte related genes. Treatment of hTSCs with lower concentrations

SSEA-4





Figure 3. Expression of the stem cell marker SSEA-4 by hTSCs cultured *in vitro* in various concentrations of PGE₂. **A:** without PGE₂ treatment; **B:** 0.01 ng/ml PGE₂; **C:** 0.1 ng/ml PGE₂; **D:** 1 ng/ml PGE₂; **E:** 10 ng/ml PGE₂; and **F:** 100 ng/ml PGE₂. hTSCs were seeded in 12-well plates, cultured with six different concentrations of PGE₂, incubated with mouse anti-human SSEA-4 primary antibody, and detected with Cy3-conjugated goat anti-mouse IgG. Nuclei were stained with Hoechst (Blue). Expression of SSEA-4 (red) is dose-dependent, with more robust expression seen in hTSCs treated with low levels of PGE₂ (**A–D**) than expression levels seen in those treated with high levels (**E, F**). Positively stained cells were also counted to calculate percentage staining (**G**) (* $p < 0.05$ with respect to hTSCs not treated with PGE₂). Bar: 100 μ m. doi:10.1371/journal.pone.0087706.g003

Stro-1





Figure 4. Expression of the stem cell marker Stro-1 by hTSCs cultured *in vitro* in medium containing various concentrations of PGE₂. A: without PGE₂ treatment; B: 0.01 ng/ml PGE₂; C: 0.1 ng/ml PGE₂; D: 1 ng/ml PGE₂; E: 10 ng/ml PGE₂; and F: 100 ng/ml PGE₂. hTSCs were seeded in 12-well plates, cultured with six different concentrations of PGE₂, incubated with mouse anti-human Stro-1, and detected with Cy3-conjugated goat anti-mouse IgG. Hoechst was used to stain nuclei (blue). Expression of Stro-1 (red) is higher in hTSCs treated with low PGE₂ concentrations (A, B) than hTSCs treated with high concentrations (E–F). Similar to SSEA-4, expression of Stro-1 is also dose-dependent. Positively stained cells were also counted to calculate percentage staining (G) (*p<0.05 in comparison with control hTSCs not treated with PGE₂). Bar: 100 μm. doi:10.1371/journal.pone.0087706.g004

(0.01, 0.1, and 1 ng/ml) of PGE₂ significantly ($p<0.05$) enhanced the expression of both collagen type I and tenascin C, two tenocyte-associated genes (Fig. 8A). However, at these lower concentrations, the expression levels of non-tenocyte associated genes PPAR γ , Sox9, and Runx2 were lower or only marginally higher than the control (Fig. 8B). On the other hand, treatment of hTSCs with higher concentrations (10 and 100 ng/ml) of PGE₂ significantly ($p<0.05$) up-regulated PPAR γ , Sox9, and Runx2 genes associated with adipogenic, chondrogenic, and osteogenic differentiation, respectively (Fig. 8B). This up-regulation corresponded with the down-regulation of collagen type I and tenascin C at 10 and 100 ng/ml PGE₂ concentrations (Fig. 8A).

Non-tendinous Tissue Formation after Implantation of PGE₂-treated hTSCs

To determine whether PGE₂-treated hTSCs underwent non-tenogenic differentiation, we subcutaneously implanted PGE₂-treated hTSCs into nude rats. We found that 4 weeks after implantation, non-tenocyte differentiation of hTSCs was more extensive in the cells treated with higher concentrations (10 and 100 ng/ml) of PGE₂ (Fig. 9E–G, and Fig. 9I–K) when compared to the hTSCs that received the lowest concentration of PGE₂ (0.1 ng/ml) (Fig. 9A–C), as evidenced by higher amounts of PPAR γ , collagen type II and osteocalcin (stained in red/pink). It appeared that more cells (black dots) were present in tissues that received hTSCs treated with high PGE₂ concentrations (Fig. 9H–L) than those that were treated with low PGE₂ concentrations (Fig. 9D). Specifically, at 100 ng/ml (Fig. 9L), numerous cells were concentrated in a specific region (triangle). The immunohis-




Figure 5. FACS analysis of SSEA-4 and Stro-1 expression in hTSCs treated with various concentrations of PGE₂. hTSCs in culture were treated with various concentrations of PGE₂. FACS analysis was performed on these cells (for details, see Methods section). It is evident that when cells were treated with 0.01 ng/ml of PGE₂ (PGE₂-0.01), more cells positively stained with SSEA-4 and Stro-1 were detected (blue dots in the P2 area) compared to control cells without PGE₂ treatment (PGE₂-0). When PGE₂ concentration increased to 1 ng/ml (PGE₂-1) and 100 ng/ml (PGE₂-100), fewer cells were actually stained with SSEA-4 and Stro-1 when compared to control cells (PGE₂-0). doi:10.1371/journal.pone.0087706.g005

tochemical observations were also confirmed by semi-quantification, which showed a significant ($P < 0.001$) dose-dependent increase in the staining of non-tenocyte associated genes with increasing amounts of PGE₂ (Fig. 9M). When compared to hTSCs treated with 0.1 ng/ml PGE₂, those treated with 10 ng/ml had a ~2 fold increase in PPAR γ , collagen type II, and osteocalcin. These increases were higher when hTSCs were treated with 100 ng/ml PGE₂ (~3 fold for PPAR γ , ~4 fold for collagen type II, and ~4 fold for osteocalcin).

Discussion

PGE₂ is one of the most abundant prostaglandins in the body, and an important causative factor of inflammation that results from tissue damage or infection. Since our previous study showed that high levels of PGE₂ (1, 10, and 100 ng/ml) decrease proliferation and induce differentiation of mouse TSCs into non-tenocytes [6], in the present study we investigated the effects of comparable and lower doses of PGE₂ (0.01 to 100 ng/ml) on hTSC proliferation and differentiation by performing cell culture and cell implantation experiments. Our results revealed a concentration-dependent biphasic effect of PGE₂ on the proliferation and differentiation of hTSCs. PGE₂ treatment of hTSCs increased cell proliferation at lower concentrations, but decreased it at higher concentrations. In particular, low levels of PGE₂ promoted the stemness of hTSCs, as evidenced by the extensive expression of stem cell markers SSEA-4 and Stro-1 in hTSCs treated with low concentrations of PGE₂. The range of PGE₂ concentrations used in this study also includes the *in vivo* physiological concentrations of PGE₂ reported in human Achilles tendons (0.8 ± 0.2 ng/ml, [10] or 54 ± 24 pg/ml [11]). It should be




Figure 6. Quantification of SSEA-4 and Stro-1 expression in hTSCs treated with various concentrations of PGE₂. Average percentage of cells expressing SSEA-4 and Stro-1 in two independent FACS experiments showed that hTSCs treated with 0.01 ng/ml of PGE₂ expressed the most SSEA-4 ($84.6 \pm 5.4\%$) and Stro-1 ($91.1 \pm 4.2\%$), which was higher than the control cells without PGE₂ treatment ($54.5 \pm 8.6\%$ for SSEA-4, and $35.8 \pm 6.5\%$ for Stro-1) and the cells treated with 1 ng/ml (SSEA-4: $23.0 \pm 2.0\%$; and Stro-1: $17.7 \pm 1.7\%$) or 100 ng/ml PGE₂ (SSEA-4: $10.6 \pm 2.1\%$, and Stro-1: $15.3 \pm 1.1\%$). SC: Stem cell; PGE₂-0: PGE₂ at 0 concentration; PGE₂-0.01: PGE₂ concentration at 0.01 ng/ml; PGE₂-1: PGE₂ concentration at 1 ng/ml; and PGE₂-100: PGE₂ concentration at 100 ng/ml. doi:10.1371/journal.pone.0087706.g006

noted that these values are likely lower due to two reasons: i) patients in these studies were at rest during these measurements and did not undergo intensive exercise, and ii) these values are average microdialysis measurements of PGE₂ concentrations over a large portion of the tendon instead of at a local site, where PGE₂ concentrations could be much higher.

The biphasic effects of PGE₂ on various tissue properties have been reported in previous studies. For example, PGE₂ has been shown to exert biphasic effects on vascularity [12]; it elicits vasodilation at low concentrations and reverses this effect at higher concentrations. Similarly, PGE₂ treatment reduced proliferation of mesenchymal stem cells (MSCs) in a dose-dependent manner (0.25 μ M PGE₂, or 88 ng/ml to 8.8 μ g/ml), with the two lowest concentrations (0.25 nM and 2.5 nM PGE₂, or 88 pg/ml to 880 pg/ml) slightly increasing MSC proliferation over baseline levels [13]. In this study, the authors demonstrated that the biphasic effect of PGE₂ was executed by differential activation of two types of protein kinase A (PKA). At low concentrations, PGE₂ activated PKA II, leading to a cascade of events that resulted in cell proliferation; at high concentrations, PGE₂ caused PKA I activation, resulting in cell cycle arrest which reduced MSC proliferation. In addition, PGE₂ was reported to have a biphasic influence on injured esophagus: at low doses PGE₂ was protective, but at high doses it damaged the esophagus, with this effect being mediated by the EP1 receptor [14]. Interestingly, the biphasic effects of PGE₂ were also reported to depend on the growth state of the tissue type. For example, PGE₂ promoted proliferation of quiescent smooth muscle cells indicated by an increase in both DNA and RNA synthesis with increasing levels of PGE₂ (10^{-10} – 10^{-5} M, or 3.5 ng/ml - 3.5 μ g/ml). However, when proliferating smooth muscle cells were treated with the same concentrations of PGE₂, DNA synthesis decreased by 48%, indicating that PGE₂ had an inhibitory effect [15].

In this study, we established the stemness of hTSCs based on three characteristics described previously for human tendon stem/progenitor cells: a) the ability to form colonies in culture; b)




Figure 7. Expression of stem cell markers Nanog and Oct-4 in hTSCs treated with various concentrations of PGE₂. Total RNA was collected from various hTSCs cultured with or without PGE₂ and used for qRT-PCR. Expression levels of Nanog and Oct-4 are more up-regulated in hTSCs treated with low concentrations of PGE₂ (0.01 and 0.1 ng/ml) than in those treated with high concentrations (1, 10, and 100 ng/ml) (* $p < 0.05$ with respect to hTSCs not treated with PGE₂). doi:10.1371/journal.pone.0087706.g007




Figure 8. Expression of tenocyte (A) and non-tenocyte (B) related genes in hTSCs treated with various concentrations of PGE₂. qRT-PCR was performed on total RNA collected from cultured hTSCs treated with PGE₂. Expression levels of tenocyte related genes, collagen type I (Collagen I) and Tenascin C, were higher in hTSCs treated with low concentrations of PGE₂ (0.01, 0.1 and 1 ng/ml) than in those treated with high concentrations (10 and 100 ng/ml) (A). However, expression levels of non-tenocyte related genes, PPAR_γ, Sox9, and Runx2, were more reduced in hTSCs treated with low (0.01, 0.1 and 1 ng/ml) than with high concentrations of PGE₂ (10 and 100 ng/ml) (B) (*p<0.05 with respect to corresponding controls that did not receive PGE₂ treatment). doi:10.1371/journal.pone.0087706.g008

expression of stem cell markers Oct-4, Nanog, and nucleostemin; and c) multi-differentiation potential [7,8]. In addition, these hTSCs assumed a cobblestone shape when grown to confluence [8]. Further, we used two stem cell markers, SSEA-4 and Stro-1, to measure the stemness of hTSCs treated with various concentrations of PGE₂. SSEA-4 and Stro-1 are highly expressed in undifferentiated stem cells and therefore are used as markers for stem cell identification. However, after differentiation, SSEA-4 is down-regulated in human embryonic stem cells [16]. Our results showing higher expression of SSEA-4 in cells treated with low levels of PGE₂ indicate that stemness is enhanced in these cells, but not in cells treated with higher levels of PGE₂. Additionally, we also found that cells treated with low levels of PGE₂ produced higher levels of stem cell-related genes (Oct-4 and Nanog) than cells treated with high levels of PGE₂. Oct-4 and Nanog are both required for the self-renewal and maintenance of stem cells in an un-differentiated state [17]. These genes were reported to downregulate the expression and activity of lineage specific factors, thereby promoting pluripotency [18]. Their downregulation, however, increased differentiation and thereby decreased the capacity of mouse embryonic stem cells for self-renewal [19–21].

This study found that higher expression levels of both Nanog and Oct-4 and corresponded low levels of non-tenocyte related genes, particularly in cells treated with low levels of PGE₂ (0.01, and 0.1 ng/ml). The results indicate maintenance of hTSCs in an undifferentiated state, at least in part through Nanog and Oct-4 suppression of adipocyte- (PPAR_γ), chondrocyte- (Sox9), and osteocyte- (Runx2) related genes. Further, lower expression levels of Nanog and Oct-4, especially in cells treated with high concentrations of PGE₂ (10 and 100 ng/ml), also corresponded to higher expression levels of non-tenocyte related genes. This effect, however, was not observed in the control cells (those without PGE₂ treatment), indicating the role high PGE₂ levels have in promoting non-tenocyte differentiation of hTSCs, which in turn reduces their stemness. Taken together, these results strongly suggest that the beneficial effects of the constitutively maintained low levels of PGE₂ may be critical for the maintenance of homeostasis in tendons *in vivo*.

hTSCs treated with higher concentrations of PGE₂ exhibited extensive expression of non-tenocyte related genes. In the *in vivo* experiment, non-tenocyte proteins PPAR_γ, collagen type II, and osteocalcin were up-regulated after implantation of hTSCs treated




Figure 9. *In vivo* expression of non-tenocyte markers PPAR_γ, collagen type II, and osteocalcin in rats implanted with hTSCs treated with various concentrations of PGE₂ and their respective hematoxylin and eosin (H&E) stained tissue sections. hTSCs cultured with three concentrations (0.1, 10, and 100 ng/ml) of PGE₂ were implanted subcutaneously into rats; later, immunohistochemical and histological analyses were performed on tissue sections. For the immunohistochemical staining, fixed tissue sections were incubated with mouse anti-human PPAR_γ antibody, mouse anti-collagen type II (Collagen II) antibody, or mouse anti-human osteocalcin antibody. Cy3-conjugated goat anti-mouse IgG was then used to detect primary binding. Nuclei were stained with Hoechst (blue). Expression levels of PPAR_γ, collagen type II, and osteocalcin (red) were lower in cells treated with 0.1 ng/ml PGE₂ (A–C) than those treated with the higher concentrations (10 and 100 ng/ml) of PGE₂ (E–G, I–K). H&E staining was also performed on tissue sections (D, H, L). More cells (black dots; see insets in D, H, and L) were observed in tissues implanted with hTSCs that had been treated with high concentrations of PGE₂ in culture (H, L). Specifically, at 100 ng/ml (L), cells were concentrated in a specific region (triangle). Additionally, semi-quantification of the stained cells was performed by counting immuno-positive cells and calculating percentage staining (M) (*p<0.05 in comparison with control hTSCs not treated with PGE₂). Bar: 100 μm. doi:10.1371/journal.pone.0087706.g009

with high levels of PGE₂. These findings suggest that PGE₂ at high concentrations could cause differentiation of TSCs to non-tenocytes; this could lead to impaired tendon healing and the formation of non-tendinous tissues in affected tendons, which would consequently reduce tendon strength. Indeed, it has been suggested that PGE₂, as a local hormone in tendons, may contribute to the development of tendinopathy [2,22–24]. In addition, prostaglandins (PGs) are known to play a pathophysiological role in the skeletal system, including contributing to the pathology of osteoporosis by enhancing bone resorption [25]. However, in the same milieu, PGs also exert a physiological role by stimulating bone formation through increased osteoblast

proliferation and differentiation. These functions of PGs are consistent with the biphasic effects of PGE₂ that maintain tendon homeostasis and lead to tendon pathology or tendinopathy.

It should be noted that when hTSCs were treated with low levels of PGE₂, tenocyte-related genes, including collagen type I and tenascin C, were highly expressed (Fig. 7A). These results suggest that PGE₂ at low concentrations may exert its effects on TSCs in two ways: promoting the stemness of TSCs, and inducing TSCs to differentiate towards tenocytes (or progenitor cells for tenocytes). TSCs in our cultures presumably consisted of two subpopulations of cells: one population consisted of early-stage stem cells expressing stem cell markers, such as Nanog, Oct-4, SSEA-4,

and Stro-1, and the other population consisted of progenitor cells, which have differentiated towards tenocytes and expressed collagen type I and tenascin C, as demonstrated in this study. In other words, low levels of PGE₂ not only promote TSC self-renewal, but also promote the differentiation of TSCs into progenitor cells for tenocytes, suggesting that low concentrations of PGE₂ cause hTSCs to undergo asymmetric differentiation. Endogenous PGE₂ has also been shown to stimulate the proliferation of human MSCs [26], protect mouse embryonic stem cells from apoptosis through EP receptor activation [27], and enhance homing, survival, and proliferation of mouse and human hematopoietic stem cells that lead to increased numbers of repopulating cells and units [28]. As tendon-specific stem cells, TSCs play an important role in the repair of injured tendons by proliferating and differentiating *in vivo*. When tendons are injured, more tenocytes are needed, and TSCs must be activated to effectively repair injured tendons. Our results indicate that the constitutive baseline levels of PGE₂, which are low, may be used to effectively expand TSCs for cell therapy of injured tendons by promoting proliferation and maintaining tendon homeostasis.

The beneficial effects of low PGE₂ levels on TSCs have several potential applications in tendon tissue engineering. Since PGE₂ at low levels can promote the stemness of TSCs, it may be used to maintain TSCs in culture. In addition, because low PGE₂ levels can accelerate TSC proliferation, they could be used to quickly expand TSC populations for the use of cell therapy to treat injured tendons. Moreover, *in vivo* tendon injuries could be potentially treated by injecting low levels of PGE₂ at the site of injury. This could enhance the healing of injured tendons because of the ability of low levels of PGE₂ to stimulate self-renewal of TSCs and promote tenogenesis. A recent study showed that low levels of PGE₂ injected into rat patellar tendons enhanced their structural properties (the ultimate load, stiffness, and elastic modulus) [29].

While this is the first study to demonstrate the biphasic effects of PGE₂ on hTSCs, the molecular mechanisms responsible for these biphasic effects are yet to be investigated. PGE₂ is known to exert its diverse biological effects through the EP receptors [14,15,30] and by differential activation of PKA types [13]. Hence, the biphasic response of hTSCs to PGE₂ observed in this study may

also involve multiple EP receptor subtypes and/or differential activation of PKA types. Also, while we have shown the beneficial effects of low PGE₂ levels on hTSCs, one limitation of the study is the use of static culture without mechanical loading applied to hTSCs. However, tendons, and therefore the TSCs *in vivo*, are constantly subjected to mechanical loading, which regulates the expression levels of collagen type I, PPAR γ , collagen type II, Sox9, and Runx2 genes. In addition, mechanical loading also increases PGE₂ levels in both patellar and Achilles tendons [9], indicating a potential interaction between mechanical loading and PGE₂. Additional studies are required to reveal the mechanisms behind this interaction. Further, we investigated only the long term effects (up to 6 days) of PGE₂ treatment on hTSCs. It is known that exercise increases PGE₂ levels in human blood only transiently, with maximum levels observed 2 hrs after exercise [31]. Therefore, it would be of interest to study the short term effects of PGE₂ on hTSCs both *in vitro* and *in vivo*.

In summary, we showed in this study that PGE₂ exerted biphasic effects on hTSCs: at low concentrations, PGE₂ enhanced their proliferation and expression of stem cell markers, whereas high concentrations of PGE₂ were detrimental to hTSCs, because they reduced their proliferation and induced non-tenocyte differentiation. These results suggest that, on one hand, low levels of PGE₂ promote tendon homeostasis by maintaining hTSCs and tenogenesis; on the other hand, high levels of PGE₂ in tendons may induce differentiation of hTSCs into non-tenocytes and thus lead to the development of the degenerative tendinopathy often seen in clinical settings.

Acknowledgments

We thank Dr. Nirmala Xavier for assistance in the preparation of this manuscript.

Author Contributions

Conceived and designed the experiments: JHW. Performed the experiments: JZ. Analyzed the data: JHW. Contributed reagents/materials/analysis tools: JHW. Wrote the paper: JZ. JHW. Supervised the experiments, analyzed and interpreted results: JHW.

References

- Meknas K, Johansen O, Steigen SE, Olsen R, Jorgensen L, et al. (2012) Could tendinosis be involved in osteoarthritis? *Scand J Med Sci Sports* 22: 627–634.
- Wang JH, Iosifidis MI, Fu FH (2006) Biomechanical basis for tendinopathy. *Clin Orthop Relat Res* 443: 320–332.
- Wang JH, Jia F, Yang G, Yang S, Campbell BH, et al. (2003) Cyclic mechanical stretching of human tendon fibroblasts increases the production of prostaglandin E2 and levels of cyclooxygenase expression: a novel *in vitro* model study. *Connect Tissue Res* 44: 128–133.
- Almekinders LC, Banes AJ, Ballenger CA (1993) Effects of repetitive motion on human fibroblasts. *Med Sci Sports Exerc* 25: 603–607.
- Langberg H, Skovgaard D, Karamouzis M, Bulow J, Kjaer M (1999) Metabolism and inflammatory mediators in the peritendinous space measured by microdialysis during intermittent isometric exercise in humans. *J Physiol* 515 (Pt 3): 919–927.
- Zhang J, Wang JH (2010) Production of PGE₂ increases in tendons subjected to repetitive mechanical loading and induces differentiation of tendon stem cells into non-tenocytes. *J Orthop Res* 28: 198–203.
- Bi Y, Ehrlich D, Kilts TM, Inkson CA, Embree MC, et al. (2007) Identification of tendon stem/progenitor cells and the role of the extracellular matrix in their niche. *Nat Med* 13: 1219–1227.
- Zhang J, Wang JH (2010) Characterization of differential properties of rabbit tendon stem cells and tenocytes. *BMC Musculoskelet Disord* 11: 10.
- Zhang J, Wang JH (2010) Mechanobiological response of tendon stem cells: implications of tendon homeostasis and pathogenesis of tendinopathy. *J Orthop Res* 28: 639–643.
- Langberg H, Boushel R, Skovgaard D, Risum N, Kjaer M (2003) Cyclooxygenase-2 mediated prostaglandin release regulates blood flow in connective tissue during mechanical loading in humans. *J Physiol* 551: 683–689.
- Alfredson H, Thorsen K, Lorentzon R (1999) In situ microdialysis in tendon tissue: high levels of glutamate, but not prostaglandin E2 in chronic Achilles tendon pain. *Knee Surg Sports Traumatol Arthrosc* 7: 378–381.
- Tang L, Loutzenhiser K, Loutzenhiser R (2000) Biphasic actions of prostaglandin E₂ on the renal afferent arteriole: role of EP(3) and EP(4) receptors. *Circ Res* 86: 663–670.
- Kleiveland CR, Kassem M, Lea T (2008) Human mesenchymal stem cell proliferation is regulated by PGE₂ through differential activation of cAMP-dependent protein kinase isoforms. *Exp Cell Res* 314: 1831–1838.
- Yamato M, Nagahama K, Kotani T, Kato S, Takeuchi K (2005) Biphasic effect of prostaglandin E2 in a rat model of esophagitis mediated by EP1 receptors: relation to pepsin secretion. *Digestion* 72: 109–118.
- Yau L, Zahrada P (2003) PGE₂ stimulates vascular smooth muscle cell proliferation via the EP2 receptor. *Molecular and Cellular Endocrinology* 203: 77–90.
- Draper J, Pigott C, Thomson J, Andrews P (2002) Surface antigens of human embryonic stem cells: changes upon differentiation in culture. *J Anat* 200: 249–258.
- Ying QL, Nichols J, Chambers I, Smith A (2003) BMP induction of Id proteins suppresses differentiation and sustains embryonic stem cell self-renewal in collaboration with STAT3. *Cell* 115: 281–292.
- Loh Y, Wu Q, Chew J, Vega V, Zhang W, et al. (2006) The Oct4 and Nanog transcription network regulates pluripotency in mouse embryonic stem cells. *Nat Genet* 38: 431–440.
- Chambers I, Silva J, Colby D, Nichols J, Nijmeijer B, et al. (2007) Nanog safeguards pluripotency and mediates germline development. *Nature* 450: 1230–1234.

20. Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, et al. (2003) The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell* 113.
21. Niwa H, Toyooka Y, Shimosato D, Strumpf D, Takahashi K, et al. (2005) Interaction between Oct3/4 and Cdx2 determines trophoctoderm differentiation. *Cell* 123: 917–929.
22. Zhang J, Wang JH (2012) BMP-2 mediates PGE(2) -induced reduction of proliferation and osteogenic differentiation of human tendon stem cells. *J Orthop Res* 30: 47–52.
23. Khan MH, Li Z, Wang JH (2005) Repeated exposure of tendon to prostaglandin-e2 leads to localized tendon degeneration. *Clin J Sport Med* 15: 27–33.
24. Riley G (2004) The pathogenesis of tendinopathy. A molecular perspective. *Rheumatology (Oxford)* 43: 131–142.
25. Raisz L (1999) Prostaglandins and bone: physiology and pathophysiology. *Osteoarthritis Cartilage* 7: 419–421.
26. Arikawa T, Omura K, Morita I (2004) Regulation of bone morphogenetic protein-2 expression by endogenous prostaglandin E2 in human mesenchymal stem cells. *J Cell Physiol* 200: 400–406.
27. Liou J, Ellent D, Lee S, Goldsby J, Ko B, et al. (2007) Cyclooxygenase-2-derived prostaglandin e2 protects mouse embryonic stem cells from apoptosis. *Stem Cells* 25: 1096–1103.
28. Hoggatt J, Singh P, Sampath J, Pelus L (2009) Prostaglandin E2 enhances hematopoietic stem cell homing, survival, and proliferation. *Blood* 113: 5444–5455.
29. Ferry S, Afshari H, Lee J, Dahners L, Weinhold P (2012) Effect of prostaglandin E2 injection on the structural properties of the rat patellar tendon. *Sports Med Arthrosc Rehabil Ther Technol* 4: 2.
30. Breyer RM, Bagdassarian CK, Myers SA, Breyer MD (2001) Prostanoid receptors: subtypes and signaling. *Annu Rev Pharmacol Toxicol* 41: 661–690.
31. Markworth JF, Vella LD, Lingard BS, Tull DL, Rupasinghe TW, et al. (2013) Human inflammatory and resolving lipid mediator responses to resistance exercise and ibuprofen treatment. *Am J Physiol Regul Integr Comp Physiol*.
32. Chen YC, Hsu HS, Chen YW, Tsai TH, How CK, et al. (2008) Oct-4 expression maintained cancer stem-like properties in lung cancer-derived CD133-positive cells. *PLoS One* 3: e2637.
33. Ling GQ, Chen DB, Wang BQ, Zhang LS (2012) Expression of the pluripotency markers Oct3/4, Nanog and Sox2 in human breast cancer cell lines. *Oncol Lett* 4: 1264–1268.
34. Minear MA, Crosslin DR, Sutton BS, Connelly JJ, Nelson SC, et al. (2011) Polymorphic variants in tenascin-C (TNC) are associated with atherosclerosis and coronary artery disease. *Hum Genet* 129: 641–654.
35. Kohjima M, Enjoji M, Higuchi N, Kotoh K, Kato M, et al. (2007) NIM811, a nonimmunosuppressive cyclosporine analogue, suppresses collagen production and enhances collagenase activity in hepatic stellate cells. *Liver International* 27: 1273–1281.
36. Zhang Y, Ba Y, Liu C, Sun G, Ding L, et al. (2007) PGC-1alpha induces apoptosis in human epithelial ovarian cancer cells through a PPARgamma-dependent pathway. *Cell Res* 17: 363–373.
37. Tew SR, Hardingham TE (2006) Regulation of SOX9 mRNA in human articular chondrocytes involving p38 MAPK activation and mRNA stabilization. *J Biol Chem* 281: 39471–39479.
38. Shui C, Spelsberg TC, Riggs BL, Khosla S (2003) Changes in Runx2/Cbfa1 expression and activity during osteoblastic differentiation of human bone marrow stromal cells. *J Bone Miner Res* 18: 213–221.