

POSTER PRESENTATION

Open Access

Synergistic effects of agonistic costimulatory antibodies adsorbed to amphiphilic poly (γ -glutamic acid) nanoparticles

Catja Freiburghaus^{1,2*}, Sissela Broos¹, Linda C Sandin³, Thomas H Tötterman³, Takami Akagi⁴, Mitsuru Akashi⁴, Carl Borrebaeck¹, Peter Ellmark^{1,2}, Malin Lindstedt¹

From Society for Immunotherapy of Cancer 28th Annual Meeting
National Harbor, MD, USA. 8-10 November 2013

Agonistic antibodies targeting costimulatory pathways for the immune system hold great potential for cancer immunotherapy. However, systemic administration of immunoactivating antibodies can be associated with side effects such as cytokine release syndrome and liver toxicity. Polymeric nanoparticles (NPs) represent an exciting approach to control the release of therapeutic antibodies and to optimize the desired immune response via selective targeting. With increased potency as objective, agonistic antibodies targeting TNF receptors were adsorbed to immune stimulating biodegradable and self-assembled polymeric nanoparticles composed of γ -glutamic acid (γ -PGA NPs). The effects of antibodies targeting CD40, CD137 and OX40 were evaluated based on T and B cell proliferation as well as on cytokine release and phenotypic maturation of antigen-presenting cells. Our results demonstrated a strong synergistic effect on human B cell proliferation of CD40 monoclonal antibodies (mAbs) carrying NPs in vitro. In addition adsorption of anti-CD40 mAb to γ -PGA NPs significantly reduced the systemic release of TNF- α , IL-6, IL-10 and IL-12, compared to treatment with the soluble mAb. Preliminary results indicate increased T cell proliferation and activation by CD137 and OX40 agonistic antibodies in combination with γ -PGA NPs. Combining NPs with agonistic antibodies for cancer immunotherapy offers intriguing opportunities for increased therapeutic efficacy and safety.

Authors' details

¹Department of Immunotechnology, Lund University, Lund, Sweden. ²Alligator Bioscience AB, Lund, Sweden. ³Department of Immunology, Genetics and Pathology, Uppsala University, Uppsala, Sweden. ⁴Department of Applied Chemistry, Graduate School of Engineering, Osaka University, Osaka, Japan.

Published: 7 November 2013

doi:10.1186/2051-1426-1-S1-P128

Cite this article as: Freiburghaus *et al.*: Synergistic effects of agonistic costimulatory antibodies adsorbed to amphiphilic poly (γ -glutamic acid) nanoparticles. *Journal for ImmunoTherapy of Cancer* 2013 **1**(Suppl 1):P128.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Department of Immunotechnology, Lund University, Lund, Sweden
Full list of author information is available at the end of the article