

## Conclusions

Our findings support that transcatheter embolotherapy should be considered, when possible, even for PAVMs with feeding artery sizes <3 mm (but not to the exclusion of other factors, like procedural risks and recurrent radiation exposure). Future research and HHT guideline iterations may need to direct their attention beyond PAVM feeding artery size as a risk factor for neurovascular complications and as a criterion for transcatheter embolotherapy application. ■

**Author disclosures** are available with the text of this letter at [www.atsjournals.org](http://www.atsjournals.org).

**Acknowledgment:** The Brain Vascular Malformation Consortium (U54NS065705) is a part of the National Center for Advancing Translational Sciences (NCATS) Rare Diseases Clinical Research Network (RDCRN) and is supported by the RDCRN Data Management and Coordinating Center (DMCC) (U2CTR002818). RDCRN is an initiative of the Office of Rare Diseases Research (ORDR), NCATS, funded through a collaboration between NCATS and National Institute of Neurological Disorders and Stroke (NINDS).

Theodora Ananiadis, B.Sc. candidate  
University of Western Ontario  
London, Ontario, Canada

Marie E. Faughnan, M.D., M.Sc.  
Dewi Clark, B.Sc.  
Vikram Prabhudesai, M.B. B.S., M.S.  
University of Toronto  
Toronto, Ontario, Canada

Helen Kim, M.P.H., Ph.D.  
University of California San Francisco  
San Francisco, California

Michael T. Lawton, M.D.  
Barrow Neurological Institute  
Phoenix, Arizona

Nicholas T. Vozoris, M.H.Sc., M.D.\*  
University of Toronto  
Toronto, Ontario, Canada

and  
ICES (formerly known as Institute of Clinical Evaluative Sciences)  
Toronto, Ontario, Canada

Brain Vascular Malformation HHT Investigator Group

ORCID ID: 0000-0003-1670-1592 (N.T.V.).

\*Corresponding author: (e-mail: [nick.vozoris@utoronto.ca](mailto:nick.vozoris@utoronto.ca)).

**Brain Vascular Malformation HHT Investigator Group:** Murali Chakinala, Marianne S. Clancy, Marie E. Faughnan, James R. Gossage, Steven W. Hetts, Vivek Iyer, Raj S. Kasthuri, Helen Kim, Timo Krings, Michael T. Lawton, Doris Lin, Hans-Jurgen Mager, Douglas A. Marchuk, Justin P. McWilliams, Jamie McDonald, Ludmila Pawlikowska, Jeffrey Pollak, Felix Ratjen, Karen Swanson, Dilini Vethanayagam, Shantel Weinsheimer, Andrew J. White, and Pearce Wilcox.

## References

- Gossage JR, Kanj G. Pulmonary arteriovenous malformations. A state-of-the-art review. *Am J Respir Crit Care Med* 1998;158:643–661.
- Circo S, Gossage JR. Pulmonary vascular complications of hereditary hemorrhagic telangiectasia. *Curr Opin Pulm Med* 2014;20:421–428.
- Faughnan ME, Granton JT, Young LH. The pulmonary vascular complications of hereditary hemorrhagic telangiectasia. *Eur Respir J* 2009;33:1186–1194.
- Faughnan ME, Palda VA, Garcia-Tsao G, Geithoff UW, McDonald J, Proctor DD, et al.; HHT Foundation International - Guidelines Working Group. International guidelines for the diagnosis and management of hereditary hemorrhagic telangiectasia. *J Med Genet* 2011;48:73–87.
- Shovlin CL, Jackson JE, Bamford KB, Jenkins IH, Benjamin AR, Ramadan H, et al. Primary determinants of ischaemic stroke/brain abscess risks are independent of severity of pulmonary arteriovenous malformations in hereditary hemorrhagic telangiectasia. *Thorax* 2008; 63:259–266.
- White RI Jr, Lynch-Nyhan A, Terry P, Buescher PC, Farnlett EJ, Charnas L, et al. Pulmonary arteriovenous malformations: techniques and long-term outcome of embolotherapy. *Radiology* 1988;169:663–669.
- Cottin V, Dupuis-Girod S, Lesca G, Cordier JF. Pulmonary vascular manifestations of hereditary hemorrhagic telangiectasia (rendu-osler disease). *Respiration* 2007;74:361–378.
- Akers AL, Ball KL, Clancy M, Comi AM, Faughnan ME, Gopal-Srivastava R, et al. Brain vascular malformation consortium: overview, progress, and future directions. *J Rare Disord* 2013;1:5.
- Shovlin CL, Guttmacher AE, Buscarini E, Faughnan ME, Hyland RH, Westermann CJ, et al. Diagnostic criteria for hereditary hemorrhagic telangiectasia (Rendu-Osler-Weber syndrome). *Am J Med Genet* 2000;91:66–67.
- Pierucci P, Lenato GM, Suppressa P, Lastella P, Triggiani V, Valerio R, et al. A long diagnostic delay in patients with hereditary hemorrhagic telangiectasia: a questionnaire-based retrospective study. *Orphanet J Rare Dis* 2012;7:33.

Copyright © 2022 by the American Thoracic Society



## Mortality among Patients with COVID-19 and Different Interstitial Lung Disease Subtypes: A Multicenter Cohort Study

To the Editor:

Although highly effective vaccines are now available for the prevention of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), this infection continues to cause substantial

morbidity and mortality in both vaccinated and unvaccinated individuals (1, 2). Some of the factors linked to poor outcomes from SARS-CoV-2 infection include advanced age, male sex, and various cardiopulmonary disorders (3).

Within the spectrum of lung diseases, emerging evidence indicates that patients with interstitial lung diseases (ILDs) have higher mortality from SARS-CoV-2 (4). Specifically, one case-control study from a large academic institution in the United States detected a nearly fourfold increase in adjusted odds of death from SARS-CoV-2 infection in patients with ILDs versus matched control subjects (5). Notably, these findings were replicated in other smaller studies, supporting the notion that patients with ILD are particularly vulnerable to this infection (6, 7). However, ILDs represent a

Ⓞ This article is open access and distributed under the terms of the Creative Commons Attribution Non-Commercial No Derivatives License 4.0. For commercial usage and reprints, please e-mail Diane Gern.

**Table 1.** Mortality risk comparison between study and control cohorts

Cohort	Patients in Cohort (after Matching)	Deceased Patients	Risk (%)	Risk Difference (%)	Confidence Interval (%)	P Value
Patient with SARS-CoV-2 without IPF	74,783	2,648	3.5	−2.0	(−2.2 to −1.7)	<0.0001
Patient with SARS-CoV-2 with IPF	74,783	4,113	5.5			
Patient with SARS-CoV-2 without RA-ILD	1,306	35	2.7	−2.9	(−4.4 to −1.4)	0.0002
Patient with SARS-CoV-2 with RA-ILD	1,306	73	5.6			
Patient with SARS-CoV-2 without SSc-ILD	5,639	111	2.0	−1.3	(−1.9 to −0.7)	<0.0001
Patient with SARS-CoV-2 with SSc-ILD	5,639	183	3.2			
Patient with SARS-CoV-2 without Sjogren's-ILD	47,327	816	1.7	0.2	(0.04 to 0.4)	0.02
Patient with SARS-CoV-2 with Sjogren's-ILD	47,327	723	1.5			
Patient with SARS-CoV-2 without HP	4,471	115	2.6	−0.7	(−1.4 to −0.04)	0.04
Patient with SARS-CoV-2 with HP	4,471	148	3.3			

*Definition of abbreviations:* HP = hypersensitivity pneumonitis; ILD = interstitial lung disease; IPF = idiopathic pulmonary fibrosis; RA = rheumatoid arthritis; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; SSc = scleroderma.

**Table 2.** Mortality risk comparison between study cohorts

Cohort	Patients in Cohort (after Matching)	Deceased Patients	Risk (%)	Risk Difference (%)	Confidence Interval (%)	P Value
Patient with SARS-CoV-2 with IPF	36,057	1,417	3.9	2.1	(1.8 to 2.3)	<0.0001
Patient with SARS-CoV-2 with Sjogren's-ILD	36,057	671	1.9			
Patient with SARS-CoV-2 with IPF	5,639	186	3.3	0.1	(−0.6 to 0.7)	0.16
Patient with SARS-CoV-2 with SSc-ILD	5,639	183	3.2			
Patient with SARS-CoV-2 with SSc-ILD	5,639	183	3.2	1.6	(1.1 to 2.2)	<0.0001
Patient with SARS-CoV-2 with Sjogren's-ILD	5,639	90	1.6			

*Definition of abbreviations:* ILD = interstitial lung disease; IPF = idiopathic pulmonary fibrosis; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; SSc = scleroderma.

heterogeneous group of disorders, and it remains to be determined whether all ILDs or just specific subtypes have higher SARS-CoV-2–related mortality.

## Methods

To address this, we performed a large, retrospective cohort study to evaluate outcomes from SARS-CoV-2 infection among patients with different ILD subtypes. Analyses were performed using data from the TriNetX Analytics Network, a global research network containing records from millions of patients (8–11). This large, multicenter database includes relevant information on diagnoses, procedures, medications, and laboratory values and incorporates patients from both the inpatient and outpatient environments. Patients included in our study were adults  $\geq 18$  years of age diagnosed with SARS-CoV-2 between the periods of January 1, 2020 to February 1, 2022. Patients with SARS-CoV-2 were identified based on diagnostic coding for coronavirus disease (COVID-19) or documentation of a positive polymerase chain reaction test result. Study cohorts included patients with one of the following ILD diagnostic subtypes: idiopathic pulmonary fibrosis (IPF), rheumatoid arthritis–ILD, scleroderma (SSc)-ILD, Sjogren's syndrome with lung involvement, and hypersensitivity pneumonitis. Control cohorts had SARS-CoV-2 but no diagnosis of ILD. Study and control cohorts underwent propensity score matching for age, sex, history of nicotine dependence, body mass index, diabetes mellitus, ischemic heart disease, hypertensive disease, and cerebrovascular disease before analyses. Cohorts were not matched for other types of lung diseases. For mortality

comparisons, we used “Deceased at 90 days” after SARS-CoV-2 diagnosis as our primary endpoint, and relative risk comparisons were performed on ILDs with more than 5,000 patients.

## Results

We identified a total of 133,526 patients with SARS-CoV-2 and a diagnosis of ILD. Overall prevalence varied among different ILD subtypes, with IPF being the most prevalent (74,783 cases), followed by Sjogren's with lung involvement (47,327) and Ssc-ILD (5,639) (Table 1). After propensity score matching, the risk of mortality was increased for all ILD subtypes (IPF, rheumatoid arthritis–ILD, SSc-ILD, and hypersensitivity pneumonitis), with the exception of ILD from Sjogren's syndrome, which had a lower overall mortality than matched control subjects (Table 1). For highly prevalent ILDs, a trend toward higher mortality risk was seen in IPF and SSc-ILD, as mortality risk for IPF and SSc-ILD were both higher than ILD in Sjogren's syndrome (Table 2).

## Discussion

ILD is recognized as a risk factor for death from SARS-CoV-2, but this study is the first to look at outcomes among patients with different ILD subtypes. Our major finding is that all ILDs increase mortality from SARS-CoV-2, with the exception of Sjogren's syndrome, which had a lower mortality than control subjects.

Specific factors contributing to higher mortality among different ILD subtypes were not identified in our study. However, it is reasonable to assume that infection may have increased mortality by

accelerating progression or causing acute exacerbations, a manifestation of ILD known to associate with substantial morbidity and mortality. It is also possible that properties intrinsic to the ILD lung contributed to worsening outcomes; this includes the potential impact of dense regions of lung fibrosis on immune cell trafficking and the role of dysfunctional alveolar type 2 cells (12) and activated myofibroblasts on lung injury and repair (13, 14).

A surprising finding in our study was that patients with Sjogren's syndrome had a reduced SARS-CoV-2 mortality. Interestingly, histopathological features of this disease are unique to other ILDs, which includes the infiltration of lymphocytes around airways and cystic dilation of distal airspaces (15, 16). Whether these structural changes somehow contribute to altering SARS-CoV-2 biology is unknown; however, we speculate that the binding of virus to epithelium may be reduced by the cystic dilation of airspaces. It is also tempting to speculate that factors specific to Sjogren's syndrome may have influenced the course of disease. For example, autoantibodies to Ro52 target a protein already linked to neutralizing viruses (17).

Our study did not detect a significant increase in mortality in IPF versus other ILDs. This was somewhat surprising, given that IPF is considered the most aggressive ILD (16). Indeed, less than half of all patients with IPF are alive at 5 years, whereas the majority of patients with SSc-ILD are alive over a similar time period (18). Our observation that mortality was similar among patients with IPF and SSc-ILD challenges traditional thinking about the unique vulnerability of patients with IPF.

Although our study has many strengths, we also recognize it has weaknesses. As with any study that relies on administrative data, we recognize our results may have been skewed by inaccuracies in diagnostic coding. Also, other confounding variables not included in our analyses may have affected the results. Moreover, the interpretation of findings is limited without details about disease severity, antiviral treatments, underlying immunosuppressive drugs, and primary cause of death. Finally, some cohorts had small numbers of patients, making it hard to generalize our results to larger populations.

In conclusion, our study suggests that mortality related to SARS-CoV-2 is increased in patients with different subtypes of ILD, highlighting the importance of prevention and early treatment in this diverse patient population. ■

**Author disclosures** are available with the text of this letter at [www.atsjournals.org](http://www.atsjournals.org).

Joy Zhao, B.S.  
Brandon Metra, M.D.  
Gautam George, M.D.  
Jesse Roman, M.D.  
Joseph Mallon, B.S.  
Baskaran Sundaram, M.D.  
Michael Li, Ph.D.  
Ross Summer, M.D.\*  
Thomas Jefferson University  
Philadelphia, Pennsylvania

ORCID IDs: 0000-0001-9288-5093 (J.Z.); 0000-0001-7668-6975 (B.M.); 0000-0003-4491-2274 (G.G.); 0000-0003-2956-9674 (J.R.); 0000-0001-9039-0396 (B.S.); 0000-0003-0351-2637 (M.L.); 0000-0003-4615-4956 (R.S.).

\*Corresponding author (e-mail: [ross.summer@jefferson.edu](mailto:ross.summer@jefferson.edu)).

## References

- Ludwig S, Zarbock A. Coronaviruses and SARS-CoV-2: a brief overview. *Anesth Analg* 2020;131:93–96.
- Cucinotta D, Vanelli M. WHO declares COVID-19 a pandemic. *Acta Biomed* 2020;91:157–160.
- Centers for Disease Control and Prevention. People with certain medical conditions. [accessed 2021 Sept 19]. Available from: <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-with-medical-conditions.html>.
- Huang H, Zhang M, Chen C, Zhang H, Wei Y, Tian J, et al. Clinical characteristics of COVID-19 in patients with preexisting ILD: a retrospective study in a single center in Wuhan, China. *J Med Virol* 2020;92:2742–2750.
- Esposito AJ, Menon AA, Ghosh AJ, Putman RK, Fredenburgh LE, El-Chemaly SY, et al. Increased odds of death for patients with interstitial lung disease and COVID-19: a case-control study. *Am J Respir Crit Care Med* 2020;202:1710–1713.
- Lee H, Choi H, Yang B, Lee SK, Park TS, Park DW, et al. Interstitial lung disease increases susceptibility to and severity of COVID-19. *Eur Respir J* 2021;58:2004125.
- Drake TM, Docherty AB, Harrison EM, Quint JK, Adamali H, Agnew S, et al.; ISARIC4C Investigators. Outcome of hospitalization for COVID-19 in patients with interstitial lung disease: an international multicenter study. *Am J Respir Crit Care Med* 2020;202:1656–1665.
- Wang L, Wang Q, Davis PB, Volkow ND, Xu R. Increased risk for COVID-19 breakthrough infection in fully vaccinated patients with substance use disorders in the United States between December 2020 and August 2021. *World Psychiatry* 2022;21:124–132.
- Khan A, Bilal M, Morrow V, Cooper G, Thakkar S, Singh S. Impact of the coronavirus disease 2019 pandemic on gastrointestinal procedures and cancers in the United States: a multicenter research network study. *Gastroenterology* 2021;160:2602–2604.e5.
- Turk MA, Landes SD, Formica MK, Goss KD. Intellectual and developmental disability and COVID-19 case-fatality trends: TriNetX analysis. *Disabil Health J* 2020;13:100942.
- Singer ME, Taub IB, Kaelber DC. Risk of myocarditis from COVID-19 infection in people under age 20: a population-based analysis. *medRxiv*; 2021 [accessed 2022 Mar 21]. Available from: <https://doi.org/10.1101/2021.07.23.21260998>.
- Bui LT, Winters NI, Chung M-I, Joseph C, Gutierrez AJ, Habermann AC, et al.; Human Cell Atlas Lung Biological Network. Chronic lung diseases are associated with gene expression programs favoring SARS-CoV-2 entry and severity. *Nat Commun* 2021;12:4314.
- Sang Y, Miller LC, Blecha F. Macrophage polarization in virus-host interactions. *J Clin Cell Immunol* 2015;6:311.
- Shirey KA, Lai W, Pletneva LM, Karp CL, Divanovic S, Blanco JC, et al. Role of the lipoxigenase pathway in RSV-induced alternatively activated macrophages leading to resolution of lung pathology. *Mucosal Immunol* 2014;7:549–557.
- Gupta N, Wikenheiser-Brokamp KA, Fischer A, McCormack FX. Diffuse cystic lung disease as the presenting manifestation of Sjogren syndrome. *Ann Am Thorac Soc* 2016;13:371–375.
- Cottin V. Significance of connective tissue diseases features in pulmonary fibrosis. *Eur Respir Rev* 2013;22:273–280.
- Burbelo PD, Teos LY, Herche JL, Iadarola MJ, Alevizos I. Autoantibodies against the immunoglobulin-binding region of Ro52 link its autoantigenicity with pathogen neutralization. *Sci Rep* 2018;8:3345.
- Wells AU, Cullinan P, Hansell DM, Rubens MB, Black CM, Newman-Taylor AJ, et al. Fibrosing alveolitis associated with systemic sclerosis has a better prognosis than lone cryptogenic fibrosing alveolitis. *Am J Respir Crit Care Med* 1994;149:1583–1590.

Copyright © 2022 by the American Thoracic Society