

Exercise Limitation in Children and Adolescents with Mild-to-Moderate Asthma

Olga Lagiou^{1,2,*}
Sotirios Fouzas^{1,2,*}
Dimosthenis Lykouras³
Xenophon Sinopidis²
Ageliki Karatza²
Kiriakos Karkoulis³
Gabriel Dimitriou²
Michael B Anthracopoulos^{1,2}

¹Pediatric Respiratory Unit, University of Patras Medical School, Patras, Greece;

²Department of Pediatrics, University of Patras Medical School, Patras, Greece;

³Department of Respiratory Medicine, University Hospital of Patras, Patras, Greece

*These authors contributed equally to this work

Background: Children with uncontrolled asthma are less tolerant to exercise due to ventilatory limitation, exercise-induced bronchoconstriction (EIB), or physical deconditioning. The contribution of these factors in children with controlled mild-to-moderate asthma is unknown.

Objective: To explore the underlying mechanisms of reduced exercise capacity in children with controlled mild-to-moderate asthma.

Methods: This was a cross-sectional study of 45 children and adolescents (age 8–18 years) with controlled mild-to-moderate asthma (asthma control test score 21–25) and 61 age-matched healthy controls. All participants completed a physical activity questionnaire and performed spirometry and cardiopulmonary exercise testing (CPET; maximal incremental protocol). Spirometric indices and CPET parameters were compared between the two groups. The effect of EIB (FEV₁ decrease >10% post CPET), ventilatory limitation and physical deconditioning on maximum oxygen uptake ($\dot{V}O_{2peak}$), was assessed by multivariable linear regression.

Results: 62.2% of children with asthma and 29.5% of controls ($P = 0.002$) were categorized as inactive. Reduced exercise capacity ($\dot{V}O_{2peak} < 80\%$) was noted in 53.3% of asthmatics and 16.4% of controls ($P < 0.001$). EIB was documented in 11.1% of participants with asthma. Physical deconditioning was noted in 37.8% of children with asthma and in 14.8% of controls ($P = 0.013$). Physical deconditioning emerged as the only significant determinant of $\dot{V}O_{2peak}$, irrespective of asthma diagnosis, body mass index, ventilatory limitation and EIB.

Conclusion: Children with controlled mild-to-moderate asthma are less tolerant to strenuous exercise than their healthy peers. The decreased exercise capacity in this population should mainly be attributed to physical deconditioning, while the contribution of ventilatory limitation and EIB is rather small.

Keywords: asthma, cardiopulmonary exercise testing, children, exercise

Plain Language Summary

Why was the study done?

Children with severe or uncontrolled asthma experience exercise intolerance due to ventilatory limitation and exercise-induced bronchoconstriction. The contribution of these factors in children with controlled mild-to-moderate asthma is unknown.

What did the researchers do and find?

In this study, 45 children and adolescents with controlled mild-to-moderate asthma and 61 age-matched healthy controls performed cardiopulmonary exercise testing. We found that children with controlled mild-to-moderate asthma were less tolerant to strenuous exercise than their healthy peers. However, their decreased aerobic capacity was mainly attributed to

Correspondence: Michael B Anthracopoulos
Pediatric Respiratory Unit, University of Patras Medical School, Patras, 26504, Greece
Tel +30 2610999716
Email manthra@otenet.gr

physical deconditioning, while the contribution of ventilatory limitation and exercise-induced bronchoconstriction was minimal.

What do these results mean?

Our findings suggest that children with mild-to-moderate asthma do not present exercise limitation due to the disease itself and, thus, should be encouraged to participate in physical activities systematically and without constraints.

Introduction

Asthma represents the most frequent chronic respiratory disorder in childhood.¹ The disease is characterized by recurrent episodes of reversible airway obstruction that differ in frequency and severity depending on the level of asthma control.² Children with controlled disease do not experience day- or night-time symptoms or limitation of their physical activity; however, exposure to triggering factors, such as viral respiratory infections, allergens, and physical exercise, may unveil asthma symptoms and lead to disease exacerbation.²

Although physical activity should be encouraged in children with asthma,³ imposed restrictions are not uncommon in this population.⁴ Such attitudes relate primarily to the parental perception that physical exercise may precipitate asthma symptoms⁵ and/or to children's belief that the disease may significantly decrease their physical performance.⁶ However, evidence of a causal relationship between asthma and reduced exercise capacity (ie the amount of physical exertion that an individual can sustain) does not exist,^{7–11} although children with exercise induced bronchoconstriction (EIB) or those with uncontrolled disease tend to be less active than their peers.^{8,10–13} The few studies in which cardiopulmonary exercise testing (CPET) was employed, suggest that a substantial percentage of children with troublesome disease have reduced exercise capacity simply due to physical deconditioning (ie a decline in physical function of the body as a result of physical inactivity and disuse).^{13–15} The combined effect of EIB, ventilatory limitation (ie a ventilatory output during exercise that closely approaches or matches maximal ventilatory capacity) and physical deconditioning on exercise capacity of pediatric patients with controlled mild-to-moderate disease, ie on the vast majority of children with asthma,^{1,2} remains largely unknown.

The aim of this study was to explore the underlying mechanisms of reduced exercise capacity in children with controlled mild-to-moderate asthma using CPET. Based on evidence from children with uncontrolled disease,¹⁵ we

hypothesized that ventilatory limitation and EIB may have minimal contribution to the level of exercise capacity in this population.

Materials and Methods

Study Design and Population

This was a cross-sectional study of Caucasian European children aged 8–18 years with mild-to-moderate asthma and age-matched healthy controls. The study was conducted at the CPET laboratory of the Pediatric Respiratory Unit of the University Hospital of Patras, Greece, between June 10 and September 6, 2019.

Participants with asthma were children who: a) attended our Outpatient Asthma Clinics regularly in the last two years, b) had at least one spirometry with FEV₁ < 80% or FEV₁ reversibility ≥12% in the previous year, c) were under controller therapy for at least six months in the previous year and, d) were classified as having well controlled, mild-to-moderate asthma (treatment steps 1 and 2), according to Global Initiative for Asthma (GINA) criteria² and the Asthma Control Test (ACT) (ie score ≥20). All had interrupted their controller medication for at least four weeks prior to CPET. Healthy controls were recruited from local public schools;¹⁶ they all had normal spirometry and no asthma diagnosis or prescription of relevant medication within the last two years. All participants had normal echocardiographic examination on the day of enrolment. Exclusion criteria for both groups were: cardiac disease, major disabilities (neurological, muscular, neurodevelopmental), lower respiratory infection in the last month or non-specific respiratory symptoms (including symptoms of allergic rhinitis) in the week prior to enrollment.

The study protocol was approved by the Ethics Committee of the University Hospital of Patras. Study participants and their parents were informed about the purpose of the study. Parental informed written consent and child's informed verbal assent were obtained at enrollment. The study was conducted in accordance with the Declaration of Helsinki.

ACT and Physical Activity Questionnaire

On the day of study visit, demographic data were collected, and height and weight were measured. The body mass index (BMI) was calculated and assessed (*z* scores) according to the International Obesity Task Force norms.¹⁷ Children with asthma were asked to complete the age-appropriate Greek version of ACT (<https://www.asthma>

controltest.com). In addition, all participants completed the Greek version of the International Physical Activity Questionnaire - Short Form (IPAQ-SF)¹⁸ to estimate the intensity, frequency, and duration of their physical activities. Children with at least 60 minutes of moderate-to-vigorous physical activity per day^{19,20} were classified as “active”, otherwise they were considered to be “inactive”. The ACT and the modified IPAQ-SF were completed in the presence of one of the investigators.

Spirometry

All participants underwent spirometry using a Micro 5000 spirometer (Medisoft, Sorinnes, Belgium) according to the ERS/ATS guidelines.²¹ Forced expiratory volume at 1 s (FEV₁), forced vital capacity (FVC), FEV₁/FVC ratio, and forced expiratory flow between 25 and 75% of FVC (FEF_{25–75}) values were assessed according to Global Lung Initiative normative data.²² Spirometry was repeated at 5, 10, 15 and 20 min after CPET; a decrease of $\geq 10\%$ in at least one of these time points was defined as EIB.

Cardiopulmonary Exercise Testing

Exercise testing was performed in an air-conditioned room at a temperature of 21–22°C and relative humidity of 50–60%, using the Ultima CPX CPET (MGC Diagnostics, Saint Paul, MN) with a cycle ergometer. Following a 3 min warm-up period of free pedaling, the imposed workload was increased in a linear ramp pattern by 15 watts/min if subject’s height was < 150 cm or by 20 watts/min if subject’s height was ≥ 150 cm, up to exhaustion.²³ Participants were instructed to maintain a constant pedaling rate of 60 \pm 5 rpm throughout the trial. Continuous electrocardiographic, blood pressure, and oxygen saturation (SpO₂; special ear-lobe sensor) monitoring was also undertaken. The test was terminated when the child was unable to maintain the pedaling speed despite encouragement or when specific adverse events (eg symptoms: chest pain, excessive dyspnea, dizziness, headache, nausea, pallor; electrocardiographic abnormalities; SpO₂ $< 92\%$) occurred.²⁴

The following variables were recorded: workload (in watts), oxygen uptake ($\dot{V}O_2$; in mL/kg/min), carbon dioxide output ($\dot{V}CO_2$; in mL/kg/min), respiratory exchange ratio (RER; $\dot{V}CO_2/\dot{V}O_2$), heart rate (HR; in bpm), and minute ventilation ($\dot{V}E$; in L/min); all parameters were obtained in a breath-by-breath mode and averaged over an 8-breath moving window. The maximum (“peak”) value of $\dot{V}O_2$, $\dot{V}CO_2$,

RER, $\dot{V}E$ and HR was defined as the average value of the last 15 s of the trial. Predicted $\dot{V}O_{2peak}$ values were calculated using the equations proposed by Cooper et al.²⁵ The ratios $\dot{V}O_{2peak}$ to maximum workload ($\dot{V}O_{2peak}/w$), $\dot{V}O_{2peak}$ to maximum HR ($\dot{V}O_{2peak}/HR$), $\dot{V}E_{peak}/\dot{V}O_{2peak}$, $\dot{V}E_{peak}/\dot{V}CO_{2peak}$, $\dot{V}E_{peak}$ to maximal voluntary ventilation (MVV, calculated as $35 \times FEV_1$) and $\dot{V}E_{peak}$ to $30 \times FEV_1$ ²⁶ were determined. The ventilation efficiency slope ($\dot{V}E/\dot{V}CO_2$ slope) was also computed. The anaerobic threshold (AT) was determined by the V-slope method, while the level of breathing discomfort at the end of the test was assessed by the modified 10-point Borg scale.

A CPET trial was considered maximal when: a) the child presented signs of maximum physical effort (sweating, fatigue), b) the HR was $\geq 85\%$ of the age-predicted value, and c) the RER was > 1.05 . Ventilatory limitation was defined as maximum $\dot{V}E_{peak}/30 \times FEV_1 > 0.85$ ²⁶ in a child with $\dot{V}O_{2peak} < 80\%$ of predicted. Participants with $\dot{V}O_{2peak} < 80\%$ of predicted and an early plateau in the $\dot{V}O_{2peak}/HR$ or a drop of $\dot{V}O_{2peak}/HR$ before peak exercise were categorized as having cardiac impairment. Physical deconditioning was defined as $\dot{V}O_{2peak} < 80\%$ of predicted, without ventilatory limitation or cardiac impairment.²⁷

Statistics

Continuous variables are presented as mean \pm SD or median with range; comparisons between the two study groups were performed with *Student’s t* or *Mann–Whitney U*-tests, respectively. The chi-square or Fisher’s exact tests were used for the comparisons of categorical variables. Univariable and multivariable linear regression was used to assess the effect of sex, age, BMI (z-score), asthma, EIB, ventilatory limitation and physical deconditioning on the achieved $\dot{V}O_{2peak}$. All predictors were first tested for collinearity and if it was present, the predictor was excluded from the model. Statistical analyses were performed with the IBM SPSS version 25 (IBM Corp., Armonk, NY).

Results

A total of 106 children participated in the study, 45 with controlled mild-moderate asthma and 61 healthy controls. The baseline characteristics of the study population are presented in [Table 1](#). There were no differences in age, height,

and weight between the two groups; children with asthma had higher BMI z-score. The median ACT score of children with asthma was 23 (range 21–25). According to the IPAQ-SF, 62.2% of the participants with asthma and 29.5% of their healthy peers were categorized as inactive ($P=0.002$; Table 1).

The children in the asthma group had lower FEV_1 , FEV_1/FVC and FEF_{25-75} as compared with controls (Table 1). At study entry, the lowest value for FEV_1 in children with asthma was 82.7% of predicted (-1.52 z-scores) and for FEV_1/FVC 0.81 (-0.97 z-scores). The lowest values for FEV_1 and FEV_1/FVC in the non-asthma group were 85.3% of predicted (-1.23 z-scores) and 0.84 (-0.29 z-scores), respectively.

All participants performed CPET that met the maximal criteria. The CPET parameters are presented in Table 2. The children with asthma achieved significantly lower mean $\dot{V}O_{2peak}$, maximal work, and $\dot{V}O_{2peak}/HR$; they also had lower $\dot{V}E_{peak}$, $\dot{V}E_{peak}/MVV$ and $\dot{V}E_{peak}/30 \times FEV_1$ (Table 2). Conversely, there were no differences between the two groups in HR_{max} , $\dot{V}O_{2peak}/work$, RER , $\dot{V}E_{peak}/\dot{V}O_{2peak}$, $\dot{V}E_{peak}/\dot{V}CO_{2peak}$, ventilation efficiency slope and level of desaturation; the Borg dyspnea scores were also comparable (Table 2).

Reduced cardiopulmonary exercise capacity ($\dot{V}O_{2peak} < 80\%$ of predicted) was noted in 53.3% of children with asthma and in 16.4% of controls ($P < 0.001$) (Figure 1). Five children, all with asthma, were diagnosed with EIB. Ventilatory limitation was noted in 13.3% of those with asthma and in 1.6% of controls ($P=0.040$); two children (4.4% of those with asthma) had both ventilatory limitation and EIB. Physical deconditioning was noted in 37.8% of participants with asthma and in 14.8% of controls ($P=0.013$) (Figure 1).

Children with asthma, and those with increased BMI or physical deconditioning were more likely to achieve a lower $\dot{V}O_{2peak}$. However, physical deconditioning emerged as the only significant determinant of cardiopulmonary exercise capacity in the multivariable regression model, irrespective of asthma diagnosis, BMI z-score, ventilatory limitation and presence of EIB (Table 3).

Discussion

In this cross-sectional study we explored the exercise capacity of children with controlled mild-moderate asthma and the underlying mechanisms that may lead to its reduction in this population. We found that

children with asthma were less exercise tolerant than their non-asthmatic peers; the main determinant of reduced exercise capacity was physical deconditioning, while the contribution of EIB and ventilatory limitation was minimal.

Studies employing CPET to assess the exercise capacity of children with asthma are sparse. Two earlier reports showed that children with newly diagnosed¹³ and those with persistent asthma¹⁴ had lower $\dot{V}O_{2peak}$ than their healthy peers; however, the underlying mechanisms of reduced exercise capacity (ie EIB, ventilatory or cardiovascular limitation) were not explored.^{13,14} A recent case-control study confirmed that children with severe refractory asthma may be less tolerant to exercise.¹⁵ Using a CPET protocol similar to ours, the investigators also showed that EIB (documented in 30% of cases), physical deconditioning (25% of cases), and a combination of the two (25% of cases) could explain the reduced exercise capacity in that cohort.¹⁵ Similar studies in adults have yielded contradictory results. Earlier reports suggest that the reduced $\dot{V}O_{2peak}$ in patients with controlled moderate-to-severe asthma cannot be attributed to airflow limitation,²⁸ while others have shown that the reduced exercise capacity is linked to small airways disease, especially in subjects with EIB.²⁹ Finally, more recent reports conclude that most adults with mild-to-moderate asthma have normal $\dot{V}O_{2peak}$ and, thus, no truly reduced exercise capacity.³⁰

To our knowledge, our study is the first to explore the mechanisms of reduced exercise capacity in children with well-controlled mild-to-moderate asthma. We found that the contribution of EIB was minimal; EIB was noted in 11.1% of participants with asthma and did not emerge as a significant determinant of $\dot{V}O_{2peak}$. In a recent meta-analysis, the pooled prevalence of EIB among children and adolescents with asthma was 46% (95% CI: 39–53%), but the estimates varied significantly depending on the method of exercise challenge and the severity of the disease (17–41% in mild versus 52–99% in severe asthmatics).³¹ However, the analyzed studies were all based on EIB-oriented protocols, ie methods of exercise challenge that differ significantly from the standard incremental CPET protocol used in our study.³² In a study methodologically comparable to ours, EIB was documented in 55% of participants with asthma and emerged as major determinant of reduced exercise capacity, but that study enrolled only subjects with severe refractory disease (ACT scores between 18 and 23).¹⁵

Table I Characteristics of the Study Groups

	Asthma Group (n=45)	Control Group (n=61)	P-value
Male gender, n (%)	32 (71.1)	49 (80.3)	0.269
Age, years	12.2 ± 2.1	12.5 ± 2.3	0.481
Height, cm	154.9 ± 13.8	157.5 ± 15.1	0.360
Weight, kg	52.0 ± 15.5	50.1 ± 13.7	0.505
BMI, kg/m ²	21.4 ± 4.5	19.9 ± 3.4	0.056
BMI, z-score	0.80 ± 0.91	0.32 ± 1.09	0.019
ACT, median (range)	23 (21–25)	NA	
Physical activity questionnaire			
Inactive, n (%)	28 (62.2)	18 (29.5)	0.002
Spirometry			
FEV ₁ , L	2.4 ± 0.6	2.9 ± 0.9	<0.001
FEV ₁ , %	91.3 ± 9.4	100.9 ± 10.5	<0.001
FEV ₁ z-score	-0.74 ± 0.81	0.08 ± 0.89	<0.001
FVC, L	2.9 ± 0.7	3.3 ± 1.1	0.019
FVC, %	90.4 ± 14.1	96.4 ± 12.2	0.021
FVC, z-score	-0.83 ± 1.21	-0.32 ± 1.04	0.022
FEV ₁ /FVC	0.87 ± 0.05	0.91 ± 0.04	<0.001
FEV ₁ /FVC z-score	0.01 ± 0.81	0.81 ± 0.83	<0.001
FEF _{25–75} , L/sec	2.1 ± 1.1	2.8 ± 1.3	0.004
FEF _{25–75} , %	80.3 ± 14.9	99.4 ± 19.6	<0.001
FEF _{25–75} , z-score	-0.94 ± 0.73	-0.04 ± 0.93	<0.001

Note: Values are mean ± SD unless stated otherwise.

Abbreviations: BMI, body mass index; ACT, asthma control test; FEV₁, forced expiratory volume at 1 s; FVC, forced vital capacity; FEF_{25–75}, forced expiratory flow at 25–75% of FVC.

The effect of ventilatory limitation in our cohort, was also small; only 13.3% of children with asthma had a $\dot{V}E_{peak}/30 \times FEV_1 \geq 0.85$,²⁶ yet without significant influence on the achieved $\dot{V}O_{2peak}$. The only other report that has examined ventilatory limitation in children with asthma using CPET showed, surprisingly, that this mechanism was absent in patients with severe refractory disease.¹⁵ The different definition for ventilation limitation (ie $\dot{V}E_{peak}/MVV \geq 0.80$) compared to $\dot{V}E_{peak}/30 \times FEV_1 \geq 0.85$ in our study, may account for this striking difference. Of note, recent evidence suggests that the criterion $\dot{V}E_{peak}/30 \times FEV_1 \geq 0.85$ is significantly more accurate than the classical $\dot{V}E_{peak}/MVV$

≥ 0.80 approach in children.²⁶ Interestingly, in 4.4% (N=2) of participants with asthma both ventilatory limitation and EIB were documented. This finding most likely confirms the presence of “breakthrough” EIB (ie bronchoconstriction that occurs during the exercise)¹² in this population. The small number of children in this group did not permit for further analysis of this intriguing finding. It should be noted that all participants had a normal echocardiographic examination on the day of enrollment, and none presented signs of cardiovascular impairment during CPET.

The most important finding of this study is that physical deconditioning, defined as reduced cardiopulmonary exercise capacity in the absence of ventilatory limitation

Table 2 CPET Parameters

	Asthma Group (n=45)	Control Group (n=61)	P-value
Maximal work, watts	136.3 ± 29.2	151.9 ± 39.0	0.021
Test time, min	10.1 ± 1.4	10.6 ± 1.7	0.075
$\dot{V}O_{2peak}$, mL/kg/min	36.1 ± 7.4	41.3 ± 6.7	<0.001
$\dot{V}O_{2peak}$, % predicted	85.2 ± 19	93.8 ± 16.5	0.015
$\dot{V}O_2$ at AT, mL/kg/min	25.1 ± 6.6	29.0 ± 6.0	0.002
AT % $\dot{V}O_{2peak}$, %	69.1 ± 9.5	70.6 ± 10.4	0.448
HRmax, bpm	188 ± 9	188 ± 10	0.917
$\dot{V}O_{2peak}/HR$, mL/beat	9.6 ± 2.3	10.9 ± 3.1	0.020
$\dot{V}O_{2peak}/work$, mL/watts×min	13.2 ± 1.2	13.4 ± 1.5	0.478
RER	1.12 (1.1–1.4)	1.16 (1.1–1.4)	0.066
$\dot{V}E$, L/min	65.1 ± 14.7	72.8 ± 19.4	0.023
$\dot{V}E/MVV$	0.79 ± 0.12	0.73 ± 0.14	0.045
$\dot{V}E/30 \times FEV_1$	0.77 ± 0.12	0.71 ± 0.13	0.041
$\dot{V}E/\dot{V}O_2$	36.4 ± 4.4	36.1 ± 4.7	0.767
$\dot{V}E/\dot{V}CO_2$	31.2 ± 3.3	31.6 ± 3.4	0.526
Ventilation efficiency slope	29.6 ± 3.4	30.2 ± 3.5	0.437
SpO ₂ decrease	2 (1–3)	2 (1–3)	0.876
Minimum SpO ₂	95 (94–98)	96 (94–98)	0.304
Borg dyspnea	7 (5–8)	7 (6–8)	0.422

Notes: Values are mean ± SD or median (range) and were compared with *t* test or Mann–Whitney U, respectively.

Abbreviations: CPET, cardiopulmonary exercise testing; AT, anaerobic threshold; HR, heart rate; RER, respiratory exchange ratio; MVV, maximal voluntary ventilation.

or cardiac impairment, was documented in 37.8% of children with asthma and in 70.8% of children with asthma and low $\dot{V}O_{2peak}$. Furthermore, physical deconditioning emerged as the only significant determinant of $\dot{V}O_{2peak}$ in our cohort, irrespective of asthma diagnosis and the presence of EIB or ventilatory limitation. Physical deconditioning also outperformed and masked the effect of BMI, which has been associated with reduced exercise capacity in other studies.^{8,11} The above finding is of particular importance, because it suggests that the decreased aerobic fitness in children with asthma^{7–15} should not be attributed to underlying cardio-respiratory mechanisms that reduce their exercise capacity, but rather to the lack of regular physical activity that leads to deconditioning. Indeed, 60.7% of our children with asthma who were categorized

as inactive were also physically deconditioned, while all children with asthma and physical deconditioning were inactive. To our knowledge, only one other study has explored the underlying mechanisms of reduced exercise capacity in children with asthma by means of CPET;¹⁵ that study focused on subjects with severe refractory asthma and reached conclusions similar to ours regarding the contribution of physical deconditioning. Of note, maximal incremental CPET has been shown to be the only reliable method to systematically assess the underlying etiology of reduced exercise capacity.³³

The relationship between asthma and physical activity in childhood remains unclear. A meta-analysis of studies that have used accelerometers concluded that children with asthma are engaged in similar amount of




Figure 1 Numbers (in boxes) and percentages (x-axis) of children with low $\dot{V}O_2$ peak, EIB, VL and physical deconditioning in the two study groups. The number of inactive participants is also presented. Comparisons were performed with chi-square or Fisher's exact test. *Two children had both EIB and VL. **Abbreviations:** EIB, exercise-induced bronchoconstriction; VL, ventilatory limitation.

physical activity as their healthy peers;⁷ subsequent studies have also confirmed these findings.^{9,11} On the other hand, children with exercise induced bronchoconstriction,^{11,12} and those with newly diagnosed,¹⁰ uncontrolled,^{10,13} or severe asthma¹³ have been found to be less active than their peers. Since our study included children and adolescents with controlled

mild-to-moderate asthma, the observed differences in physical activity between cases and controls were unexpected. A likely explanation is that the children of our cohort who were less active reported realistic estimates regarding the frequency and duration of their physical activities, as they were aware that an objective measure of cardiovascular fitness (ie the CPET) would follow.

Table 3 Determinants of Cardiopulmonary Exercise Performance*

Factor	Unadjusted Effect		Adjusted Effect R ² 0.455	
	Beta	P	Beta	P
Male gender	0.089 (-0.11 to 0.28)	0.367	0.044 (-0.20 to 0.11)	0.567
Age	0.143 (-0.05 to 0.34)	0.143	0.005 (-0.15 to 0.16)	0.949
BMI, z-score	-0.247 (-0.44 to -0.06)	0.011	-0.113 (-0.28 to 0.05)	0.173
Asthma	-0.236 (-0.42 to -0.05)	0.015	-0.007 (-0.16 to 0.17)	0.932
EIB	-0.025 (-0.22 to 0.17)	0.801	-0.057 (-0.22 to 0.11)	0.497
Ventilatory limitation	-0.018 (-0.21 to 0.17)	0.853	-0.108 (-0.28 to 0.06)	0.213
Physical deconditioning	-0.645 (-0.79 to 0.50)	<0.001	-0.653 (-0.82 to 0.49)	<0.001

Notes: Values are standardized linear regression coefficients *beta* with 95% confidence intervals in parenthesis. The unadjusted effect (univariable models) refers to the effect of each factor separately. The adjusted effect (multivariable model) refers to the combined effect of these factors when adjusted for each other. The achieved power of the multivariable model was 99.99%. * $\dot{V}O_2$ peak % predicted.

Abbreviations: BMI, body mass index; EIB, exercise-induced bronchoconstriction.

The fact that the IPAQ-SF classification (ie active versus inactive) was in accordance with the percentage of children categorized as fit or unfit by CPET analysis (Figure 1), renders further support to the above hypothesis and to the validity of our results.

Physical activity is of great importance for children because it promotes cardiovascular health, improves cognitive function, and establishes the foundations for lifelong health and well-being.^{19,20} Patients with asthma may have additional benefits, since exercise training reduces airway inflammation and bronchial hyperresponsiveness, thus improving respiratory symptoms and quality of life.³⁴ Moreover, the physically active children with asthma have lower body fat and are less likely to develop obesity.³⁵ Unfortunately, despite these benefits, imposed exercise restrictions are not uncommon in this population,⁴ and relate primarily to parental and/or children's perception that more strenuous activities may precipitate asthma symptoms and affect physical performance, thus leading to devaluation by the social environment.⁶ In this regard, the findings of the present study are of importance, since we show that children with mild-to-moderate asthma are less tolerant to exercise due to physical deconditioning and not because of underlying cardio-respiratory mechanisms which limit their exercise capacity. Although the cross-sectional design of our study does not permit us to fully explore the direction of causality, the assumption that children with asthma may have adopted a less active lifestyle emerges as more plausible; such an indolent lifestyle may progressively lead to physical deconditioning and further deterioration of their exercise capacity which may, in turn, feed into a vicious cycle. Thus, regular physical activity, such as training and participation in sports, should be encouraged in all children with mild-to-moderate asthma. The importance of a thorough evaluation of the exercise capacity in these children, ideally by CPET, should also be highlighted.

Inevitably, our study has some limitations. First, the level of physical activity was not assessed by objective tools (eg accelerometers). The IPAQ-SF is—in principle—a subjective method; however, a systematic analysis of physical activity was not within the purposes of the present study. Second, measurements of allergic airway inflammation were not employed and, thus, it is not possible to assess its possible contribution. However, the study was performed in the summer months when the weather in Greece is typically hot

and dry and the prevalence of respiratory allergies is extremely low. Moreover, as already discussed, the standard incremental CPET protocol may not be appropriate to assess the presence of EIB.³² Nevertheless, in contrast to similar studies,¹⁵ our cohort consisted of children with well-controlled mild-to-moderate asthma; therefore, the low prevalence of EIB reported herein is most likely valid. Finally, this study was not designed to assess the underlying causes that may lead children with mild-to-moderate asthma to inactivity and physical deconditioning. These children may be reluctant to participate in physical activities due to the belief that the disease has affected their performance or that exercise may precipitate asthma symptoms, thus leading to stigmatization by their social environment. Parents, on the other hand, may not encourage children with asthma to participate in physical activities due to the misunderstanding that exercise may be harmful or due to lack of appropriate specialist guidance. Future studies should focus on these important issues, in order to uncover possible targets for individual- and family-oriented management strategies.

Conclusions

Children and adolescents with controlled mild-to-moderate asthma seem to be less capable of strenuous exercise than their healthy peers. However, the decreased exercise capacity in this population is mainly attributed to physical deconditioning, while the contribution of ventilatory limitation and/or EIB is minimal. Therefore, pediatric patients with mild-to-moderate asthma should be encouraged to participate in physical activities systematically and without constraints.

Abbreviations

CPET, cardiopulmonary exercise testing; EIB, exercise-induced bronchoconstriction; ACT, asthma control test; IPAQ-SF, international physical activity questionnaire - short form.

Data-Sharing Statement

Data supporting the findings of this manuscript may be available from the corresponding author on reasonable request.

Acknowledgments

We would like to thank all participating children and their families for their enthusiastic participation in this study.

Author Contributions

All authors made a significant contribution to the work reported, whether that is in the conception, study design, execution, acquisition of data, analysis and interpretation, or in all these areas; took part in drafting, revising or critically reviewing the article; gave final approval of the version to be published; have agreed on the journal to which the article has been submitted; and agree to be accountable for all aspects of the work.

Funding

This research is financed by the Hellenic Foundation for Research and Innovation (H.F.R.I.) through the “1st Call for H.F.R.I. Scholarships to PhD Candidates” to promote actions that support new scientists. Scholarship Recipient: Dr. Olga Lagiou (# 129322/I2). The publication of this paper was financed by the University of Patras.

Disclosure

The authors declare no conflicts of interest in relation to the present work.

References

- Global Asthma Network. The global asthma report; 2018. Available from: www.globalasthmareport.org. Accessed August 16, 2021.
- Global Initiative for Asthma. Global Initiative for Asthma: global strategy for asthma management and prevention (Updated 2021); 2020. Available from: www.ginasthma.org. Accessed August 16, 2021.
- Hughes D. Childhood asthma and exercise. *Paediatr Child Health*. 2014;19(9):467–468. doi:10.1093/pch/19.9.467
- Lang DM, Butz AM, Duggan AK, Serwint JR. Physical activity in urban school-aged children with asthma. *Pediatrics*. 2004;113(4):e341–346. doi:10.1542/peds.113.4.e341
- Dantas FM, Correia MA, Silva AR, Peixoto DM, Sarinho ES, Rizzo JA. Mothers impose physical activity restrictions on their asthmatic children and adolescents: an analytical cross-sectional study. *BMC Public Health*. 2014;14(1):287–294. doi:10.1186/1471-2458-14-287
- Williams B, Powell A, Hoskins G, Neville R. Exploring and explaining low participation in physical activity among children and young people with asthma: a review. *BMC Fam Pract*. 2008;9(1):40–51. doi:10.1186/1471-2296-9-40
- Cassim R, Koplin JJ, Dharmage SC, et al. The difference in amount of physical activity performed by children with and without asthma: a systematic review and meta-analysis. *J Asthma*. 2016;53(9):882–892. doi:10.1080/02770903.2016.1175474
- Sousa AW, Cabral AL, Martins MA, Carvalho CR. Daily physical activity in asthmatic children with distinct severities. *J Asthma*. 2014;51(5):493–497. doi:10.3109/02770903.2014.888571
- Matsunaga NY, Oliveira MS, Morcillo AM, Ribeiro JD, Ribeiro MAGO, Toro AADC. Physical activity and asthma control level in children and adolescents. *Respirology*. 2017;22(8):1643–1648. doi:10.1111/resp.13093
- Pike KC, Griffiths LJ, Dezateux C, Pearce A. Physical activity among children with asthma: cross-sectional analysis in the UK millennium cohort. *Pediatr Pulmonol*. 2019;54(7):962–969. doi:10.1002/ppul.24314
- Anthracopoulos MB, Fouzas S, Papadopoulos M, et al. Physical activity and exercise-induced bronchoconstriction in Greek schoolchildren. *Pediatr Pulmonol*. 2012;47(11):1080–1087. doi:10.1002/ppul.22620
- van der Kamp MR, Thio BJ, Tabak M, Hermens HJ, Driessen J, van der Palen J. Does exercise-induced bronchoconstriction affect physical activity patterns in asthmatic children? *J Child Health Care*. 2020;24(4):577–588. doi:10.1177/1367493519881257
- Vahlkvist S, Pedersen S. Fitness, daily activity and body composition in children with newly diagnosed, untreated asthma. *Allergy*. 2009;64(11):1649–1655. doi:10.1111/j.1398-9995.2009.02081.x
- Villa F, Castro AP, Pastorino AC, et al. Aerobic capacity and skeletal muscle function in children with asthma. *Arch Dis Child*. 2011;96(6):554–559. doi:10.1136/adc.2011.212431
- Faleiro RC, Mancuzo EV, Lanza FC, et al. Exercise limitation in children and adolescents with severe refractory asthma: a lack of asthma control? *Front Physiol*. 2021;11:620736. doi:10.3389/fphys.2020.620736
- Frima ES, Theodorakopoulos I, Gidaris D, et al. Lung function variability in children and adolescents with and without asthma (LUV Study): protocol for a prospective, nonrandomized, clinical trial. *JMIR Res Protoc*. 2020;9(8):e20350. doi:10.2196/20350
- Cole TJ, Lobstein T. Extended international (IOTF) body mass index cut-offs for thinness, overweight and obesity. *Pediatr Obes*. 2012;7(4):284–294. doi:10.1111/j.2047-6310.2012.00064.x
- Papathanasiou G, Georgoudis G, Papandreou M, et al. Reliability measures of the short International Physical Activity Questionnaire (IPAQ) in Greek young adults. *Hellenic J Cardiol*. 2009;50(4):283–294.
- Piercy KL, Troiano RP, Ballard RM, et al. Physical Activity Guidelines for Americans. 2nd ed. Washington DC: U.S. Department of Health and Human Services; 2018: 46–54
- NHS. Physical activity guidelines for children and young people; 2019. Available from: <https://www.nhs.uk/live-well/exercise/physical-activity-guidelines-children-and-young-people/>. Accessed August 16, 2021.
- Miller MR, Hankinson J, Brusasco V, et al. ATS/ERS Task Force. Standardisation of spirometry. *Eur Respir J*. 2005;26(2):319–338. doi:10.1183/09031936.05.00034805
- Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH. ERS Global Lung Function Initiative. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J*. 2012;40(6):1324–1343. doi:10.1183/09031936.00080312
- Godfrey S, Davies CT, Wozniak E, Barnes CA. Cardio-respiratory response to exercise in normal children. *Clin Sci*. 1971;40(5):419–431. doi:10.1042/cs0400419
- Paridon SM, Alpert BS, Boas SR, et al; American Heart Association Council on Cardiovascular Disease in the Young, Committee on Atherosclerosis, Hypertension, and Obesity in Youth. Clinical stress testing in the pediatric age group: a statement from the American Heart Association Council on Cardiovascular Disease in the Young, Committee on Atherosclerosis, Hypertension, and Obesity in Youth. *Circulation*. 2006;113(15):1905–1920. doi:10.1161/CIRCULATIONAHA.106.174375
- Cooper DM, Weiler-Ravell D, Whipp BJ, Wasserman K. Aerobic parameters of exercise as a function of body size during growth in children. *J Appl Physiol Respir Environ Exerc Physiol*. 1984;56(3):628–634. doi:10.1152/jappl.1984.56.3.628
- Pianosi PT, Smith JR. Ventilatory limitation of exercise in pediatric subjects evaluated for exertional dyspnea. *Front Physiol*. 2019;10:20. doi:10.3389/fphys.2019.00020
- American Thoracic Society; American College of Chest Physicians. ATS/ACCP Statement on cardiopulmonary exercise testing. *Am J Respir Crit Care Med*. 2003;167(2):211–277. doi:10.1164/rccm.167.2.211

28. Clark CJ, Cochrane LM. Assessment of work performance in asthma for determination of cardiorespiratory fitness and training capacity. *Thorax*. 1988;43:745–749. doi:10.1136/thx.43.10.745
29. Chrysanthopoulos C, Maksud MG, Funahashi A, Hoffmann RG, Barboriak JJ. An assessment of cardiorespiratory adjustments of asthmatic adults to exercise. *J Allergy Clin Immunol*. 1979;63:321–327. doi:10.1016/0091-6749(79)90126-X
30. Garfinkel SK, Kesten S, Chapman KR, Rebuck AS. Physiologic and nonphysiologic determinants of aerobic fitness in mild to moderate asthma. *Am Rev Respir Dis*. 1992;145:741–745. doi:10.1164/ajrccm/145.4_Pt_1.741
31. de Aguiar KB, Anzolin M, Zhang L. Global prevalence of exercise-induced bronchoconstriction in childhood: a meta-analysis. *Pediatr Pulmonol*. 2018;53(4):412–425. doi:10.1002/ppul.23951
32. Stensrud T, Carlsen KH. Can one single test protocol for provoking exercise-induced bronchoconstriction also be used for assessing aerobic capacity? *Clin Respir J*. 2008;2(1):47–53. doi:10.1111/j.1752-699X.2007.00030.x
33. Goddard T, Sonnappa S. The role of cardiopulmonary exercise testing in evaluating children with exercise induced dyspnoea. *Paediatr Respir Rev*. 2021;38:24–32. doi:10.1016/j.prrv.2020.08.002
34. Eichenberger PA, Diener SN, Kofmehl R, Spengler CM. Effects of exercise training on airway hyperreactivity in asthma: a systematic review and meta-analysis. *Sports Med*. 2013;43(11):1157–1170. doi:10.1007/s40279-013-0077-2
35. Lu KD, Manoukian K, Radom-Aizik S, Cooper DM, Galant SP. Obesity, asthma, and exercise in child and adolescent health. *Pediatr Exerc Sci*. 2016;28(2):264–274. doi:10.1123/pes.2015-0122

Journal of Asthma and Allergy

Dovepress

Publish your work in this journal

The Journal of Asthma and Allergy is an international, peer-reviewed open-access journal publishing original research, reports, editorials and commentaries on the following topics: Asthma; Pulmonary physiology; Asthma related clinical health; Clinical immunology and the immunological basis of disease; Pharmacological interventions and

new therapies. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/journal-of-asthma-and-allergy-journal>