

5-(4-Hydroxy-3-methoxybenzyl)-1,3-thiazolidine-2,4-dione monohydrate

Li-Yan Xiong,^a Ting-Fang Wang,^b Li-Ping Zheng,^b Chuan Zhang^{a*} and Feng-Chun Wang^c

^aNew Drug Research Center, School of Pharmacy, Second Military Medical University, Shanghai 200433, People's Republic of China, ^bSchool of Pharmacy, Fujian University of Traditional Chinese Medicine, Fuzhou 350108, People's Republic of China, and ^cDepartment of Medicine, The General Hospital of Beijing Military Command, Beijing 100700, People's Republic of China. Correspondence e-mail: zhangchuan@smmu.edu.cn


Received 22 November 2010; accepted 29 November 2010

Key indicators: single-crystal X-ray study; $T = 293$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.043; wR factor = 0.119; data-to-parameter ratio = 13.0.

In the title compound, $\text{C}_{11}\text{H}_{11}\text{NO}_4\text{S}\cdot\text{H}_2\text{O}$, the five-membered thiazolidine ring is nearly planar, with a maximum deviation of 0.010 (2) Å. The dihedral angle between the thiazolidine and benzene rings is 49.16 (9)°. Intermolecular O—H...O and N—H...O hydrogen bonding is present in the crystal structure.

Related literature

For the therapeutic and pharmacological properties of thiazolidinediones, see: Day (1999); Spiegelman (1998). For the synthesis of the title compound, see: Madhavan *et al.* (2002); Shoda *et al.* (1983). For related structures, see: Divjaković *et al.* (1991); Yathirajan *et al.* (2005).



Experimental

Crystal data

$\text{C}_{11}\text{H}_{11}\text{NO}_4\text{S}\cdot\text{H}_2\text{O}$
 $M_r = 271.28$
 Monoclinic, $P2_1/n$
 $a = 10.684$ (4) Å
 $b = 8.151$ (3) Å
 $c = 14.747$ (5) Å
 $\beta = 99.657$ (4)°

$V = 1266.0$ (8) Å³
 $Z = 4$
 Mo $K\alpha$ radiation
 $\mu = 0.27$ mm⁻¹
 $T = 293$ K
 $0.15 \times 0.12 \times 0.10$ mm

Data collection

Bruker SMART 1000 CCD area-detector diffractometer
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.960$, $T_{\max} = 0.974$

4985 measured reflections
 2226 independent reflections
 1902 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.047$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.119$
 $S = 1.05$
 2226 reflections
 171 parameters

H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 0.22$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.31$ e Å⁻³

Table 1
 Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N3}-\text{H3}\cdots\text{O2}^{\text{i}}$	0.86	2.03	2.886 (2)	174
$\text{O4}-\text{H4A}\cdots\text{O5}^{\text{ii}}$	0.82	1.87	2.685 (2)	171
$\text{O5}-\text{H5A}\cdots\text{O3}$	0.82 (5)	2.19 (5)	2.962 (2)	156 (4)
$\text{O5}-\text{H5A}\cdots\text{O4}$	0.82 (5)	2.37 (4)	2.947 (2)	127 (4)
$\text{O5}-\text{H5B}\cdots\text{O1}^{\text{iii}}$	0.85 (3)	1.97 (3)	2.795 (3)	163 (3)

Symmetry codes: (i) $-x + 1, -y + 2, -z + 2$; (ii) $-x + \frac{1}{2}, y + \frac{1}{2}, -z + \frac{1}{2}$; (iii) $-x + 1, -y + 2, -z + 1$.

Data collection: SMART (Bruker, 2003); cell refinement: SAINT (Bruker, 2003); data reduction: SAINT; program(s) used to solve structure: SHELXTL (Sheldrick, 2008); program(s) used to refine structure: SHELXTL; molecular graphics: SHELXTL; software used to prepare material for publication: SHELXTL.

This study was supported by the Science and Technology Commission of Shanghai special purpose for modernization of traditional Chinese medicine in 2008 (No 08DZ1970802) and the National Basic Research Program of China (No 2006CB504100 and 2009CB521907).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU5102).

References

- Bruker (2003). SAINT and SMART. Bruker AXS Inc., Madison, Wisconsin, USA.
 Day, C. (1999). *Diabet. Med.* **16**, 179–192.
 Divjaković, V., Popov-Pergal, K., Pergal, M. & Klement, U. (1991). *Acta Cryst.* **C47**, 1760–1761.
 Madhavan, G. R., Chakrabarti, R., Vikramadithyan, R. K., Mamidi, R. N. V. S., Balraju, V., Rajesh, B. M., Misra, P., Kumar, S. K. B., Lohray, B. B., Lohraya, V. B. & Rajagopalan, R. (2002). *Bioorg. Med. Chem.* **10**, 2671–2680.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Shoda, T., Mizuno, K., Hirata, T., Maki, Y. & Kawamatsu, Y. (1983). *Chem. Pharm. Bull.* **31**, 560–569.
 Spiegelman, B. M. (1998). *Diabetes*, **47**, 507–514.
 Yathirajan, H. S., Rai, K. M. L., Gaonkar, S. L., Narasegowda, R. S., Prabhuswamy, B. & Bolte, M. (2005). *Acta Cryst.* **E61**, o245–o246.

supplementary materials

Acta Cryst. (2011). E67, o16 [doi:10.1107/S1600536810049895]

5-(4-Hydroxy-3-methoxybenzyl)-1,3-thiazolidine-2,4-dione monohydrate

L.-Y. Xiong, T.-F. Wang, L.-P. Zheng, C. Zhang and F.-C. Wang

Comment

Thiazolidinediones (TZDs), which are known to sensitize tissues to insulin, have been developed and clinically used as antidiabetic agents. They have been shown to reduce plasma glucose, lipid, and insulin levels, and used for the treatment of type 2 diabetes (Day, 1999; Spiegelman, 1998). Prompted by the activity of TZDs, we have synthesized the title compound to study its crystal structure.

The asymmetric unit contains a 5-(4-hydroxy-3-methoxybenzyl)thiazolidine-2,4-dione molecule and a solvate water molecule (Fig. 1). The geometric parameter of the title compound are to its related structures (Divjakovic *et al.*, 1991; Yathirajan *et al.*, 2005). The dihedral angle between the thiazolidinedione ring [S1/C2/N3/C4/C5] and the benzene ring [C7–C12] is 49.16 (9)°. In the crystal packing (Fig. 2), the molecules are linked *via* intermolecular N1—H1···O2 hydrogen bonds. In addition, the molecule is connected to the water molecule by O5—H5···O1, O5—H5···O4, O5—H5···O3 and O4—H4···O5 hydrogen bonds which generate a three dimensional network (Table 1).

Experimental

A mixture of 2,4-tThiazolidinedione (3.51 g, 0.03 mol), 4-hydroxy-3-methoxybenzaldehyde (4.56 g, 0.03 mol), acetic acid (0.18 g, 0.003 mol) and piperidine (0.26 g, 0.003 mol) in toluene (60 ml) was refluxed for 5 h with continuous removal of water. The reaction mixture was cooled to room temperature and the resultant crystalline compound was filtered and washed with water and dried to afford the (*Z*)-5-(4-hydroxy-3-methoxybenzylidene)thiazolidine-2,4-dione. Yield=7.33 g, 97.3%. To a solution of (*Z*)-5-(4-hydroxy-3-methoxybenzylidene) thiazolidine-2,4-dione (4 g, 0.016 mol) in 1,4-dioxane (400 ml), hydrogenated in the presence of 10% Pd/C (1.0 g) at 60 psi for 24 h. The mixture was filtered through a bed of Celite. The filtrate was evaporated under reduced pressure and purified by column chromatography using 50:1 CH₂Cl₂/MeOH to afford the title compound as yellowish solid. Yield = 1.96 g, 48.6% (Madhavan *et al.*, 2002; Shoda *et al.*, 1983). Crystallization of the product was carried out by dissolving the product in 10 ml a solvent mixture of MeOH and water (4:1) at room temperature.

Refinement

Water H atoms were located in a difference Fourier map and refined isotropically. Other H atoms were positioned geometrically and refined using the riding-model approximation with C—H = 0.93–0.97, O—H = 0.82 and N—H = 0.86 Å, and $U_{\text{iso}}(\text{H}) = 1.5U_{\text{eq}}(\text{C},\text{O})$ for methyl H and hydroxy H atoms and $1.2U_{\text{eq}}(\text{C},\text{N})$ for the others.

Figures




Fig. 1. View of the asymmetric unit of the compound showing the atom-labelling scheme. Displacement ellipsoids are drawn at the 30% probability level. H atoms are presented as small spheres of arbitrary radius.




Fig. 2. Crystal packing of the title compound. Intermolecular hydrogen bonds are shown as dashed lines.

5-(4-Hydroxy-3-methoxybenzyl)-1,3-thiazolidine-2,4-dione monohydrate

Crystal data

$C_{11}H_{11}NO_4S \cdot H_2O$

$M_r = 271.28$

Monoclinic, $P2_1/n$

Hall symbol: -P 2yn

$a = 10.684 (4) \text{ \AA}$

$b = 8.151 (3) \text{ \AA}$

$c = 14.747 (5) \text{ \AA}$

$\beta = 99.657 (4)^\circ$

$V = 1266.0 (8) \text{ \AA}^3$

$Z = 4$

$F(000) = 568$

$D_x = 1.423 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 889 reflections

$\theta = 2.8\text{--}27.5^\circ$

$\mu = 0.27 \text{ mm}^{-1}$

$T = 293 \text{ K}$

Block, yellow

$0.15 \times 0.12 \times 0.10 \text{ mm}$

Data collection

Bruker SMART 1000 CCD area-detector diffractometer

Radiation source: fine-focus sealed tube graphite

Detector resolution: $10.0 \text{ pixels mm}^{-1}$

ϕ and ω scans

Absorption correction: multi-scan (SADABS; Sheldrick, 1996)

$T_{\min} = 0.960$, $T_{\max} = 0.974$

4985 measured reflections

2226 independent reflections

1902 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.047$

$\theta_{\max} = 25.0^\circ$, $\theta_{\min} = 2.2^\circ$

$h = -12 \rightarrow 12$

$k = -9 \rightarrow 6$

$l = -16 \rightarrow 17$

Refinement

Refinement on F^2

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.043$

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

$wR(F^2) = 0.119$

$S = 1.05$

2226 reflections

171 parameters

0 restraints

H atoms treated by a mixture of independent and constrained refinement

$$w = 1/[\sigma^2(F_o^2) + (0.0681P)^2 + 0.2943P]$$

$$\text{where } P = (F_o^2 + 2F_c^2)/3$$

$$(\Delta/\sigma)_{\max} = 0.001$$

$$\Delta\rho_{\max} = 0.22 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.31 \text{ e } \text{\AA}^{-3}$$

Special details

Experimental. 1H NMR (300 MHz, CDCl₃): δ 8.72(bar, 1H, N—H), 6.87–6.71 (m, 3H, 8-H, 11-H, 12-H), 5.46 (bar, 1H, 10-OH), 4.47–4.51 (m, 1H, 5-H), 3.83 (s, 3H, 9-OCH₃), 3.46 (dd, 1H, $j=14.4, 4.2$, 3-H), 3.06 (dd, 1H, $j=14.1, 9.6$, 3-H). 13C NMR (300 MHz, CDCl₃): δ 174.5, 167.5, 147.8, 145.7, 132.7, 123.1, 116.7, 114.9, 57.2, 56.1, 36.2. MS(ESI) m/z calc. for C₁₁H₁₁NO₄S 253.27, found [M–1]⁺ 252.15. m.p. 109–110°C

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.49420 (5)	1.06186 (7)	0.70922 (3)	0.0523 (2)
N3	0.51317 (16)	1.07172 (19)	0.88568 (11)	0.0430 (4)
H3	0.5303	1.1072	0.9413	0.052*
C2	0.5427 (2)	1.1623 (3)	0.81421 (15)	0.0529 (5)
C4	0.45618 (17)	0.9243 (2)	0.86643 (12)	0.0383 (4)
C5	0.43481 (18)	0.8857 (2)	0.76418 (12)	0.0403 (4)
H5	0.4868	0.7902	0.7545	0.048*
C6	0.29661 (18)	0.8453 (3)	0.72695 (13)	0.0448 (5)
H6A	0.2706	0.7526	0.7606	0.054*
H6B	0.2440	0.9383	0.7370	0.054*
C7	0.27553 (18)	0.8048 (2)	0.62553 (13)	0.0407 (4)
C8	0.31489 (17)	0.6537 (2)	0.59516 (12)	0.0386 (4)
H8	0.3523	0.5768	0.6378	0.046*
C9	0.29866 (16)	0.6178 (2)	0.50237 (12)	0.0360 (4)
C10	0.24649 (18)	0.7353 (2)	0.43822 (12)	0.0408 (4)
C11	0.2075 (2)	0.8830 (2)	0.46857 (14)	0.0512 (5)
H11	0.1718	0.9613	0.4261	0.061*
C12	0.2206 (2)	0.9167 (2)	0.56175 (14)	0.0504 (5)
H12	0.1918	1.0163	0.5812	0.061*
C13	0.3765 (2)	0.3430 (3)	0.52648 (16)	0.0541 (5)
H13A	0.3955	0.2494	0.4917	0.081*

supplementary materials

H13B	0.3128	0.3141	0.5625	0.081*
H13C	0.4520	0.3781	0.5665	0.081*
O1	0.5944 (2)	1.2929 (2)	0.82373 (12)	0.0877 (7)
O2	0.42480 (14)	0.83322 (17)	0.92352 (9)	0.0483 (4)
O3	0.33054 (14)	0.47280 (16)	0.46520 (9)	0.0485 (4)
O4	0.23570 (15)	0.69314 (17)	0.34819 (9)	0.0537 (4)
H4A	0.2034	0.7691	0.3160	0.080*
O5	0.36833 (18)	0.4211 (2)	0.27286 (12)	0.0558 (4)
H5A	0.336 (4)	0.444 (5)	0.318 (3)	0.134 (16)*
H5B	0.376 (3)	0.518 (4)	0.253 (2)	0.085 (10)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0699 (4)	0.0527 (4)	0.0342 (3)	-0.0171 (2)	0.0079 (2)	0.0023 (2)
N3	0.0578 (10)	0.0384 (9)	0.0326 (8)	-0.0079 (7)	0.0065 (7)	-0.0046 (6)
C2	0.0715 (14)	0.0423 (11)	0.0453 (11)	-0.0124 (10)	0.0112 (10)	-0.0012 (9)
C4	0.0432 (9)	0.0353 (10)	0.0352 (9)	-0.0008 (7)	0.0033 (7)	-0.0011 (7)
C5	0.0510 (10)	0.0365 (10)	0.0332 (9)	-0.0018 (8)	0.0060 (8)	-0.0021 (8)
C6	0.0518 (11)	0.0454 (11)	0.0374 (10)	-0.0063 (9)	0.0079 (8)	-0.0067 (8)
C7	0.0465 (10)	0.0381 (10)	0.0369 (10)	-0.0074 (8)	0.0050 (8)	-0.0055 (8)
C8	0.0441 (9)	0.0368 (10)	0.0337 (9)	-0.0031 (8)	0.0029 (7)	0.0046 (7)
C9	0.0409 (9)	0.0300 (9)	0.0367 (9)	-0.0029 (7)	0.0051 (7)	-0.0018 (7)
C10	0.0509 (10)	0.0363 (10)	0.0328 (9)	-0.0020 (8)	0.0006 (8)	-0.0018 (7)
C11	0.0727 (14)	0.0357 (10)	0.0405 (10)	0.0059 (10)	-0.0042 (9)	0.0021 (8)
C12	0.0679 (13)	0.0344 (10)	0.0466 (11)	0.0016 (9)	0.0028 (10)	-0.0078 (8)
C13	0.0650 (13)	0.0390 (11)	0.0599 (13)	0.0097 (10)	0.0149 (10)	0.0102 (10)
O1	0.1467 (19)	0.0578 (11)	0.0617 (11)	-0.0545 (12)	0.0262 (11)	-0.0083 (8)
O2	0.0667 (9)	0.0429 (7)	0.0340 (7)	-0.0108 (7)	0.0047 (6)	0.0033 (6)
O3	0.0718 (9)	0.0344 (7)	0.0391 (7)	0.0089 (7)	0.0084 (6)	0.0019 (6)
O4	0.0836 (10)	0.0421 (8)	0.0312 (7)	0.0092 (7)	-0.0022 (7)	-0.0009 (6)
O5	0.0797 (11)	0.0438 (9)	0.0440 (8)	0.0040 (8)	0.0106 (8)	-0.0066 (7)

Geometric parameters (\AA , $^\circ$)

S1—C2	1.751 (2)	C8—H8	0.9300
S1—C5	1.815 (2)	C9—O3	1.370 (2)
N3—C4	1.356 (2)	C9—C10	1.395 (3)
N3—C2	1.366 (3)	C10—O4	1.358 (2)
N3—H3	0.8600	C10—C11	1.373 (3)
C2—O1	1.197 (3)	C11—C12	1.385 (3)
C4—O2	1.211 (2)	C11—H11	0.9300
C4—C5	1.520 (2)	C12—H12	0.9300
C5—C6	1.523 (3)	C13—O3	1.424 (2)
C5—H5	0.9800	C13—H13A	0.9600
C6—C7	1.511 (3)	C13—H13B	0.9600
C6—H6A	0.9700	C13—H13C	0.9600
C6—H6B	0.9700	O4—H4A	0.8200
C7—C12	1.370 (3)	O5—H5A	0.82 (5)

C7—C8	1.399 (3)	O5—H5B	0.85 (3)
C8—C9	1.382 (3)		
C2—S1—C5	92.79 (9)	C9—C8—C7	120.60 (17)
C4—N3—C2	118.11 (16)	C9—C8—H8	119.7
C4—N3—H3	120.9	C7—C8—H8	119.7
C2—N3—H3	120.9	O3—C9—C8	125.52 (16)
O1—C2—N3	123.5 (2)	O3—C9—C10	114.77 (15)
O1—C2—S1	125.58 (18)	C8—C9—C10	119.71 (17)
N3—C2—S1	110.93 (15)	O4—C10—C11	124.07 (17)
O2—C4—N3	124.38 (17)	O4—C10—C9	116.61 (16)
O2—C4—C5	123.37 (17)	C11—C10—C9	119.31 (17)
N3—C4—C5	112.25 (16)	C10—C11—C12	120.76 (18)
C4—C5—C6	112.19 (15)	C10—C11—H11	119.6
C4—C5—S1	105.90 (13)	C12—C11—H11	119.6
C6—C5—S1	113.61 (13)	C7—C12—C11	120.64 (19)
C4—C5—H5	108.3	C7—C12—H12	119.7
C6—C5—H5	108.3	C11—C12—H12	119.7
S1—C5—H5	108.3	O3—C13—H13A	109.5
C7—C6—C5	112.25 (15)	O3—C13—H13B	109.5
C7—C6—H6A	109.2	H13A—C13—H13B	109.5
C5—C6—H6A	109.2	O3—C13—H13C	109.5
C7—C6—H6B	109.2	H13A—C13—H13C	109.5
C5—C6—H6B	109.2	H13B—C13—H13C	109.5
H6A—C6—H6B	107.9	C9—O3—C13	118.00 (15)
C12—C7—C8	118.92 (18)	C10—O4—H4A	109.5
C12—C7—C6	120.66 (18)	H5A—O5—H5B	98 (3)
C8—C7—C6	120.40 (17)		

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N3—H3 \cdots O2 ⁱ	0.86	2.03	2.886 (2)	174
O4—H4A \cdots O5 ⁱⁱ	0.82	1.87	2.685 (2)	171
O5—H5A \cdots O3	0.82 (5)	2.19 (5)	2.962 (2)	156 (4)
O5—H5A \cdots O4	0.82 (5)	2.37 (4)	2.947 (2)	127 (4)
O5—H5B \cdots O1 ⁱⁱⁱ	0.85 (3)	1.97 (3)	2.795 (3)	163 (3)

Symmetry codes: (i) $-x+1, -y+2, -z+2$; (ii) $-x+1/2, y+1/2, -z+1/2$; (iii) $-x+1, -y+2, -z+1$.

Fig. 1




Fig. 2

