

Detection of Cytoplasmic and Nuclear Circular RNA via RT-qPCR

Ke-En Tan^{1, Sa}, Wei Lun Ng^{1, Sb}, Chee-Kwee Ea^{1, Sc}, and Yat-Yuen Lim^{1, *}

¹Institute of Biological Sciences, Faculty of Science, Universiti Malaya, Kuala Lumpur, Malaysia

^{Sa}Current address: Department of Biochemistry and Molecular Biology, Mayo Clinic College of Medicine and Science, Rochester, MN, USA

^{Sb}Current address: Programme in Emerging Infectious Diseases, Duke-NUS Medical School, Singapore, Singapore

^{Sc}Current address: Department of Molecular Biology, University of Texas Southwestern Medical Center, Dallas, TX, USA

*For correspondence: yatyuen.lim@um.edu.my

Abstract

Circular RNA (circRNA) is an intriguing class of non-coding RNA that exists as a continuous closed loop. With the improvements in high throughput sequencing, biochemical analysis, and bioinformatic algorithms, studies on circRNA expression became abundant in recent years. However, functional studies of circRNA are still limited. Subcellular localization of circRNA may provide some clues in elucidating its biological functions by performing subcellular fractionation assay. Notably, circRNAs that are predominantly found in the cytoplasm are more likely to be involved in post-transcriptional gene regulation, e.g., acting as miRNA sponge, whereas nuclear-retained circRNAs are predicted to play a role in transcriptional regulation. Subcellular fractionation could help researchers to narrow down and prioritize downstream experiments. The majority of the currently available protocols describe the steps for subcellular fractionation followed by western blot analysis for protein molecules. Here, we present a protocol for the subcellular fractionation of cells to detect circRNA via RT-qPCR with divergent primers. Moreover, detailed steps for the generation of specific circRNAs-enriched cDNA included in this protocol will enhance the amplification and detection of low-abundance circRNAs. This will be useful for researchers studying low-abundance circRNAs.


Key features

- This protocol builds upon the method developed by Gagnon et al. (2014) and extends its application to circRNA study.
- Protocol for amplification of low levels of circRNA expression.
- Analysis takes into consideration the ratio of cytoplasmic RNA concentration to nuclear RNA concentration.

Keywords: Subcellular fractionation, RT-qPCR, Circular RNA, circRNA, Cytoplasmic circRNA, Nuclear circRNA, Low-abundance circRNA

This protocol is used in: Sci. Rep. (2021), DOI: 10.1038/s41598-021-93781-w

Graphical overview



Background

Circular RNA (circRNA) is an intriguing class of non-coding RNA that is formed through a unique mechanism known as backsplicing, in which the 5' and 3' termini are covalently joined (Jeck and Sharpless, 2014). Due to the absence of free termini in the circular structure, circRNA can easily escape from hydrolysis by numerous cellular exonucleases. CircRNAs can be derived from exons, introns, or both, with a great diversity in length. Emerging evidence has shown that circRNAs are involved in many different biological processes and human diseases by functioning as miRNA sponges, RNA-binding protein sponges, transcriptional regulators, mRNA traps, regulators of assembly and transport of cellular proteins, as well as templates for translation.

Several computational methods, such as CIRI2, CIRCexplorer2, and find_circ, have been developed to identify circRNA by employing alignment-based strategies to recognize the back-spliced junction (BSJ), which is a unique feature of circRNAs (Szabo and Salzman, 2016; Cai et al., 2020). To confirm the presence of a circRNA identified *in silico*, an experimental approach is applied to validate the computationally predicted circRNAs (Zhang et al., 2016).

Reverse-transcription PCR (RT-PCR) is commonly used for the validation of circRNAs identified by RNA sequencing using divergent primers. In this approach, circRNA is converted to cDNA via reverse transcription (RT) with the presence of random hexamer or gene-specific reverse primers. Priming with random hexamers offers flexibility by ensuring RT of all RNA sequences, but gene-specific RT enhances the sensitivity of detecting the target of interest by converting the target of interest—instead of transcribing all RNA in the mix—into cDNA. Priming with gene-specific primers could enhance the detection of circRNA of interest, especially for those present in low abundance (Bustin and Nolan, 2004).

The conventional RT-PCR assay using convergent primers is unable to differentiate circRNAs from their linear transcript when the linear genome is used as template for primer design. Divergent primers are needed for the RT-PCR, as they allow direct detection of BSJ and quantification of circRNA. Divergent primers are a pair of outward-facing primers that allow amplification of circRNAs and not linear mRNA with the same sequence in cDNA and not gDNA (Figure 1). Sanger sequencing is usually performed on the PCR product amplified with the divergent primers to confirm the presence of a BSJ, as to rule out any incorrect amplification or artifact. Quantitative RT-PCR (RT-qPCR) can be used to relatively quantify the expression of circRNA across a panel of samples.




Figure 1. Primer design for circular RNA (circRNA) detection. Divergent primers are a pair of outward-facing primers that allow amplification of circRNA backsplice junction (BSJ), whereas convergent primers are a pair of inward-facing primers that allow amplification of linear mRNA from prepared cDNA.

Combining subcellular fractionation and RT-qPCR with divergent primers may provide some hints on circRNA's biological functions. For example, circRNAs that are predominantly found in the cytoplasm are more likely to be involved in post-transcriptional gene regulation, whereas nuclear-retained circRNAs are predicted to play a role in transcription regulation. Currently available protocols are mainly for protein work, describing the steps for subcellular fractionation followed by western blot analysis, which is not entirely suitable for RNA work. Instead, this protocol allows the downstream detection of circRNA through RT-qPCR with divergent primers. A detailed protocol for the generation of specific circRNAs-enriched cDNA will also be included to better detect low-abundance circRNAs.

Materials and reagents

Biological material

1. GM12878 cell line

Reagents

1. Dulbecco's phosphate-buffered saline (PBS) (Fisher Scientific, catalog number: BP399-1)
2. Trypsin-EDTA solution (0.5%) (Gibco, catalog number: 15400054)
3. Tissue culture media: Roswell Park Memorial Institute (RPMI) 1640 medium (Gibco, catalog number: 11875093)
4. Fetal bovine serum (FBS) (Gibco, catalog number: 10270106)
5. Penicillin-Streptomycin (Gibco, catalog number: 15140122)
6. Trypan blue (Alfa Aesar, catalog number: A1860)
7. Tris-Base (Fisher Scientific, catalog number: BP152-5)
8. Potassium chloride (KCl) (Fisher Scientific, catalog number: BP366-1)
9. Magnesium chloride ($MgCl_2$) (Fisher Scientific, catalog number: BP214-500)
10. DL-Dithiothreitol (DTT) (Sigma-Aldrich, catalog number: D0632)
11. NP-40 (Sigma-Aldrich, catalog number: 9016-45-9)
12. Ribonucleoside vanadyl complexes (NEB, catalog number: S1402S)
13. Sodium acetate (Fisher Scientific, catalog number: BP333-500)
14. Molecular grade ethanol (Merck, catalog number: 64-17-5)
15. TRIzol (Life Technologies, catalog number: 15-596-018)
16. Chloroform:isoamyl alcohol (1:24) (Fisher Scientific, catalog number: BP1752I-400)
17. Molecular grade isopropanol (Fisher Scientific, catalog number: BP2618-1)

18. RNase-free water (Macherey Nigel, catalog number: 740378)
19. 10× M-MuLV RT buffer (NEB, catalog number: #M0253)
20. DNase I (NEB, catalog number: #M0303)
21. M-MuLV reverse transcriptase (NEB, catalog number: #M0253)
22. RNase inhibitor (NEB, catalog number: #M0314)
23. dNTP mix (Thermo Scientific, catalog number: R0192)
24. Random hexamer (Invitrogen, catalog number: N8080127)
25. Primers (IDT, origin: USA)
26. SYBR Fast qPCR Master Mix (KAPA Biosystem, catalog number: KK4602)

Solutions

1. Complete media for GM12878 cell line (see Recipes)
2. Hypotonic buffer (see Recipes)
3. RNA precipitation solution (RPS) (see Recipes)

Recipes

1. Complete media for GM12878 cell line

Reagent	Stock concentration	Final concentration	Volume (mL)
RPMI media			395
FBS	100%	20%	100
Penicillin-Streptomycin	10,000 U/mL	100 U/mL	5
Total			500

2. Hypotonic buffer

Reagent	Stock concentration	Final concentration	Volume (μL)
Tris (pH 7.5)	100 mM	10 mM	500
KCl	100 mM	10 mM	500
MgCl ₂	150 mM	1.5 mM	50
DTT	100 mM	0.5 mM	25
NP-40	10%	0.075%	37.5
Ribonucleoside vanadyl complexes	200 mM	2 mM	50
H ₂ O			3837.5
Total			5,000

3. RNA precipitation solution (RPS)

Reagent	Stock concentration	Final concentration	Volume (mL)
Sodium acetate (pH 5.5)	3 M	150 mM	0.5
Ethanol	99.5%		9.5
Total			10

Final concentration of each component is 0.15 M sodium acetate (pH 5.5) in ethanol.

Keep at -20 °C for storage.

Equipment

1. Refrigerated tabletop centrifuge (15 or 50 mL conical tube adaptors) (Thermo Scientific, model: Multifuge X1R)
2. Refrigerated benchtop centrifuge (1.5 mL tube rotor) (Eppendorf, model: 5424R)

3. Light microscope (Olympus, origin: USA)
4. Vortexer (Biosan, model: V-1 plus)
5. Micropipettors (Gilson, origin: USA)
6. NanoDrop 2000c UV-Vis Spectrophotometer (Thermo Scientific, origin: USA)
7. Applied Biosystems Veriti Dx 96-Well Thermal Cycler PCR Thermocycler (Applied Biosystem, origin: USA)
8. Bio-Rad Connect Real-Time PCR System (Bio-Rad, origin: USA)

Procedure

A. Subcellular fractionation

1. Cell harvesting from culture plates
 - a. Grow the cells to a confluency of ~80%. Rinse the cells with 1× PBS.
 - b. For adherent cells, detach the cells with Trypsin-EDTA. For suspension cells, proceed to step A1d.
 - c. Add an equal volume of complete media to stop the trypsin proteolysis.
 - d. Collect the cell suspension in a conical tube and centrifuge at 200× g for 5 min.
 - e. Remove the supernatant and resuspend the cell pellet with 1× PBS.
 - f. Perform cell counting and aliquot 1 × 10⁶ cells into a 1.5 mL microcentrifuge tube.
 - g. Centrifuge the tube at 200× g for 5 min and remove the supernatant.
 - h. Keep the cell pellet on ice prior to the next step.
2. RNA isolation from cytoplasmic and nuclear fractions of cells
 - a. Resuspend the cell pellet gently with 100 μL of ice-cold hypotonic buffer and incubate on ice for 5 min.
Note: Volume of the hypotonic buffer can be adjusted based on the size of cell pellet.
 - b. Check for membrane lysis with trypan blue. Proceed with centrifugation at 500× g for 10 min at 4 °C until > 90% of cells are lysed. Otherwise, increase the incubation time for step A2a until > 90% of cells are lysed.
Note: Do not overlyse the cells.
 - c. Collect the supernatant (cytoplasmic fraction) and wash the pellet (nuclear fraction) with 300 μL of ice-cold hypotonic buffer three times with centrifugation at 500× g for 5 min at 4 °C.
 - d. Add 1 mL of RPS to the cytoplasmic fraction and incubate at -20 °C for at least 1 h.
 - e. Vortex the cytoplasmic fraction in RPS and centrifuge at 18,000× g for 15 min at 4 °C.
 - f. Discard the supernatant and rinse the pellet with 70% (v/v) ice-cold ethanol.
 - g. Add 1 mL of TRIzol to the semi-dry nuclear and cytoplasmic pellets followed by the addition of 10 μL of 0.5 M EDTA.
 - h. Heat both fractions at 65 °C until the pellet dissolves with vortexing.
 - i. Cool the samples to room temperature and add 200 μL of chloroform:isoamyl alcohol (1:24).
 - j. Vortex the samples and centrifuge at 18,000× g for 10 min at room temperature.
 - k. Transfer the aqueous supernatant into a clean microcentrifuge tube.
 - l. Add an equal volume of isopropanol and incubate at -20 °C for at least 1 h.
 - m. Vortex the samples and centrifuge at 18,000× g for 15 min at room temperature.
 - n. Wash the pellet with 70% (v/v) ethanol and centrifuge at 18,000× g for 5 min.
 - o. Air-dry the pellet for 5–10 min.
 - p. Dissolve the air-dried RNA pellet in 30 μL of RNase-free water.
 - q. Quantify the RNA concentration and purity with NanoDrop 2000c UV-Vis spectrophotometer.

B. cDNA synthesis

1. DNase I treatment for isolated RNA
 - a. Prepare the RNA in PCR tube for DNase I treatment as shown in Table 1.

Table 1. Components used for DNase I treatment

Components	Stock concentration	Final concentration	Volume (μL)
MuLV RT Buffer	10×	1×	1.4
DNase I		1 U	0.5
RNA		500–2,000 ng	x
H ₂ O			Top up to 14

- b. Incubate the tube in a thermal cycler at 37 °C for 30 min.
2. Reverse transcription (RT)
 - a. Prepare the RT reaction as shown in Table 2.

Table 2. Components used for reverse transcription

Components	Stock concentration	Final concentration	Volume (μL)	
			Random hexamer	Primer specific
MuLV RT Buffer	10×	1×	0.6	0.6
dNTP mix	10 mM	0.5 mM	1	1
Random hexamer	50 mM	2.5 mM	1	–
Reverse primers	10 μM	0.5 μM	–	0.5 for each primer
RNase inhibitor		2 U	0.5	0.5
Reverse transcriptase		10 U	0.5	0.5
H ₂ O			2.4	Top up to 6
Subtotal			6	6
DNase I–treated RNA			14	14
Total			20	20

- b. Perform the cDNA conversion on a thermal cycler with the setup of 42 °C for 60 min and follow by heat inactivation at 65 °C for 20 min.

Notes:

- i. If more than four pairs of primer sets (targeting linear mRNAs and/or circRNAs) are needed for primer-specific cDNA conversion, prepare 5 μM primer mix and add 1 μL into the RT reaction.
- ii. One may start with random hexamer for the cDNA conversion and switch to primer-specific cDNA conversion if the amplification of the targeted circRNA BSJs is weak, absent, or unspecific.
- iii. Random hexamer-generated cDNA allows amplification of any circRNA target of interest at any one time, while primer-specific-generated cDNA only allows amplification of the planned circRNA targets and does not allow any amplification of any unplanned circRNA targets.
- iv. Caution is needed when generating circRNA-specific cDNA, as the remaining unused reverse primer in the RT reaction could possibly interrupt or participate in qPCR process and lead to undesired amplification from linear mRNA.

C. Quantitative PCR (qPCR)

1. Prepare the PCR reaction as shown in Table 3.

Table 3. Components used for SYBR Green RT-qPCR

Components	Stock concentration	Final concentration	Volume (μL)
SYBR Fast qPCR Master Mix	2×	1×	5
Forward primer	10 μM	0.1 μM	0.1
Reverse primer	10 μM	0.1 μM	0.1
6-fold diluted cDNA			3

RNase-free water	1.8
Total	10

- Perform the qPCR on a real time thermal cycler with the cycling parameter of 95 °C for 3 min, followed by 40 cycles of 95 °C for 2 s and 60 °C for 20 s.
- Calculate the relative expression of each gene based on the equation below:

$$(a) \text{Relative expression in cytoplasm} = \frac{2^{(Cq_{cyto} - Cq_{nuc})}}{\text{RNA ratio}}$$

$$(b) \text{Relative expression in cytoplasm} = \frac{2^{(Cq_{nuc} - Cq_{cyto})}}{\text{RNA ratio}}$$

RNA ratio is the ratio of cytoplasmic RNA concentration to nuclear RNA concentration eluted in a similar volume of water. Gene expression for a cytoplasmic marker is calculated using formula (a), whereas gene expression of a nuclear marker is calculated using formula (b).

Data analysis

Representative data

Subcellular fractionation was performed using GM12878 cells, which is a lymphoblastoid cell line that contains Epstein-Barr Virus (EBV). The concentration and purity of each fraction were measured using NanoDrop 2000c (Table 4). The cytoplasmic *RPL30* and nuclear *MALAT1* transcripts were used as positive controls to indicate the purity of cytoplasmic and nuclear fractions, respectively. As shown in Figure 2, *RPL30* was enriched in the cytoplasm, whereas *MALAT1* was enriched in the nuclear fraction, indicating that subcellular fractionation was successfully performed. In this representative data, the linear EBV latent membrane protein 2 (*LMP-2*) transcripts were found in both cytoplasmic and nuclear fractions. The subcellular localization pattern of circ*LMP-2_e5* is similar to that of linear *LMP-2A* and *LMP-2B*.

Note that the protocol described here requires a real-time PCR machine to generate the data, which may not be available in all laboratories. In such a case, conventional agarose gel-based RT-PCR can be conducted to obtain a semi-quantitative result.

Table 4. Concentration and purity of cytoplasmic and nuclear fraction

	Concentration (ng/μL)	260/280	260/230
Cytoplasmic fraction 1	1,279	1.96	1.95
Cytoplasmic fraction 2	1,242	1.93	1.74
Nuclear fraction 1	566	1.76	0.71
Nuclear fraction 2	430.3	1.87	1.46

1 and 2 represent biological replicates.




Figure 2. Subcellular localization of circLMP-2_e5. RT-qPCR analysis showed *LMP-2* and *circLMP-2_e5* were localized in both nucleus and cytoplasm of GM12878 cells. Data was normalized to the RNA yield ratio and represents the mean \pm SD of two independent experiments. This result was published in Figure 3C in Tan et al. (2021).

Validation of protocol

This protocol or parts of it has been used and validated in the following research article:

Tan et al. (2021). Identification and characterization of a novel Epstein-Barr Virus-encoded circular RNA from *LMP-2* Gene. *Scientific Reports* (Figure 3, panel C).

Acknowledgments

The work was supported financially by the Ministry of Higher Education Malaysia via Fundamental Research Grant Scheme (FRGS/1/2017/SKK08/UM/02/11) and University of Malaya High Impact Research Grant (UM.C/625/1/HIR/MOE/CHAN/02/07).

The original research paper in which this protocol was described and validated is Tan et al. (2021).

Competing interests

The authors declare no competing interests.

References

- Bustin S. A. and Nolan, T. (2004). [Pitfalls of quantitative real-time reverse-transcription polymerase chain reaction](#). *J. Biomol. Tech.* 15(3): 155–166.
- Cai, Z., Fan, Y., Zhang, Z., Lu, C., Zhu, Z., Jiang, T., Shan, T. and Peng, Y. (2020). [VirusCircBase: a database of virus circular RNAs](#). *Briefings Bioinf.* 22(2): 2182–2190.
- Gagnon, K. T., Li, L., Janowski, B. A. and Corey, D. R. (2014). [Analysis of nuclear RNA interference in human cells by subcellular fractionation and Argonaute loading](#). *Nat. Protoc.* 9(9): 2045–2060.
- Jeck, W. R. and Sharpless, N. E. (2014). [Detecting and characterizing circular RNAs](#). *Nat. Biotechnol.* 32(5): 453–461.

- Szabo, L. and Salzman, J. (2016). [Detecting circular RNAs: bioinformatic and experimental challenges](#). *Nat. Rev. Genet.* 17(11): 679–692.
- Tan KE, Ng WL, Marinov GK, Yu KH, Tan LP, Liau ES, Goh SY, Yeo KS, Yip KY, Lo KW et al. (2021). Identification and characterization of a novel Epstein-Barr Virus-encoded circular RNA from LMP-2 Gene. *Sci Rep.* 2021 Jul 13;11(1):14392.
- Zhang, Y., Yang, L. and Chen, L. L. (2016). [Characterization of Circular RNAs](#). In: Feng, Y. and Zhang, L. (Eds.). *Long Non-Coding RNAs* (pp. 215–227). Methods in Molecular Biology. Humana Press, New York.