

MEETING ABSTRACT

Open Access

Use of platelet gel after surgery for wound breakdown prevention in recurrent vulvar cancer: case report

M Morelli^{1*}, A Dominijanni², S Del Negro¹, A Brescia², F Cariati¹, P Puzzonio², F Zullo¹

From de Senectute: Age and Health Forum
Catanzaro, Italy. 5-7 December 2009

Background

Vulvar carcinomas comprise almost 5% of all malignant tumors of the female genital tract [1]. Primary therapeutic approach is considered a radical surgery [2]. Infection and wound breakdown are the most common complications, these occur in approximately 40-60% of patients [3]. Objective to value the efficacy of platelet gel as support for reconstructive surgery in recurrent vulvar cancer.

Materials and methods

Five old caucasian women with a history of vulvar cancer, referred to our Department, were treated for recurrent vulvar cancer. Before the vulvar surgery reconstruction, an application of platelet gel was performed.

Results

Our data about wound healing, compared to results of previous cases of vulvar recurrence treated by our department, immediately show a reduction of complications, early healing and patient mobilization with less hospital stay, minimizing the costs. Other advantages of this new technique are the facility of application, no influence on operative time and low cost.

Conclusions

In consideration of the most frequent complications that are reported after surgical treatment of vulvar recurrence, as difficult and longer healing due to frequent necrosis and breakdown of wounds, we had experimented with the use of platelet gel [3]. In fact, thanks to its properties of regenerating and repairing tissues, platelet

gel is being used ever more frequently in reconstructive surgery [4]. The biological properties of platelet gel are basically due to the high concentrations of growth factors (GF) [4].

Author details

¹Department of Obstetrics and Gynecology, University 'Magna Graecia', Catanzaro, Italy. ²Servizio di Immunoematologia e Medicina Trasfusionale, Hospital 'Pugliese-Ciaccio', Catanzaro, Italy.

Published: 19 May 2010

References

1. Durdević S, Hadzić B, Petrović D: Radical vulvectomy with inguino-femoral lymphadenectomy in the surgical treatment of vulvar carcinoma. *Med Pregl* 2000, **53**:607-12.
2. Hacker NF: Current management of early vulvar cancer. *Ann Acad Med Singapore* 1998, **27**:688-92.
3. Gaarenstroom KN, Kenter GG, Trimbos JB, Agous I, Amant F, Peters AA, Vergote I: Postoperative complication after vulvectomy and inguinofemoral lymphadenectomy using separate groin incisions. *Int J Gynecol cancer* 2003, **13**:522-7.
4. van Lindert AC, Symons EA, Damen BF, Heintz AP: Wound healing after radical vulvectomy and inguino-femoral lymphadenectomy experience with granulocyte colony stimulating factor (filgrastim, r-metHuG-CSF). *Eur J Obstet Gynecol Reprod Biol* 1995, **62**:217-9.

doi:10.1186/1471-2318-10-S1-A67

Cite this article as: Morelli et al.: Use of platelet gel after surgery for wound breakdown prevention in recurrent vulvar cancer: case report. *BMC Geriatrics* 2010 **10**(Suppl 1):A67.

¹Department of Obstetrics and Gynecology, University 'Magna Graecia', Catanzaro, Italy