



Published in final edited form as:

*Oncogene*. 2014 February 27; 33(9): 1135–1147. doi:10.1038/onc.2013.49.

## Aberrant IKK $\alpha$ and IKK $\beta$ cooperatively activate NF- $\kappa$ B and induce EGFR/AP1 signaling to promote survival and migration of head and neck cancer

Liesl K. Nottingham<sup>1,2</sup>, Carol H. Yan<sup>1,2,3</sup>, Xiping Yang<sup>1</sup>, Han Si<sup>1</sup>, Jamie Coupar<sup>1</sup>, Yansong Bian<sup>1</sup>, Tsu-Fan Cheng<sup>1</sup>, Clint Allen<sup>1,4</sup>, Pattatheyl Arun<sup>1</sup>, David Gius<sup>1,5</sup>, Lenny Dang<sup>1,6</sup>, Carter Van Waes<sup>1,7</sup>, and Zhong Chen<sup>1,7</sup>

<sup>1</sup>Tumor Biology Section, Head and Neck Surgery Branch, National Institute on Deafness and Other Communication Disorders, NIH, Bethesda, MD, 20892, USA

<sup>3</sup>Howard Hughes Medical Institute-NIH Research Scholars Program

<sup>4</sup>Clinical Research Training Program supported jointly by NIH and Pfizer Inc

<sup>5</sup>Department of Radiation Oncology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

### Abstract

The Inhibitor- $\kappa$ B Kinase-Nuclear Factor- $\kappa$ B (IKK-NF- $\kappa$ B) and Epidermal Growth Factor Receptor-Activator Protein-1 (EGFR-AP-1) pathways are often co-activated and promote malignant behavior, but the underlying basis for this relationship is unclear. Resistance to inhibitors of IKK $\beta$  or EGFR is observed in head and neck squamous cell carcinomas (HNSCC). Here, we reveal that both IKK $\alpha$  and  $\beta$  contribute to nuclear activation of canonical and alternate NF- $\kappa$ B/REL family transcription factors, and overexpression of signal components enhancing co-activation of the EGFR-AP1 pathway. We observed that IKK $\alpha$  and IKK $\beta$  exhibit increased protein expression, nuclear localization and phosphorylation in HNSCC tissues and cell lines. Individually, IKK activity varied amongst different cell lines, but overexpression of both IKKs induced the strongest NF- $\kappa$ B activation. Conversely, siRNA knockdown of both IKKs significantly decreased nuclear localization and phosphorylation of canonical RELA and I $\kappa$ B $\alpha$ , and alternative p52 and RELB subunits. Knockdown of both IKKs more effectively inhibited NF- $\kappa$ B activation, broadly modulated gene expression, and suppressed cell proliferation and migration. Global expression profiling revealed that NF- $\kappa$ B, cytokine, inflammatory response, and growth factor signaling are among the top pathways and networks regulated by IKKs. Importantly,

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: [http://www.nature.com/authors/editorial\\_policies/license.html#terms](http://www.nature.com/authors/editorial_policies/license.html#terms)

Correspondence: Zhong Chen, MD, PhD; Phone: 301-435-2073; Fax: 301-594-4643; [chenz@nidcd.nih.gov](mailto:chenz@nidcd.nih.gov) (ZC); Head and Neck Surgery Branch, NIDCD/NIH, 10/5D55, MSC-1419, Bethesda, MD 20892-1419. Or Carter Van Waes, MD, PhD; Phone: 301-402-4216; Fax: 301-402-1140; [vanwaesc@nidcd.nih.gov](mailto:vanwaesc@nidcd.nih.gov); Head and Neck Surgery Branch, NIDCD/NIH, 10/5D55, MSC-1419, Bethesda, MD 20892-1419.

<sup>2</sup>Contribute equally as the first author.

<sup>6</sup>Current affiliation: Agios Pharmaceuticals, Cambridge, MA.

<sup>7</sup>Contributed equally as senior authors.

### Conflicts of interests

Lenny Dang was an employee at Millennium Pharmaceuticals, Inc. Other authors have no conflict of interest.

IKK $\alpha$  and IKK $\beta$  together promoted the expression and activity of TGF $\alpha$ , EGFR, and AP1 transcription factors cJun, JunB, and Fra1. Knockdown of AP1 subunits individually decreased 8/15 (53%) of IKK-targeted genes sampled, and similarly inhibited cell proliferation and migration. Mutations of NF- $\kappa$ B and AP1 binding sites abolished or decreased IKK-induced *IL-8* promoter activity. Compounds such as wedelactone with dual IKK inhibitory activity, and geldanamycins that block IKK $\alpha/\beta$  and EGFR pathways were more active than IKK $\beta$ -specific inhibitors in suppressing NF- $\kappa$ B activation and proliferation, and inducing cell death. We conclude that IKK $\alpha$  and IKK $\beta$  cooperatively activate NF- $\kappa$ B and EGFR/AP1 networks of signaling pathways, and contribute to the malignant phenotype and the intrinsic or acquired therapeutic resistance of HNSCC.

## Keywords

IKK $\alpha$ ; IKK $\beta$ ; NF- $\kappa$ B; AP1; head and neck cancers

## Introduction

The inhibitor-kappaB kinases (IKKs) mediate signal phosphorylation-induced degradation of inhibitor kappaBs (I $\kappa$ Bs), a critical step in releasing the NF-kappaB (NF- $\kappa$ B) family of transcription factors for cytoplasmic-nuclear translocation, DNA binding, and target gene transactivation (1, 2). The activation of the canonical NF- $\kappa$ B pathway is triggered by pro-inflammatory signals through a heterotrimeric complex of the IKK $\alpha$  and IKK $\beta$  catalytic subunits and the non-catalytic subunit, IKK $\gamma$ . This complex can phosphorylate and degrade I $\kappa$ Bs and promote nuclear translocation and phosphorylation of RELA/NF- $\kappa$ B1 (p65/p50) (1, 2). An alternative pathway, regulated by IKK $\alpha$  homodimers, induces processing of the precursor p100 to NF- $\kappa$ B2 (p52) to translocate with RELB (1, 2).

Aberrant activation of IKKs, NF- $\kappa$ B subunits, and their regulated gene programs have been implicated in the pathogenesis of many cancers, including head and neck squamous cell carcinomas (HNSCC) (3, 4). Blocking IKK-mediated I $\kappa$ B $\alpha$  degradation and NF- $\kappa$ B activation has been shown to inhibit aberrant gene expression, malignant phenotypes, and therapeutic resistance in pre-clinical HNSCC models (5, 6). Earlier experiments suggested that the IKK $\beta$  subunit was essential for canonical NF- $\kappa$ B activation through mediation of I $\kappa$ B degradation (1, 2), while IKK $\alpha$  was dispensable for I $\kappa$ B phosphorylation (7). The importance of IKK $\beta$  and proteasome in the canonical NF- $\kappa$ B pathway lead to drug development of bortezomib, a proteasome inhibitor that blocks I $\kappa$ B degradation and RELA activation, and MLN-120b, an inhibitor of IKK $\beta$  kinase activity (8, 9). However clinical trials using bortezomib in HNSCC showed limited clinical efficacy, due to incomplete targeting of other NF- $\kappa$ B/REL subunits, and other signaling pathways such as EGFR and AP1, which were shown to contribute to the resistance to bortezomib and radiation in HNSCC (10–12). Similarly, our preliminary studies showed that IKK $\beta$  inhibitors had modest effects *in vitro* and in pre-clinical HNSCC xenograft models (J. Ricker and J. Friedman, unpublished observations). Together, these findings suggested that drugs targeting IKK $\beta$ -mediated activation of NF- $\kappa$ B alone are insufficient, and the hypothesis that IKK $\alpha$  may contribute to canonical and/or alternative NF- $\kappa$ B/REL activation, and promotion

of the malignant phenotype. The few prior studies of IKK $\alpha$  in SCC have emphasized its potential role as a tumor suppressor, reporting increased malignant phenotype with decrease in its expression in poorly differentiated human SCC and in IKK $\alpha$  KO mice (13, 14). However, we have observed that increased IKK $\alpha$  with IKK $\beta$  are prevalent in the majority of differentiated HNSCC. How both IKKs together contribute to activation of the NF- $\kappa$ B pathway(s) and promoting the malignant phenotype in these cancers, remain incompletely understood.

Concurrent with NF- $\kappa$ B activation, the EGFR/AP1 signaling pathway is another important contributor of HNSCC pathogenesis. EGFR and its ligand TGF $\alpha$  are overexpressed in more than 90% HNSCC (15–17). The *EGFR* and *TGF $\alpha$*  genes may be amplified, but are more often overexpressed at the transcriptional level, by unknown mechanism(s) in HNSCC (18). Together, EGFR and TGF $\alpha$  form an autocrine signaling loop with the MAPK/ERK pathway and activate AP1 transcription factors to promote malignancy (15, 19). The AP1 transcription factor family of Jun and Fos contains 7 members: cJun, JunB, JunD, cFos, FosB, FosL(Fra1) and FosL2(Fra2) (19). We previously observed that co-activation of AP1 in HNSCC, either constitutively or induced through upstream signaling by TNF- $\alpha$ , EGFR, and MAPK/ERK, or tobacco carcinogens, is mediated through cJun, JunB and Fra1 (15, 16, 20). Nuclear co-activation of NF- $\kappa$ B/RELS, and overexpression of EGFR-AP1, was also subsequently linked in oral premalignant lesions and HNSCC tumor tissues by immunostaining (21), but the basis for this association remains undefined. Evidence for cross-talk and co-activation between NF- $\kappa$ B and AP1 signaling has been observed in skin, breast, and other solid tumors (22–24). To date, the role of IKK $\alpha$  and  $\beta$  in mechanisms of activation of the classical and alternative NF- $\kappa$ B pathways, and EGFR/AP1 signaling in human HNSCC remain unclear.

In this study, we observed that both IKK $\alpha$  and IKK $\beta$  are overexpressed and activated in most HNSCC tumors and cell lines. Cooperatively, they activated NF- $\kappa$ B/REL family members, cross-regulated the expression and activities of EGFR/AP1, and promoted proliferation and migration of HNSCC. Consistent with defined genetic studies, dual chemical inhibitors of IKKs, or IKKs and EGFR/AP1 more effectively inhibited NF- $\kappa$ B activation, cellular proliferation, and survival. Together, our data suggest the critical roles of IKKs as co-regulators of both the canonical and alternative NF- $\kappa$ B pathways, and mediators of NF- $\kappa$ B and EGFR/AP1 signaling cross-talk that promote pathogenesis of HNSCC.

## Results

### Aberrant expression and phosphorylation of IKK $\alpha$ and IKK $\beta$ in HNSCC tumor specimens and cell lines

IKK $\alpha$  and IKK $\beta$  expression, phosphorylation and localization in HNSCC were examined by immunohistochemical staining (IHC) of frozen sections of 18 HNSCC and 5 matched mucosa samples (Fig. 1A). Increased staining of IKK $\alpha$ , IKK $\beta$  and phospho-IKK $\alpha/\beta$  was observed throughout malignant epithelia compared to mucosa, which exhibited limited IKK staining in basilar layers (Fig. 1A). The staining intensity and localization of IKKs in tumors were quantified by histoscore. Over 90% exhibited predominantly nuclear IKK $\alpha$  staining, IKK $\beta$  was detected in both nucleus and cytoplasm, and phosphorylated-IKK staining was

more intense in the cytoplasm (Fig. 1B). The staining specificity of the antibody for p-IKK $\alpha$ / $\beta$  was confirmed by immunofluorescence staining after knockdown of IKK $\alpha$ , IKK $\beta$  or both, in HNSCC line UMSCC1 (data not shown). Furthermore, IKK $\alpha$  and IKK $\beta$  protein expression and phosphorylation was increased in the whole cell and nuclear extracts of 9 HNSCC lines (UM-SCC) derived from patients with aggressive clinical courses [survival 15 months,](25), compared to primary human keratinocytes (HeKa, Fig. 1C). Thus, these HNSCC cell lines exhibited both IKK $\alpha$  and  $\beta$  expression and p-IKK $\alpha$ / $\beta$ , consistent with most HNSCC specimens.

### **Overexpression and activation of IKK $\alpha$ or IKK $\beta$ enhance I $\kappa$ B degradation and NF- $\kappa$ B signaling in HNSCC cell lines**

To study the functional roles of IKKs, we overexpressed wild type (WT) or genetic mutant constructs of IKKs in UM-SCC lines. IKK $\alpha$  and IKK $\beta$  can be phosphorylated at the Serine phospho-acceptor sites, S176/S180 or S177/181, which were substituted with Alanines (SS->AA) to prevent IKK activation, or with Glutamate (SS->EE) to produce constitutively active IKKs. Catalytically inactive kinases (KA mutants) were produced by Lysine44 substitution with Alanine (K44A, Fig. 2A, upper panels) (26, 27). Transfection of all IKK expression vectors increased corresponding IKK protein expression (Fig. 2A, lower panels). Overexpression of IKK $\alpha$  or  $\beta$  EE increased kinase activity and decreased I $\kappa$ B $\alpha$  luciferase fusion protein reporter by >90% (Fig. 2B). Conversely, overexpression of kinase-inactive KA plasmids increased I $\kappa$ B $\alpha$ , with IKK $\beta$  KA mutant exhibiting a stronger dominant negative effect (Fig. 2B). A higher basal level of NF- $\kappa$ B activity was observed in UM-SCC11B compared to UM-SCC1. IKK WT and IKK EE individually increased NF- $\kappa$ B activation, with IKK $\beta$  having a greater effect in UM-SCC1 and IKK $\alpha$  in UM-SCC11B (Fig 2C, D). Co-expression of IKK $\alpha$  and IKK $\beta$  WT or EE resulted in the highest NF- $\kappa$ B activation in UM-SCC1, but not in UM-SCC11B (Fig. 2C, D). In both cell lines, IKK $\alpha$  AA had a greater effect in attenuation of NF- $\kappa$ B activity, whereas IKK $\beta$  KA inhibited NF- $\kappa$ B more than IKK $\alpha$  KA (Fig. 2C, D). Together, the data suggest differential but combinatory roles of IKK $\alpha$  and IKK $\beta$  in mediating I $\kappa$ B $\alpha$  degradation and NF- $\kappa$ B activation in different HNSCC lines.

### **Knockdown of IKK $\alpha$ or IKK $\beta$ alone or in combination by siRNA inhibits NF- $\kappa$ B subunit expression, I $\kappa$ B $\alpha$ degradation and NF- $\kappa$ B activation**

To define the roles of endogenous IKK $\alpha$  and IKK $\beta$  on NF- $\kappa$ B/REL activation, we knocked down both IKK subunits individually and in combination using siRNAs. We first examined the effects of IKK knockdown in UM-SCC1 cells treated with TNF- $\alpha$ , an inducer of canonical signaling which is detected in human HNSCC tumor specimens (28). The efficacy and specificity of the knockdowns were verified in cytoplasmic and nuclear fractions in UM-SCC1 (Fig. 3A). Canonical subunit RELA was modestly inhibited by IKK $\alpha$  or  $\beta$  knockdown, and strongly inhibited by dual IKK knockdown in both the cytoplasm and nucleus. A similar stepwise reduction from individual to dual IKK inhibition was observed for cytoplasmic phospho-RELA (p-RELA) at Serine536, while individual IKK knockdown enhanced nuclear p-RELA relative to basal or dual knockdown. The NF- $\kappa$ B1 subunit involved in the canonical pathway, p105 and its processed product p50, were also reduced after IKK $\alpha$  or  $\beta$  individual knockdown, while the strongest effects were observed after

knockdown of both. We detected expression of alternate NF- $\kappa$ B p100 protein predominantly localized in the cytoplasm, while processed p52 and RelB predominantly localized in the nuclear fraction (Fig. 3A). Interestingly, expression of the NF- $\kappa$ B subunit p100, and its processed product, p52, were suppressed more by IKK $\beta$  knockdown, while dual knockdown was the most effective in inhibiting the nuclear fraction. These protein expression data for p100/p52 can be largely explained by the effects of IKK siRNAs on *p100/p52* mRNA, as blocking IKK $\beta$  exhibited a stronger inhibitory effect on p100/p52 mRNA expression than knockdown of IKK $\alpha$ , while knockdown of both IKKs exhibited the strongest effects (Fig. 3B). Knockdown of IKKs did not significantly affect *RELA* gene expression (Fig. 3B). These data are consistent with previous observations that the canonical pathway is reported to regulate transcription of alternate pathway subunits NF- $\kappa$ B2/p100 precursor of p52 (29, 30). In addition, IKK $\alpha$  knockdown preferentially inhibited nuclear RelB along with p-p52 at Ser222 (Fig. 3A), a site implicated in p52 dimer and heterodimer formation recently identified by Perkins and colleagues (31, 32). Further, under TNF $\alpha$  stimulation, we did not detect significant induction of p-p100 (ser866/870), a marker for alternate IKK $\alpha$  pathway signal mediated processing, in the cytoplasm or nucleus (data not shown). p-I $\kappa$ B $\alpha$ , the downstream target of canonical IKK activation, was more strongly inhibited by IKK $\beta$ -specific or combinatory knockdowns in the cytoplasmic or nuclear fraction. Together, the greater effects of dual inhibition upon the cytoplasmic-nuclear translocation and phosphorylation of NF- $\kappa$ B subunits and I $\kappa$ B $\alpha$  are consistent with predominantly canonical IKK $\alpha$ / $\beta$ -dependent activation of these subunits in response to inflammatory signaling by TNF- $\alpha$ .

We next examined the functional effects of IKK knockdown under Lymphotoxin  $\beta$  (LT $\beta$ ) stimulation (Fig. 3C and D), which is a major upstream signal for alternative NF- $\kappa$ B pathway activation, and detected in lymph nodes to which HNSCC metastasize (33). LT $\beta$  induced increased p-p100 in both cytoplasm and nucleus in UM-SCC1. As a consequence, LT $\beta$  increased p52 in the nucleus. Interestingly, knockdown of IKK $\beta$  decreased p-p100 and p52 more significantly than knockdown of IKK $\alpha$ , while knockdown of both IKKs blocked p-p100 induced with LT $\beta$  treatment in both cytoplasm and nucleus. These observations are consistent with above findings, where knockdown of *IKK $\beta$*  exhibited a stronger inhibitory effect on p100/p52 mRNA and total protein expression than knockdown of *IKK $\alpha$* , and knockdown of both IKKs exhibited the strongest effects. Together, our results provide evidence that in HNSCC, the upstream signaling from both IKKs regulate downstream NF- $\kappa$ B subunits. The effect of IKK $\beta$  on the p100/p52 is predominantly through its regulatory function controlling *p100/p52* gene expression and processing, and is not solely dependent on IKK $\alpha$ -mediated phosphorylation of p-100, consistent with previous findings in other cell types (34–36).

To assay the functional effect of IKK knockdowns on I $\kappa$ B and NF- $\kappa$ B activity, we used I $\kappa$ B $\alpha$ -Luciferase fusion protein (Fig. 3E) and NF- $\kappa$ B reporter constructs (Fig. 3F). Dual IKK knockdown had a greater effect than individual knockdowns in different HNSCC cell lines UM-SCC1 (Fig. 3 E, F), UM-SCC11B (Fig. 3E) and UM-SCC6 (data not shown). Together, these data indicate that both IKK $\alpha$  and IKK $\beta$  individually and together contribute to I $\kappa$ B $\alpha$  degradation and NF- $\kappa$ B reporter activation. We examined the relative contribution of

canonical and alternate NF- $\kappa$ B/REL subunits on NF- $\kappa$ B reporter activity, which demonstrated significant effects of p65, p105/p50 and RELB knockdown (Nottingham L, unpublished observations).

### Dual knockdown of IKKs suppress a broader gene expression program involving NF- $\kappa$ B, cytokines and growth factor signaling pathways

Dual knockdown of IKKs showed significant down-regulation of genes compared to individual IKK knockdown at 48 hours using ~23K human gene profiling (Fig. 4A). Representative down-regulated genes are shown in Supplemental Fig. 1. IPA annotation indicates the genes enriched in cytokine-, immune response-, and NF- $\kappa$ B related pathways (top), and growth factor signaling pathways (bottom, Fig. 4B, Supplemental Table 2). The interactive networks, established using gene components in cytokines and NF- $\kappa$ B related pathways (Fig. 4C), or growth factor signaling pathways (Fig. 4D), reveal potential interactive relationships amongst different molecules and pathways.

Among the knockdown targets, we validated 20 genes by real-time RT-PCR, with the strongest effects were observed by dual IKK knockdown (Fig. 5A). These include known canonical NF- $\kappa$ B modulated genes such as *I $\kappa$ B $\alpha$* , *IL-1* and *IL-8*, and the non-canonical *NF- $\kappa$ B2/p52* subunit gene, established above. Several genes involved in the EGFR/AP1 pathways were suppressed, prompting knockdown experiments with *FOSL1*, *cJun*, or *JunB* (Fig 5B). Eight of fifteen (53%) of IKK target genes were co-modulated by AP1 family members, suggesting overlapped as well as distinct gene targets between IKKs and AP1 signaling pathways. Many of the IKK or AP1 target genes are implicated in cell proliferation and migration, according to Ingenuity Pathway Analysis annotations (Supplemental Table 2). Genes in the ERK/p38/MAPK and PI3K/AKT pathways involved in cell proliferation, include *TGF $\alpha$* , *HBEGF*, *EGFR*, *Jun*, *JunB*, *FOSL*, *EGR1*, *cyclinD1*, and *IGFBPs*. Genes involved in cell migration included those in the HGF and TGF $\beta$  signaling pathways, such as *EGR1*, *ETS*, *Jun*, *SERPINE1*, *MMPs*, *SMADs* and their family members. Together, these signaling pathways regulated by IKKs and AP1 family members have previously been implicated as individually important molecules in promoting the malignant phenotype of HNSCC and other cancers.

### Knockdown of IKKs and RELA inhibit EGFR/AP1 expression as well as AP1 and IL-8 promoter activities

To further investigate the specific effects of IKKs and NF- $\kappa$ B signaling on the EGFR/AP1 pathway, we examined the effects of IKK knockdowns on EGFR/AP1 protein expression by Western blot (Fig. 6A). Individual and dual IKK knockdown, affected expression of EGFR, cJun, JunB and Fra-1/FOSL proteins in whole cell lysates, consistent with that observed for mRNA expression shown in Fig. 5A. To further investigate the role of canonical IKK mediated regulation of transcription factor RELA in EGFR/AP1 gene regulation, we knocked down RELA by siRNA. A significant decrease in EGFR, JunB and FOSL(Fra1) expression were observed (Fig. 6B), establishing an important role for RELA in co-expression of these EGFR/AP-1 pathway components. Consistent with the above observations, knockdown of IKKs suppressed AP1 reporter activity at 48 and 72 hours (Fig. 6C and data not shown). Further, wild-type IKK-induced proximal IL-8 promoter activity

(-133bp from the transcription starting site) was completely abolished by a point mutation at the NF- $\kappa$ B site (-83bp), and partially blocked by a point mutation at the AP1 site (-128bp, Fig. 6D).

### **Knockdown of IKKs and target AP1 genes inhibits cell proliferation and migration**

The biological roles of IKK $\alpha$  and IKK $\beta$  on proliferation and migration were examined, and demonstrated that dual knockdown of IKKs more significantly inhibited cellular proliferation (Fig. 7A). Knockdown of IKK $\beta$  exhibited a greater inhibition of cellular migration than IKK $\alpha$ , while the dual knockdown demonstrated the strongest effect (Fig. 7B and C). We examined the effect of knockdown of IKK target AP1 subunits on cell proliferation and migration. Knockdown of FOSL and JunB individually, but not cJun, significantly inhibited cell proliferation (Fig. 7D; data not shown). Knockdown of all AP1 subunits individually suppressed cell migration, with cJun exhibiting the strongest effects (Figure 7E and F). The data suggest that knockdown of both IKKs and target AP1 subunits differentially affect cell growth and migration.

### **Drugs targeting IKK $\beta$ , dual IKK, and IKK/EGFR/AP1 components differentially inhibit NF- $\kappa$ B activity and cell survival**

To explore the therapeutic implications of the above findings, we examined the effects of small molecules available for preclinical or clinical studies capable of inhibiting IKK $\beta$  alone (MLN120b, SC-514) (37), both IKKs (Wedelactone), (38), or IKK $\alpha$  and EGFR/AP1 signaling (Hsp90 inhibitor and geldanamycin derivatives, 17-AAG, 17-DMAG) (39). We found that 17-AAG preferentially inhibited IKK $\alpha$  expression in UM-SCC1, and reduced a slower migrating IKK $\beta$  band in both cell lines (Fig. 8A). In contrast, MLN-120b most selectively inhibited IKK $\beta$  in UM-SCC11B cells. Pan-IKK inhibitor Wedelolactone attenuated IKK $\alpha$  expression in both cell lines, and the slower migrating band of IKK $\beta$  and phospho-p65 in UM-SCC11B cells (Fig. 8A). Hsp90 inhibitor, 17-DMAG, more potently inhibited the NF- $\kappa$ B reporter than MLN-120b in UM-SCC1 cells, while UM-SCC11B cells were sensitive to both drugs (Fig. 8B). 17-DMAG and Wedelolactone more potently inhibited cell proliferation in both cell lines, compared to IKK $\beta$  selective inhibitors, MLN-120b and SC-514 (Supplemental Fig. 2). Both 17-AAG and Wedelolactone are also strong inducers for cell death as demonstrated by the increased sub-G0 DNA fragmentation and cell cycle arrest at the G2M phase, decreased cell density and increased cell blebbing and detachment (Fig. 8C, and Supplemental Fig. 3A, B). MLN-120b and SC-514 showed cytostatic effects and an increased vacuolization (Supplemental Fig. 3B). We confirmed that IKK $\alpha$  wt and EE mutants rescued NF- $\kappa$ B reporter activation in cells treated with IKK $\beta$  selective inhibitor MLN120b, when compared to broader inhibitors such as 17AAG (data not shown). Altogether, these data suggest that inhibitors with dual IKK $\alpha/\beta$  or broader activity exhibit more potent effects on IKK and NF- $\kappa$ B protein expression, NF- $\kappa$ B activation, and induction of cell cycle arrest and cell death, compared to selective-IKK $\beta$  inhibitors.

## Discussion

In this study, we provide evidence supporting a model whereby both IKK $\alpha$  and IKK $\beta$  cooperatively activate NF- $\kappa$ B, as well as enhance EGFR/AP1 signaling, to promote the malignant phenotype of HNSCC (Fig. 8D). IKK $\alpha$  and IKK $\beta$  were aberrantly overexpressed in most tumor tissues and cell lines, indicating their co-expression is prevalent in oncogenesis in differentiated HNSCC (Fig. 1). IKK $\alpha$  was previously implicated as a tumor suppressor of differentiation and IKK $\beta$ -mediated NF- $\kappa$ B and EGFR/AP-1 activation in a smaller subset of undifferentiated human HNSCC and knockout mice (13, 24, 40). However, when co-expressed, modulating IKK $\alpha$  and IKK $\beta$  in combination induced the strongest effects on NF- $\kappa$ B activity, although either were found to dominantly modulate NF- $\kappa$ B in different cell lines (Fig. 2). IKK $\alpha$  and IKK $\beta$  were both shown to mediate nuclear translocation and activation of canonical NF- $\kappa$ B/REL subunits (Fig. 3A). While IKK $\alpha$  predominately affected nuclear RELB and p-p52 as expected, IKK $\beta$  predominantly modulated overall expression of p100/p52 mRNA and protein available for nuclear activation (Fig. 3A, B). Pathway and network analyses indicated a crosstalk between IKK/NF- $\kappa$ B and the EGFR/AP1 signaling pathways (Fig. 4), which was shown to be modulated through the expression and functions of NF- $\kappa$ B targets and EGFR/AP1 molecules (Fig 5–7). Knockdown of AP1 subunits affected an overlapping subset of IKK target genes, and inhibited cell proliferation and migration (Fig. 5 and 7). Dual targeting of IKKs either by siRNA or small molecule inhibitors with broader activities, showed greater inhibitory effects on cell proliferation, migration, cell cycle arrest, and apoptosis (Fig. 7 and 8). Our data strongly suggest that both IKK $\alpha/\beta$  are important modulators of signaling upstream of NF- $\kappa$ B activation, and cross-talk promoting co-activation of the EGFR/AP1 pathway, important in pathogenesis and therapeutic targeting of HNSCC.

Previously, the IKK $\beta$  subunit of the canonical IKK complex was believed to be the predominant kinase responsible for I $\kappa$ B-dependent NF- $\kappa$ B activation (7), and promotion of tumorigenesis and metastasis. By utilizing IKK genetic mutant IKK constructs, we demonstrate that the specific impacts and variable potencies of IKK $\alpha$  and IKK $\beta$  activity differ, but cooperate in NF- $\kappa$ B activation in HNSCC. Consistent with the previous paradigm, overexpressing kinase dead mutant IKK $\beta$  KA had a relatively greater effect compared with IKK $\alpha$  KA in blocking I $\kappa$ B $\alpha$  degradation or NF- $\kappa$ B activation, but constitutively active IKK $\alpha$  or  $\beta$  EE mutants both promoted I $\kappa$ B degradation and NF- $\kappa$ B activation (Fig. 2B–D), indicating IKK $\alpha$  potentiates activation. The activating effects of overexpression of IKK $\alpha$  WT and EE mutants, or inhibitory effects of IKK $\alpha$  AA mutants, together further support the role of signal activated IKK $\alpha$  in potentiating IKK $\beta$  activation of NF- $\kappa$ B. These results are consistent with previous reports that IKK $\alpha$  can enhance IKK $\beta$  activity, though IKK $\beta$  is the stronger, direct kinase for downstream canonical NF- $\kappa$ B activation (1, 26). Interestingly, UMSSC11B demonstrated a stronger effect of IKK $\alpha$  WT or EE overexpression (Fig. 2C, D), but greater sensitivity to IKK $\beta$  inhibitor MLN-120b (Fig. 8A and B), consistent with its constitutively predominant IKK $\beta$  activation.

Further results modulating IKK $\alpha$ , IKK $\beta$ , or both by siRNA support the hypothesis that endogenous IKK $\alpha$  and IKK $\beta$  cooperatively regulate nuclear canonical and alternative NF- $\kappa$ B/REL subunits and functional NF- $\kappa$ B reporter activation. As expected, either IKK $\alpha$  or  $\beta$

partially attenuated canonical nuclear RELA and NF- $\kappa$ B reporter activity, while dual inhibition had the most potent effects (Fig. 3A, E, F). TNF- $\alpha$  and LT $\beta$  are often expressed by infiltrating cells in HNSCC and lymph nodes, where HNSCC preferentially metastasize (28, 41). Interestingly, while overall nuclear RELA was partly inhibited by knockdown of either IKK under TNF- $\alpha$  treatment, IKK-dependent phosphorylation of nuclear RELA ser536 was enhanced unless both were knocked down, a phenomenon observed when IKK $\alpha$  or  $\beta$  compensate upon knockout of either kinase in mice and human cell lines (40, 42). Overall and nuclear p100 and p52 were also inhibited with IKK $\beta$  siRNA (Fig. 3A), which is consistent with prior findings that IKK $\beta$  strongly regulates TNF- $\alpha$ -induced p100/p52 gene expression (Fig. 3B) (29, 43). LT $\beta$ , as an alternate pathway inducer, was shown to induce detectable p-p100 and nuclear p52, but IKK $\beta$  and dual knockdown still attenuated p-p100 and nuclear p52 protein (Fig. 3C, D). Recent observations suggest that expression, processing and nuclear localization of p100/p52 may be more dependent on canonical IKK activation and I $\kappa$ B $\alpha$  degradation than previously appreciated (34–36, 43–45). With canonical signal activation, Scheidereit and colleagues found that p105 and p100 interact and are co-processed to p50 and p52, and that canonically regulated I $\kappa$ B can co-modulate nuclear translocation of RELB, potentially modifying previous paradigms separating canonical and alternative pathways (33). Consistent with this possibility, our data show similar effects of IKK knockdowns on p105/p50 and p100/p52 (Fig 3A). Together, our results indicate that dual inhibition of IKK $\alpha$  and  $\beta$  has greatest effects on expression, processing, nuclear localization, and functional activation of canonical and alternate NF- $\kappa$ B/RELBs in HNSCC.

Although our data clearly demonstrate that both IKK $\alpha$  and IKK $\beta$  co-modulate the NF- $\kappa$ B pathway and expression of many genes, IKK $\alpha$  contains a nuclear localization signal, and is capable of modulating gene expression through histone H3 modification, important for NF- $\kappa$ B dependent, and independent gene expression, such as Fos after stimulation with EGF (46). We found that IKK $\alpha$  was localized in both nucleus and cytoplasm of human HNSCC, and IKK $\alpha$ ,  $\beta$  and p65 co-activated EGFR and AP-1 subunit gene expression (Fig. 5). This is in apparent contrast with non-malignant epidermis of IKK $\alpha$  knockout mice, where IKK $\alpha$  can repress an autocrine loop of EGFR, ERK, and EGFR ligands, potentially through NF- $\kappa$ B independent mechanisms (24). Further studies suggested that where IKK $\alpha$  is decreased, this may promote carcinogenesis primarily through loss of its dual function as a tumor suppressive component of an otherwise intact TGF $\beta$  pathway (14). However, for the majority of HNSCC tumors and cell lines, we have recently shown that TGF $\beta$  function is attenuated via decreased TGF $\beta$  receptor expression or downstream responsiveness, and IKK $\alpha$  is overexpressed together with enhancement of NF- $\kappa$ B, as well as EGFR expression and pathway activation (47–49).

Through global gene analysis, we identified a broad gene program regulated by IKKs individually or in combination (Fig. 4, 5, Supplemental Fig. 1). Interestingly, IKK modulated genes were enriched in several growth factor signaling pathways, including EGFR/AP1 signaling often co-activated and important in the malignant phenotype in HNSCC (Fig. 4B, lower panel, Fig. 5A). Knockdown of AP1 subunits revealed that AP1 co-modulates a major subset of the IKK regulated genes. Previously, IKK or NF- $\kappa$ B-modulated expression of AP1 family members was suggested by studies in LPS stimulated B cells,

monocytes, and dendritic cells (50–52), as well as in IKK $\alpha$  knockout MEFs (46, 53). However, the role and means by which IKK modulates the EGFR/AP1 pathway in HNSCC tumorigenesis has not previously been well elucidated. Our data in established HNSCC cell lines support a model, whereby activated IKKs and RELA can induce co-expression of TGF- $\alpha$ , EGFR and downstream Jun and FosL(Fra1) family members, and co-activation of AP1. Interestingly, the three AP1 subunits identified in this study are the same ones we had previously shown to be co-activated with NF- $\kappa$ B in HNSCC (20). In addition, our data showed that AP1 family members exhibited differential functions to modulate cellular functions, such as proliferation and migration (Fig 7D–F). Our data suggest that FosL(Fra1) cooperates with cJun or JunB to promote migration and inflammatory cytokine production (54, 55), extending findings beyond a previous study highlighting cJun's role in promoting cellular proliferation (19). Knockdown of IKK $\beta$  and cJun exhibited strong effects on cell migration (Fig. 7), consistent with knockdown of IKK $\beta$  affecting cJun mRNA and protein expression more significantly than IKK $\alpha$  (Fig. 5 and 6). Our data are also supported by recent evidence, that activated NF- $\kappa$ B signaling contributes to resistance of EGFR-targeted therapy (56). Together with our study, this suggests that concurrent suppression of both IKK/NF- $\kappa$ B and EGFR/AP1 signaling could enhance therapeutic targeting of HNSCC.

We found that small molecule inhibitors with broader activity against both IKKs (wedelactone), or IKKs and EGFR (Hsp90 inhibitors), had more potent anti-proliferative and cytotoxic activity than IKK $\beta$  specific inhibitors (MLN120b and SC-514; Fig. 7C, Supplemental Fig. 2 and 3), consistent with the combined targeting of IKK $\alpha$  and IKK $\beta$  with siRNA. Geldanamycin derivatives inhibit the heat shock protein 90kDa (HSP90), a molecular chaperone preferentially targeting IKK $\alpha$  degradation over IKK $\beta$ , and inducing apoptosis in different cancers (39). Although the chemical drugs, 17-AAG/17-DMAG have broader ranges of targets compared to specific siRNA targeting, inhibition of IKK $\alpha$  activity was confirmed by the significant suppression of NF- $\kappa$ B activation induced by transfection with WT and EE IKK $\alpha$  mutants, while IKK $\alpha$  rescued NF- $\kappa$ B inhibitory effects of IKK $\beta$  selective inhibitor MLN120 (data not shown). Also, 17AAG significantly decreased IKK $\alpha$  protein levels (Fig 8A), consistent with the drug's known ability to destabilize IKK $\alpha$  and increase its degradation rate. Wedelolactone also inhibits both IKK $\alpha$  and IKK $\beta$  through modulation of caspases and I $\kappa$ B phosphorylation (38). In contrast, IKK $\beta$  specific inhibitors MLN-120b and SC514 exhibited limited efficacy when evaluated in preclinical studies of multiple cancers (37, 57), consistent with our observations (Fig. 8C, Supplemental Fig. 2 and 3, and data not shown). Our data have implications for dual IKK $\alpha$  and  $\beta$  inhibition in chemo- or radiation sensitization. IKK knockdown may sensitize cancer cells to chemotherapy-induced apoptosis (58), while Hsp90 inhibitors analogous to 17-AAG/17-DMAG increase radiation sensitivity both *in vitro* and *in vivo* in HNSCC (manuscript in preparation). Our current study challenges the conventional idea of focusing heavily on IKK $\beta$ -mediated signaling for development of anti-tumor drugs, which have not lead to expected efficacies in preclinical studies. The current inhibitors targeting NF- $\kappa$ B signaling are based on previously observed roles of IKK $\beta$  and the classical NF- $\kappa$ B pathway in transgenic or knock-out mouse models (7). However, the lack of efficacy of these inhibitors in solid epithelial tumors suggest that blockade of IKK $\beta$  maybe be insufficient. Our data suggest the underlying reasons for the lack of therapeutic efficacy, and support our

hypothesis that blocking both classical NF- $\kappa$ B pathway mediated by IKK $\alpha/\beta$  and the alternative pathway mediated by IKK $\alpha$  produces an additive and more complete inhibition of down stream genes and malignant phenotypes.

## Materials and methods

### Immunohistochemical analysis of HNSCC tissue specimens, Western blot, Reporter gene assay, MTT assay, Flow cytometry, Wound healing assay

See Supplemental Materials and Methods.

### Cell lines and culture

University of Michigan head and neck squamous cell carcinoma (UM-SCC) cell lines were from T.E. Carey (University of Michigan, Ann Arbor, MI), and cultured under standard growth conditions. Cell DNA was sent for sequence genotyping in 2008 and fall 2010 to compare and verify their unique origin from original stocks, as recently described (25). The 9 loci analyzed included D3S1358, D5S818, D7S820, D8S1179, D13S317, D18S51, D21S11, FGA, vWA and amelogenin. Human epidermal keratinocytes were obtained from (Invitrogen, Carlsbad, CA) and harvested at less than passage 5.

### siRNAs and plasmids

siRNAs were obtained from IDT (Coralville, IA), multiple siRNAs were individually tested, and three duplexes of siRNAs with best knockdown efficiency and specificity for each target were selected and pooled for the experiments (Supplemental Table 1). IKK expression and mutant constructs were kindly provided by Dr. Ulrich Siebenlist (NIH/NIAID, Bethesda, MD) and Dr. Michael Karin (UC San Diego), consisting of IKK $\alpha$  and IKK $\beta$  wild type (WT), phosphoacceptor-mutant (SS->AA, Serine to Alanine), constitutively activated (SS->EE, Serine to Glutamic Acid), and kinase dead (K44A, Lysine to Alanine) vectors (26, 27).

### Microarray and gene expression analysis

Microarray expression profiling was performed using Illumina Beadchip Array, and data were normalized and summarized in GenomeStudio software (Illumina). Genes with more than 50% decreased expression after IKK $\alpha$  and/or IKK $\beta$  siRNA knockdown were selected and compared to scramble siRNA treated cells, and analyzed in statistical software "R". To display the common and unique down-regulated genes in different treated samples, 3-way Venn diagram was made using VennDiagram package from Bioconductor (<http://www.bioconductor.org>). Ingenuity Pathway Analysis (IPA) was used to detect the significantly enriched signaling pathways within down-regulated genes, and two networks were established using genes involved in depicted pathways.

### Chemotherapeutic drugs

17-AAG (17-allylamino 17-demethoxygeldanamycin, tanespimycin) and 17-DMAG (17-Dimethylaminoethylamino-17-demethoxygeldanamycin, alvespimycin), exhibit very similar *in vitro* activity, but have different solubility and toxicity profiles *in vivo* (59). The drugs were synthesized by the Radiation Oncology Branch at NCI/NIH, and kindly provided by

Dr. David Guis. SC-514 is a selective and reversible inhibitor of I $\kappa$ B kinase 2 (IKK2) (IC<sub>50</sub> = 3–12  $\mu$ M) that displays greater than 10-fold selectivity over 28 other kinases (Cayman Chemical). Wedelolactone (a natural compound isolated from the herb *E. prostrata*) was obtained from Biomol (Plymouth Meeting, PA). MLN-120b was provided by Millennium Pharmaceuticals, Inc under a Material Transfer Agreement.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

This project is supported by NIDCD Intramural projects Z01-DC-00016, Z01-DC-00073 and Z01-DC-00074. The authors wish to thank Drs. Michael Karin (UC San Diego) and Ulrich Siebenlist (NIH/NIAID) for kindly providing the IKK plasmids; Dr. Neil Perkins (Newcastle University, UK) for kindly providing anti-p-p52 antibody; Drs. Michael Karin, Christophe Cataisson (NCI/NIH) and Claire Sauvageot (Millennium Pharmaceuticals, Inc) for critique of the manuscript.


## References

- Hacker H, Karin M. Regulation and function of IKK and IKK-related kinases. *Sci STKE*. 2006; 2006(357):re13. Epub 2006/10/19. [PubMed: 17047224]
- Scheidereit C. IkappaB kinase complexes: gateways to NF-kappaB activation and transcription. *Oncogene*. 2006; 25(51):6685–705. Epub 2006/10/31. [PubMed: 17072322]
- Karin M, Greten FR. NF-kappaB: linking inflammation and immunity to cancer development and progression. *Nat Rev Immunol*. 2005; 5(10):749–59. Epub 2005/09/22. [PubMed: 16175180]
- Van Waes C. Nuclear factor-kappaB in development, prevention, and therapy of cancer. *Clin Cancer Res*. 2007; 13(4):1076–82. Epub 2007/02/24. [PubMed: 17317814]
- Duffey DC, Chen Z, Dong G, Ondrey FG, Wolf JS, Brown K, et al. Expression of a dominant-negative mutant inhibitor-kappaB $\alpha$  of nuclear factor-kappaB in human head and neck squamous cell carcinoma inhibits survival, proinflammatory cytokine expression, and tumor growth in vivo. *Cancer Res*. 1999; 59(14):3468–74. Epub 1999/07/23. [PubMed: 10416612]
- Sunwoo JB, Chen Z, Dong G, Yeh N, Crowl Bancroft C, Sausville E, et al. Novel proteasome inhibitor PS-341 inhibits activation of nuclear factor-kappa B, cell survival, tumor growth, and angiogenesis in squamous cell carcinoma. *Clin Cancer Res*. 2001; 7(5):1419–28. Epub 2001/05/15. [PubMed: 11350913]
- Delhase M, Hayakawa M, Chen Y, Karin M. Positive and negative regulation of IkappaB kinase activity through IKKbeta subunit phosphorylation. *Science*. 1999; 284(5412):309–13. Epub 1999/04/09. [PubMed: 10195894]
- Adams J. The proteasome: a suitable antineoplastic target. *Nature reviews Cancer*. 2004; 4(5):349–60. Epub 2004/05/04. [PubMed: 15122206]
- Nagashima K, Sasseville VG, Wen D, Bielecki A, Yang H, Simpson C, et al. Rapid TNFR1-dependent lymphocyte depletion in vivo with a selective chemical inhibitor of IKKbeta. *Blood*. 2006; 107(11):4266–73. Epub 2006/01/28. [PubMed: 16439676]
- Allen C, Saigal K, Nottingham L, Arun P, Chen Z, Van Waes C. Bortezomib-induced apoptosis with limited clinical response is accompanied by inhibition of canonical but not alternative nuclear factor- $\kappa$ B subunits in head and neck cancer. *Clin Cancer Res*. 2008; 14(13):4175–85. Epub 2008/07/03. [PubMed: 18593997]
- Chen Z, Ricker JL, Malhotra PS, Nottingham L, Bagain L, Lee TL, et al. Differential bortezomib sensitivity in head and neck cancer lines corresponds to proteasome, nuclear factor-kappaB and activator protein-1 related mechanisms. *Mol Cancer Ther*. 2008; 7(7):1949–60. Epub 2008/07/23. [PubMed: 18645005]

12. Argiris A, Duffy A, Kummar S, Simone NL, Arai Y, Kim S, et al. Early tumor progression associated with enhanced EGFR signaling with bortezomib, cetuximab, and radiotherapy for head and neck cancer. *Clin Cancer Res*. 2011 Epub 2011/07/14.
13. Liu B, Park E, Zhu F, Bustos T, Liu J, Shen J, et al. A critical role for I kappaB kinase alpha in the development of human and mouse squamous cell carcinomas. *Proc Natl Acad Sci U S A*. 2006; 103(46):17202–7. Epub 2006/11/03. [PubMed: 17079494]
14. Marinari B, Moretti F, Botti E, Giustizieri ML, Descargues P, Giunta A, et al. The tumor suppressor activity of IKKalpha in stratified epithelia is exerted in part via the TGF-beta antiproliferative pathway. *Proc Natl Acad Sci U S A*. 2008; 105(44):17091–6. Epub 2008/10/30. [PubMed: 18957551]
15. Lee TL, Yeh J, Van Waes C, Chen Z. Epigenetic modification of SOCS-1 differentially regulates STAT3 activation in response to interleukin-6 receptor and epidermal growth factor receptor signaling through JAK and/or MEK in head and neck squamous cell carcinomas. *Mol Cancer Ther*. 2006; 5(1):8–19. Epub 2006/01/25. [PubMed: 16432158]
16. Pernas FG, Allen CT, Winters ME, Yan B, Friedman J, Dabir B, et al. Proteomic signatures of epidermal growth factor receptor and survival signal pathways correspond to gefitinib sensitivity in head and neck cancer. *Clin Cancer Res*. 2009; 15(7):2361–72. Epub 2009/03/26. [PubMed: 19318490]
17. Sharafinski ME, Ferris RL, Ferrone S, Grandis JR. Epidermal growth factor receptor targeted therapy of squamous cell carcinoma of the head and neck. *Head & neck*. 2010; 32(10):1412–21. Epub 2010/09/18. [PubMed: 20848399]
18. Grandis JR, Tweardy DJ. Elevated levels of transforming growth factor alpha and epidermal growth factor receptor messenger RNA are early markers of carcinogenesis in head and neck cancer. *Cancer Res*. 1993; 53(15):3579–84. Epub 1993/08/01. [PubMed: 8339264]
19. Shaulian E, Karin M. AP-1 as a regulator of cell life and death. *Nature cell biology*. 2002; 4(5):E131–6. Epub 2002/05/04. [PubMed: 11988758]
20. Ondrey FG, Dong G, Sunwoo J, Chen Z, Wolf JS, Crowl-Bancroft CV, et al. Constitutive activation of transcription factors NF-(kappa)B, AP-1, and NF-IL6 in human head and neck squamous cell carcinoma cell lines that express pro-inflammatory and pro-angiogenic cytokines. *Molecular carcinogenesis*. 1999; 26(2):119–29. Epub 1999/10/03. [PubMed: 10506755]
21. Mishra A, Bharti AC, Saluja D, Das BC. Transactivation and expression patterns of Jun and Fos/ AP-1 super-family proteins in human oral cancer. *Int J Cancer*. 2010; 126(4):819–29. Epub 2009/08/05. [PubMed: 19653276]
22. Park E, Liu B, Xia X, Zhu F, Jami WB, Hu Y. Role of IKKalpha in skin squamous cell carcinomas. *Future oncology*. 2011; 7(1):123–34. Epub 2010/12/23. [PubMed: 21174543]
23. Ndlovu MN, Van Lint C, Van Wesemael K, Callebert P, Chalbos D, Haegeman G, et al. Hyperactivated NF-(kappa)B and AP-1 transcription factors promote highly accessible chromatin and constitutive transcription across the interleukin-6 gene promoter in metastatic breast cancer cells. *Molecular and cellular biology*. 2009; 29(20):5488–504. Epub 2009/08/19. [PubMed: 19687301]
24. Liu B, Xia X, Zhu F, Park E, Carbajal S, Kiguchi K, et al. IKKalpha is required to maintain skin homeostasis and prevent skin cancer. *Cancer cell*. 2008; 14(3):212–25. Epub 2008/09/06. [PubMed: 18772111]
25. Yang X, Lu H, Yan B, Romano RA, Bian Y, Friedman J, et al. DeltaNp63 versatily regulates a Broad NF-kappaB gene program and promotes squamous epithelial proliferation, migration, and inflammation. *Cancer Res*. 2011; 71(10):3688–700. Epub 2011/05/18. [PubMed: 21576089]
26. Yamamoto Y, Yin MJ, Gaynor RB. IkappaB kinase alpha (IKKalpha) regulation of IKKbeta kinase activity. *Molecular and cellular biology*. 2000; 20(10):3655–66. Epub 2000/04/25. [PubMed: 10779355]
27. Zandi E, Rothwarf DM, Delhase M, Hayakawa M, Karin M. The IkappaB kinase complex (IKK) contains two kinase subunits, IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappaB activation. *Cell*. 1997; 91(2):243–52. Epub 1997/11/05. [PubMed: 9346241]


28. Younes F, Quartey EL, Kiguwa S, Partridge M. Expression of TNF and the 55-kDa TNF receptor in epidermis, oral mucosa, lichen planus and squamous cell carcinoma. *Oral Dis.* 1996; 2(1):25–31. Epub 1996/03/01. [PubMed: 8957934]
29. Madge LA, May MJ. Classical NF-kappaB activation negatively regulates noncanonical NF-kappaB-dependent CXCL12 expression. *J Biol Chem.* 2010; 285(49):38069–77. Epub 2010/10/07. [PubMed: 20923761]
30. Basak S, Hoffmann A. Crosstalk via the NF-kappaB signaling system. *Cytokine Growth Factor Rev.* 2008; 19(3–4):187–97. Epub 2008/06/03. [PubMed: 18515173]
31. Barre B, Perkins ND. The Skp2 promoter integrates signaling through the NF-kappaB, p53, and Akt/GSK3beta pathways to regulate autophagy and apoptosis. *Mol Cell.* 2010; 38(4):524–38. Epub 2010/06/02. [PubMed: 20513428]
32. Barre B, Perkins ND. Phosphorylation of the p52 NF-kappaB subunit. *Cell Cycle.* 2010; 9(24):4774–5. Epub 2010/12/15. [PubMed: 21150331]
33. Wolf MJ, Seleznik GM, Zeller N, Heikenwalder M. The unexpected role of lymphotoxin beta receptor signaling in carcinogenesis: from lymphoid tissue formation to liver and prostate cancer development. *Oncogene.* 2010; 29(36):5006–18. Epub 2010/07/07. [PubMed: 20603617]
34. Madge LA, Kluger MS, Orange JS, May MJ. Lymphotoxin-alpha 1 beta 2 and LIGHT induce classical and noncanonical NF-kappa B-dependent proinflammatory gene expression in vascular endothelial cells. *J Immunol.* 2008; 180(5):3467–77. Epub 2008/02/23. [PubMed: 18292573]
35. Savinova OV, Hoffmann A, Ghosh G. The Nfkb1 and Nfkb2 proteins p105 and p100 function as the core of high-molecular-weight heterogeneous complexes. *Mol Cell.* 2009; 34(5):591–602. Epub 2009/06/16. [PubMed: 19524538]
36. Scheidereit C. NF-kappaB and DNA Damage. *Keystone Symposia, NF-kappaB Signaling and Biology: From Bench to Bedside*; March, 2012; Whistler, British Columbia, Canada.
37. Lam LT, Davis RE, Pierce J, Hepperle M, Xu Y, Hottelet M, et al. Small molecule inhibitors of IkappaB kinase are selectively toxic for subgroups of diffuse large B-cell lymphoma defined by gene expression profiling. *Clin Cancer Res.* 2005; 11(1):28–40. Epub 2005/01/27. [PubMed: 15671525]
38. Kobori M, Yang Z, Gong D, Heissmeyer V, Zhu H, Jung YK, et al. Wedelolactone suppresses LPS-induced caspase-11 expression by directly inhibiting the IKK complex. *Cell Death Differ.* 2004; 11(1):123–30. Epub 2003/10/04. [PubMed: 14526390]
39. Hertlein E, Wagner AJ, Jones J, Lin TS, Maddocks KJ, Towns WH 3rd, et al. 17-DMAG targets the nuclear factor-kappaB family of proteins to induce apoptosis in chronic lymphocytic leukemia: clinical implications of HSP90 inhibition. *Blood.* 2010; 116(1):45–53. Epub 2010/03/31. [PubMed: 20351313]
40. Hu Y, Baud V, Delhase M, Zhang P, Deerinck T, Ellisman M, et al. Abnormal morphogenesis but intact IKK activation in mice lacking the IKKalpha subunit of IkappaB kinase. *Science.* 1999; 284(5412):316–20. Epub 1999/04/09. [PubMed: 10195896]
41. Ammirante M, Luo JL, Grivennikov S, Nedospasov S, Karin M. B-cell-derived lymphotoxin promotes castration-resistant prostate cancer. *Nature.* 2010; 464(7286):302–5. Epub 2010/03/12. [PubMed: 20220849]
42. Lam LT, Davis RE, Ngo VN, Lenz G, Wright G, Xu W, et al. Compensatory IKKalpha activation of classical NF-kappaB signaling during IKKbeta inhibition identified by an RNA interference sensitization screen. *Proc Natl Acad Sci U S A.* 2008; 105(52):20798–803. Epub 2008/12/24. [PubMed: 19104039]
43. Basak S, Shih VF, Hoffmann A. Generation and activation of multiple dimeric transcription factors within the NF-kappaB signaling system. *Molecular and cellular biology.* 2008; 28(10):3139–50. Epub 2008/02/27. [PubMed: 18299388]
44. Fusco AJ, Savinova OV, Talwar R, Kearns JD, Hoffmann A, Ghosh G. Stabilization of RelB requires multidomain interactions with p100/p52. *J Biol Chem.* 2008; 283(18):12324–32. Epub 2008/03/07. [PubMed: 18321863]
45. Hinz M, Arslan SC, Scheidereit C. It takes two to tango: IkappaBs, the multifunctional partners of NF-kappaB. *Immunol Rev.* 2012; 246(1):59–76. Epub 2012/03/23. [PubMed: 22435547]

46. Anest V, Cogswell PC, Baldwin AS Jr. IkappaB kinase alpha and p65/RelA contribute to optimal epidermal growth factor-induced c-fos gene expression independent of IkappaBalpha degradation. *J Biol Chem*. 2004; 279(30):31183–9. Epub 2004/05/25. [PubMed: 15155743]
47. Cohen J, Chen Z, Lu SL, Yang XP, Arun P, Ehsanian R, et al. Attenuated transforming growth factor beta signaling promotes nuclear factor-kappaB activation in head and neck cancer. *Cancer Res*. 2009; 69(8):3415–24. Epub 2009/04/09. [PubMed: 19351843]
48. Bian Y, Hall B, Sun ZJ, Molinolo A, Chen W, Gutkind JS, et al. Loss of TGF-beta signaling and PTEN promotes head and neck squamous cell carcinoma through cellular senescence evasion and cancer-related inflammation. *Oncogene*. 2012; 31(28):3322–32. Epub 2011/11/01. [PubMed: 22037217]
49. Freudlsperger C, Bian Y, Contag Wise S, Burnett J, Coupar J, Yang X, et al. TGF-beta and NF-kappaB signal pathway cross-talk is mediated through TAK1 and SMAD7 in a subset of head and neck cancers. *Oncogene*. 2012 Epub 2012/05/30.
50. Krappmann D, Wegener E, Sunami Y, Esen M, Thiel A, Mordmuller B, et al. The IkappaB kinase complex and NF-kappaB act as master regulators of lipopolysaccharide-induced gene expression and control subordinate activation of AP-1. *Molecular and cellular biology*. 2004; 24(14):6488–500. Epub 2004/07/01. [PubMed: 15226448]
51. Gomard T, Michaud HA, Tempe D, Tholon K, Pelegrin M, Piechaczyk M. An NF-kappaB-dependent role for JunB in the induction of proinflammatory cytokines in LPS-activated bone marrow-derived dendritic cells. *PLoS one*. 2010; 5(3):e9585. Epub 2010/03/12. [PubMed: 20221401]
52. Carayol N, Chen J, Yang F, Jin T, Jin L, States D, et al. A dominant function of IKK/NF-kappaB signaling in global lipopolysaccharide-induced gene expression. *J Biol Chem*. 2006; 281(41):31142–51. Epub 2006/08/18. [PubMed: 16914552]
53. Anest V, Hanson JL, Cogswell PC, Steinbrecher KA, Strahl BD, Baldwin AS. A nucleosomal function for IkappaB kinase-alpha in NF-kappaB-dependent gene expression. *Nature*. 2003; 423(6940):659–63. Epub 2003/06/06. [PubMed: 12789343]
54. Zerbini LF, Wang Y, Cho JY, Libermann TA. Constitutive activation of nuclear factor kappaB p50/p65 and Fra-1 and JunD is essential for deregulated interleukin 6 expression in prostate cancer. *Cancer Res*. 2003; 63(9):2206–15. Epub 2003/05/03. [PubMed: 12727841]
55. Debinski W, Gibo DM. Fos-related antigen 1 (Fra-1) pairing with and transactivation of JunB in GBM cells. *Cancer Biol Ther*. 2011; 11(2):254–62. Epub 2010/11/23. [PubMed: 21088499]
56. Bivona TG, Hieronymus H, Parker J, Chang K, Taron M, Rosell R, et al. FAS and NF-kappaB signalling modulate dependence of lung cancers on mutant EGFR. *Nature*. 2011; 471(7339):523–6. Epub 2011/03/25. [PubMed: 21430781]
57. Yemelyanov A, Gasparian A, Lindholm P, Dang L, Pierce JW, Kisseljev F, et al. Effects of IKK inhibitor PS1145 on NF-kappaB function, proliferation, apoptosis and invasion activity in prostate carcinoma cells. *Oncogene*. 2006; 25(3):387–98. Epub 2005/09/20. [PubMed: 16170348]
58. Bednarski BK, Ding X, Coombe K, Baldwin AS, Kim HJ. Active roles for inhibitory kappaB kinases alpha and beta in nuclear factor-kappaB-mediated chemoresistance to doxorubicin. *Mol Cancer Ther*. 2008; 7(7):1827–35. Epub 2008/07/23. [PubMed: 18644995]
59. Egorin MJ, Lagattuta TF, Hamburger DR, Covey JM, White KD, Musser SM, et al. Pharmacokinetics, tissue distribution, and metabolism of 17-(dimethylaminoethylamino)-17-demethoxygeldanamycin (NSC 707545) in CD2F1 mice and Fischer 344 rats. *Cancer Chemother Pharmacol*. 2002; 49(1):7–19. Epub 2002/02/22. [PubMed: 11855755]



**Figure 1. Aberrant expression of IKK $\alpha$  and IKK $\beta$  proteins in HNSCC tumor specimens and cell lines**

A, Immunohistochemical analysis of IKK $\alpha$ , IKK $\beta$ , and phospho-IKK in representative tumor specimens and matched mucosa. H&E, pan-cytokeratin and rabbit IgG isotype stains served as controls. Magnification: 100X and 400X. B, Intensities of immunohistochemical staining of nuclear and cytoplasmic IKKs were semi-quantified and presented as histoscores of 18 HNSCC specimens. C, Overexpression of IKK $\alpha$ , IKK $\beta$  and phospho-IKK proteins in whole cell and nuclear lysates of 9 UM-SCC lines were compared with normal human primary keratinocytes (HeKa).  $\beta$ -actin and Ponceau-S staining as loading controls.



**Figure 2. Overexpression of IKK $\alpha$  or  $\beta$  mutant vectors differentially modulates I $\kappa$ B- $\alpha$  and NF- $\kappa$ B mediated reporter activities**

A, Schematic depiction of functional domains of IKK $\alpha$  and IKK $\beta$  (upper). KD, Kinase domain; NSL, Nuclear localization signal; ULD, Ubiquitin-like domain; LZ, Leucine zipper; HLH, Helix-loop-helix; NBD, Nemo binding domain. K44, Lysine44; S176/S180, Serine176 and 180; S177/S181, Serine177 and 181. Western blot analysis of IKK $\alpha$  or IKK $\beta$  expression of control (Con), wild type (WT), phospho-resistant (AA), constitutively activated (EE), or kinase inactive (KA) IKK plasmids in whole cell lysates of UM-SCC1 cells 48h post-transfection (lower).  $\beta$ -actin as loading control. B, Relative I $\kappa$ B $\alpha$ -luciferase expression levels with EE or KA mutant of IKK $\alpha$  or IKK $\beta$  vector in UM-SCC1 cells 48h post-transfection. Statistical significance (t-test, \* $p$ <0.001, + $p$ <0.05) compared to the control. Experiments were performed in triplicate and presented as a mean with standard deviation. Relative NF- $\kappa$ B reporter activity measured 48 hrs after overexpression of mutant


IKK plasmids in UM-SCC1 (C) and UM-SCC11B cells (D). Statistically significant differences between individual IKK constructs (\*) or dual IKK overexpression (\*\*) with the control plasmid (t-test,  $p < 0.05$ ).

Author Manuscript

Author Manuscript

Author Manuscript


Author Manuscript



**Figure 3. A combinatory effect of dual IKKα and IKKβ knockdown by siRNA in blocking NF-κB signaling molecule expression and reporter activity**


A, Knockdown efficiency of IKKα and IKKβ individually or in combination, and effects on downstream NF-κB subunits RelA, p-RelA (ser536), p105, p50, RelB, p100, p52, p-p52 and p-IκBα (ser32/36) after knockdown for 48 hours and treatment with TNF-α for 1 hour. β-tubulin and Histone1 served as loading and fractionation controls. B, Knockdown of IKKs affected mRNA expression of NF-κB p100/p52 subunits. UM-SCC1 cells were transfected with siRNA of IKKs under the same experimental condition as shown above, and RNAs were harvested and relative fold-change in mRNA expression was measured via quantitative

RT-PCR. Starred (\*) values indicate a p-value<0.05 by t-test vs. negative control. p-p100 and p52 subunits in cytoplasmic (C) or nuclear (D) fraction were tested after IKK knockdown for 44 hours and treatment with lymphotoxin beta ( $\alpha 1/\beta 2$ ; 100ng/ml) for 4 hours. E, Greater I $\kappa$ B $\alpha$  luciferase fusion protein was observed in the double IKK knockdown than individual knockdowns. F, Conversely, the greater combinatory inhibitory effect of NF- $\kappa$ B reporter activity was observed in the IKK double knockdown over the single knockdowns in UM-SCC1 (left), UM-SCC11B (right), and UM-SCC6 (data not shown). Statistical significance of t-test, \* single knockdowns compared to negative control: p<0.01; \*\* double knockdowns compared to negative control: p<0.001; ++ dual knockdowns compared to single knockdowns: p<0.001.




**Figure 4. Array profiling of gene expression after IKK single or dual knockdowns by siRNA in UM-SCC1 cells**

A, Venn-diagram of down-regulated gene expressions and overlaps in IKK $\alpha$ , IKK $\beta$ , IKK $\alpha$  & IKK $\beta$  knockdown samples compared with control scramble siRNA transfected cells. B, Enriched signaling pathways were annotated among down-regulated genes, and p value (-log) depicted the significance. Ingenuity Pathway Analysis (IPA) constructed networks using the genes involved in the cytokine and NF $\kappa$ B pathways (C), and the growth factor signaling pathways (D).




**Figure 5. Knockdowns of IKKs and AP1 subunits down-regulate a partially overlapped gene program involved in NF- $\kappa$ B and EGFR/AP1 signaling pathways**

A, Relative fold-change in mRNA expression after IKK single or double siRNA transfection of UM-SCC1 for 48 hrs were measured by quantitative RT-PCR (white, IKK $\alpha$ ; grey, IKK $\beta$ ; black, IKK $\alpha$ +IKK $\beta$ ). B, Relative fold-change in mRNA expression after knockdown of AP1 subunits individually by siRNA in UM-SCC1 for 48 hrs were measured by quantitative RT-PCR (white, FOSL1; grey, cJun; black, JunB).  $p < 0.05$  by t-test vs. controls (\*).




**Figure 6. Knockdowns of IKKs and NF- $\kappa$ B RelA subunits modulated EGFR and AP1 expression and activity**

A, Western blots of UM-SCC1 whole cell lysates showed decreased EGFR and AP1 subunit protein expression after IKK knockdowns. Histone1 as the loading control. B, Relative fold-change in gene expression of EGFR and AP1 subunit molecules after knockdown of NF- $\kappa$ B RelA(p65) in UM-SCC1 for 48 hours. C, Decreased AP1 reporter activity after knockdown of IKKs individually or in combination. D, Overexpressed wild-type IKKs significantly increased IL-8 promoter (-133bp) activity. Point mutation of NF- $\kappa$ B (-83bp) or AP1 (-128bp) binding site abolished or inhibited IL-8 reporter activity.  $p < 0.05$  by t-test vs. controls (\*), or vs. IL-8 promoter activity when overexpressing IKKs (#).



**Figure 7. Knockdown of IKKs or AP1 subunits by siRNAs potently inhibit cell proliferation and migration**

A, IKK $\alpha$  and IKK $\beta$  double knockdown by siRNAs potently inhibited cell proliferation over single IKK knockdowns by MTT assay. Statistical significance of t-test, \* single knockdown:  $p < 0.05$ ; +double knockdown:  $P < 0.01$ . B, UM-SCC1 cells were transfected with IKK specific siRNAs for 48 hrs. Scratches were made on cell monolayers and wound closure was monitored at 14, 20, and 30 hrs. C, Quantification of wound healing with statistical significance. The distances between the wounds were measured and significant differences by individual IKK knockdowns were examined by t-test (\*,  $p < 0.05$ ), or in dual IKK knockdown (+,  $p < 0.01$ ). D, Knockdown of FOSL and JunB individually by siRNAs inhibited cell proliferation by MTT assay. Statistical significance of t-test, \*  $p < 0.05$ . E, AP1 subunits were individually knocked down for 48 hrs. Scratches were made and wound closure was monitored at 11, and 15 hrs under serum free condition. F, Quantification of wound healing with statistical significance. The distances between the wounds were measured and significant differences were examined by t-test. \*,  $p < 0.05$ .



**Figure 8. Drugs targeting IKK $\alpha$  and IKK $\beta$  inhibit NF- $\kappa$ B activity, and promote cell cycle arrest and apoptosis of UM-SCC cells**

A, UM-SCC1 and UM-SCC11B cells were treated with 17-AAG (50nM), MLN-120b (40 $\mu$ M), Wedelolactone (20 $\mu$ M) for 24hrs. IKK $\alpha$ , IKK $\beta$ , and phospho-p-65 S536 protein were tested in whole cell lysates by Western blots.  $\beta$ -actin as the loading control. B, Inhibition of NF- $\kappa$ B activity in UM-SCC1 or UM-SCC11B cells transfected with constitutively activated IKK $\alpha$  or IKK $\beta$  EE expression vector then treated with various doses of 17-DMAG or MLN-120b for 48hrs. Data calculated from triplicates and presented as mean with standard deviation. \*Statistical significance compared to controls (t-test, p<0.05).

C, Flow cytometry was performed to test cell-cycle and DNA fragmentation in sub-G0/G1 phase (% cell death) by UM-SCC1 and UM-SCC11B (data not shown) 24 hrs post treatment with 17-AAG (50nM), Wedelolactone (20 $\mu$ M), MLN-120b (40 $\mu$ M), and SC-514 (50 $\mu$ M, Supplemental Fig. 3A). D, Schematic of classical and alternative NF- $\kappa$ B pathway depicting the specific activation of IKK/NF- $\kappa$ B subunits as targets by each chemical inhibitors and the cross-induction of EGFR/AP1 expression and signaling.