

# What to do with the equipoise?

Wiebe G. Knol<sup>1\*</sup>, Grigorios Papageorgiou<sup>1,2</sup>, and Mostafa M. Mokhles<sup>1</sup>

<sup>1</sup>Department of Cardiothoracic Surgery, Erasmus Medical Center, Room Na-2624, P.O. Box 2040, 3000 CA Rotterdam, The Netherlands; and <sup>2</sup>Department of Biostatistics, Erasmus Medical Center, Room Na-2624, P.O. Box 2040, 3000 CA Rotterdam, The Netherlands

Online publish-ahead-of-print 24 April 2019

**This commentary refers to ‘Coronary computed tomography angiography for heart team decision-making in multivessel coronary artery disease’, by C. Collet et al., *Eur Heart J* 2018;39:3689–3698.**

After the rise of coronary computed tomography angiography (cCT-A) in diagnosing coronary artery disease, the aim shifts towards treatment decision-making in revascularization of these patients. The recent SYNTAX-III ‘Revolution’ trial<sup>1</sup> evaluated the appropriateness of cCTA for this purpose as compared to invasive coronary angiography. We received the results of this elegantly designed trial with great interest. They found high agreement between heart team decisions on percutaneous coronary intervention (PCI) or coronary artery bypass graft (CABG) surgery in patients with complex coronary disease. However, some questions remain about the analysis of their results.

The authors chose the SYNTAX-II score-based recommendation as primary endpoint measurement, rather than the subsequent heart team’s decision. Although a decision between PCI/CABG is still needed in case of equipoise in predicted mortality, we appreciate the value of the SYNTAX-II score. It eliminates bias by physicians or clinical factors unaccounted for in the SYNTAX-II score. This score, therefore, is indeed a more reliable tool for the comparison.

However, the authors merged the PCI-group and ‘equipoise between PCI/CABG’-group. We could not find an explanation for this. One could argue that in case of similar predicted mortality, the less invasive option would be preferred, hence the combination with PCI. Nevertheless in both study arms a majority (106 patients) in the equipoise-group were selected for CABG. Consequently, it would be more sensible to combine the equipoise- and CABG-group. Furthermore, combining categories in a non-trivial  $3 \times 3$  agreement

table will increase or decrease the Cohen’s kappa coefficient.<sup>2</sup> Therefore, if the SYNTAX-II recommendations are used, agreement using all three categories separately should be calculated. Since the categories can be viewed as ordinal data, a weighted kappa may be considered to assess inter-rater agreement.<sup>3</sup>

Secondly, addition of cCTA derived fractional flow reserve (CT-FFR) reduced the overestimation of the anatomical SYNTAX-score, changing the decision based on cCT-A alone in 7% of patients. Unfortunately, the authors did not quantify whether this change improved the agreement.

Finally, although trivial, we could not reproduce the significant *P*-values for ‘current smoking’ and ‘diabetes mellitus’ in *Table 4*.

We believe that implementation of CT-FFR can improve the decision-making process. It is important that the boundaries of these possibilities continue to be explored, and we endorse the authors’ effort in this regard. In order to fully appreciate the SYNTAX-III trial, we hope to receive more insight in the outcomes of this clinically important and indeed revolutionary trial.

**Conflict of interest:** none declared.

## References

1. Collet C, Onuma Y, Andreini D, Sonck J, Pompilio G, Mushtaq S, La Meir M, Miyazaki Y, de Mey J, Gaemperli O, Ouda A, Maureira JP, Mandry D, Camenzind E, Macron L, Doenst T, Teichgraber U, Sigusch H, Asano T, Katagiri Y, Morel MA, Lindeboom W, Pontone G, Luscher TF, Bartorelli AL, Serruys PW. Coronary computed tomography angiography for heart team decision-making in multivessel coronary artery disease. *Eur Heart J* 2018;39:3689–3698.
2. Warrens MJ. Cohen’s kappa can always be increased and decreased by combining categories. *Stat Methodol* 2010;7:673–677.
3. Cohen J. Weighted kappa: nominal scale agreement with provision for scaled disagreement or partial credit. *Psychol Bull* 1968;70:213–220.

\* Corresponding author. Tel: +31625540273, Email: [w.knol@erasmusmc.nl](mailto:w.knol@erasmusmc.nl)

© The Author(s) 2019. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)