

Anesthetic approach to Niemann–Pick Type C patient for dental treatment

Sir,

Niemann–Pick disease is an autosomal recessive lipid storage disease, characterized with differentiating levels of hepatosplenomegaly and progressive psychomotor retardation (this disease exhibiting heterogenic symptoms has four subtypes; Type A and B, in which sphingomyelinase values are normal, and Type C and D, in which sphingomyelinase values are close to normal values). Disease emerges with early childhood ataxia and progressive dementia, and the most evident features are early childhood hepatosplenomegaly, vertical supranuclear ophthalmoplegia, ataxia, dysarthria, mental-motor retardation, and seizures.^[1-3] In this report, we present our anesthetic approach to an uncooperative 3-year-old female patient with hepatosplenomegaly, mental-motor retardation, developmental retardation, deglutition, speech deficiency, ataxia, and seizures who admitted to our clinic in collaboration with our pedodontics clinic for preventive dentistry practice.

Preventive measures such as topical fluoride application to teeth and incision of the excess mucosa over the left upper and lower primary second molars to obtain eruption was planned, and the patient was consulted to our department for the surgical procedure and the need of sufficient mouth opening under anesthesia for the uncooperative patient. Mallampati was Grade 3, sternomental and thyromental distances were short. After 6 h of fasting, sedoanalgesia with monitored anesthesia care was planned because of the presence of difficult intubation criteria. Patient was taken to the operation room without premedication, was monitored and oxygenized 2 L/min. Five minutes after EMLA® application to the left-hand dorsal side, peripheral intravenous cannulation was obtained. Metoclopramide, to minimize vomiting possibility by increasing lower esophagus sphincter pressure, and ranitidine, to neutralize stomach acid was administered. Afterward, 0.5 mg midazolam and 20 mg ketamine were administered intravenously. After termination of the 20 min operation, patient was observed for 10 min and sent to service while hemodynamically stable.

Our choice of anesthetic in addition to local infiltration anesthesia (Ultracain®) was ketamine, which has analgesic, hypnotic, and amnestic effects but does not repress pharyngeal and laryngeal reflexes and does not evoke cardiovascular and respiratory depression.^[3] The potential effect of ketamine of elevating convulsion threshold was prevented with midazolam. We are in the opinion that this method, which did not require additional anesthetic methods, was free of side effects and safe for the 3-year-old child with Niemann–Pick Type C disease.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Ayşe Hande Arpacı

Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Ankara University, Besevler, Ankara, Turkey

Address for correspondence: Dr. Ayşe Hande Arpacı, Department of Oral and Maxillofacial Surgery, Faculty of Dentistry, Ankara University, Besevler, Ankara, Turkey.
E-mail: handarpaci@yahoo.com

REFERENCES

1. Miao N, Lu X, O'Grady NP, Yanjanin N, Porter FD, Quezado ZM. Niemann-pick disease type C: Implications for sedation and anesthesia for diagnostic procedures. *J Child Neurol* 2012;27:1541-6.
2. Lee SY, Lee HJ, Kim SH, Jeong YJ, Jin HK, Bae JS, *et al.* Two siblings with adolescent/adult onset Niemann-Pick disease type C in Korea. *J Korean Med Sci* 2016;31:1168-72.
3. Gazal G, Fareed WM, Zafar MS, Al-Samadani KH. Pain and anxiety management for pediatric dental procedures using various combinations of sedative drugs: A review. *Saudi Pharm J* 2016;24:379-85.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

Access this article online	
Quick Response Code:	Website: www.jmsjournal.net
	DOI: 10.4103/1735-1995.200276

How to cite this article: Arpacı AH. Anesthetic approach to Niemann–Pick Type C patient for dental treatment. *J Res Med Sci* 2017;22:25.