



Secondary metabolites from endophytic fungi: Production, methods of analysis, and diverse pharmaceutical potential

Vivek Kumar Singh¹ · Awanish Kumar¹

Received: 22 January 2023 / Accepted: 22 May 2023
© The Author(s), under exclusive licence to Springer Nature B.V. 2023

Abstract

The synthesis of secondary metabolites is a constantly functioning metabolic pathway in all living systems. Secondary metabolites can be broken down into numerous classes, including alkaloids, coumarins, flavonoids, lignans, saponins, terpenes, quinones, xanthenes, and others. However, animals lack the routes of synthesis of these compounds, while plants, fungi, and bacteria all synthesize them. The primary function of bioactive metabolites (BM) synthesized from endophytic fungi (EF) is to make the host plants resistant to pathogens. EF is a group of fungal communities that colonize host tissues' intracellular or intercellular spaces. EF serves as a storehouse of the above-mentioned bioactive metabolites, providing beneficial effects to their hosts. BM of EF could be promising candidates for anti-cancer, anti-malarial, anti-tuberculosis, antiviral, anti-inflammatory, etc. because EF is regarded as an unexploited and untapped source of novel BM for effective drug candidates. Due to the emergence of drug resistance, there is an urgent need to search for new bioactive compounds that combat resistance. This article summarizes the production of BM from EF, high throughput methods for analysis, and their pharmaceutical application. The emphasis is on the diversity of metabolic products from EF, yield, method of purification/characterization, and various functions/activities of EF. Discussed information led to the development of new drugs and food additives that were more effective in the treatment of disease. This review shed light on the pharmacological potential of the fungal bioactive metabolites and emphasizes to exploit them in the future for therapeutic purposes.

Keywords Endophytic fungi · Secondary metabolites · Characterization · Bioactive · Therapeutics

1 Introduction

In recent years, the world population has faced many challenges in the field of healthcare. Its associated problem, including various diseases like cancer, diabetes, cardiovascular disorder, multiple drug resistance, etc. are major defiance for the human fraternity (Qadri et al. 2021; WHO 2020). Even in the era of vaccines and drugs, infectious diseases are responsible for the high mortality rate. Diseases like diarrhea cause the death of around 525 000 children under five years of age (Tesfaye et al. 2020). Around 33 million populations are infected with HIV and reported to be two million deaths last decade (Laga et al. 2015). *Mycobacterium tuberculosis* is recorded to infect around one-third

of the world's population. In tropical and subtropical areas, underprivileged populations are infected with malaria, accounting for about one million deaths (Cárdenas et al. 2021). Similarly, COVID-19 (Pokhrel and Chhetri 2021) and other pandemics as monkeypox (Bhattacharya et al. 2022), immunosuppressive disorders, and the appearance of highly virulent viruses open a new discussion for future treatment and remedies to address those problems (Chatelain 2015). An international health agency prioritizes the conservation of natural resources to create novel therapeutics. Plants are major natural resources and have therapeutic potential that has been tracked down thousand years ago and utilized for the treatment of many diseases (Abel 2013). Today around 40% of modern medicines are plant-derived because of fewer side effects. Plants generate various secondary metabolites, which may be broken down into the chemical categories of phenolics, terpenes, and alkaloids (Sharma and Singh 2021). Secondary metabolites secreted by the plant during different stress and developmental stages make them competitive in their inhabitant (Pang et al. 2021). These small molecules

✉ Awanish Kumar
drawanishkr@gmail.com

¹ Department of Biotechnology, National Institute of Technology, Raipur (CG), Raipur 492010, Chhattisgarh, India

exert a significant impact on plants themselves and also on humans and other organisms (Teoh 2016).

Plants are the natural inhabitant of microorganisms as they form a symbiotic association with them to accomplish one and all requirements. Endophytes (endosymbiotic microbes colonize in plants) and microbes and their bioactive metabolites are important natural sources for promising therapeutic agents. The therapeutic potential of endophytes and their metabolites as biotherapeutic agents has garnered a lot of interest (Xia et al. 2022). Endophytes are associated with the healthy tissue of the plant (Sadrati et al. 2013) and reside in the interior sections of the plant, such as the root, the stem, the petiole, and other components that are referred to as endophytes (Ma et al. 2013; Deepthi et al. 2018). In 1898, Vogl was the first person who reported that endophytic mycelium was present in the grass seeds of *Latium teinutentuin* (Waghunde et al. 2021). Around 300,000 plant species, each individual having one or more endophytes and having one to hundred strains of endophytes varying according to the host system (Gao et al. 2018). They got much attention when it became apparent that endophytes can produce bioactive secondary metabolites with varied molecular structures, which are barely impersonated by synthetic chemistry (Mengistu 2020). Endophytes play a crucial role in plant development, and survival, and regulate some defense mechanisms. They are formed to be a composite equilibrium to achieve the host boundary and build a mutualistic association with the host (Alam et al. 2021; Nanda et al. 2019). During stressed conditions in the plant, endophytes secreted a range of secondary metabolites in plant cells and incorporated them into the stressed pathways (Nanda et al. 2019). Secretion of stress-controlled molecules like gibberellins (GB), cytokinin (CIS), salicylic acid (SA), and indole-3-acetic acid (IAA) enhance plant growth and development and is responsible for many physiological changes in the plant (Shekhawat et al. 2021; Duc et al. 2018; Shaffique et al. 2022).

The biochemical and pharmaceutical industries rely on endophytic fungi as a source of new therapeutic biomolecules that could be immune-suppressant compounds, anti-cancer drugs, plant growth promoters, anti-microbial volatiles, insecticides, anti-oxidants, and antibiotics, offer significant potential for application in medicine. Furthermore, endophytic microorganisms can lessen a plant's capacity to endure nutritional deficiency, high temperatures, salt, trace metals, and water scarcity (Eid et al. 2021). Amylase, cellulase, lyase, and laccase are important enzymes that have significant industrial applications and endophytes play a role in their synthesis (Sharma et al. 2021). Research suggested that the fungal endophytes found to be heterotrophic organisms with various life cycles in natural ecosystems form a symbiotic relationship with plants. The fossil record also reported that endophytic fungi and host plants were associated for 400 million years and intimately involved in the

ecology, proliferation, fitness, and steer the evolution of their life (Krings et al. 2007). Plant systems change their habits from water to terrestrial atmosphere with many challenges like nutrient-deficient soil, high carbon dioxide varying temperature conditions, and water availability. Fungal endophytes provided tolerance to the plant during adverse conditions and fixed them in the soil. In the evolutionary period, fungal endophytes evolved themselves in the plant environment through an alteration of genetic behavior, uptake of the DNA, and started producing secondary metabolites (Arora and Ramawat 2017). At least one plant harbour one or more fungal endophytes, especially woody plants containing hundreds of species of fungal endophytes. The fungal endophytes are found in different geographical and climate regions, fin-ray ubiquitously distributed, and rich in species diversity. The abundance of fungal endophytes with great extent, ubiquitous nature, diversity, and wide range of ecological functions are shown to be greatly adapted for plants under worldwide distribution and selective pressure (Rodriguez et al. 2009).

All these findings indicate that fungal endophytic population colonization inside the host tissue confers tolerance under specific environmental stresses condition and is responsible for the survival of plants. Extensive research on fungal endophytes (FE) found their crucial role in abiotic and biotic stress tolerance, nutrient supply, growth, and plant development. When searching for natural products mediated by endophytes, researchers may explore EF. It is well established that many bioactive metabolites (BM) with potential pharmacological effects are produced by EF (Tiwari and Bae 2022). The bioactive compounds produced by EF albeit receiving much less attention. The development and production of biomass need a thorough understanding of endophyte ecology, bioactive components, and the bio-transformation of substrates. Considering the information available on EF, the article is aimed to review the sampling, optimization, production, and extraction of the secondary metabolites from EF and the pharmacological relevance of the different bioactive metabolites produced by endophytic fungi associated with plants (Fig. 1).

2 Sampling and optimization of media for the production of BM from EF

The first steps in sampling and isolation of bioactive molecules from EF are collecting plant material having EF from different geographical areas and pre-processing the plant material, which may involve surface cleaning, slicing, and selecting media. Surface sterilizing agents like mercuric chloride (HgCl_2), ethanol, etc., are used to wash and surface sterilize the plant components (leaves, stems, seeds, etc.) (Bisht et al. 2016). They are subsequently




Fig. 1 Entry and Colonization of the fungal endophytes in response to biotic (herbivores and pathogens) and abiotic (drought, salinity, heavy metals, and others) stresses and production of secondary

metabolites to modulate plant defense, plant growth, and functional role in the pharmacology

diced and cultured in LB media on a PDA (potato dextrose agar) plate. Hyphal tips are harvested from the fungus and then transferred to PDA slants, where they are tested for bioactive secondary metabolites after an extended period of incubation (Sharma et al 2016). It is vital to build a suitable cultivation system for commercial use because EF can produce many biologically active metabolites. Endophytes can be grown through liquid-submerged or solid-state fermentation (SSF). EF fermentation is fruitful, continuous, and environmentally beneficial. Submerged culture produces mycelial biomass and bioactive metabolites faster. It needs less time and has fewer contamination risks. Extensive research has been done on bioactive compounds produced by endophytic fungus in submerged fermentation. In liquid fermentation, temperature, pH, aeration, and agitation affect secondary metabolite production. In addition, each of these different characteristics has been optimized, which has led to an increase in the overall production of BM (Debbab et al. 2013; Brader et al 2014). Various researchers used liquid-submerged fermentation

to manufacture myriads of BM. These include antibiotics, anti-oxidants, pigments, enzymes, etc. (Mrudula and Murugammal 2011; Patil et al. 2016). Similarly, cellulase enzyme production from *Pestalotiopsis* sp (Chen et al. 2011) and glucoamylase from *Aspergillus flavus* (Karim et al 2017) has been performed previously. Vimal and Kumar 2022 reported optimized production of medically important L-asparaginase enzyme under solid-state and submerged fermentation from agricultural wastes. Similarly, microorganisms are cultured on Wheat bran, Cajanuscajan (red gram), Phaseolus mungo (mung bean), and Glycine max (soybean) bran in solid-state fermentation (SSF). Chitosan production by *A. terreus* was reported by submerged fermentation in the optimized condition (Abo Elsoud et al. 2023). SSF from fungal cultures provides various advantages over submerged fermentation to manufacture bioactive chemicals in the food, agricultural, and pharmaceutical industries. Furthermore, this includes comparatively improved productivity, greater product concentrations, and simple equipment requirements for the

fermentation process of BM (Patil et al. 2016). Lovastatin, a potent medication for decreasing blood cholesterol, was extracted from the healthy tissues of *Taxusbaccata* by an endophytic fungus, *A. niger* PN2, using SSF with wheat bran as the substrate (Raghunath et al. 2012). Further, we discuss the production and extraction of the SM from EF.

3 Production and extraction of the secondary metabolites (SM) from endophytic fungi (EF)

Biomolecules like polysaccharides, polypeptides, unsaturated fatty acids, and glycoproteins are commonly known as elicitors. EF produces the elicitors or signaling molecules stimulated by the bioactive phytochemical accumulation in plants. Some of them provide defense to plants against disease-causing organisms. They also stimulate the production of several phytochemicals, including alkaloids, flavonoids, terpenoids, saponins, and phenols (Chandran et al. 2020). The oligosaccharide components of *Colletotrichum gloeosporioides*' crude endophytic mycelium have been demonstrated to stimulate artemisinin synthesis in hairy root cultures of *Artemannua* (Hussain et al. 2015). The bacterial culture was incubated at 32 °C in broth for 36 h, whereas fungal cultures were incubated at 28 °C for two weeks with 150 rpm shaking. Several solvents were employed alone or in combination to extract metabolites. Ethyl acetate, methanol, dichloromethane, hexane, and ethanol are routinely used to extract metabolites from the culture broth. The solubility of the desired component determines the extraction solvent. Equal amounts of solvents were added to the filtrate and agitated for 10 min until two transparent immiscible layers appeared. The extracted compounds were separated from the solvent using a funnel. The solvent was evaporated and the compound was dried in a rotator vacuum evaporator to produce the crude metabolite (Bhardwaj et al. 2015). The crude extract was diluted with dimethyl sulphoxide and kept at 4 °C. Phytochemical screening was conducted to look for alkaloids, saponins, tannins, flavonoids, steroids, sugars, and cardiac glycosides (Mathew et al. 2012). The isolation and characterization methods of secondary metabolites isolated from fungal endophytes are discussed in Table 1 and in this section; we discuss some high throughput methods precisely used for analysis of SM from EF. Chromatographic methods such as TLC and HPLC were employed to get the secondary metabolite extract as pure as feasible. The recovered fractions are usually analyzed by the use of gas chromatography-mass spectrometry (GC-MS), Fourier transform infrared (FTIR), and nuclear magnetic resonance (NMR). NMR and MS are the principal techniques exploited

in the structural characterization of BM. According to Madhusudhan et al. (2015), X-ray diffraction (XRD) is also promising for crystalline biomolecules.

3.1 TLC

TLC analysis was used to extract the bioactive components of *Pestalotiopsis neglecta* BAB-5510, a fungal endophyte that was isolated from the leaves of *Cupressus torulosa* D. Don. Two distinct fractions were observed on the silica gel TLC plates after being developed in dichloromethane and methanol at a ratio of 90:10, with the second fraction having an R_f value of 0.79. TLC was employed to separate the extracted secondary metabolites synthesized by fungal endophytes, which have been isolated from *Mentha piperita*. The R_f values of the metabolites that were isolated from bacterial endophytes were found to be quite comparable to those that were obtained by the TLC chromatogram.

3.2 GC-MS

According to the findings of gas chromatography performed on *Pestalotiopsis neglecta* BAB-5510, the most important active compounds of *Pestalotiopsis sp.* BAB-5510 are nonadecane (19.74%), 1,2,3-propanetriol, 1-acetate (17.21%), bis (2-Ethylhexyl) phthalate (14.41%), and 4 Hpyran-4-one, 2,3-dihydro-3,5-d (Bunaciu et al. 2015). The GC-MS spectra revealed that these metabolites were terpenes, more notably cinnamaldehyde, cinnamyl alcohol, and eugenol (Kumar et al. 2017) and GC-MS is particularly suitable to identify volatile organic compounds.

3.3 HPLC & LC-MS

Silica gel column chromatography and high-performance liquid chromatography were used to purify BMs. The *Taxus cuspidate* culture media was treated with di-chloromethane to extract taxol, which was then purified and quantified by employing HPLC. Finally, the structure of taxol was confirmed by utilizing LC-MS and H-NMR spectroscopy (Sharma et al. 2016). Ethyl acetate was used to extract vinblastine and vincristine from the fungus endophyte *Fusarium oxysporum*.

3.4 NMR

To evaluate the molecular masses of the purified compounds, electrospray ionization mass spectrometry (ESI-MS) and tandem mass spectrometry (MS-MS) followed by nuclear magnetic resonance (NMR) analysis utilized by Kumar et al. 2013. They performed isolation, purification, and characterization of Vinblastine and Vincristine from EF *Fusarium oxysporum* isolated from *Catharanthus roseus*.

Table 1 Isolation, characterization, and their functional role of secondary metabolites isolated from fungal endophytes

Endophytes	Compound	Host plant	Genome size of Endophytes (Mb)	Class of compound	Activity/Function	Yield	Growth media/extraction medium	Method of purification	Characterization	Reference
<i>Alternaria alternata</i>	Taxol	<i>Taxaschinensis</i> var. <i>mairei</i>	33.67	Alkaloids	Anticancer, Cytotoxicity,	84.50 mg/mL	PDB medium	HPLC	LC-MS, H NMR	(Shankar Naik 2019)
<i>Fusarium oxysporum</i>	Vincristine	<i>Catharanthus roseus</i>	52.60	Alkaloids	Anticancer	67 µg/L	Methanol extraction PDB medium Ethyl acetate extraction	TLC and HPLC	UV-Vis spectroscopy, ESI-MS/MS, and 1 H NMR	(Kumar et al. 2013)
<i>Penicillium citrinum</i>	Huperzine A	<i>Huperziaserrata</i>	32.19	Alkaloids	Treating for Alzheimer's disease (AD)	1.38 mg/L	PDB Medium Chloroform & Methanol extraction	TLC and HPLC	ESI-MS, LC-MS, H NMR	(Thi Minh et al. 2019)
<i>Nectria haematococca</i>	Quercetin	<i>Zingiberis monnii</i> (J. Graham)	54.43	flavonoid glycosides	Anti-oxidant	5.82 ± 0.2 mg/gm	PDB Medium Ethyl acetate extraction	HPLC	ESI-MS and MS-MS	(Das et al. 2017)
<i>Fusarium solani</i>	Camptothecin	<i>Camptotheca acuminata</i>	51.74	Quinolone Alkaloids	Anticancer	(1.2 to 152 fold)	PDB medium Chloroform and methanol (4:1 v/v)	TLC and HPLC	ESI-MS	(Kaur et al. 2020)
<i>Bisogniauxia cylindrospora</i>	Isofraxidin	<i>Rice</i> (<i>Oryza sativa</i>)	-	Polyketides	Antibacterial activity, Anti-oxidant, Anticancer	2.5 mg/1.5 KG	PDA medium Methanol extraction	TLC	1H NMR, ESI-MS	(Wu et al. 2019)
<i>Phomopsis</i> sp.	Gallic acid	<i>Acer ginnala</i>	-	Phenol	anti-oxidant, anti-inflammatory, anti-microbial, anti-cancer	29.25 mg/gm	PDA medium Methanol extraction	HPLC	ESI-NMR	(Qi et al. 2009)
<i>Alternaria</i> sp.	Berberin	<i>Coptis chinensis</i>	-	Isoquinoline alkaloid	Antibacterial, antidiabetic, antihypertensive,	9.313 µg/gm	PDA medium Methanol extraction	TLC, HPLC	MS-NMR	(Zhang et al. 2016)
<i>Colletotrichum gloeosporioides</i>	Piperine	<i>Piper nigrum</i>	57.60	Alkaloids	anti-diabetic, antidiarrheal, anti-oxidant, antibacterial	-	PDA medium Ethyl acetate extraction	HPLC	HPLC and LCMS	(Chithra et al. 2014)
<i>Phomopsis vexans</i>	Lovastatin	<i>Solanum xanthocarpum</i>	59.78	Polyketide	Controlling the blood cholesterol level	550 mg/L	PDA, Czapek Dox broth medium, Ethyl acetate extraction	TLC, HPLC	FT-IR, UV, C NMR, and LC-MS analyses	(Parthasarathy and Sathiyabama 2015)
<i>Fusarium oxysporum</i>	Diosgenin	<i>Dioscorea zingiberensis</i>	52.60	triterpenoids	Antitumor, Anti-inflammatory	5.21 mg/L	PDA medium Ethyl acetate, methanol extraction	HPLC	GC-MS, LC-MS	(Biswas et al. 2020)

Table 1 (continued)

Endophytes	Compound	Host plant	Genome size of Endophytes (Mb)	Class of compound	Activity/Function	Yield	Growth media/extraction medium	Method of purification	Characterization	Reference
<i>Epicoccum nigrum</i>	Hypericin	<i>Hypericum perforatum</i>	34.73	Benzopyrenes	Antidepressants, antitumors and antiviral	320.4 ng/mg	PDB medium Ethyl acetate, extraction	HPLC	HPLC-HRMS, HRMS/MS	(Vigneshwari et al. 2019)
<i>Fusarium oxysporum</i>	Ginkgolide B	<i>Ginkgo biloba</i>	52.60	Diterpenoids	Anti-inflammatory, Antiallergic	0.2 mg/mL	PDB medium Ethyl acetate, extraction	TLC	HPLC/ESI-MS and 13C-NMR	(Cui et al. 2012)
<i>Colletotrichum coccodes</i>	Tyrosol	<i>Houttuynia cordata Thunb</i>	50.24	Phenolic glycosides	Anti-microbial	2.3 mg/mL	PDB medium Ethyl acetate, extraction	TLC	1H NMR-MS	(Talukdar et al. 2021)
<i>Alternaria</i> spp.	Emodin	<i>Hypericum perforatum</i>	33.67	Hydroxyanthraquinones	Antibacterial, Antiulcer, Anti-inflammatory, Anticancer, and Antinociceptive	20.8 ng/mg	PDB medium Ethyl acetate, extraction	HPLC	HPLC-HRMS, HRMS/MS	(Vigneshwari et al. 2019)
<i>Penicillium frequentans</i>	Curcumin	<i>Curcuma wenyujin</i>	-	phytopolyphenol	anti-microbial, anti-cancer, anti-inflammatory and anti-oxidant	12.6 mg/5.213 g	PDA medium Czapek medium chloroform-methanol extraction	silica column chromatography, TLC	ESI-MS, 1H NMR	(Yan et al. 2014)
<i>Annulohypoxylon boveri</i> var. <i>microspora</i>	Cinnamic acid	<i>Oryza sativa</i>	-	Phenolic acid	anti-oxidant and antibacterial activities,	1.5 mg/2.8 gm	PDA & RGY medium, N-butanol, chloroform-ethyl acetate extraction	TLC	UV, ESI-MS, 1H NMR	(Cheng et al. 2011)

4 Pharmaceutical application of the endophytic fungal secondary metabolites

EFs are eminent for their competence in synthesizing a distinct variety of pharmacologically significant chemicals with immense therapeutic promise, including antiviral, antifungal, antibacterial, antitumor, and anti-cancer activity. Various EFs are potential sources of plant growth factors and hormones. Some endophytes have been demonstrated to release a broad spectrum of extracellular enzymes, such as phosphatase enzyme, which converts insoluble phosphates to soluble forms for easier assimilation by plants. Secondary metabolites of EFs have been demonstrated to strengthen the host's immune system, reducing the severity of infections and the damage caused by pathogenic microorganisms (Sharma et al. 2021). The biocontrol systems of plants are responsible for the production of various kinds of BMs, which shield plants against potentially lethal diseases and stimulate their development (Santos et al. 2018; Hardoim et al. 2015). Their pharmaceutical role was experienced against various diseases (Table 1) and they are discussed below.

4.1 Antibiotics

Antibiotic synthesis through metabolic pathways is widely regarded as an effective strategy for protecting plants against illness. Phytopathogens can be inhibited by various bioactive substances and out of them few have been researched (Suryanarayanan 2013; Daguerre et al. 2017). Endophytes produced diverse metabolites, most of which have anti-microbial properties. These metabolites include alkaloids, flavonoids, peptides, phenols, polyketides, quinones, steroids, and terpenoids (Lugtenberg et al. 2016; Fadiji and Babalola 2020). DAPG, also known as 2, 4-diacetyl phloroglucinol, is a phenolic antibiotic with a broad spectrum of activity and has shown *Pseudomonas* spp. It contributes to the biological control of plant diseases, particularly soil-borne (Bonaterra et al. 2022). The novel alkaloid altersetin was isolated from the endophyte *Alternaria* spp. showing substantial antibacterial activity against various pathogenic gram-positive bacteria. Fungal endophytes isolated from *Artemisia annua* have been shown to inhibit the development of most phytopathogenic organisms in vitro by secreting n-butanol and ethylacetate (Fadiji and Babalola 2020). In addition, *pseudomonads* spp. generate cyclic lipopeptides (CLPs) amphiphilic molecules with chains of 7–25 amino acids that act as biosurfactants and are significant in biological control because of their favourable competitive capability with numerous groups of microorganisms (Flury et al. 2017; Bonaterra et al. 2022).

4.2 Siderophore

Micronutrient metals including nickel, copper, zinc, and iron are essential for soil plants and microorganisms, however, their bio-availability is often low due to environmental factors (Satapute et al. 2019). Reduced bioavailability of Fe(III) directly results from forming insoluble oxyhydroxide phases in response to harsh environmental conditions. Due to a lack of iron, plants develop chlorosis and have lessened metabolic activity and biomass. In response to various stresses, plants, and microbes have evolved a chelation strategy to increase metal availability (Chowdappa et al. 2020). Endophytes secreted small molecules called siderophores capable of chelating iron and increasing the bioavailability of the iron molecule to the plant (Yadav 2018). The iron in the soil can be dissolved with the assistance of secreted siderophores, which have a strong affinity for the substrate and the potential to assimilate it. The bacterial iron-siderophore complex makes iron accessible for plant development while simultaneously reducing the acquisition of iron by phytopathogens, which restricts phytopathogen proliferation (Santos et al. 2018). Despite this, the chemical composition of different microorganisms' siderophores can be somewhat distinct. For instance, bacterial hydroxamates are composed of hydroxylated and acylated alkylamines, whereas fungal hydroxamates are composed of hydroxylated and acylated ornithine groups. Chelating compounds can be obtained from endophytic fungi derived from *Cymbidium aloifolium*. This medicinal orchid can secrete exogenous siderophores and form stable complexes with the metal ion Fe^{3+} . These metabolites have the great potential to act as antibacterial siderophores. In a similar vein, a hydroxamate-type siderophore obtained from *P. crysogenum* was found to possess potent antibacterial activities against some of the most virulent phytopathogens, hence safeguarding peanuts and rice plants (Chowdappa et al. 2020). To evaluate the inhibitory efficacy of exogenous deferoxamine-B and siderophores-exochelin MS (a pentapeptide derivative) against methicillin-resistant *Staphylococcus aureus*, metallo-lactamase producers *Acinetobacter baumannii*, and *Pseudomonas aeruginosa*, disc diffusion, micro broth dilution, and turbidimetric growth tests were utilized (Gokarn and Pal 2018). The combination of siderophores and antibiotics was effective against the drug-resistant isolates. They can treat antibiotic-resistant bacteria and acute iron intoxications such as hemochromatosis by producing sideromycins. Evidence suggests they help to treat malaria (Chowdappa et al. 2020), bioremediating mercury (Pietro-Souza et al. 2020), and other plant diseases. According to some research, microorganisms' sensitivity to oxidative stress is lowered by siderophores. They can be used in cosmetics, cancer therapy, combat fish infections, and against bacterial/fungal phytopathogens (Chowdappa et al. 2020; Peralta et al. 2016).

4.3 Hydrolyzing enzyme

Endophytes make lytic enzymes like amylases, lipases, cellulases, pectinases, proteases, phosphatases, hemicellulases, chitinases, and 1, 3-glucanases (Mishra et al. 2019), which help them form symbiotic relationships with host plants and control of plant pathogens. Cellulase produced by endophytic fungi such as *Epicoccum nigrum*, *Trichoderma asperellum*, and *Alternaria longipes* has been shown to suppress the development of *Epicoccum sorghinum*, *Alternaria alternata*, *Fusarium thapsinum*, and *Curvularia lunata* in vitro by hydrolyzing the cell wall (Fadiji and Babalola 2020). The biocontrol efficacy against tall fescue leafspot and sugar beet damping-off was reduced after mutations in the 1,3-glucanase genes of *Lysobacteren zymogenes*. The lytic enzymes produced by *Streptomyces* are also effective against cocoa witch broom (Gao et al. 2018). Lytic enzymes produced by endophytes are often more robust and can function on a broader pH, temperature, and pressure range as compared to enzymes produced using traditional chemical catalysts (Tiwari 2015). These features also bode well for the commercial use of these enzymes in the food, detergent, paper, pharmaceutical, energy, and biofuel industry (Rana et al. 2019) because endophytic amylase hydrolyzes starch by accelerating the creation of 1,4 glycosidic bonds (Tiwari 2015). Bacterial pectinases, which depolymerize pectin linkages, are also widely utilized and have many applications in juice and food industries as well as in paper/pulp production, composting, recycling, etc. (Haile and Ayele 2022).

4.4 Growth-promoting hormones

Plant growth stimulation is bolstered, and phytohormones produced by fungal endophytes alter the plant's shape and form. This quality of endophytes has helped them progress in sustainable agriculture (Fadiji and Babalola 2020). Phytohormones or plant growth regulators are chemical substances that control, impede, or accelerate growth promotion and development of plants at low concentrations (Damam et al. 2016). Many phytohormones, including auxins, gibberellins, abscisic acid, cytokinins, ethylene, strigolactones, brassinosteroids, and jasmonates are produced by endophytic microorganisms (Santoyo et al. 2016; Shahzad et al. 2016). The primary function of indole-3-acetic acid (IAA) in plants is to stimulate cell growth and division. The IAA generated by the bacteria in symbiosis facilitates nutrient availability, increases root exudation, and encourages the growth of adventitious and lateral roots. The bacteria of the genera *Azospirillum*, *Herbaspirillum*, *Azotobacter*, *Alcaligenes*, *Pseudomonas*,

Enterobacter, *Klebsiella*, *Burkholderia*, *Pantoea*, *Rhizobium*, *Bacillus*, *Rhodococcus*, *Acetobacter*, etc., are familiar endophytic IAA producers (Eid et al. 2021). IAA synthesis also increases the size of bacterial cell walls, speeds up the release of exudates, and makes more of the nutrients that help other beneficial bacteria thrive in the rhizosphere more accessible. As a result, IAA is the primary effector molecule of endophytic bacteria involved in phytostimulation, pathogenicity, and plant–microbe interaction (Etesami et al. 2015). A specific form of bacterial endophyte contains 1-aminocyclopropane-1-carboxylate deaminase, which allows it to convert the ethylene precursor ACC into ammonia and beta-ketobutyrate. (Eid et al. 2021). Due to the release of ACC deaminases, these bacterial endophytes can stimulate plant development in nitrogen-limiting environments. Additional benefits include a more robust immune system and increased resistance to abiotic stress. In the case of sorghum plants, *Pseudomonas brassicacearum* (SVB6R1) increases the expression of ACC deaminase, increasing the plant's resistance to salt stress (Gamalero et al. 2020).

4.5 Nutrients

The expansion of agricultural production is facilitated by biofortification, which simplifies the uptake of nutrients by plants. Fungal endophytes can colonize plant structures such as roots, stems, and leaves and they are less likely to be outcompeted than microbes that live in the soil, therefore, they can be utilized as a replacement to boost the plant's ability to fix nitrogen (Santos et al. 2018). The presence of many microorganisms in the soil allows for the breakdown of insoluble phosphate and thus phosphorus is made available to the plants (Alori et al. 2017). In addition to their involvement in the release of organic acids into the soil, endophytes are also involved in the solubilization of phosphate complexes and the conversion of these compounds into the ortho-phosphates that plants absorb more readily. $\text{Ca}_3(\text{PO}_4)_2$ solubility was observed for poplar endophyte strains WP5 and WP42, and subsequent solubilization tests validated this observation (Kandel et al. 2017; Khan et al. 2015; Varga et al. 2020). Selenobacteria is a plant endophyte that can draw selenium out of the soil and transmit it to the host plant, where it can be used to promote plant development. Selenium biofortification in Glycine max may be enhanced by the endophyte *Paraburkholderia megapolitana* (MGT9) during drought (Trivedi et al. 2020). Zn solubilizing endophytes (such as *B. subtilis* DS-178 and *Arthrobacter sp.* DS-179) improve the translocation and enrichment of Zn to grain in specific Wheat genotypes (Singh et al. 2018).

4.6 Phytoremediation

As the global economy has become more industrialized over the past century, various anthropogenic chemicals have been released into the environment. These include polycyclic aromatic hydrocarbons (PAHs), petroleum hydrocarbons (PHC), halogenated hydrocarbons, salt, solvents, pesticides, and heavy metals (Bisht et al. 2015). Microbe-assisted phytoremediation is a method that deals with these issues. Due to endophytes' closer relationship with their host plants, phytoremediation may be enhanced (Li et al. 2012). It has been discovered that many endophytes can withstand high concentrations of heavy metals and degrade organic pollutants. Because of their prolonged exposure to the high metal concentrations stored in the hyperaccumulators of these plants, the endophytes associated with these plants may have become more resistant to the negative consequences of metal exposure (Aishwarya et al. 2014). Furthermore, it has been demonstrated that releasing low-molecular-mass organic acids by specific endophytes might boost heavy-metal mobilization. For example, the organic acids produced by endophytes caused the pH of a solution to fall when the water-soluble Pb concentration increased. When the concentration of water-soluble Pb rose, the pH of the solution decreased because of the organic acids generated by endophytes (Yongpisanphop et al. 2020).

4.7 Anti-cancer activity

Cancer is the second largest cause of mortality worldwide due to its high incidence rate. Malignant cells kill 15 million people annually, and the number is rising. Cancer can be treated with safe, biocompatible, less toxic, and more resistant natural chemicals from endophytic organisms. These natural chemicals are cancer treatment alternatives to chemotherapeutic medicines. These natural chemicals are anti-cancer and can control many malignancies. Due to their abundance, they can be employed to treat cancer. Endophytic fungi including *Taxomyces andreanae*, *Seimatoantlerium nepalense*, *Alternaria alternate*, and *Chaetomellaphigera* have been linked to the production of the anti-cancer drug paclitaxel (Kousar et al. 2022). Paclitaxel treats Kaposi's sarcoma, prostate, lung, and ovarian cancer (Weaver 2014). It binds to tubulin and inhibits depolymerization during cell division (Leung and Cassimeris 2019). Taxol isolated from sick Chilli plant fruits has shown cytotoxic activity against human cell lines MCF-7, HLK-210, and HL-251. Endophytes *Sinopodophyllum hexandrum* and *Dysosma veitchii* generate podophyllotoxin which is used to treat leukemia, testicular, prostate, lung, and ovarian cancer (Leung and Cassimeris 2019). Camptothecin is an effective cytotoxic compound for the treatment of solid tumors of the liver, bladder, lungs, and ovaries. A study found that

camptothecin (extracted from *A. niger*) caused apoptosis when given to colon cancer cell lines, and cell death occurs at dosages as low as 7.8 mg/L and as high as 1000 mg/L with the highest and lowest cell viability occurring at concentrations of 11.85 and 65.13%, respectively (Aswani and Soundhari 2018). *Fusarium oxysporum* generates the anti-cancer compound vinblastine in *Cathranthus roseus*, which is helpful against lymphoblastic leukemia and cancer cell lines HepG-2 at 7.48 g/mL (Kousar et al. 2022).

4.8 Immunosuppressive activity

An immediate quench strategy to overcome graft rejection and autoimmune diseases; researchers have been looking for a substance that might dampen the immune system (Rajamanikyam et al. 2017). Fungal endophytes can synthesize some substances with immunosuppressive effects (Egbuta et al. 2017). Synthetic chemical immunosuppressive medications have serious side effects because of the time they took to treat diseases. Infection risks like hyperlipidemia, nephrotoxicity, hypertension, and neurotoxicity have side effects from long-term usage of chemical immunosuppressive medications (Hošková et al. 2017). A recent study has revealed that immunosuppressive medicines produced from fungal endophytes are highly effective. Sydoxanthones A and B, 13-O-acetylsydowinin B, colutellin A, methyl peniphenone, dibenzofurane, xanthone derivatives, subglutinols A and B, peniphenone, lipopeptide, benzophenone derivatives, (-) mycousnine, etc. are used in the treatments. There was speculation that cyclosporin A, an immunosuppressant medication, originated in the fungus *Tolypocladium flatum* (El-Gowelli and El-Mas 2015). cyclosporin A, an extract from the endophytic soil fungus *Trichoderma polysporum*, was discovered as a critical immunosuppressive agent. Endophytic fungus *Fusarium subglutinatum* was discovered in *Tripterygium wilfordii*, where it produces noncytotoxic diterpene pyrenes and immunosuppressive Subglutinol A and B (Vasundhara et al. 2016; Adeleke et al. 2021).

4.9 Anti-diabetic activity

Recent research has suggested that endophytic fungus might be a source of substances with anti-diabetic properties. The endophytic fungus *Nigrospora oryzae* in *Combretum dolichopetalum* leaves has been proven to lower fasting blood sugar in diabetic mice when its purified components are administered (Uzor et al. 2017). These chemicals include abscisic acid, 70-hydroxy abscisic acid, and 4-des-hydroxyl alter-solanol A. Indrianingsih and Tachibana (2017) showed that 8-hydroxy-6,7-dimethoxy-3-methyl isocoumarins exhibit strong glucosidase inhibitory effect and are produced by the endophytic fungus *Xylariaceae* spp. in the stem of *Quercus gilva* Blume. Oral administration of glucose and alloxan to

Wistar albino rats accompanied with *Salvadoraoleoides* extracts of *Phoma* spp. and *Aspergillus* spp. induced anti-diabetic and hypolipidemic effects (Ezekwesili and Ogbunugafor 2015). Endophytic fungi isolated from medicinal plants such as *Rauwolfia densiflora* and *Leucas ciliata* have been tested as a potential therapeutics for diabetes through bioprospecting. There is evidence that compounds generated from *Fusarium* spp. and *Alternaria* spp. exhibit anti-diabetic action, suggesting that these fungal endophytes may serve as a source of multifunctional therapies (Adeleke et al. 2021).

4.9.1 Anti-malarial activity

Due to the rapid spread of anti-drug resistance malaria parasites in recent years, there is an urgent need for novel malaria therapy drugs. It was shown that the anti-malarial activity of two endophytic fungi, munumbicins E-4 and E-5, was twice as potent as that of chloroquine (Fadiji and Babalola 2020). The endophyte *Diaporthemiriciae* is responsible for producing the secondary metabolite epoxy cytochalasin H. This compound exhibits robust anti-malarial suppression against a strain of *Plasmodium falciparum* resistant to chloroquine (Ferreira et al. 2017). Ateba et al. (2018) found that the endophyte species *Paecilomyces lilcinus* and *Penicillium janthinellum* are great resources for new compounds that are active against *Plasmodium falciparum* and show potential in the treatment of malaria. It has been proven that various endophytic fungi, in addition to *Aspergillus niger*, *Fusarium* spp., and *Nigrospora* spp. can produce bioactive compounds with an antiplasmodial effect against *Plasmodium falciparum* (Kaushik et al. 2014).

4.9.2 Antituberculosis

Tuberculosis (TB), an infection of the lungs caused by *Mycobacterium tuberculosis*, is a worldwide health concern. It has lasted for centuries and is one of the world's most devastating illnesses. The World Health Organization (WHO) estimates that 10 million individuals are currently living with TB. Ending the TB pandemic by 2030 is one of the health aims of the sustainable development goals of the United Nations. *M. tuberculosis* has been proven to acquire resistance to numerous synthetic medications. To this end, it is crucial to keep looking for new, natural antimycobacterial medicines that may kill mycobacteria without causing resistance. Bioprospecting for fungal endophytes as a cure for TB is an exciting field of study since many fungal metabolites are naturally antimycobacterial. *Azadirachta indica* and *Parthenium hysterophorus* fungal endophytes have been found to exhibit antibacterial effects against TB (Mane et al. 2017). *Phomopsis* spp., an endophytic fungus isolated from *Garcinia* spp. is responsible for the production of Phomoxanthone A and B, which have been shown to suppress the development of *M. tuberculosis*

(Kumar et al. 2017). *Alternaria alternata* and *Phomopsis* spp., two fungal endophytes isolated from Thai medicinal plants, have been shown to generate 3-nitro propionic acid and tenuazonic acid, exhibit indecisive action against *M. tuberculosis* (H37Ra) (Kumar et al. 2017; Deshmukh et al. 2015). Endophytic fungus *Phomopsis* spp. isolated from *Garcinia adulcis* generates bioactive metabolites with anti-tuberculosis potential (Kumar et al. 2017). These metabolites include phomoeamide and phomonitroester. Benzopyran, diaporthone A and B are bioactive chemicals produced by *Diaporthe* spp. They are associated with the leaves of *Pandanus amaryllifolius* and suppress aggressive strains of *M. tuberculosis* (Chepkirui and Stadler 2017).

4.9.3 Antiviral

Evidence suggests that endophytic fungi can create antiviral drugs that are effective against a wide range of viruses, including HIV (Farooq et al. 2016), human CMV (Raekiansyah et al. 2017), Dengue virus (Liu et al. 2019), and influenza A (H1N1) virus (Ambele et al. 2020). Antiviral activity has been found in two novel substances, cytonic acid A and cytonic acid B, which are isolated from *Cytospora* spp. With the use of mass spectrometry and nuclear magnetic resonance, the structures of pteridines isomers were conferred, leading to the identification of novel inhibitors of the protease activity of human cytomegalovirus. Fungal endophytes in the phyllosphere (leaves) of an oak tree (*Quercus coccifera*) generate the antiviral chemical Hinnuliquinone, which has been linked to the inhibition of HIV-1 protease activity (Adeleke et al. 2021). *Alternaria tenuissima* QUE1Se is an endophytic fungus that generates altertoxins, a substance with potent anti-HIV-1 action. In addition to emerimidine (A, B), dehydroaustin, austinol, aspernidine (A, B), austin, emeriphenolicins (A, D), and acetoxyldehydroaustin, many other compounds isolated from *Emericella* spp. (HKZJ) have been found to have antiviral activity against the influenza A virus (H1N1) (Raekiansyah et al. 2017). The antiviral activity of most medicinal plant mixtures is relatively high, even in their crudest forms. Antiviral activity in certain actinomycetes has been demonstrated (Fadiji and Babalola 2020). The antiviral chemical 2-(furan-2-yl)-6-(2S, 3S, 4-trihydroxybutyl) pyrazine was initially isolated from the plant species *Jishengella endophytica* 161,111. This chemical is adequate to combat the spread of the influenza A (H1N1) virus (Fadiji and Babalola 2020; Raekiansyah et al. 2017).

4.9.4 Other pharmacological potentials of the endophytic fungal secondary metabolites

In addition to their potential uses in food, agriculture, medicine, and cosmetics, the bioactive metabolites found in endophytes are excellent pharmaceutical sources for

treating various disease conditions (Shukla et al. 2014). The metabolites generated by fungal endophytes contain a wide variety of functional groups including alkaloids, flavonoids, terpenoids, phenolic acids, quinones, steroids, benzopyranones, tannins, tetralones, and chinones (Gouda et al. 2016). Half of all deaths worldwide are attributable to infectious and parasitic disorders. It has been established that endophytes are the origin of a wide variety of commercially accessible bioactive chemicals and secondary metabolites. New compounds produced by endophytic microbes have shown promise as antibiotics, anti-inflammatory agents, cancer treatments, immunosuppressants, anti-diabetic activity, anti-malarial activity, and even insecticides (Fadiji and Babalola 2020).

5 Conclusion

The purpose of this article is to gather updated knowledge on secondary metabolites of endophytic fungi, their production, methods of analysis, pharmaceutical potential, and application. Microorganisms that are endophytic to a plant provide benefits to the host plant and stimulate plant growth via several direct and indirect mechanisms of action. Fungal endophytes represent an inexhaustible reservoir of pharmacologically essential compounds. Endophytic fungi are an essential component for the production of novel biomolecules for the biochemical and pharmaceutical industries. The fungal endophytes are in a pivotal position in producing certain enzymes, such as amylase, cellulase, laccase, lyase, etc., that have significant commercial and pharmaceutical applications. Several promising pharmaceutical lead molecules have been reported and derived from endophytic fungi. Because they produce physiologically active metabolites that are immune suppressants, anti-cancer agents, promote plant growth, anti-microbial volatiles, anti-oxidants, and antibiotics. Future insights are necessary to understand more about dynamic fungal endophytes, host interactions, and molecular players of fungal endophytes involved in producing biopharmaceuticals of human interest. Finally, the use of metagenomics in combination with next-generation sequencing technologies is anticipated to unlock Endophytes have the potential to offer at least partial, if not total, answers to critical concerns such as rising strain on the global food supply, climate change, environmental degradation, and therapeutics. As a result, the biology of endophytes has to be investigated more if we are going to reap the advantages of these organisms in the fields of agriculture, industry, and medicine. Information on bioactive natural compounds found in endophytic fungi has not been explored effectively. The reservoir of pharmacologically important chemicals that may be found in fungal endophytes is almost endless. To get a better understanding of the dynamic fungal endophytes, the host relationships, and the

molecular actors of fungal endophytes that are engaged in the production of biopharmaceuticals of human interest, more research is required. In conclusion, the use of metagenomics in conjunction with next-generation sequencing technologies is projected to open a wide variety of hitherto undiscovered pools of antimicrobials that are released by endophytic microorganisms that have not yet been farmed. Endophytes have just come to be seen as an important contributor to the overall pool of biological variety and numerous bioactive molecules secreted by as-yet-uncultivated endophytic microbes are not explored. The resources that endophytes give may be used for a variety of purposes, including formulations and bioprospecting. Because of this, further research into the biology of endophytes is required if we want to benefit from the presence of these organisms in the sectors of agriculture, industry, and medicine. The discovery of bioactive natural chemicals discovered in endophytic fungus has had an indelible impact on the treatment of a variety of diseases, including cancer, diabetes, and neurological conditions. With the use of cutting-edge biotechnology like genetic engineering and the microbial fermentation process, these microbial resources may be better utilized for human benefit.

Acknowledgements The authors thank the parent institute NIT Raipur (CG), India, for continuous support and assistance during scientific writing.

Declarations

Ethics declarations Not Required.

Consent for publication Not applicable.

Conflict of interest The authors declare no conflict of interest.

References

- Abel EL (2013) Marihuana: the first twelve thousand years. Springer Science & Business Media.
- Abo Elsoud MM, Mohamed SS, Selim MS, Sidkey NM (2023) Characterization and Optimization of Chitosan Production by *Aspergillus terreus*. Arab J Sci Eng 48:93–106. <https://doi.org/10.1007/s13369-022-07163-z>
- Adeleke BS, Babalola OO, Glick BR (2021) Plant growth-promoting root-colonizing bacterial endophytes. Rhizosphere 20:100433. <https://doi.org/10.1016/j.rhisp.2021.100433>
- Aishwarya S, Venkateswarlu N, Chandramouli K et al (2014) Role of endophytic fungi in the restoration of heavy metal contaminated soils. Indo Am j Pharm Sci 4:5427–5436
- Alam B, Li J, Gě Q, Khan MA, Gōng J, Mehmood S, Yuán Y, Gōng W (2021) Endophytic fungi: from symbiosis to secondary metabolite communications or vice versa? Front Plant Sci 12:791033. <https://doi.org/10.3389/fpls.2021.791033>
- Alori ET, Glick BR, Babalola OO (2017) Microbial Phosphorus Solubilization and Its Potential for Use in Sustainable Agriculture.

- Front Microbiol 8:971. <https://doi.org/10.3389/fmicb.2017.00971>
- Ambele CF, Ekesi S, Bisseleua HDB et al (2020) Entomopathogenic fungi as endophytes for biological control of subterranean termite pests attacking cocoa seedlings. *J Fungi* 6:126. <https://doi.org/10.3390/jof6030126>
- Arora J, Ramawat KG (2017) An introduction to endophytes. In: Maheshwari D. (eds) *Endophytes: Biology and Biotechnology. Sustainable Development and Biodiversity*, vol 15. Springer, Cham. https://doi.org/10.1007/978-3-319-66541-2_1
- Aswani A, Soundhari C (2018) Production of camptothecin from endophytic fungi and characterization by high-performance liquid chromatography and anticancer activity against colon cancer cell line. *Asian J Pharm Clin Res* 11:166. <https://doi.org/10.22159/ajpcr.2018.v11i3.18921>
- Ateba J, Togheue R, Awantu A et al (2018) Antiplasmodial properties and cytotoxicity of endophytic fungi from symphonia globulifera (clusiaceae). *J Fungi* 4:70. <https://doi.org/10.3390/jof4020070>
- Bhardwaj A, Sharma D, Jadon N, Agrawal PK (2015) Antimicrobial and phytochemical screening of endophytic fungi isolated from spikes of *Pinus roxburghii*. *Arch Clin Microbiol* 6(3):1
- Bhattacharya M, Dhama K, Chakraborty C (2022) Recently spreading human monkeypox virus infection and its transmission during COVID-19 pandemic period: A travelers' prospective. *Travel Med Infect Dis* 49:102398. <https://doi.org/10.1016/j.tmaid.2022.102398>
- Bisht S, Pandey P, Bhargava B et al (2015) Bioremediation of polycyclic aromatic hydrocarbons (PAHs) using rhizosphere technology. *Braz J Microbiol* 46:7–21. <https://doi.org/10.1590/S1517-838246120131354>
- Bisht R, Sharma D, Agrawal PK (2016) Antagonistic and antibacterial activity of endophytic fungi isolated from needle of *Cupressus torulosa* D. Don. *Asian J Pharm Clin Res* 9:282–288
- Biswas D, Nazir R, Biswas P et al (2020) Endophytic sources of diosgenin, a natural steroid with multiple therapeutic values. *S Afr J Bot* 134:119–125. <https://doi.org/10.1016/j.sajb.2020.04.009>
- Bonaterre A, Badosa E, Daranas N et al (2022) Bacteria as biological control agents of plant diseases. *Microorganisms* 10:1759. <https://doi.org/10.3390/microorganisms10091759>
- Brader G, Compant S, Mitter B et al (2014) Metabolic potential of endophytic bacteria. *Curr Opin Biotechnol* 27:30–37. <https://doi.org/10.1016/j.copbio.2013.09.012>
- Bunaciu AA, Udriștioiu EG, Aboul-Enein HY (2015) X-Ray Diffraction: Instrumentation and Applications. *Crit Rev Anal Chem* 45:289–299. <https://doi.org/10.1080/10408347.2014.949616>
- Cárdenas G, Salgado P, Laura-Foronda E, Popoca-Rodríguez I, Delgado-Hernández RD, Rojas R, Palacios E (2021) Neglected and (re-)emergent infections of the CNS in low-/middle-income countries. *Infez Med* 29(4):513–525. <https://doi.org/10.53854/liim-2904-3>
- Chandran H, Meena M, Barupal T, Sharma K (2020) Plant tissue culture as a perpetual source for production of industrially important bioactive compounds. *Biotechnol Rep* 26:e00450. <https://doi.org/10.1016/j.btre.2020.e00450>
- Chatelain E (2015) Chagas disease drug discovery: Toward a New Era. *SLAS Discov* 20:22–35. <https://doi.org/10.1177/1087057114550585>
- Chen H-Y, Xue D-S, Feng X-Y, Yao S-J (2011) Screening and production of ligninolytic enzyme by a marine-derived fungal *Pestalotiopsis* sp. J63. *Appl Biochem Biotechnol* 165:1754–1769. <https://doi.org/10.1007/s12010-011-9392-y>
- Cheng M-J, Wu M-D, Hsieh S-Y et al (2011) Secondary metabolites from the endophytic fungus *Annulohyphoxylon boveri* var. *microspora* BCRC 34012. *Chem Nat Compd* 47:536–540. <https://doi.org/10.1007/s10600-011-9990-5>
- Chepkirui C, Stadler M (2017) The genus *Diaporthe*: a rich source of diverse and bioactive metabolites. *Mycol Prog* 16:477–494. <https://doi.org/10.1007/s11557-017-1288-y>
- Chithra S, Jasim B, Sachidanandan P et al (2014) Piperine production by endophytic fungus *Colletotrichum gloeosporioides* isolated from *Piper nigrum*. *Phytomedicine* 21:534–540. <https://doi.org/10.1016/j.phymed.2013.10.020>
- Chowdappa S, Jagannath S, Konappa N et al (2020) Detection and characterization of antibacterial siderophores secreted by endophytic fungi from *Cymbidium alofolium*. *Biomolecules* 10:1412. <https://doi.org/10.3390/biom10101412>
- Cui Y, Yi D, Bai X et al (2012) Ginkgolide B produced endophytic fungus (*Fusarium oxysporum*) isolated from *Ginkgo biloba*. *Fito-terapia* 83:913–920. <https://doi.org/10.1016/j.fitote.2012.04.009>
- Daguette Y, Edel-Hermann V, Steinberg C (2017) Fungal genes and metabolites associated with the biocontrol of soil-borne plant pathogenic fungi. In: Mérillon J-M, Ramawat KG (eds) *Fungal metabolites*. Springer International Publishing, Cham, Switzerland, pp 33–104
- Damam M, Kaloori K, Gaddam B et al (2016) Plant growth promoting substances (phytohormones) produced by rhizobacterial strains isolated from the rhizosphere of medicinal plants. *Int J Pharm Sci Rev Res* 37:130–136
- Das M, Prakash HS, Nalini MS (2017) Antioxidative properties of phenolic compounds isolated from the fungal endophytes of *Zingiber nimmonii* (J. Graham) Dalzell. *Front Biol (Beijing)* 12:151–162. <https://doi.org/10.1007/s11515-016-1441-z>
- Debbab A, Aly AH, Proksch P (2013) Mangrove derived fungal endophytes – a chemical and biological perception. *Fungal Divers* 61:1–27. <https://doi.org/10.1007/s13225-013-0243-8>
- Deepthi VC, Sumathi S, Faisal M et al (2018) Isolation and identification of endophytic fungi with anti-microbial activities from the leaves of *Elaeocarpus sphaericus* (gaertn.) K. Schum. and *Myristica fragrans* houtt. *Int J Pharm Sci Res* 9:2783–2791. [https://doi.org/10.13040/IJPSR.0975-8232.9\(7\).2783-91](https://doi.org/10.13040/IJPSR.0975-8232.9(7).2783-91)
- Deshmukh SK, Verekar SA, Bhavé SV (2015) Endophytic fungi: a reservoir of antibacterials. *Front Microbiol* 5:715. <https://doi.org/10.3389/fmicb.2014.00715>
- Duc NH, Csintalan Z, Posta K (2018) Arbuscular mycorrhizal fungi mitigate negative effects of combined drought and heat stress on tomato plants. *Plant Physiol Biochem* 132:297–307. <https://doi.org/10.1016/j.plaphy.2018.09.011>
- Egbuta M, Mwanza M, Babalola O (2017) Health risks associated with exposure to filamentous fungi. *Int J Environ Res Public Health* 14:719. <https://doi.org/10.3390/ijerph14070719>
- Eid AM, Fouda A, Abdel-Rahman MA et al (2021) Harnessing bacterial endophytes for promotion of plant growth and biotechnological applications: An overview. *Plants* 10:935. <https://doi.org/10.3390/plants10050935>
- El-Gowelli HM, El-Mas MM (2015) Central modulation of cyclosporine-induced hypertension. *Naunyn Schmiedeberg Arch Pharmacol* 388:351–361. <https://doi.org/10.1007/s00210-014-1074-1>
- Etesami H, Alikhani HA, Hosseini HM (2015) Indole-3-acetic acid (IAA) production trait, a useful screening to select endophytic and rhizosphere competent bacteria for rice growth promoting agents. *MethodsX* 2:72–78. <https://doi.org/10.1016/j.mex.2015.02.008>
- Ezekwesili CN, Ogbunugafor HA (2015) Blood glucose lowering activity of five Nigerian medicinal plants in alloxan-induced diabetic Wistar albino rats. *Anim Res Int* 12:2150–2158. <https://doi.org/10.4314/ARI.V12I2>
- Fadiji AE, Babalola OO (2020) Elucidating mechanisms of endophytes used in plant protection and other bioactivities with multifunctional prospects. *Front Bioeng Biotechnol* 8:467. <https://doi.org/10.3389/fbioe.2020.00467>

- Farooq T, Hameed A, Rehman K et al (2016) Antiretroviral agents: Looking for the best possible chemotherapeutic options to conquer HIV. *Crit Rev Eukaryot Gene Expr* 26:363–381. <https://doi.org/10.1615/CritRevEukaryotGeneExpr.2016018255>
- Ferreira MC, Cantrell CL, Wedge DE et al (2017) Antimycobacterial and antimalarial activities of endophytic fungi associated with the ancient and narrowly endemic neotropical plant *Vellozia gigantea* from Brazil. *Mem Inst Oswaldo Cruz* 112:692–697. <https://doi.org/10.1590/0074-02760170144>
- Flury P, Vesga P, Péchy-Tarr M, Aellen N, Dennert F, Hofer N, Kupferschmied KP, Kupferschmied P, Metla Z, Ma Z, Siegfried S, de Weert S, Bloemberg G, Höfte M, Keel CJ, Maurhofer M (2017) Antimicrobial and Insecticidal: Cyclic Lipopeptides and Hydrogen Cyanide Produced by Plant-Beneficial *Pseudomonas* Strains CHA0, CMR12a, and PCL1391 Contribute to Insect Killing. *Front Microbiol* 8:100. <https://doi.org/10.3389/fmicb.2017.00100>
- Gamalero E, Favale N, Bona E et al (2020) Screening of bacterial endophytes able to promote plant growth and increase salinity tolerance. *Appl Sci* 10:5767. <https://doi.org/10.3390/app10175767>
- Gao H, Li G, Lou H-X (2018) Structural diversity and biological activities of novel secondary metabolites from endophytes. *Molecules* 23:646. <https://doi.org/10.3390/molecules23030646>
- Gokarn K, Pal R (2018) Activity of siderophores against drug-resistant Gram-positive and Gram-negative bacteria. *Infect Drug Resist* 11:61–75. <https://doi.org/10.2147/IDR.S148602>
- Gouda S, Das G, Sen SK, Shin HS, Patra JK (2016) Endophytes: a treasure house of bioactive compounds of medicinal importance. *Front Microbiol* 7:1538. <https://doi.org/10.3389/fmicb.2016.01538>
- Haile S, Ayele A (2022) Pectinase from microorganisms and its industrial applications. *Sci World J* 2022:1–15. <https://doi.org/10.1155/2022/1881305>
- Hardoim PR, van Overbeek LS, Berg G et al (2015) The hidden world within plants: Ecological and evolutionary considerations for defining functioning of microbial endophytes. *Microbiol Mol Biol Rev* 79:293–320. <https://doi.org/10.1128/MMBR.00050-14>
- Hořková L, Málek I, Kopkan L, Kautzner J (2017) Pathophysiological mechanisms of calcineurin inhibitor-induced nephrotoxicity and arterial hypertension. *Physiol Res* 66(2):167–180. <https://doi.org/10.33549/physiolres.933332>
- Hussain H, Root N, Jabeen F, Al-Harrasi A, Ahmad M, Mabood F, Hassan Z, Shah A, Green IR, Schulz B (2015) Microsphaerol and seimatorone: two new compounds isolated from the endophytic fungi, *Microsphaeropsis* sp. and *Seimatosporium* sp. *Chem Biodivers* 12(2):289–294
- Indrianingsih AW, Tachibana S (2017) α -Glucosidase inhibitor produced by an endophytic fungus, *Xylariaceae* sp. QGS 01 from *Quercus gilva* Blume. *Food Sci Human Wellness* 6:88–95. <https://doi.org/10.1016/j.fshw.2017.05.001>
- Kandel S, Joubert P, Doty S (2017) Bacterial endophyte colonization and distribution within plants. *Microorganisms* 5:77. <https://doi.org/10.3390/microorganisms5040077>
- Karim KMR, Husaini A, Tasnim T (2017) Production and characterization of crude glucoamylase from newly isolated *Aspergillus flavus* NSH9 in liquid culture. *Am J Biochem Mol Biol* 7:118–126. <https://doi.org/10.3923/ajbmb.2017.118.126>
- Kaur P, Kumar V, Singh R et al (2020) Biotechnological strategies for production of camptothecin from fungal and bacterial endophytes. *S Afr J Bot* 134:135–145. <https://doi.org/10.1016/j.sajb.2020.07.001>
- Kaushik NK, Murali TS, Sahal D, Suryanarayanan TS (2014) A search for antiparasitoid metabolites among fungal endophytes of terrestrial and marine plants of southern India. *Acta Parasitol* 59(4):745–57. <https://doi.org/10.2478/s11686-014-0307-2>
- Khan A, Khan S, Khan MA et al (2015) The uptake and bioaccumulation of heavy metals by food plants, their effects on plants nutrients, and associated health risk: a review. *Environ Sci Pollut Res* 22:13772–13799. <https://doi.org/10.1007/s11356-015-4881-0>
- Kousar R, Naeem M, Jamaludin MI et al (2022) Exploring the anticancer activities of novel bioactive compounds derived from endophytic fungi: Mechanisms of action, current challenges and future perspectives. *Am J Cancer Res* 12:2897
- Krings M, Taylor TN, Hass H et al (2007) Fungal endophytes in a 400-million-yr-old land plant: infection pathways, spatial distribution, and host responses. *New Phytol* 174:648–657. <https://doi.org/10.1111/j.1469-8137.2007.02008.x>
- Kumar M, Mathur A (2017) First study on anti-microbial activities of endophytes isolated from aerial parts of *Mentha Piperita*. *Int J Sci Res Publ* 7:90–99
- Kumar A, Patil D, Rajamohanam PR, Ahmad A (2013) Isolation, purification and characterization of vinblastine and vincristine from endophytic fungus *fusarium oxysporum* isolated from *catharanthus roseus*. *PLoS One* 8:e71805. <https://doi.org/10.1371/journal.pone.0071805>
- Kumar G, Chandra P, Choudhary M (2017) Endophytic fungi: A potential source of bioactive compounds. *Chem Sci Rev Lett* 6:2373–2381
- Laga V, Vasilyev A, Lapovok I et al (2015) HIV type 1 subtype A1 dominates in Armenia. *Curr HIV Res* 13:219–225. <https://doi.org/10.2174/1570162X13666150407142834>
- Le Thi Minh T, Thi Hong Hoang A, Le Thi Bich T et al (2019) Isolation of endophytic fungi and screening of Huperzine A-producing fungus from *Huperzia serrata* in Vietnam. *Sci Rep* 9:16152. <https://doi.org/10.1038/s41598-019-52481-2>
- Leung JC, Cassimeris L (2019) Reorganization of paclitaxel-stabilized microtubule arrays at mitotic entry: roles of depolymerizing kinesins and severing proteins. *Cancer Biol Ther* 20:1337–1347. <https://doi.org/10.1080/15384047.2019.1638678>
- Li H-Y, Wei D-Q, Shen M, Zhou Z-P (2012) Endophytes and their role in phytoremediation. *Fungal Divers* 54:11–18. <https://doi.org/10.1007/s13225-012-0165-x>
- Liu S-S, Jiang J-X, Huang R et al (2019) A new antiviral 14-nordrimane sesquiterpenoid from an endophytic fungus *Phoma* sp. *Phytochem Lett* 29:75–78. <https://doi.org/10.1016/j.phytol.2018.11.005>
- Lugtenberg BJJ, Caradus JR, Johnson LJ (2016) Fungal endophytes for sustainable crop production. *FEMS Microbiol Ecol* 92:fiw194. <https://doi.org/10.1093/femsec/fiw194>
- Ma L, Cao YH, Cheng MH et al (2013) Phylogenetic diversity of bacterial endophytes of *Panax notoginseng* with antagonistic characteristics towards pathogens of root-rot disease complex. *Antonie Van Leeuwenhoek* 103:299–312. <https://doi.org/10.1007/s10482-012-9810-3>
- Madhusudhan M, Bharathi T, Prakash H (2015) Isolation and purification of bioactive metabolites from fungal endophytes— a review. *Curr Biochem Eng* 2:111–117. <https://doi.org/10.2174/2212711902999150706114209>
- Mane R, Shinde M, Wagh P (2017) Isolation of Endophytic microorganisms as a source of novel secondary metabolite producers against tuberculosis. *Int J Sci Res Technol* 3:1267–1269
- Mathew BB, Jatawa SK, Tiwari A (2012) Phytochemical analysis of Citrus limonum pulp and peel. *Int J Pharm Pharm Sci* 4:369–71
- Mathew BB, Jatawa SK, Tiwari A (2012) Phytochemical analysis of citrus limonum pulp and peel. *Int J Pharm Pharm Sci* 4(2)
- Mengistu AA (2020) Endophytes: Colonization, behaviour, and their role in defense mechanism. *Int J Microbiol* 2020:1–8. <https://doi.org/10.1155/2020/6927219>
- Mishra R, Kushveer JS, Revanthbabu P, Sarma VV (2019) Endophytic fungi and their enzymatic potential. In: Singh B (ed) *Advances in endophytic fungal research: present status and future challenges*. Springer, Cham, pp 283–337

- Mrudula S, Murugammal R (2011) Production of cellulase by *Aspergillus niger* under submerged and solid state fermentation using coir waste as a substrate. *Braz J Microbiol* 42:1119–1127. <https://doi.org/10.1590/S1517-83822011000300033>
- Nanda S, Mohanty B, Joshi RK (2019) Endophyte-mediated host stress tolerance as a means for crop improvement. *Reference Series in Phytochemistry* 677–701. https://doi.org/10.1007/978-3-319-76900-4_28-1
- Pang Z, Chen J, Wang T, Gao C, Li Z, Guo L, Xu J, Cheng Y (2021) Linking Plant Secondary Metabolites and Plant Microbiomes: A Review. *Front Plant Sci* 12:621276. <https://doi.org/10.3389/fpls.2021.621276>
- Parthasarathy R, Sathiyabama M (2015) Lovastatin-producing endophytic fungus isolated from a medicinal plant *Solanum xanthocarpum*. *Nat Prod Res* 29:2282–2286. <https://doi.org/10.1080/14786419.2015.1016938>
- Patil RH, Patil MP, Maheshwari VL (2016) Bioactive secondary metabolites from endophytic fungi. *Studies in Natural Products Chemistry* 189–205. <https://doi.org/10.1016/B978-0-444-63601-0.00005-3>
- Peralta DR, Adler C, Corbalán NS et al (2016) Enterobactin as Part of the Oxidative Stress Response Repertoire. *PLoS One* 11:e0157799. <https://doi.org/10.1371/journal.pone.0157799>
- Pietro-Souza W, de Campos Pereira F, Mello IS et al (2020) Mercury resistance and bioremediation mediated by endophytic fungi. *Chemosphere* 240:124874. <https://doi.org/10.1016/j.chemosphere.2019.124874>
- Pokhrel S, Chhetri R (2021) A literature review on impact of COVID-19 pandemic on teaching and learning. *High Educ Future* 8:133–141. <https://doi.org/10.1177/2347631120983481>
- Qadri H, Shah AH, Mir M (2021) Novel strategies to combat the emerging drug resistance in human pathogenic microbes. *Curr Drug Targets* 22:1424–1436. <https://doi.org/10.2174/1389450121666201228123212>
- Qi F-H, Jing T-Z, Wang Z-X, Zhan Y-G (2009) Fungal endophytes from *Acer ginnala* Maxim: isolation, identification and their yield of gallic acid. *Lett Appl Microbiol* 49:98–104. <https://doi.org/10.1111/j.1472-765X.2009.02626.x>
- Raekiansyah M, Mori M, Nonaka K et al (2017) Identification of novel antiviral of fungus-derived brefeldin A against dengue viruses. *Trop Med Health* 45:32. <https://doi.org/10.1186/s41182-017-0072-7>
- Raghunath R, Radhakrishna A, Angayarkanni J, Palaniswamy M (2012) Production and cytotoxicity studies of lovastatin from *Aspergillus niger* PN2: an endophytic fungi isolated from *Taxus baccata*. *Int J Appl Biol Pharm* 3(3):342–351
- Rajamanikyam M, Vadlapudi V, amanchy R, Upadhyayula SM, (2017) Endophytic Fungi as Novel Resources of natural Therapeutics. *Brazilian Archives of Biology and Technology* 60:1–26. <https://doi.org/10.1590/1678-4324-2017160542>
- Rana KL, Kour D, Sheikh I et al (2019) Endophytic Fungi: biodiversity, ecological significance, and potential industrial applications. *Recent Advancement in White Biotechnology Through Fungi* 1–62. https://doi.org/10.1007/978-3-030-10480-1_1
- Rodriguez RJ, White JF Jr, Arnold AE, Redman RS (2009) Fungal endophytes: diversity and functional roles. *New Phytol* 182:314–330. <https://doi.org/10.1111/j.1469-8137.2009.02773.x>
- Sadrati N, Daoud H, Zerroug A et al (2013) Screening of antimicrobial and antioxidant secondary metabolites from endophytic fungi isolated from wheat (*triticum durum*). *J Plant Prot Res* 53:128–136. <https://doi.org/10.2478/jppr-2013-0019>
- Santos MLD, Berlitz DL, Wiest SLF et al (2018) Benefits Associated with the Interaction of Endophytic Bacteria and Plants. *Brazilian Archives of Biology and Technology* 61:1–11. <https://doi.org/10.1590/1678-4324-2018160431>
- Santoyo G, Moreno-Hagelsieb G, del Carmen Orozco-Mosqueda Ma, Glick BR (2016) Plant growth-promoting bacterial endophytes. *Microbiol Res* 183:92–99. <https://doi.org/10.1016/j.micres.2015.11.008>
- Satapute P, Kamble MV, Adhikari SS, Jogaiah S (2019) Influence of triazole pesticides on tillage soil microbial populations and metabolic changes. *Sci Total Environ* 651:2334–2344. <https://doi.org/10.1016/j.scitotenv.2018.10.099>
- Shaffique S, Khan MA, Wani SH et al (2022) A review on the role of endophytes and plant growth promoting rhizobacteria in mitigating heat stress in plants. *Microorganisms* 10:1286. <https://doi.org/10.3390/microorganisms10071286>
- Shahzad R, Waqas M, Khan AL et al (2016) Seed-borne endophytic *Bacillus amyloliquefaciens* RWL-1 produces gibberellins and regulates endogenous phytohormones of *Oryza sativa*. *Plant Physiol Biochem* 106:236–243. <https://doi.org/10.1016/j.plaphy.2016.05.006>
- Shankar Naik B (2019) Developments in taxol production through endophytic fungal biotechnology: a review. *Orient Pharm Exp Med* 19:1–13. <https://doi.org/10.1007/s13596-018-0352-8>
- Sharma D, Pramanik A, Agrawal PK (2016) Evaluation of bioactive secondary metabolites from endophytic fungus *Pestalotiopsis neglecta* BAB-5510 isolated from leaves of *Cupressus torulosa* D. Don. *3 Biotech* 6:210. <https://doi.org/10.1007/s13205-016-0518-3>
- Sharma H, Rai AK, Dahiya D et al (2021) Exploring endophytes for *in vitro* synthesis of bioactive compounds similar to metabolites produced *in vivo* by host plants. *AIMS Microbiol* 7:175–199. <https://doi.org/10.3934/microbiol.2021012>
- Sharma P, Singh SP (2021) Role of the endogenous fungal metabolites in the plant growth improvement and stress tolerance. In: *Fungi Bio-prospects in sustainable agriculture, environment and nanotechnology*, vol. 3. Elsevier, pp 381–401
- Shekhawat K, Saad MM, Sheikh A, Mariappan K, Al-Mahmoudi H, Abdulhakim F, Eida AA, Jalal R, Masmoudi K, Hirt H (2021) Root endophyte induced plant thermotolerance by constitutive chromatin modification at heat stress memory gene loci. *EMBO Rep* 22(3):e51049. <https://doi.org/10.15252/embr.202051049>
- Shukla ST, Habbu PV, Kulkarni VH et al (2014) Endophytic microbes: a novel source for biologically/pharmacologically active secondary metabolites. *Asian J Pharmacol Toxicol* 2:1–6
- Singh D, Geat N, Rajawat MVS et al (2018) Prospecting endophytes from different Fe or Zn accumulating wheat genotypes for their influence as inoculants on plant growth, yield, and micronutrient content. *Ann Microbiol* 68:815–833. <https://doi.org/10.1007/s13213-018-1388-1>
- Suryanarayanan TS (2013) Endophyte research: going beyond isolation and metabolite documentation. *Fungal Ecol* 6:561–568. <https://doi.org/10.1016/j.funeco.2013.09.007>
- Talukdar R, Padhi S, Rai AK, Masi M, Evidente A, Jha DK, Cimmino A, Tayung K (2021) Isolation and Characterization of an Endophytic Fungus *Colletotrichum coccodes* Producing Tyrosol From *Houttuynia cordata* Thunb. Using ITS2 RNA Secondary Structure and Molecular Docking Study. *Front Bioeng Biotechnol* 9:650247. <https://doi.org/10.3389/fbioe.2021.650247>
- Teoh ES (2015) Secondary Metabolites of Plants. *Medicinal Orchids of Asia* 5:59–73. https://doi.org/10.1007/978-3-319-24274-3_5
- Tesfaye TS, Magarsa AU, Zeleke TM (2020) Moderate to severe diarrhea and associated factors among under-five children in Wonago District, South Ethiopia: A Cross-sectional study. *Pediatric Health Med Ther* 11:437–443. <https://doi.org/10.2147/PHMT.S266828>
- Tiwari K (2015) The future products: Endophytic fungal metabolites. *J Biodivers, Bio Prospect Dev* 2:2–7

- Tiwari P, Bae H (2022) Endophytic fungi: KEY insights, emerging prospects, and challenges in natural product drug discovery. *Microorganisms* 10:360. <https://doi.org/10.3390/microorganisms10020360>
- Trivedi G, Patel P, Saraf M (2020) Synergistic effect of endophytic selenobacteria on biofortification and growth of Glycine max under drought stress. *S Afr J Bot* 134:27–35. <https://doi.org/10.1016/j.sajb.2019.10.001>
- Uzor PF, Osadebe PO, Nwodo NJ (2017) Antidiabetic activity of extract and compounds from an endophytic fungus *Nigrospora oryzae*. *Drug Res* 67:308–311. <https://doi.org/10.1055/s-0042-122777>
- Varga T, Hixson KK, Ahkami AH, Sher AW, Barnes ME, Chu RK, Battu AK, Nicora CD, Winkler TE, Reno LR, Fakra SC, Antipova O, Parkinson DY, Hall JR, Doty SL (2020) Endophyte-promoted phosphorus solubilization in populus. *Front Plant Sci* 11:567918. <https://doi.org/10.3389/fpls.2020.567918>
- Vasundhara M, Baranwal M, Kumar A (2016) *Fusarium tricinctum*, An endophytic fungus exhibits cell growth inhibition and antioxidant activity. *Indian J Microbiol* 56:433–438. <https://doi.org/10.1007/s12088-016-0600-x>
- Vigneshwari A, Rakk D, Németh A et al (2019) Host metabolite producing endophytic fungi isolated from *Hypericum perforatum*. *PLoS One* 14:e0217060. <https://doi.org/10.1371/journal.pone.0217060>
- Vimal A, Kumar A (2022) optimized production of medically significant enzyme L-asparaginase under submerged and solid-state fermentation from agricultural wastes. *Curr Microbiol* 379(12):394. <https://doi.org/10.1007/s00284-022-03095-x>
- Waghunde RR, Shelake RM, Pramanik D et al (2021) Application of cereal and vegetable endophytes in plant health management. In: *Microbiomes and Plant Health*. Elsevier, pp 29–51
- Weaver BA (2014) How Taxol/paclitaxel kills cancer cells. *Mol Biol Cell* 25:2677–2681. <https://doi.org/10.1091/mbc.e14-04-0916>
- World Health Organization (2020) WHO package of essential non-communicable (PEN) disease interventions for primary health care. NCD Management-Screening, Diagnosis and Treatment 77. <https://www.who.int/publications/i/item/9789240009226>
- Wu M-D, Cheng M-J, Lin R-J et al (2019) Chemical constituents of the fungus *biscogniauxia cylindrospora*. *Chem Nat Compd* 55:924–926. <https://doi.org/10.1007/s10600-019-02848-8>
- Xia Y, Liu J, Chen C et al (2022) The multifunctions and future prospects of endophytes and their metabolites in plant disease management. *Microorganisms* 10:1072. <https://doi.org/10.3390/microorganisms10051072>
- Yadav AN (2018) Biodiversity and biotechnological applications of host-specific endophytic fungi for sustainable agriculture and allied sectors. *Acta Sci Microbiol* 1:01–05. <https://doi.org/10.31080/ASMI.2018.01.0044>
- Yan J, Qi N, Wang S et al (2014) Characterization of secondary metabolites of an endophytic fungus from *Curcuma wenyujin*. *Curr Microbiol* 69:740–744. <https://doi.org/10.1007/s00284-014-0647-z>
- Yongpisanphop J, Babel S, Kurisu F et al (2020) Isolation and characterization of Pb-resistant plant growth promoting endophytic bacteria and their role in Pb accumulation by fast-growing trees. *Environ Technol* 41:3598–3606. <https://doi.org/10.1080/09593330.2019.1615993>
- Zhang FH, Xiang JH, Cui WX et al (2016) (2016) Isolation and identification of berberine from endophytic fungi HL-Y-3. *China J Chin Mater Med*. <https://doi.org/10.4268/cjcmm20161609>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.