

## EPV0368

### A cross sectional observational study on the effect of music on the anxiety state of patients admitted for COVID 19 in a tertiary care hospital in New Delhi

N.A. Jose\*, R. Prakash, P. Bansal and M. Khan

Hamdard Institute of Medical Sciences & Research, Psychiatry,  
New Delhi, India

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1237

**Introduction:** COVID 19 has led to dramatic changes in the lives of people leading to an increase in stress and anxiety. Music intervention is a non-medicated method for relieving anxiety. This current study aims to understand whether music can be effectively used to alleviate anxiety in admitted COVID-19 patients.

**Objectives:** To study the effect of music on anxiety in patients admitted for COVID-19 in a tertiary care hospital in New Delhi

**Methods:** 34 patients (17 females & 17 males) were randomly divided into two groups, a control (N=17) and a music group (N=17). Vitals of all the patients were noted. Patients of the music group were asked to listen to relaxing instrumental for 30 minutes, while patients of the control group were asked to relax for 30 minutes. Vitals of all the patients were noted again. Patients were asked to fill State Trait Anxiety Inventory (STAI) before and after intervention.

**Results:** The post-intervention mean scores of STAI after the music session were lower in the music group than the control group [95.06 (SD 8.5)) versus 102.37 (SD 10.3)]. The differences in mean values of pre- to post-intervention changes between both groups after music session were statistically significant.

**Conclusions:** Our findings suggest that listening to music lowers anxiety. As music is non-invasive and free of side-effects we recommend that music intervention service should be used to improve health care quality.

**Disclosure:** No significant relationships.

**Keywords:** Covid-19; Anxiety; music

## EPV0369

### Ms. Noemi Monika Szeifert

N. Szeifert<sup>1\*</sup>, S. Szilágyi<sup>1</sup> and Á. Schmelowszky<sup>2</sup>

<sup>1</sup>Peterfy Alexander Hospital, Crisis Intervention And Psychiatry, Budapest, Hungary and <sup>2</sup>University of Eötvös Lóránd, Clinical Psychology And Addictology, Budapest, Hungary

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1238

**Introduction: Working in COVID – 19 Health Care Units – Psychological impacts (PTSD, Depression, Anxiety Disorders)** Working in the frontline during COVID-19 has put under extreme psychological and physical pressure the health care workers. The severe psychological symptoms can emerge on short, mid and long term as well. Our research is focusing on the psychological impacts of front line health care workers.

**Objectives:** In our lecture, we summarize our research made in the National Traumatology Center, Budapest, Hungary among health care workers who worked in the front and in the second line during the most severe period of the pandemic in Hungary, and discuss probable risk factors for PTSD and chronic psychological distress related to COVID-19. Furthermore we make an overview on the most frequently used coping skills for dealing with the psychological stress caused by the pandemic among the health care workers. The sample taking was processed in July 2021, after 3 month the 3rd COVID – 19 wave officially ended in Hungary.

**Methods:** 123 health care workers completed the online survey anonymously, included Beck Depression Inventory, Spielberger Anxiety State Inventory, PTSD Checklist, Lazarus Coping Scale.

**Results:** 23% of HCW reported symptoms of depression, nearly 25% dealing with sleeping disorders and 38% with chronic fatigue, 42,50 % HCW suffering from anxiety disorders, 22% HCW experiencing symptoms of PTSD.

**Conclusions:** With our results we would like to raise awareness of the challenges and severe psychological consequences that these colleagues, our national heroes we can say, are facing after working in COVID-19 Health Care Units.

**Disclosure:** No significant relationships.

**Keywords:** COVID-19 Health Care Workers; Psychological distress; vicarious traumatization; PTSD

## EPV0370

### Prevalence and socio-demographic correlates of psychological health problems in inhabitants of the West Pomeranian Voivodeship during the COVID-19 outbreak

A. Cybulska<sup>1\*</sup>, K. Rachubińska<sup>2</sup>, D. Schneider-Matyka<sup>1</sup> and E. Grochans<sup>1</sup>

<sup>1</sup>Pomeranian Medical University, Department Of Nursing, Szczecin, Poland and <sup>2</sup>Pomeranian Medical University in Szczecin, Department Of Nursing, Szczecin, Poland

\*Corresponding author.

doi: 10.1192/j.eurpsy.2022.1239

**Introduction:** Psychological health problems, especially emotional disorders, have become an important topic of considerations for many scientists, because the epidemiology of these disorders is strongly influenced by stressful events, such as the SARS-CoV-2 coronavirus pandemic.

**Objectives:** The aim of this study was to evaluate selected parameters of psychosocial functioning as well as socio-demographic correlates of depression, anxiety, sleep disorders and perceived stress among the residents of the West Pomeranian Voivodeship.

**Methods:** An online questionnaire was completed by 323 participants, in whom the parameters of psychosocial functioning were assessed: symptoms of depression (PHQ-9), anxiety (GAD-7), severity of sleep disorders (AIS) and perceived stress (Perceived Stress Scale).

**Results:** The majority of the respondents (75.2%) scored high on the Perceived Stress Scale, almost half of the respondents (47.1%)