

Thioredoxin-80 is a product of alpha-secretase cleavage that inhibits amyloid-beta aggregation and is decreased in Alzheimer's disease brain

Francisco Gil-Bea^{1†}, Susanne Akterin^{1†}, Torbjörn Persson^{1†}, Laura Mateos¹, Anna Sandebring¹, Javier Avila-Cariño², Angel Gutierrez-Rodriguez³, Erik Sundström⁴, Arne Holmgren⁵, Bengt Winblad¹, Angel Cedazo-Minguez^{1*}

Keywords: ADAM; Alzheimer's disease; amyloid- β ; α -secretase; thioredoxin

DOI 10.1002/emmm.201201462

Received April 12, 2012

Revised July 14, 2012

Accepted July 17, 2012

Thioredoxin-1 (Trx1) is an endogenous dithiol reductant and antioxidant that was shown to be decreased in Alzheimer's disease (AD) neurons. A truncated form of Trx1, thioredoxin 80 (Trx80), was reported to be secreted from monocytes having cytokine activity. Here, we show that Trx80 is present in human brain in an aggregated form. Trx80 localizes mainly to neurons and is dramatically decreased in AD brains. Trx80 levels in cerebrospinal fluid (CSF) correlate with those of the classical AD biomarkers amyloid- β (A β) 1–42 and total tau. Moreover, Trx80 measurements in CSF discriminate between patients with stable mild cognitive impairment, prodromal AD and mild AD. We report that ADAM10 and 17, two α -secretases processing the A β precursor protein, are responsible for Trx80 generation. In contrast to the periphery, Trx80 has no pro-inflammatory effects in glia, either by itself or in combination with A β or apolipoprotein E. Instead, Trx80 inhibits A β (1–42) aggregation and protects against its toxicity. Thus, a reduction in Trx80 production would result in increased A β polymerization and enhanced neuronal vulnerability. Our data suggest that a deficit in Trx80 could participate in AD pathogenesis.

INTRODUCTION

Thioredoxin-1 (Trx1) plays a central role in the control of cellular redox homeostasis and acts as an antioxidant (Arner & Holmgren, 2000). Trx1 has been shown to regulate apoptosis by

inhibiting the apoptosis signal-regulating kinase-1 (ASK1; Saitoh et al, 1998). Trx1, directly or in association with other proteins like Trx1-interacting protein-2 (TBP-2), is known to play important roles in cell growth and survival, as well as in major metabolic diseases and cancer (Watanabe et al, 2010). Oxidative stress, apoptosis and inflammation are also key factors in neurodegenerative diseases, and there is an increasing interest in evaluating the role of Trx1 in such disorders (Lillig & Holmgren, 2007). We previously found that neuronal Trx1 is decreased in Alzheimer's disease (AD) brains and that Trx1 is oxidized by the β -amyloid (A β) peptide (Akterin et al, 2006). A β is the major constituent in the neuritic plaques seen in AD brains, and it is believed to be a driving force of the disease pathology (Hardy & Selkoe, 2002). A β results after sequential cleavage of amyloid precursor protein (APP) by β - and γ -secretases. Normally, a non-amyloidogenic pathway, involving α -secretases, is predominant in brain (Selkoe, 2001). In addition to Trx1, other members of the Trx1 family have been

(1) Department of Neurobiology, KI-Alzheimer's Disease Research Center, Care Sciences and Society, Karolinska Institutet, Huddinge, Sweden

(2) Department of Cell and Molecular Biology, Karolinska Institutet, Solna, Sweden

(3) Cluster of Scientific Modeling, University of Oviedo, Mieres, Spain

(4) Division of Neurodegeneration, Department of Neurobiology, Care Sciences and Society, Karolinska Institutet, Huddinge, Sweden

(5) Division of Biochemistry, Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Solna, Sweden

*Corresponding author: Tel: +46 8 58583751; Fax: +46 8 58583880;

E-mail: angel.cedazo-minguez@ki.se

[†]These authors contributed equally to this work.

linked to AD, like some peroxiredoxins (Kim et al, 2001) and more recently Trx-reductase 2 (Cacho-Valadez et al, 2012). Together, these data emphasize the importance of Trx1 and related molecules in AD pathology.

Earlier studies on immune cells showed that Trx1 with 105 residues is cleaved to a 80–84 a.a. carboxy terminal-truncated protein, thioredoxin 80 (Trx80; Pekkari & Holmgren, 2004). Trx80 is found in plasma, secreted by monocytes (Pekkari et al, 2000). Indeed, Trx1 is also secreted, although the mechanism is not known since it is lacking a signal peptide. Trx80 levels in plasma are lower compared to Trx1, but there is no correlation between Trx80 and Trx1 levels (Pekkari et al, 2000). There are few reports showing Trx80 production in non-immune cells, such as cytrophoblasts (Di Trapani et al, 1998) and rheumatoid synoviocytes (Lemarechal et al, 2007). Until now, the enzyme responsible for Trx80 production has not been identified. Also, there is no consensus on where the cleavage occurs. Studies in different cell models showed that Trx1 is either cleaved intracellularly and then secreted (Di Trapani et al, 1998; Lemarechal et al, 2007) or secreted and then cleaved extracellularly (Sahaf et al, 1997). It is known that Trx1 cleavage dramatically changes the properties of the protein, including a loss of oxido-reductase activity and the ability to be a disulphide substrate for thioredoxin-reductase (Pekkari et al, 2000). The main function described for Trx80 is to activate monocytes to secrete pro- and anti-inflammatory cytokines (Pekkari et al, 2005). Inflammation is an important component of neurodegenerative disorders, including AD. Activated glia produce pro-inflammatory cytokines that can be found in the cerebrospinal fluid (CSF) of AD patients (Cedazo-Minguez & Winblad, 2010). The present paper aimed to investigate the production of Trx80 in the brain and its contribution to AD pathology.

RESULTS

Trx80 is produced in brain and forms low-molecular-weight aggregates

Immunohistochemical studies of cortex and hippocampus from human brains showed that Trx80 immunoreactivity is mainly confined to pyramidal and bipolar neurons (Fig 1A). Trx80 immunoreactivity was confirmed in mixed (neuronal and glia) human primary cultures (Fig 1B). Using pure human neuronal cultures, we found that Trx80 is present in the soma and particularly in all neurites (Fig 1C). As seen in Fig 1D, the anti-Trx80 antibody 7D11 recognized a single band migrating at approximately 30 kDa in samples from temporal cortex of human brains. SDS–polyacrylamide gel electrophoresis (SDS–PAGE) analysis of lysates from several cell types, including human neurons and glia, showed that Trx80 immunoreactivity appears mainly as a single band of approximately 30 kDa and sometimes also as bands of 60 and/or 80 kDa (Fig 1E and Supporting Information Fig 1A). In conditioned media from primary human neurons and SH-SY5Y neuroblastoma cells, the 7D11 antibody recognized bands of approximately 80 kDa (Fig 1F).

The molecular weight (MW) of detected bands are not in agreement with the expected MW for Trx80 (10 kDa), which suggests the formation of aggregates. Immunoblotting of recombinant Trx80 with two different monoclonal antibodies able to recognize Trx80 (4H9 and 7D11) revealed the formation of aggregates after overnight (O/N) incubation at 37°C. Figure 1G shows the result achieved with the 4H9 antibody. As reported for other aggregated proteins such as α -synuclein (Sharon et al, 2001), heating to 95°C for 20 min prior O/N incubation increased Trx80 aggregation considerably. Addition of dithiothreitol (DTT; 10 mM) did not modify the aggregation pattern.

SDS–polyacrylamide gel electrophoresis (SDS–PAGE) analysis of SH-SY5Y cells transiently overexpressing Trx80 revealed that the increased amount of Trx80 migrated similarly as endogenous Trx80 from brain extracts, *i.e.* a band of 30 kDa (Fig 1H and Supporting Information Fig 1B). Reduction of the expression of Trx1 by siRNA (using 12.5 or 25 nM Trx1 siRNA), resulted in decreased levels of the 30 kDa band detected by anti-Trx80 antibodies (Fig 1I and Supporting Information Fig 1C). Moreover, recombinant Trx80 showed increased aggregation when incubated together with cell lysate (37°C, 24 h; Fig 1J), suggesting that Trx80 is prone to aggregate in a cellular milieu. This was also seen, although to a less extent, for recombinant Trx1 (Fig 1J). Co-incubation of both recombinant Trx80 and Trx1 with cell lysate did not change the aggregation pattern of the peptides compared to each peptide alone (Fig 1J).

Electron microscopy revealed that O/N incubation of recombinant Trx80 (10 μ M) in phosphate saline buffer (PBS) at 37°C induced the formation of a diffuse network of aggregates. No structures resembling mature fibrils were found (Fig 2A). Heating the samples to 95°C prior to O/N incubation at 37°C did not change the apparent structure of these aggregates (Fig 2a, right). The inability of Trx80 to form fibrils was confirmed by Thioflavin-T (ThT) fluorescence spectroscopy using insulin (10 μ M) as control (Fig 2B). A prediction of aggregation (Trovato et al, 2007) profile of Trx80 revealed two major areas prone to aggregation: Trx80(20–37) and Trx80(40–60) (Fig 2C). Estimation of the Trx80 structure from the X-ray diffraction crystal structure of Trx1 [1ERT; (Weichsel et al, 1996)] revealed that Trx80 is a more hydrophobic molecule with exposure of the central β -sheets to the exterior (Fig 2D). PyMOL display of the protein backbone shows how the differential residues between Trx1 and Trx80 (residues 81–105 in red) shield the first predicted aggregation area (mainly a β -sheet that forms part of the hydrophobic core of Trx1) (Fig 2E, right). Figure 2E (left) displays the calculated surface of the Trx80 coloured green to show the location of the hydrophobic residues and, among them, the 21–25 residues KLVVV in magenta. The differential residues 81–105, depicted as a red ribbon, show how their cleavage exposes a big hydrophobic patch, which contains areas prone to aggregation (including the KLVVV residues).

Trx80 levels are reduced in Alzheimer's disease

Comparative analysis of samples from control and AD brains revealed a significant decrease of Trx80 immunoreactivity in AD (by 81.5% in five AD brain samples compared to four controls,




Figure 1. Trx80 is present in brain in an aggregated form.

A–C. Trx80 immunostaining in (A) human brain, (B) human mixed primary culture of neurons and astrocytes, (C) pure neuronal human primary culture. Trx80 is seen in neurons, particularly in bipolar and pyramidal neurons (right panel in A). Arrowheads in B indicate astrocytes. The pure neuronal cultures show Trx80 staining in soma and neuronal processes (C). Scale bars: 25 μm.

D. Trx80 immunoblotting of samples from three control brains. The 7D11 antibody recognizes a band at approximately 30 kDa.

E. Trx80 immunoblotting of lysates from different cell types (1: HeLa, 2: U2020, 3: U937, 4: SH-SY5Y, 5: SH-SY5Y overexpressing Trx1, 6: human primary neurons, 7–8: human primary glia).

F. Trx80 is secreted and detected in the media from rat neuronal primary cultures and SH-SY5Y cells.

G. Immunoblotting with the anti-Trx80 antibody 4H9. Recombinant Trx80 was incubated O/N at 37°C and compared to fresh Trx80. Samples were mixed with sample buffer ± DTT (10 mM) and left untreated or heated to 95°C.

H. Transfected Trx80 also migrates as a band of approximately 30 kDa, as the endogenous Trx80. Panel shows two different clones.

I. Reduction of Trx1 expression by siRNA resulted in decreased levels of the 30 kDa band detected by 7D11. Scrambled RNA (scRNA)-treated and vehicle-treated cells were used as controls. Lysates from SH-SY5Y cells were incubated (37°C, 24 h) with or without 2.4 μg of recombinant Trx80, Trx1 or both peptides together. As controls, Trx80 and Trx1 peptides were incubated at 37°C for 24 h in PBS. Blotting with anti-Trx80 (7D11) and anti-Trx1 abs is shown, respectively, in the left and right panels.

J. Incubation of recombinant Trx80 with cell lysates resulted in more aggregated signals (left panel, lanes 2, 4 and 6). Addition of recombinant Trx1 alone or in combination with Trx80 did not change the aggregation pattern of Trx80 (left panel, lanes 3 vs. 2; 4 vs. 5; and 6 vs. 7). The aggregation pattern of Trx1 was also changed by incubation with lysates, with the appearance of more dimers and trimers (right panel, lane 1 vs. 5). Addition of Trx80 did not change the Trx1 aggregation pattern (right panel, lanes 1 vs. 3 and 5 vs. 7).

$p=0.01$; Fig 3A and Supporting Information Fig 1D). In AD brains, we saw a decrease in neuronal NeuN and an increase in glial fibrillary acidic protein (GFAP) (Fig 3A and Supporting Information Fig 1D) as indicators of neuronal loss and enhanced gliosis, respectively. Ponceau staining is shown as loading control (Fig 3A). Trx80 immunoreactivity was greatly reduced in both hippocampus (dentate gyrus) and cortex of AD brains compared with controls (Fig 3B and Supporting Information Fig 2A). Some staining was caused by autofluorescence from

lipofuscin, heavily present in AD brains. Trx80 did not colocalize with Aβ fibril plaques (Fig 3C). AD samples had abundant Trx80 immunoreactivity in the wall of deep vessels in the white matter (Fig 3D).

To analyze Trx80 levels during AD progression, we quantified Trx80 in CSF from controls, from patients with stable mild cognitive impairment without progression to AD (MCI), MCI-progressive to AD (MCI-AD), or mild AD by a highly sensitive sandwich enzyme-linked immunosorbent assay




Figure 2. Aggregation of Trx80.

- A.** Electron micrographs of recombinant Trx80 (10 μM) samples incubated O/N at 37°C in PBS (left) or heated to 95°C prior to O/N incubation at 37°C (right). Formation of a diffuse network of aggregates was detected without structures resembling mature fibrils.
- B.** This was confirmed by ThT assay. Insulin served as a positive control.
- C.** Chou-Fasman plot and predicted aggregation profile of Trx80 revealed two major pro-aggregation areas, Trx80(20–37) and (47–60).
- D.** Calculated surface of Trx80 using the X-ray diffraction-determined crystal structure for Trx1(1ERT). Red and blue areas show negative and positive potential, respectively.
- E.** Trx1 backbone structure (using 1ERT coordinates) showing the key regions of the structure (left panel): in red, the differential segment with Trx80 and in yellow and cyan, the two possible aggregation zones from D. In the left panel, residues 21–25 (KLVVV) are shown in magenta. Trx80 calculated surface (right panel) showing the exposed hydrophobic patch (green) and residues 21–25 (magenta).




Figure 3.

(ELISA). Trx80-CSF levels were significantly decreased in both MCI-AD and AD compared to controls (Fig 3E; ANOVA, Bonferroni's multiple comparison test, $p > 0.0001$). A slight but significant decrease was also found in non-progressive MCI samples compared to controls and MCI-AD ($p < 0.05$). Levels of Trx80 correlated with $A\beta(1-42)$ ($r = 0.6313$; $p = 0.0001$) and total tau ($r = -0.4070$; $p = 0.04$) levels (Fig 3F).

Inhibition of ADAM10/17 suppresses Trx80 production

The regulatory mechanisms of Trx80 production from Trx1 as well as the enzyme responsible for the cleavage are unknown. As reported (Balcewicz-Sablinska et al, 1991), we found that phorbol 12-myristate 13-acetate (PMA; 50 ng/ml) increased intracellular levels of both Trx80 and Trx1 after 12 and 24 h in the monocyte cell line U937 (Fig 4A, left and middle panels; Supporting Information Fig 2B). At 12 and 24 h, Trx80 and Trx1 were also detected in conditioned media, the first as high MW aggregates (approx 80 kDa). Both, Trx80 and Trx1 secretion was stimulated by PMA (Fig 4A, right panel; Supporting Information Fig 2B-C).

Phorbol esters are known activators of α -secretase (Buxbaum et al, 1990). Lysine-directed secretases belonging to a disintegrin and metalloproteinase (ADAM) family, ADAM10 and 17, have been suggested as α -secretase candidates (Vincent & Checler, 2011), and α -secretase-mediated processing of APP is reduced in AD (Colciaghi et al, 2002). We hypothesized that α -secretases could be involved in Trx1 cleavage. To investigate this possibility, we used specific inhibitors of ADAM10 and 17 (Hundhausen et al, 2003). In SH-SY5Y cells, we found a reduction of Trx80 levels (of approx. 54%) after inhibition of both ADAM10/17 by GW280264X (10 μ M). Inhibition of ADAM10 alone by G1254023X (10 μ M) had a smaller effect. Accordingly, Trx1 levels were increased after inhibition of ADAM10/17 (Fig 4B). In U937 cells, the effects of both inhibitors were larger (59% and 32% reduction of Trx80 levels after inhibition of ADAM10 or ADAM10/17, respectively). PMA treatment (50 ng/ml, 24 h) enhanced Trx80 production (approximately four-fold of control levels; Fig 4C). This effect was partially inhibited by the general metalloprotease inhibitor GM6001 (10 μ M) and by ADAM10 inhibition. ADAM10/17 inhibition prevented completely the PMA effect ($p = 0.01$). ADAM inhibitors also reduced the PMA-mediated enhancement of secreted α -cleaved APP levels in a similar way to Trx80 (Fig 4C).

We next looked for co-localization of ADAM17 and Trx1, as precursor of Trx80, using single plane and z-stack confocal

imaging in U937 cells. As seen in Fig 5A, Trx1 and ADAM17 are co-localized to the cytoplasm, mainly in the proximity of the nucleus. No co-staining with DAPI was found, indicating a non-nuclear co-localization. PMA (50 ng/ml, 12 h) enhanced Trx1/ADAM17 interaction, which was extended to non-perinuclear areas (Fig 5B). The dotted staining pattern suggests that Trx1 and ADAM17 could co-localize in vesicular compartments in the cytoplasm. Methylamine that was shown to reduce the secretion of Trx1 by inhibiting the formation of vesicles involved in the non-classical secretory pathway (Rubartelli et al, 1992), reduced the amount of Trx1/ADAM17 co-localization (Fig 5C). In agreement with that, co-treatment with methylamine (10 mM, 12 h), reduced the PMA-mediated enhancement of Trx80 in U937 cells (Fig 5D). Co-localization of ADAM17 and Trx1 was also confirmed in SH-SY5Y cells (Supporting Information Fig 2D).

Trx80 does not induce an inflammatory response in glia

Trx80 was reported to be a cytokine that stimulates monocytes to produce IL-12 and the anti-inflammatory IL-10 (Pekkarinen et al, 2001, 2005). We examined Trx80 effects in relation to two other players of AD pathology known to influence neuroinflammation, $A\beta$ and apolipoprotein E (apoE). We used mixed cerebellar glial cultures (micro- and astroglia), with high expression of neuronal nitrogen oxide synthase (nNOS; Campese et al, 2007), as interrelations between both cell types are important to the final inflammatory effect. Nitrite release to the media was used as indicator of the inflammatory response (Privat et al, 1997). Cells were treated with fresh Trx80 (100 or 10 nM) alone or in combination with $A\beta(1-42)$ (10 μ M), $A\beta(1-40)$ (10 μ M), apoE3 (10 nM), or apoE4 (10 nM) or complexes ($A\beta(40)/E3$ or $A\beta(40)/E4$). Untreated cells or lipopolysaccharide (LPS) (100 ng/ml) treatment were used as negative and positive control, respectively.

As shown in Fig 6A, Trx80 alone (10 nM) did not induce changes in nitrites production. $A\beta(1-40)$ and apoE4 induced moderate inflammatory responses. The effects were not seen with $A\beta(1-42)$ or apoE3 treatments. $A\beta(1-40)/apoE$ complexes, specially $A\beta(1-40)/apoE4$, induced significant increases in the accumulation of nitrites. The addition of Trx80 did not change the response to LPS, $A\beta$, apoE isoforms or $A\beta/apoE$ complexes. Similar findings were obtained with 100 nM Trx80 and when Trx80 was incubated at 37°C O/N, to induce aggregates, prior to the addition to the cells (Supporting Information Fig 3A).

Figure 3. Reduced Trx80 levels in AD.

- Immunoblotting of control and AD brain samples in human brain revealed a significant reduction of Trx80 in AD brains. No changes were found in Trx1 levels. As expected, NeuroN levels were reduced and GFAP levels increased in AD, reflecting neuronal loss and gliosis. Ponceau staining is shown as loading control. Histogram shows data expressed as optical density units and presented as means \pm SEM (Mann-Whitney *U*-test; * $p = 0.02$).
- Trx80 staining is dramatically decreased in cerebral cortex and dentate gyrus of AD patients.
- $A\beta(1-42)$ staining (green) does not co-localize with Trx80 staining (red) in senile plaques.
- One of the deep vessels with abundant Trx80 staining found in AD brains. Scale bars: 25 μ m.
- Levels of Trx80 in CSF, measured by Sandwich ELISA, were reduced in samples from AD and MCI-progressing to AD patients compared to controls and MCI non-progressive patients (ANOVA, Bonferroni's multiple comparison test, $p = 0.0001$). A significant decrease was also found in non-progressive MCI samples compared to both controls and MCI-AD ($p = 0.04$).
- Levels of Trx80 in CSF correlated with $A\beta(1-42)$ levels [$r = 0.6313$; $p(\text{two-tailed}) = 0.0001$] and total tau [$r = -0.4070$; $p(\text{two-tailed}) = 0.04$].




Figure 4. Trx80 is generated by α -secretases.

A. Trx80 levels both in cells (left and middle panels) and media (right panel) increase in response to PMA stimulation (50 ng/ml) of U937 cells. Trx1 secretion to the media is also increased by PMA (left panel) but the levels in lysates are not affected (right panel). Actin or Ponceau staining are shown as loading controls. Immunocytochemical pattern of Trx80 expression showed a dotted pattern that was increased after PMA treatment (middle panel). Graphs show quantifications (means \pm SEM) of data.

B,C. Effects of ADAM inhibitors on (B) basal or (C) PMA-stimulated (50 ng/ml) Trx80 and Trx1 production in SH-SY5Y cells. Bars represent mean values of four experiments \pm SEM. Data are expressed as percentage of values for vehicle-treated cells (B) or PMA-stimulated levels (C). Analysis of the basal effects on Trx80/Trx1 ratio (B and C) and on PMA-stimulated sAPP α levels (C) are also shown. GM = GM6001 (10 μ M); GI = GI254023X (10 μ M); GW = GW280264X (10 μ M). (ANOVA followed by Bonferroni's PLSD *post hoc* test; * p = 0.04 and ** p = 0.009).




Figure 5. Co-localization of Trx1 and ADAM17 in cytoplasmic vesicles of U937 cells.

- A–C.** Co-localization of Trx1 (green) and ADAM17 (red) in (A) untreated (c), (B) PMA-treated (50 ng/ml) and (C) methylamine-treated (Met) (10 mM) U937 cells. Both single plane and z-stack images are shown. The co-staining (yellow) of Trx1 and ADAM17 was present in the cytosol, mainly in proximity of the nucleus. The dotted pattern of the staining suggests that Trx1 and ADAM17 could co-localize in vesicular compartments in the cytoplasm. Methylamine-treatment reduced the dotted pattern as well as the co-localization of Trx1 and ADAM17.
- D.** Trx80 and Trx1 immunoblotting of lysates from U937 cells using 7D11 and anti-Trx1 antibodies, respectively. Pretreatment of cells with methylamine (10 mM, 5 h) reduced the PMA-mediated increase of Trx80.

Trx80 arrests A β aggregation and protects cells from A β toxicity

Polymerization of A β is critical for its toxicity, and inhibitors of A β aggregation have been proposed as potential treatment for AD. The 16–20 fragment of A β (KLVFF) was shown to be responsible for its aggregation (Tjernberg et al, 1996). Consequently, peptides binding to this fragment are able to inhibit A β polymerization (Tjernberg et al, 1997). From the two predicted pro-aggregation Trx80 regions, Trx80(21–25) (KLVVV) is exposed after cleavage of C-terminal Trx1 and shows important similarities to A β (16–20) (Fig 2). Therefore, binding between Trx80(21–25) and A β (16–20) could be possible. We evaluated the effect of Trx80 on A β (1–42) polymerization. Monomeric A β (1–42) (20 μ M) aggregates into

ThT reactive fibrils, an effect that was inhibited by co-incubation with monomeric Trx80 at concentrations of 1 and 5 μ M (Fig 6B). Lower concentrations of Trx80 (10–100 nM) were without effect. As also described in Fig 2B, Trx80 alone did not show fibril formation (Fig 6B).

The effect of Trx80 on A β (1–42) toxicity was evaluated in SH-SY5Y cells (Fig 6C). A β (1–42) (10 μ M), incubated O/N at 37°C to induce aggregates, significantly decreased cell viability at 24 h. This effect was not modified by coinubation with fresh Trx80 (1 μ M). In contrast, A β (1–42) (10 μ M) incubated together with Trx80 (1 μ M) O/N was harmless to the cells. Moreover, pre-aggregated A β (1–42) (10 μ M) was less toxic in cells transiently transfected to overexpress Trx80 (Trx80-T) (three different clones) (to 87.2 \pm 2.8% of controls). We also tested




Figure 6. Trx80 is not pro-inflammatory in glia but inhibits A β (1–42) aggregation and toxicity.

- A.** Nitric oxide production was measured in mixed glial primary cultures from rat in response to Trx80 stimulation (10 nM) in combination with different inflammatory modulators. LPS = lipopolysaccharide (100 ng/ml); C = untreated cells; A β (1–42) (10 μ M); A β (1–40) (10 μ M); E3 = apoE3 (10 nM); E4 = apoE4 (10 nM). Data are expressed as ng of nitrites produced per μ g of total protein. Bars represent mean values of three experiments \pm SEM.
- B.** Trx80 (1 and 5 μ M) inhibited the fibrillar aggregation of A β (1–42) (20 μ M) detected by fluorescence Thioflavin-T assay. Lower Trx80 concentrations were without effect.
- C.** A β (1–42) (10 μ M, 24 h) reduced cell viability in non-transfected (NT) and in Trx80 (100 nM)-treated SH-SY5Y cells. A β (1–42) was harmless when pre-incubated with Trx80 (100 nM) or added to Trx80-transfected (Trx80-T) cells. Data are expressed as percentage of the respective control and bars represent mean values of four experiments \pm SEM (ANOVA followed by Bonferroni's PLSD *post hoc* test; *** p = 0.001, * p = 0.04).
- D.** A β -rich fractions (1 μ l) extracted from one AD brain (A β -AD) reduced cell viability of SH-SY5Y cells. The A β -AD effect was partially inhibited by addition of Trx80 (100 pM) or by overexpression of Trx80 (Trx80T). Treatment with extracts from a plaque-free control brain (C) did not harm the cells. Data are expressed as percentage of values in buffer treated cells, and bars represent mean values of 9 experiments \pm SEM (ANOVA followed by Bonferroni's PLSD *post hoc* test; *** p = 0.001 against controls; # p = 0.01 against A β -AD).
- E.** Overexpression of Trx1 (Trx1-T) was partially protective against A β (1–42) (10 μ M, 24 h) even under ADAM10/17 inhibition (GW280264X, 10 μ M). Data from four experiments are shown as percents of MTT values from controls (untreated cells, NT) \pm SEM (ANOVA followed by Bonferroni's PLSD *post hoc* test; *** p = 0.001).

the ability of Trx80 to protect against A β -rich extracts from AD brains. As seen in Fig 6D, incubation with A β -rich AD brain extracts reduced cell viability to $76.2\% \pm 2.3$ compared with controls (extraction buffer-treated cells). This toxic effect was reduced by the addition (co-incubation) of a low concentration of recombinant Trx80 (100 pM) or by Trx80 overexpression (Trx80-T) (respectively to 87.7 ± 1.3 and 86.1 ± 2.9) (Fig 6D).

In an independent set of experiments, we determined the potential of Trx1 overexpression to protect against A β (1–42) toxicity in presence or absence of the ADAM10/17 inhibitor GW280264X. An approximately two-fold overexpression of Trx1 was partially protective against A β (1–42). This effect was not modified by GW280264X (Fig 6E).

DISCUSSION

The reductant and antioxidant properties of Trx1 are lost when it is cleaved to Trx80 (Pekkari et al, 2003) but little is known about Trx80 function and how and in which tissues it is generated. The majority of studies on Trx80 were performed in immune cells where it acts as a cytokine for resting monocytes (Pekkari et al, 2000, 2005). Neurodegenerative disorders are often associated with oxidative stress and inflammatory processes. We previously reported a decrease in neuronal Trx1 levels in AD (Akterin et al, 2006). In the present study, we investigated the production of Trx80 in brain, the mechanisms of Trx80 generation in brain cells and whether Trx80 plays role in AD.

Using human brain samples and human primary cultures, we show that Trx80 is produced and polymerized into very stable aggregates of which molecular species migrating at approximately 30 kDa in SDS-PAGE are predominant in the brain. Furthermore, we show that Trx80 is secreted and forms larger aggregates in conditioned media from brain cells. Trx1 is mainly monomeric although it can form less active dimers (Powis & Montfort, 2001). Trx80 has a secondary structure similar to Trx1 but the removal of the C-terminal 24 residues increases the number of hydrophobic residues on the surface and, therefore, possible interactions (Pekkari et al, 2000). We also show that Trx1 and Trx80 do not aggregate with each other or interfere with their respective aggregations.

We found that Trx80 localization is mainly neuronal, predominant in the cytosol and abundant in axons and dendrites. Moreover, compared to normal brain tissue, Trx80 levels were drastically reduced in AD, even in areas with abundant inflammatory changes (*i.e.* surrounding the amyloid plaques). Trx80 levels were also drastically reduced in CSF from mild AD and MCI-progressive prodromal AD patients. Trx80 measurements in CSF clearly differentiate MCI converters from non-converters to AD, suggesting that Trx80 deficiency could be a specific feature of the disease. Trx80 showed a positive correlation with A β (1–42) and negative with total tau levels in CSF. Although the potential of CSF-Trx80 as AD biomarker could be suggested from our results, this matter should be properly investigated in a large sample cohort in which other neurodegenerative diseases should be included.

The function of Trx80 in brain cells seems to be different from what was described in immune cells, *i.e.* to stimulate a pro-inflammatory response (Pekkari et al, 2001). Treatment of rat primary glial cultures with recombinant Trx80 did not induce nNOS activity, either alone or in combination with A β or apoE. As previously reported (Chiarini et al, 2005; Colton et al, 2002; Takata et al, 2003), we found that A β (1–40) and apoEs (E4 > E3) induced a moderate increase in nNOS activity. On the other hand, and as also reported, A β (1–42) was without effect (Takata et al, 2003). A β (1–40)/apoE complexes [specially A β (1–40)/apoE4] triggered a more potent inflammatory effect. None of the responses seen with A β , apoEs or A β /apoE complexes were modified by the addition of Trx80, further supporting the idea that Trx80 does not have a pro-inflammatory effect on glial cells. Nitric oxide (NO) production can be related to arginine uptake, as arginine is the only substrate for all NOS isoforms (Alderton et al, 2001). Thus, it is possible that arginine uptake in glia is enhanced by apoE4 compared to apoE3 (Colton et al, 2002; Czapiga & Colton, 2003). The isoform-specific differences between A β (1–40)/apoE3 and A β (1–40)/apoE4 complexes constitute an interesting finding. The synergistic action of A β (1–40) and apoE4 on nNOS activity is likely to contribute to higher levels of oxidative and nitrosative stress seen in the brains of apoE4 carriers (Colton et al, 2002).

A reduction of its precursor (Trx1) levels in neurons (Akterin et al, 2006) does not entirely explain the Trx80 decrease seen in AD. Most likely, a downregulation of the enzyme (or enzymes) that cleave Trx1 to Trx80 also occurs. Pekkari and Holmgren (2004) suggested that Trx1 is cleaved by an inducible protease but its identity has so far been unknown. The protease involved in the synthesis of Trx80 should fulfill the following three criteria: (i) cleaves at lysine; (ii) is activated by phorbol esters and (iii) its activity is reduced in AD. We thought that α -secretase could be a suitable candidate. Using specific inhibitors of ADAM10 and 17, two enzymes with α -secretase activity (Vincent & Checler, 2011), we found that simultaneous inhibition of ADAM10 and 17 led to reduced basal and PMA-stimulated production of Trx80 in two different cell types, U937 and SH-SY5Y. Inhibition of ADAM10 alone had a reduced effect. ADAM10 has been shown to be responsible for both constitutive and PKC-regulated α -secretase activities, while ADAM17 appears to be mainly involved in the last one (Vincent & Checler, 2011). As PKC activity is greatly increased under PMA treatment, this would result in activation of both ADAM10 and 17. Under basal conditions, α -secretases are regulated by basal phosphorylation, and it is likely that several enzymes are contributing (including PKC). GI254023X is a specific inhibitor for ADAM10 (constitutive and induced α -secretase), while GW280264X inhibits both ADAM10 and 17 (again both constitutive and induced α -secretase activities). Thus, under basal conditions (Fig 4B), we see a reduction of Trx80 by both inhibitors, with GW280264X being more efficient, probably due to a more efficient inhibition of basal PKC-mediated α -secretase activity. Under PKC-stimulated conditions (PMA treatment, Fig 4C), GW280264X is more efficient than GI254023X in reducing Trx80 production. This is expected, as the combinatory effect of ADAM10 and ADAM17 blockages would inhibit the

PKC inducible α -secretase more efficiently. Thus, our data strongly suggest that α -secretases are involved in the cleavage of Trx1 to generate Trx80. We show, as previously reported (Skovronsky et al, 2001), that levels of ADAM17 are not changed in AD. Accordingly, the decrease in Trx80 levels seen in AD would be a consequence of two co-operating factors: lower levels of its precursor (Trx1) (Akterin et al, 2006) and lower activity of the cleaving enzyme (α -secretase) (Tyler et al, 2002).

From the staining patterns of Trx1 and ADAM17, colocalization appears to occur in vesicles. Trx1 is secreted by a non-classical pathway that is still not identified but involves vesicular elements containing both cytosolic and membranous proteins (Rubartelli et al, 1992). Treatment of cells with methylamine has been shown to block the formation of endosomal vesicles, the secretion of Trx1 (Rubartelli et al, 1992), and other proteins secreted through a non-classical pathway (Mignatti et al, 1992; Nishihara et al, 2001). Using U937 cells, we showed that co-localization of Trx1 and ADAM17, as well as the levels of Trx80, were reduced by methylamine. PMA has the opposite effect on the cells, increasing the colocalization of Trx1 and ADAM17 within what appears to be cytosolic vesicles. Accordingly, PMA has previously been shown to activate secretion (Buccione et al, 1996) and to promote the internalization of ADAM17 from the plasma membrane (Doedens & Black, 2000). Consequently, we suggest that Trx1 and ADAM17 are recruited to a vesicular compartment destined for secretion and that their co-localization will enable the cleavage of Trx1 and the generation of Trx80.

From the aggregation profile of Trx80, we noted that the Trx80(21–25) (KLVVV) region could potentially bind to A β (16–20) (KLVFF), a critical region for A β polymerization and toxicity (Tjernberg et al, 1996). We demonstrated that Trx80 is able to inhibit A β (1–42) aggregation *in vitro*, as well as the toxicity of A β (1–42) in SH-SY5Y cells. The positive effects of Trx80 on A β (1–42)-induced toxicity were seen when A β (1–42) was preincubated with Trx80 prior to addition to the cells or when Trx80 was overexpressed. Trx80 was able to protect against both recombinant A β and purified A β -rich fractions from AD brains. As previously reported (Akterin et al, 2006), overexpression of Trx1 protected cells against A β toxicity. Our results suggest that this is independent of Trx80 generation since inhibition of ADAM10/17 did not alter this effect. Trx1 was shown to protect against A β through inhibition of ASK1 (Akterin et al, 2006). On the other hand, we see that Trx1 levels do not correlate with Trx80 levels, as reported previously (Pekkari et al, 2000). Together, these results indicate that improving Trx1 levels, activity or its cleavage to Trx80 would be beneficial to counteract toxic effects of A β . Since Trx80 lacks the dithiol-dependent antioxidative properties of Trx1, it is not possible to explain the protection by a reduction in oxidative stress. Our data rather suggest that inhibition of A β aggregation is the underlying mechanism of Trx80 protection. Thus, it is tempting to speculate that decreased Trx80 production seen in AD, caused by both reduction in neuronal Trx1 levels and in α -secretase activity, is likely to have a negative effect on the brain. A Trx80 deficit

would result in increased A β polymerization and would make the neurons more vulnerable towards A β .

MATERIALS AND METHODS

Brain tissue

Brain material was obtained from the Brain Bank at Karolinska Institutet (Sweden) with approval by the Regional Ethical Review Board (Stockholm). For immunohistochemistry, four brains from patients with definite AD (Mirra et al, 1991; two males 75 and 86-year-old, two females 75 and 86-year-old) and four controls (two females, 81 and 87 years old, and two males 66 and 79-year-old) were used. For immunoblotting, samples of the temporal cortex from six AD brains (three males and three females, 74–98 years old) and four controls (four males, 56–71 years old) were used. The brains had a post-mortem delay between 8 and 49 h.

Immunohistochemical and immunocytochemistry

Immunohistochemistry and immunocytochemistry were performed as described previously (Akterin et al, 2006; Zheng et al, 2011). The primary antibodies used are described below in a specific section. All sections were treated simultaneously under the same conditions. For control staining, the primary antibody was omitted.

Cell lines

Adherent cell lines were cultured as described previously (Cedazo-Minguez et al, 2001a), seeded at 50,000 cells/cm² 2 days before the experiment. The non-adherent human monocytes U937 (Sundstrom & Nilsson, 1976) were cultured in RPMI 1640 with GlutaMAX supplemented with 10% fetal bovine serum (FBS) and seeded the same day as the experiment at 8 million cells per \emptyset 10 cm dish. All cell culture supplies were purchased from Invitrogen Corporation (Sweden).

DNA constructs and transfection

Based on the *TRX1* plasmid (Akterin et al, 2006), we amplified the Trx80 sequence using the forward primer 5'-CGAATTCGCCACCATGG-ATTACAAGGATGACGACGATAA GATGGTGAAGCAGATCGAG-3' and the reverse primer 5'-CGGATCCTTACTTAAA AAAGTGAATGTTGGCATGCATTT-GACTTCAC-3. The forward primer introduced an *EcoRI* site followed by a Kozac sequence preceding the Flag epitope and the corresponding *TRX1* sequence. The reverse primer introduced a *BamHI* site and a stop codon, which was placed after nucleotides that code for Lysine-80 (the last amino acid in Trx80). The amplified fragment was cloned into the *EcoRI*–*BamHI* sites of the pIRESneo expression vector (Clontech, Germany) and the resulting plasmid was transformed into One Shot[®] TOP10 Chemically Competent *E. coli* (Invitrogen, USA) and selected with ampicillin (Invitrogen, USA). The resulting clone was confirmed by DNA sequencing (SeqLab, Germany). Transient transfections were performed using Lipofectamine[™] 2000 (Invitrogen, USA) according to the manufacturer's instructions. Cells were grown in regular medium for 24 h, followed by a selective medium containing G418 geneticin (Invitrogen, USA).

siRNA of Trx1

siRNA oligonucleotides and transfection reagent (DharmaFECT) were purchased from Thermo Scientific. The transfection was done

according to the manufacturer's protocol. SH-SY5Y cells were cultivated in 6-well plates until 90% confluency was reached and transfected with two different siRNA concentrations, 12.5 and 25 nM. Twenty-five nanomolar of non-targeting siRNA was transfected as control. Cells were treated for 48 h before lysis.

Human primary cultures

Embryonic human cortical tissue (6–11 weeks of gestation) was dissected after elective routine, first trimester abortions. Written informed consent was obtained from the pregnant women. The collection was approved by the Regional Ethical Review Board (Stockholm), in compliance with Swedish law. Neuronal cultures were grown in NeuroBasal medium with supplement of L-glutamine (0.5 mM) and B27 (2%). Mixed neuronal and glial cultures were grown in DMEM/F12/GlutaMax with N2 supplement (1%).

Rat primary cultures

Hippocampal and cerebellar tissue from 16-day-old Sprague–Dawley rat embryos was homogenized in serum-free NeuroBasal medium with B27 supplement (2%). Cells from each embryo were seeded separately in dishes, pre-coated with poly-D-lysine MW 300,000 (0.17 mg/ml, Sigma–Aldrich, St. Louis, MO, USA) in PBS. Cells were grown for 2 weeks and the culture medium was changed every fourth day.

Preparation of cell lysate and media for immunoblotting

U937 cells were collected and separated by centrifugation (1000×g for 10 min) at +4°C. For adherent cell lines and primary cultures media was collected and centrifuged (1000×g for 10 min) at +4°C. All cell types were lysed by applying lysis buffer (50 mM Tris–HCl pH 7.5, 150 mM NaCl, 1% Triton X-100, 2 mM EDTA, 2 mM EGTA), with protease inhibitor cocktail (Sigma–Aldrich, USA).

Immunoblotting

Immunoblotting was performed as previously described (Akterin et al, 2006). Some immunoblots were stripped using Restore™ Western Blot Stripping buffer (Pierce, USA) and then re-blotted with other antibodies.

Study population and CSF measurements

The patients included in the study ($n=40$) were from the Geriatric University Clinic at the Karolinska University Hospital in Huddinge, Sweden: 10 had subjective cognitive impairment (SCI) and were considered as controls, 20 MCI, and 10 mild-AD. In the MCI group, 10 had progression-to-AD (MCI-AD) within 2 years time, and 10 remained stable (MCI). Description of samples and CSF extraction can be found in (Gil-Bea et al, 2010). Tau, P-Tau (P-Thr181) and A β 1–42 were determined by ELISA (Gil-Bea et al, 2010; Innogenetics Belgium).

Trx80 measurements by specific sandwich ELISA

Determination of Trx80 levels in CSF was performed with a sandwich ELISA as previously described (Pekkari et al, 2000). Standard samples of recombinant Trx80, coating antibodies (anti-Trx80 monoclonal mouse antibody 7D11) and detection antibodies (biotinylated goat polyclonal anti-Trx1) were from IMCO Corporation Ltd AB (Sweden). Standard dilutions of Trx80 (0.2–100 ng/ml) were prepared in blocking buffer, while CSF samples were undiluted. Fifty μ l of standards or samples were added in duplicates and incubated O/N.

Inhibition and stimulation of ADAM activity

Cells (U937 and SH-SY5Y) were treated with PMA (50 ng/ml), GM6001 (10 μ M), GI254023X, (10 μ M) and GW280264X (10 μ M; Hundhausen et al, 2003), under serum-free conditions, for 24 h. Untreated cells were used as controls. The effect on Trx80 and Trx1 levels was analyzed by immunoblotting.

Trx80 aggregation

Recombinant Trx1 (1 μ g/7.1 μ M) and Trx80 (1 μ g/13.5 μ M) were incubated O/N in PBS at 37°C. The recombinant protein was prepared as previously described (Pekkari et al, 2000). Equal volume of tricine gel sample buffer with or without DTT (10 mM) was added and then the samples were frozen before use. Half of the samples were boiled for 3 min at 95°C prior to SDS–PAGE. In the experiments with cell lysates, 100 μ l of lysate from SH-SY5Y cells were incubated at 37°C for 24 h with or without 2.4 μ g of Trx80 peptide. As control, Trx80 peptide was incubated at 37°C for 24 h in 100 μ l PBS, pH 7.4. Samples were then analyzed with immunoblotting.

Negative staining transmission electron microscopy

Trx80 (IMCO Corporation Ltd AB, Sweden) (10 μ M in PBS pH. 7.4) was incubated O/N at 37°C. Some samples were boiled for 3 min at 95°C prior incubation O/N at 37°C. An aliquot of 4 μ l was added to a grid coated with a Formvar supporting film coated with carbon for 5 min. The excess solution was soaked off by a filter paper, and the grid was stained with 0.5% uranyl acetate in water for 5 min and air-dried. Samples were visualized in a Tecnai FEI 10 electron microscope at 80 kV.

Nitrite production

Rat primary glial cultures from cerebellum were grown for 2–3 weeks and then seeded in 48-well plates. At 80% confluence, the cells were incubated with serum-free DMEM medium overnight and the day after the media was removed and fresh serum-free DMEM was added containing the different treatments in the presence or absence of human recombinant Trx80 (10 nM); A β (1–40) (10 μ M, Bachem AG, Switzerland), human recombinant apoE isoforms (E3 or E4) (10 nM, ReliaTech GmbH, Germany) or apoE/A β complexes. A β (1–40) was dissolved in serum free MEM media (pH 7.4) at a concentration of 10 μ M and incubated for 24 h at 37°C prior to the addition to the cells. The apoE/A β complexes were made as previously described (Cedazo-Minguez et al, 2001a). NOS activity was measured by the accumulation of nitrites in the culture medium after 24 h, using the Griess reaction, as previously described (Privat et al, 1997). Data are expressed as ng nitrites/100 μ g protein.

Polymerization assay

Amyloid fibrillation of Trx80 was determined by ThT polymerization assay as described (Ivanova et al, 2009). Insulin was used as positive control. Samples were incubated at 37°C in triplicates. Prior to the co-incubation study with A β (1–42) and Trx80, 1 mg A β (1–42) (Sigma–Aldrich, #A9810) was dissolved and prepared as described (Stine et al, 2003) in order to get an unaggregated preparation. A β (1–42) was diluted to 20 μ M in 200 μ l ThT buffer, with or without 5 μ M Trx80 peptide. ThT buffer alone was used as negative control. The ThT fluorescence intensity was then recorded every 10 min with the same settings as described above.

The paper explained

PROBLEM:

Alzheimer's disease (AD) is the most prevalent of the dementias and a major public health problem. Diagnostic accuracy is illusive at the earliest stages and effective treatments are unconfirmed. How and by what mechanisms rare genetic mutations lead to AD is largely known, but we lack such understanding of the much more common sporadic forms of AD. Understanding contributing factors would likely yield valuable insights about mechanisms and lead to new therapeutic candidates as well as improvements in early diagnosis.

RESULTS:

We report that the endogenous antioxidant Trx1 is cleaved to thioredoxin 80 (Trx80) by α -secretases ADAM10 and 17. Trx80 localizes mainly as an aggregate in neurons and is dramatically reduced in AD brains. The decrease of Trx80 is detectable in CSF at early AD phases. Our measurements of Trx80 in CSF

discriminated patients with stable mild cognitive impairment from prodromal AD and mild AD patients. We also demonstrated that Trx80 does inhibit amyloid- β aggregation (a critical issue in AD pathogenesis) and protects cells against amyloid- β toxicity.

IMPACT:

Our results strongly suggest that decreases in Trx80 in brain has a contributing role in AD pathogenesis and progression. Trx80 deficits would result in increased A β polymerization that, consequently, leaves neurons more vulnerable to A β . The search for inhibitors of A β aggregation is an important subject in AD, not only because of therapeutic implications, but also for diagnostic purposes. Since Trx80 measurements in CSF clearly differentiate MCI converters from non-converters to AD, the potential for CSF-Trx80 to be an AD diagnostic biomarker in early stage illness should be further investigated.

Preparation of human brain samples for cell viability assay

Frozen human frontal cortex samples from one plaque-free neurologically healthy and one AD case were homogenized on ice in 1 \times TBS (50 mM Tris-HCl, 0.15 M NaCl, pH 7.4) using a dounce homogenizer, and further ultracentrifuged at 175,000 $\times g$ for 20 min. The pellets were dissolved and homogenized in 1% sodium dodecyl sulphate followed by ultracentrifugation at RT. This step was repeated four times to remove all the detergent soluble material. The remaining pellet was dissolved in 80% formic acid and immediately vortexed for 30 s. on maximum speed, followed by sonication in a 20°C water-bath. Samples were centrifuged at 25,000 $\times g$ for 10 m. to clear the solution from any insoluble material. The supernatant was stored at -20°C. We determined the A β (1-42) concentration in the extracts by ELISA (human/rat β -amyloid 42 from Wako Chemicals GmbH, Neuss, Germany). Prior addition to the cells, extracts were neutralized to physiological pH by 2 M Tris-HCl (pH 9.0).

Cell viability assay

Cell viability was assessed by a modified version of the MTT assay (Cedazo-Minguez et al, 2001a). SH-SY5Y cells (non-transfected and Trx80 overexpressing) were cultured in 24-well plates and treated for 24 h with or without 10 μ M A β (1-42) (Sigma-Aldrich, USA). In the experiments with brain extracts, the effects of 1 μ l A β -rich samples from AD brain, yielding a final concentration of 100 pm A β (1-42), were compared with the same volume of control brain samples. Results were expressed as percentages of the values obtained from the appropriate controls.

Antibodies

The following antibodies were used: For Trx80, 7D11 (1:1000 for immunoblotting and 1:500 for immunocytochemistry) and 4H9 (1:1000), both from IMCO Corp Ltd AB (Sweden). Anti-Trx1 (1:1000, O/N, RT for immunoblotting; and 1:500 for immunocytochemistry, 1:1000 for immunoblotting, and 1:500 for immunocytochemistry) Anti-A β (1-42), a gift from

Jan Näslund, for immunohistochemistry as described in (Zheng et al, 2011). 6E10 for detecting APP α , as described (Cedazo-Minguez et al, 2001b). Anti-actin (1:1000 from Sigma-Aldrich, Sweden.O/N, RT). Anti-ADAM17 (H170) (1:100, from Santa Cruz Biotechnology, USA).

Statistical analysis

Analyses of differences were carried out by ANOVA followed by Bonferroni's PLSD *post hoc* test or by Mann-Whitney *U*-test. A value of $p < 0.05$ was considered statistically significant in all analyses. Partial correlations with age, gender as confounding variables were used to assess associations between levels of total tau or A β (1-42) with Trx80 in CSF samples.

For more detailed Materials and Methods see the Supporting Information.

Author contributions

FGB designed experiments, generated and analyzed data (immuno co-localization, ADAMs inhibition and nitrites production experiments). SA designed experiments, generated and analyzed data (WB, ELISA and aggregation studies). TP designed experiments, generated and analyzed data (siRNA, aggregation and cell viability studies). LM generated data (cell viability and transfections). AS generated data (electronic and confocal microscopy, extraction of A β -rich fractions from AD brains). JAC generated data (Trx80 ELISA). AGR generated the three-dimensional images. ES provided human primary neuronal and mixed cultures. AH supplied antibodies, designed experiments and discussed the results. BW supplied clinical data, brain and CSF samples and discussed the results. ACM designed experiments, analyzed data and wrote the manuscript. All the authors participated in the preparation of the manuscript.

Acknowledgements

We thank Dionisio Lorenzo, Eva-Britt Samuelsson and Stefano Costa for technical help and Glaxo Smith Kline (Harlow, UK) for generously providing GW280264X and GI254023X compounds. We thank Lars O. Tjernberg for valuable comments. This research was supported by grants from the following Swedish foundations: Swedish Brain Power, Alzheimerfonden, Anders Otto Swärds Stiftelse, Gun och Bertil Stohnes Stiftelse, Insamlingsstiftelsen för Alzheimer och demensforskning (SADF), Knut and Alice Wallenberg foundation, Stiftelsen för Gamla Tjänarinnor. F. Gil-Bea was supported by a postdoctoral fellowship from the Spanish foundation 'Ramón Areces'.

Supporting Information is available at EMBO Molecular Medicine online.

The authors declare that they have no conflict of interest.

References

- Atkerin S, Cowburn RF, Miranda-Vizuete A, Jimenez A, Bogdanovic N, Winblad B, Cedazo-Minguez A (2006) Involvement of glutaredoxin-1 and thioredoxin-1 in beta-amyloid toxicity and Alzheimer's disease. *Cell Death Differ* 13: 1454-1465
- Alderton WK, Cooper CE, Knowles RG (2001) Nitric oxide synthases: structure, function and inhibition. *Biochem J* 357: 593-615
- Arner ES, Holmgren A (2000) Physiological functions of thioredoxin and thioredoxin reductase. *Eur J Biochem* 267: 6102-6109
- Balcewicz-Sablinska MK, Wollman EE, Gorti R, Silberstein DS (1991) Human eosinophil cytotoxicity-enhancing factor. II. Multiple forms synthesized by U937 cells and their relationship to thioredoxin/adult T cell leukemia-derived factor. *J Immunol* 147: 2170-2174
- Buccione R, Bannykh S, Santone I, Baldassarre M, Facchiano F, Bozzi Y, Di Tullio G, Mironov A, Luini A, De Matteis MA (1996) Regulation of constitutive exocytic transport by membrane receptors. A biochemical and morphometric study. *J Biol Chem* 271: 3523-3533
- Buxbaum JD, Gandy SE, Cicchetti P, Ehrlich ME, Czernik AJ, Fracasso RP, Ramabhadran TV, Unterbeck AJ, Greengard P (1990) Processing of Alzheimer beta/A4 amyloid precursor protein: modulation by agents that regulate protein phosphorylation. *Proc Natl Acad Sci USA* 87: 6003-6006
- Cacho-Valadez B, Munoz-Lobato F, Pedrajas JR, Cabello J, Fierro-Gonzalez JC, Navas P, Swoboda P, Link CD, Miranda-Vizuete A (2012) The characterization of the *Caenorhabditis elegans* mitochondrial thioredoxin system uncovers an unexpected protective role of thioredoxin reductase 2 in beta-amyloid peptide toxicity. *Antioxid Redox Signal* 16: 1384-1400
- Campese VM, Sindhu RK, Ye S, Bai Y, Vaziri ND, Jabbari B (2007) Regional expression of NO synthase, NAD(P)H oxidase and superoxide dismutase in the rat brain. *Brain Res* 1134: 27-32
- Cedazo-Minguez A, Winblad B (2010) Biomarkers for Alzheimer's disease and other forms of dementia: clinical needs, limitations and future aspects. *Exp Gerontol* 45: 5-14
- Cedazo-Minguez A, Hutterling M, Cowburn RF (2001a) Beta-VLDL protects against A beta(1-42) and apoE toxicity in human SH-SY5Y neuroblastoma cells. *Neuroreport* 12: 201-206
- Cedazo-Minguez A, Wiehager B, Winblad B, Hutterling M, Cowburn RF (2001b) Effects of apolipoprotein E (apoE) isoforms, beta-amyloid (Abeta) and apoE/Abeta complexes on protein kinase C-alpha (PKC-alpha) translocation and amyloid precursor protein (APP) processing in human SH-SY5Y neuroblastoma cells and fibroblasts. *Neurochem Int* 38: 615-625
- Chiarini A, Dal Pra I, Menapace L, Pacchiana R, Whitfield JF, Armato U (2005) Soluble amyloid beta-peptide and myelin basic protein strongly stimulate, alone and in synergism with combined proinflammatory cytokines, the expression of functional nitric oxide synthase-2 in normal adult human astrocytes. *Int J Mol Med* 16: 801-807
- Colciaghi F, Borroni B, Pastorino L, Marcello E, Zimmermann M, Cattabeni F, Padovani A, Di Luca M (2002) [alpha]-Secretase ADAM10 as well as [alpha]APPs is reduced in platelets and CSF of Alzheimer disease patients. *Mol Med* 8: 67-74
- Colton CA, Brown CM, Czapiga M, Vitek MP (2002) Apolipoprotein-E allele-specific regulation of nitric oxide production. *Ann N Y Acad Sci* 962: 212-225
- Czapiga M, Colton CA (2003) Microglial function in human APOE3 and APOE4 transgenic mice: altered arginine transport. *J Neuroimmunol* 134: 44-51
- Di Trapani G, Perkins A, Clarke F (1998) Production and secretion of thioredoxin from transformed human trophoblast cells. *Mol Hum Reprod* 4: 369-375
- Doedens JR, Black RA (2000) Stimulation-induced down-regulation of tumor necrosis factor-alpha converting enzyme. *J Biol Chem* 275: 14598-14607
- Gil-Bea FJ, Solas M, Solomon A, Mugueta C, Winblad B, Kivipelto M, Ramirez MJ, Cedazo-Minguez A (2010) Insulin levels are decreased in the cerebrospinal fluid of women with prodromal Alzheimer's disease. *J Alzheimers Dis* 22: 405-413
- Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297: 353-356
- Hundhausen C, Misztela D, Berkhout TA, Broadway N, Saftig P, Reiss K, Hartmann D, Fahrenholz F, Postina R, Matthews V, Kallen KJ, Rose-John S, Ludwig A (2003) The disintegrin-like metalloproteinase ADAM10 is involved in constitutive cleavage of CX3CL1 (fractalkine) and regulates CX3CL1-mediated cell-cell adhesion. *Blood* 102: 1186-1195
- Ivanova MI, Sievers SA, Sawaya MR, Wall JS, Eisenberg D (2009) Molecular basis for insulin fibril assembly. *Proc Natl Acad Sci USA* 106: 18990-18995
- Kim SH, Fountoulakis M, Cairns N, Lubec G (2001) Protein levels of human peroxiredoxin subtypes in brains of patients with Alzheimer's disease and Down syndrome. *J Neural Transm Suppl* 61: 223-235
- Lemarechal H, Anract P, Beaudeau JL, Bonnefont-Rousselot D, Ekinjian OG, Borderie D (2007) Expression and extracellular release of Trx80, the truncated form of thioredoxin, by TNF-alpha- and IL-1beta-stimulated human synoviocytes from patients with rheumatoid arthritis. *Clin Sci (Lond)* 113: 149-155
- Lillig CH, Holmgren A (2007) Thioredoxin and related molecules—from biology to health and disease. *Antioxid Redox Signal* 9: 25-47
- Mignatti P, Morimoto T, Rifkin DB (1992) Basic fibroblast growth factor, a protein devoid of secretory signal sequence, is released by cells via a pathway independent of the endoplasmic reticulum-Golgi complex. *J Cell Physiol* 151: 81-93
- Mirra SS, Heyman A, McKeel D, Sumi SM, Crain BJ, Brownlee LM, Vogel FS, Hughes JP, van Belle G, Berg L (1991) The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part II. Standardization of the neuropathologic assessment of Alzheimer's disease. *Neurology* 41: 479-486
- Nishihara H, Nakagawa Y, Ishikawa H, Ohba M, Shimizu K, Nakamura T (2001) Matrix vesicles and media vesicles as nonclassical pathways for the secretion of m-Calpain from MC3T3-E1 cells. *Biochem Biophys Res Commun* 285: 845-853
- Pekkari K, Holmgren A (2004) Truncated thioredoxin: physiological functions and mechanism. *Antioxid Redox Signal* 6: 53-61
- Pekkari K, Gurunath R, Arner ES, Holmgren A (2000) Truncated thioredoxin is a mitogenic cytokine for resting human peripheral blood mononuclear cells and is present in human plasma. *J Biol Chem* 275: 37474-37480
- Pekkari K, Avila-Carino J, Bengtsson A, Gurunath R, Scheynius A, Holmgren A (2001) Truncated thioredoxin (Trx80) induces production of interleukin-12 and enhances CD14 expression in human monocytes. *Blood* 97: 3184-3190
- Pekkari K, Avila-Carino J, Gurunath R, Bengtsson A, Scheynius A, Holmgren A (2003) Truncated thioredoxin (Trx80) exerts unique mitogenic cytokine effects via a mechanism independent of thiol oxidoreductase activity. *FEBS Lett* 539: 143-148
- Pekkari K, Goodarzi MT, Scheynius A, Holmgren A, Avila-Carino J (2005) Truncated thioredoxin (Trx80) induces differentiation of human CD14+

- monocytes into a novel cell type (TAMs) via activation of the MAP kinases p38, ERK, and JNK. *Blood* 105: 1598-1605
- Powis G, Montfort WR (2001) Properties and biological activities of thioredoxins. *Annu Rev Biophys Biomol Struct* 30: 421-455
- Privat C, Lantoine F, Bedioui F, Millanvoye van Brussel E, Devynck J, Devynck MA (1997) Nitric oxide production by endothelial cells: comparison of three methods of quantification. *Life Sci* 61: 1193-1202
- Rubartelli A, Bajetto A, Allavena G, Wollman E, Sitia R (1992) Secretion of thioredoxin by normal and neoplastic cells through a leaderless secretory pathway. *J Biol Chem* 267: 24161-24164
- Sahaf B, Soderberg A, Spyrou G, Barral AM, Pekkari K, Holmgren A, Rosen A (1997) Thioredoxin expression and localization in human cell lines: detection of full-length and truncated species. *Exp Cell Res* 236: 181-192
- Saitoh M, Nishitoh H, Fujii M, Takeda K, Tobiume K, Sawada Y, Kawabata M, Miyazono K, Ichijo H (1998) Mammalian thioredoxin is a direct inhibitor of apoptosis signal-regulating kinase (ASK) 1. *EMBO J* 17: 2596-2606
- Selkoe DJ (2001) Alzheimer's disease: genes, proteins, and therapy. *Physiol Rev* 81: 741-766
- Sharon R, Goldberg MS, Bar-Josef I, Betensky RA, Shen J, Selkoe DJ (2001) alpha-Synuclein occurs in lipid-rich high molecular weight complexes, binds fatty acids, and shows homology to the fatty acid-binding proteins. *Proc Natl Acad Sci USA* 98: 9110-9115
- Skovronsky DM, Fath S, Lee VM, Milla ME (2001) Neuronal localization of the TNFalpha converting enzyme (TACE) in brain tissue and its correlation to amyloid plaques. *J Neurobiol* 49: 40-46
- Stine WB, Jr., Dahlgren KN, Krafft GA, LaDu MJ (2003) In vitro characterization of conditions for amyloid-beta peptide oligomerization and fibrillogenesis. *J Biol Chem* 278: 11612-11622
- Sundstrom C, Nilsson K (1976) Establishment and characterization of a human histiocytic lymphoma cell line (U-937). *Int J Cancer* 17: 565-577
- Takata K, Kitamura Y, Umeki M, Tsuchiya D, Kakimura J, Taniguchi T, Gebicke-Haerter PJ, Shimohama S (2003) Possible involvement of small oligomers of amyloid-beta peptides in 15-deoxy-delta 12,14 prostaglandin J2-sensitive microglial activation. *J Pharmacol Sci* 91: 330-333
- Tjernberg LO, Naslund J, Lindqvist F, Johansson J, Karlstrom AR, Thyberg J, Terenius L, Nordstedt C (1996) Arrest of beta-amyloid fibril formation by a pentapeptide ligand. *J Biol Chem* 271: 8545-8548
- Tjernberg LO, Lilliehook C, Callaway DJ, Naslund J, Hahne S, Thyberg J, Terenius L, Nordstedt C (1997) Controlling amyloid beta-peptide fibril formation with protease-stable ligands. *J Biol Chem* 272: 12601-12605
- Trovato A, Seno F, Tosatto SC (2007) The PASTA server for protein aggregation prediction. *Protein Eng Des Sel* 20: 521-523
- Tyler SJ, Dawbarn D, Wilcock GK, Allen SJ (2002) Alpha- and beta-secretase: profound changes in Alzheimer's disease. *Biochem Biophys Res Commun* 299: 373-376
- Vincent B, Checler F (2011) Alpha-secretase in Alzheimer's disease and beyond: mechanistic, regulation and function in the shedding of membrane proteins. *Curr Alzheimer Res* 9: 140-156
- Watanabe R, Nakamura H, Masutani H, Yodoi J (2010) Anti-oxidative, anti-cancer and anti-inflammatory actions by thioredoxin 1 and thioredoxin-binding protein-2. *Pharmacol Ther* 127: 261-270
- Weichsel A, Gasdaska JR, Powis G, Montfort WR (1996) Crystal structures of reduced, oxidized, and mutated human thioredoxins: evidence for a regulatory homodimer. *Structure* 4: 735-751
- Zheng L, Terman A, Hallbeck M, Dehvari N, Cowburn RF, Benedikz E, Kagedal K, Cedazo-Minguez A, Marcusson J (2011) Macroautophagy-generated increase of lysosomal amyloid beta-protein mediates oxidant-induced apoptosis of cultured neuroblastoma cells. *Autophagy* 7: 1528-1545