


BMJ Open Is in situ simulation in emergency medicine safe? A scoping review

Jennifer Truchot,^{1,2,3,4} Valérie Boucher,^{4,5} Winny Li,⁶ Guillaume Martel,¹ Eva Jouhair,^{1,4} Éliane Raymond-Dufresne,^{1,2} Andrew Petrosoniak,^{6,7} Marcel Emond ^{1,2,4,5}

To cite: Truchot J, Boucher V, Li W, *et al.* Is in situ simulation in emergency medicine safe? A scoping review. *BMJ Open* 2022;**12**:e059442. doi:10.1136/bmjopen-2021-059442

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2021-059442>).

Received 22 November 2021
Accepted 30 June 2022

ABSTRACT

Objectives To provide an overview of the available evidence regarding the safety of in situ simulation (ISS) in the emergency department (ED).

Design Scoping review.

Methods Original articles published before March 2021 were included if they investigated the use of ISS in the field of emergency medicine.

Information sources MEDLINE, EMBASE, Cochrane and Web of Science.

Results A total of 4077 records were identified by our search strategy and 2476 abstracts were screened. One hundred and thirty full articles were reviewed and 81 full articles were included. Only 33 studies (40%) assessed safety-related issues, among which 11 chose a safety-related primary outcome. Latent safety threats (LSTs) assessment was conducted in 24 studies (30%) and the cancellation rate was described in 9 studies (11%). The possible negative impact of ISS on real ED patients was assessed in two studies (2.5%), through a questionnaire and not through patient outcomes.

Conclusion Most studies use ISS for systems-based or education-based applications. Patient safety during ISS is often evaluated in the context of identifying or mitigating LSTs and rarely on the potential impact and risks to patients simultaneously receiving care in the ED. Our scoping review identified knowledge gaps related to the safe conduct of ISS in the ED, which may warrant further investigation.

INTRODUCTION

Emergency medicine (EM) is a complex and challenging specialty requiring the mastery of numerous technical and non-technical skills. Most emergency departments (ED) are overcrowded, chaotic environments with reported rates of errors up to 10%.¹ Regular simulation is needed to establish and maintain skills competency for ED professionals.^{2,3} However, recent literature suggests a bias towards simulations for rare and unexpected cases (cardiac arrest, difficult airway, disaster medicine and/or rare cases) rather than for more common, routine cases.⁴⁻⁶ The retention rate of acquired knowledge and skills can also be quite low.^{7,8}

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ First complete overview of safety of conducting in situ simulation (ISS) in emergency medicine.
- ⇒ The methodology includes a detailed search strategy, clear inclusion and exclusion criteria, duplicate screening and data extraction by independent reviewers.
- ⇒ The main limitation is the widespread design variations of the included studies and of the terminology and concepts related to ISS.
- ⇒ Isolating the impact of ISS from various potential confounding factors remains a major challenge.
- ⇒ Our study identified major knowledge gaps related to the safe conduct of ISS in emergency medicine, which require further investigation.

In situ simulation (ISS) offers an optimal training solution for these practical EM-related issues.⁹ ISS is defined as simulation taking place in the participant's everyday work environment.^{10,11} The ED environment is very different from a simulation centre and this can greatly impact the learning process.¹² The value of ISS is predicated on principles from the 'situated learning theory' which states that learning is closely linked to the context of the experience.¹² ISS is a 'point of care' type of simulation training used to respond to specific local teaching and training needs and, as a result, may enhance patient safety by exposing latent safety threats (LSTs) and trialling mitigation strategies.^{10,13}

As with any intervention (or any clinical research project),^{14,15} there are potential harms linked to ISS including inadvertent use of simulation equipment/medications for real patient care.¹⁶⁻¹⁸ There is a potential for tension between the present and future state of ED care related to the delivery of ISS. During an ISS session, patients in the ED may be at risk of negative impacts because of the misdirection of resources towards the ISS. The potential risks associated with ISS and the published no go criteria seem to have been reached empirically through



© Author(s) (or their employer(s)) 2022. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Marcel Emond;
marcel.emond@fmed.ulaval.ca

expert consensus.^{16–18} The main objective of this scoping review is to provide an overview of the available evidence regarding the safety of ISS in the ED. Our secondary objective is to explore the benefits of ISS on all levels of Kirkpatrick's pyramid in EM.^{19 20}

METHODS

Design

We followed published guidance for conducting a scoping review.²¹

The results of this scoping review are reported as per the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines.²²

Patient and public involvement

Patients or the public were not involved in the design or the conduct of this study.

Eligibility criteria

We used broad inclusion criteria to present a comprehensive overview of ISS in EM, without any language or date limitations. Original studies were included if they investigated the use of ISS in the ED. Specifically, this comprised studies pertaining to clinician education, system/process evaluation, patient outcomes and patient safety.

We included single-group pretest, non-randomised and randomised studies, parallel-group and cross-over designs, and studies of 'adjuvant' instruction in which simulation was added to other instructions common to all learners.

Protocols, commentaries, conference abstracts, posters and correspondences were excluded. This review excludes systematic reviews or meta-analyses and editorial-style reviews. However, those reviews (invited, narrative or systematic), editorials and letters to the editor were scrutinised and citation tracking was conducted to retrieve potentially relevant studies.

Search strategy and information sources

An experienced librarian designed a search strategy (online supplemental eAppendix 1) to search MEDLINE, EMBASE, Cochrane and Web of Science, using keywords specific to each database from inception to January 2020. The search was updated in March 2021.

Since there is no MESH keyword for 'ISS', a manual search was performed in our four targeted databases.

Selection process

All identified studies were exported to Endnote V.X9 (Clarivate Analytics, Philadelphia, Pennsylvania, USA) and duplicates were then removed. Four authors (JT, WL, GM and EJ) independently assessed the eligibility of each study. Title, abstract and full text were first screened and then cross-referenced between reviewers. Disagreements were resolved by consensus. Any unresolved disagreement was discussed with a third researcher with experience in the field of simulation studies (ER-D).




Figure 1 The validated Kirkpatrick pyramid used to rank educational interventions.

Given the nature of a scoping review, risk of bias and quality appraisals were not performed.²¹

Data collection

Four independent reviewers completed data extraction, ensuring a double reviewing for each article.

Outcomes

The main outcome, safety, is defined in this study as the absence of incidents, accidents or a state with the minimal acceptable level of risk. This definition of safety has been used in other simulation studies.^{23–25} We selected specific criteria such as wait time, patients leaving without being seen or accident reports, which are based on literature on safety metrics in EM.^{26 27} We explored two safety concepts: 'ongoing' safety of patients being managed while ISS was ongoing in the ED (measured by ISS cancellation rate, accident reports, ED median wait time, number of patients who left without being seen) and 'future patient' safety with more long-term benefits from ISS (measured by LST evaluation). We sought to explore the impact of ISS on patients' safety throughout time. Indeed, future patients may benefit from ISS if its output translates into improved skills, clinical care and patient safety.

Secondary outcomes were assessed using the validated Kirkpatrick pyramid, which is a tool used to rank educational interventions on a scale of 1–5 (figure 1).¹⁹ Because of the specificity of ISS, we also distinguished studies that used ISS to teach and train from those that aimed to explore procedures and systems.

Data synthesis

Descriptive statistics and percentages were used to describe our results. No statistical analysis, quantitative meta-analysis or pooling was conducted due to the heterogeneous study designs and methodologies.

RESULTS

Study selection

A total of 4077 records were identified by our initial search strategy. After removing 1601 duplicates, 2476 titles and abstracts were screened. Of those, 2366 did not meet our eligibility criteria. A second search was conducted in March 2021 after which seven original articles were added to our review. A total of 81 full articles were analysed (online supplemental eAppendix 2).

General characteristics of included studies

All reviewed articles were published in English between 2006 and 2021 in specialised journals (81%, n=66).

Most included studies used a prospective design (94%; n=76), 47 of which were observational (58%). Only 3 studies were randomised controlled trials and 28 were pre/post intervention studies (35%). For the most part, the included studies were single centre (80%; n=65) and quantitative (67%; n=54). A high majority included inter-professional participants, defined as ≥ 2 professions (84%; n=68).

The clinical topics explored in the studies included ED care (77%; n=62) and extrahospital care settings (such as HEMS, prehospital care, etc) (22%; n=18). Most studies focused on ISS training for advanced life support/basic life support (n=44, 54%). The remainder focused on a variety of clinical topics including trauma (n=35, 43%) and airway management (n=21, 26%). Three studies particularly focused on COVID-19 (4%). Very few studies used consistent scenarios and as a result more detailed thematic analysis was not performed. Ethical approval was obtained for 51 studies (63%). The full characteristics of the included studies are presented in [table 1](#).

Is ISS safe for patients in EM?

We found that 33 studies (40%) evaluated safety issues, among which 12 (15%) chose a safety-related primary outcome ([table 2](#)).

Ongoing safety

As for ongoing safety of patients during ISS, none of the included studies assessed the following: (1) The number of patients in ED who left without being seen during the period of the ISS training, or (2) The impact of ISS on patient wait times and (3) Safety-related quality parameters in the ED.

No studies included official institutional accident reports, but three studies included surveys on accident reports.^{28–30} No accident was reported during ISS.^{10 31 32} The cancellation rate of ISS was described in nine studies (11%).^{10 31–38} One study included analyses on feasibility (based on department status, such as patient waiting to be seen) in their secondary outcomes. The authors showed that 3 ISS were delayed, and none were cancelled but no other patient-related outcome was presented.³⁹

The impact of ISS on real ED patients was assessed in two studies. In one multicentre clinical trial, ISS was used to prepare and facilitate the identification and

Table 1 General characteristic of studies (n=81)

Study characteristics	n (%)
Study design	
Pre/post design	28 (35)
Comparative	
≥ 2 groups	16 (20)
1 group	55 (68)
Prospective	76 (94)
Retrospective	4 (5)
Randomised controlled trial	3 (4)
Single centre	65 (80)
Quantitative	54 (67)
Qualitative	17 (21)
Mixed	9 (11)
Participants	
Mean no of participants per study \pm SD (min; max)	88.9 \pm 66 (4; 398)
Medical students	0 (0)
Residents	4 (4.9)
Emergency department teams	62 (76.5)
Nurses and nursing students	2 (2.5)
Physicians in practice	3 (3.7)
Paramedics	2 (2.5)
Interprofessional	68 (84)
Clinical topics	
ALS, BLS	44 (54)
Trauma	35 (43)
Intubation	21 (26)
Pediatrics	22 (27)
COVID-19	3 (4)
Goal of ISS	
to train skills	34 (49)
to assess skills or a procedure	24 (35)
to train and assess	16 (18.5)
NA	7 (9)
ALS, advanced life support; BLS, basic life support; ISS, in situ simulation; NA, not available.	

mitigation of threats to study participation and patient safety.²⁶ In this case, ISS was not really used to teach and directly improve skills but rather to prepare teams for the implementation of a clinical trial. As an ongoing research project can also jeopardise the safety of other patients by redirecting attention on the study and not on real patients, safety was analysed with parameters such as latent threats identification, mitigation and protocol errors and deviation. Patterson *et al's* work on LST identification also addressed the impact of ISS training on patient care,¹⁰ which was assessed using a questionnaire. Four participants out of 118 reported that the simulation

Table 2 Safety

Safety-related parameters	n (%)	References
Safety assessment	33 (40)	Amiel <i>et al.</i> ²⁸ Abulebda <i>et al.</i> ²⁹ Chan <i>et al.</i> ³¹ Fan <i>et al.</i> ⁴⁵ Hamman <i>et al.</i> ⁵⁹ Hamman <i>et al.</i> ⁶⁰ Hargestam <i>et al.</i> ⁶¹ Hunt <i>et al.</i> ⁴⁴ Kerner <i>et al.</i> ⁶² Kobayashi <i>et al.</i> ⁵² Kobayashi <i>et al.</i> ⁶³ Mondrup <i>et al.</i> ³³ O'Leary <i>et al.</i> ⁶⁴ Patterson <i>et al.</i> ¹⁰ Patterson <i>et al.</i> ⁶⁵ Petrosoniak <i>et al.</i> ³⁰ Shrestha <i>et al.</i> ³⁴ Sørensen <i>et al.</i> ⁶⁶ Theilen <i>et al.</i> ⁴¹ Ullman <i>et al.</i> ⁶⁷ Walsh <i>et al.</i> ⁶⁸ Whitfill <i>et al.</i> ⁶⁹ Zimmermann <i>et al.</i> ⁷⁰ Bradley <i>et al.</i> ⁷¹ Paltved <i>et al.</i> ³² Bredmose <i>et al.</i> ³⁵ Couto <i>et al.</i> ³⁶ Geis <i>et al.</i> ⁷² Lakissian <i>et al.</i> ⁷³ Shah <i>et al.</i> ³⁷ Wong <i>et al.</i> ⁷⁴ Petrosoniak <i>et al.</i> ⁷⁵ Aljahany <i>et al.</i> ⁷⁶
Quantitative data such as number of departures without being seen, median waiting time	0	
Accident reports	3 (4)	Chan <i>et al.</i> ³¹ Patterson <i>et al.</i> ¹⁰ Paltved <i>et al.</i> ³²
Cancellation rate	9 (11)	Chan <i>et al.</i> ³¹ Mondrup <i>et al.</i> ³³ Patterson <i>et al.</i> ¹⁰ Shrestha <i>et al.</i> ³⁴ Paltved <i>et al.</i> ³² Bredmose <i>et al.</i> ³⁵ Couto <i>et al.</i> ³⁶ Shah <i>et al.</i> ³⁷ Auerbach <i>et al.</i> ³⁸

Continued

Table 2 Continued

Safety-related parameters	n (%)	References
LST evaluation	24 (25)	Chan <i>et al.</i> ³¹ Hamman <i>et al.</i> ⁵⁹ Hamman <i>et al.</i> ⁶⁰ Hunt <i>et al.</i> ⁴⁴ Kerner <i>et al.</i> ⁶² Kobayashi <i>et al.</i> ⁵² Kobayashi <i>et al.</i> ⁶³ O'Leary <i>et al.</i> ⁶⁴ Patterson <i>et al.</i> ¹⁰ Shrestha <i>et al.</i> ³⁴ Theilen <i>et al.</i> ⁴¹ Ullman <i>et al.</i> ⁶⁷ Walsh <i>et al.</i> ⁶⁸ Whitfill <i>et al.</i> ⁶⁹ Zimmermann <i>et al.</i> ⁷⁰ Bradley <i>et al.</i> ⁷¹ Couto <i>et al.</i> ³⁶ Geis <i>et al.</i> ⁷² Lakissian <i>et al.</i> ⁷³ Shah <i>et al.</i> ³⁷ Wong <i>et al.</i> ⁷⁴ Petrosoniak <i>et al.</i> ⁷⁵ Aljahany <i>et al.</i> ⁷⁶ Zern <i>et al.</i> ⁷⁷ Shrestha <i>et al.</i> ⁷⁸ Gray <i>et al.</i> ⁴²
Others (survey, qualitative analysis)	5 (6.3)	Amiel <i>et al.</i> ²⁸ Abulebda <i>et al.</i> ²⁹ Petrosoniak <i>et al.</i> ³⁰ Wong <i>et al.</i> ⁷⁴ Sørensen <i>et al.</i> ⁷⁹
LST, latent safety threat.		

was disruptive or affected the participants in a negative way. These are the only studies exploring the impact of ISS on patients being managed in the ED.

Future safety of patients

The LST assessment was conducted in 24 studies (30%).

Kirkpatrick evaluation level

Most studies employed Kirkpatrick-related endpoints (n=75; 92%). Fourteen studies (17%) reported comparison with a preintervention assessment or a control group using knowledge as the outcome (KP1), mostly through a pre/post design. The assessment of KP2 level was provided for 10 studies (12%), and 27 (33%) using ISS as an intervention in EM to assess a level 3 outcome.

Twenty-two studies assessed a level 4 outcome. We considered LST identification as a level 4, because of the direct impact of this methodology on patient care. This classification was decided on consensus among the authors. For instance, in the study by Gray *et al.* (38), the authors showed that the identification of various LSTs regarding massive blood transfusion led to improved patient outcomes (measured by a decrease in the delays between massive

haemorrhage protocol activation and blood component administration). We identified three studies with a direct impact on patient outcomes (morbidity, mortality, length of stay). Using a pre–post study design, Steinemann *et al* demonstrated a significant improvement in mean team-work scores of real patient care in addition to a 16% reduction in the mean resuscitation time in the ED.⁴⁰ The other study that included a patient outcome analysis examined the effect of introducing a paediatric medical emergency team combined with regular ISS training on patient and system-level outcomes.⁴¹ Even though the decrease in paediatric intensive care unit mortality was nonsignificant, the authors found an association between ISS and a reduction of healthcare-related costs (level 5). Using a prospective observational design, Theilen *et al* showed that regular ISS training of ED healthcare professionals was associated with a reduction in costs and overall patient mortality. Another more recent study showed direct patient outcome improvement.⁴² The authors showed that a novel ISS-based quality improvement (QI) intervention for blood component administration in bleeding trauma patients led to a 21% mean reduction in time between massive haemorrhage protocol activation and blood component administration. The impact of changing the institution’s massive haemorrhage protocol after the identification of LSTs during ISS was explored in this study using a pre/post design.

No studies showed a negative impact when exploring the improvement of knowledge or skills. The efficacy of ISS was always superior when compared with any another training method or to a control group. We were unable to categorise 10 of our included studies (12.3%) within the 5 levels of KP, either because the studies assessed guideline adherence^{38 43} or because ISS was employed to assess a process or a system (table 3).^{44 45}

Indeed, for 24 studies (29%), ISS was used to explore and assess a procedure, an organisation, or guideline adherence and not to teach or train. Seven studies (8.6%) assessed the feasibility of ISS itself without exploring learning or safety outcomes.

DISCUSSION

This scoping review assessed the safety of ISS in EM.

ISS and patient safety

Conducting ISS within the chaotic and busy ED environment may raise concerns for the ongoing care of real patients. However, we found that few studies evaluated the impact of ISS on the safety of actual ED patients. Our scoping review focused on the available evidence regarding two safety concepts: ongoing ED patient safety during ISS, and the safety of future patients (long-term benefits of ISS). It is important for educators and ED professionals to be aware of the associated risks of ISS. Specific, rigorous guidelines are needed to help create a framework aiming to enhance ISS ‘ongoing’ safety. Surprisingly, we did not find any studies evaluating the

Table 3 Outcomes

Kirkpatrick level	n (%)	References
KP 1: satisfaction, feelings and perceptions	14	Bischof <i>et al.</i> ⁸⁰ Burke <i>et al.</i> ⁸¹ Couto <i>et al.</i> ¹³ Davison <i>et al.</i> ⁸² Gangadharan <i>et al.</i> ⁸³ Gundrosen <i>et al.</i> ⁸⁴ Hunt <i>et al.</i> ⁴⁴ Mannenbach <i>et al.</i> ⁸⁵ Meurling <i>et al.</i> ⁸⁶ O’Leary <i>et al.</i> ⁸⁷ Shrestha <i>et al.</i> ³⁴ Sørensen <i>et al.</i> ⁸⁸ Ullman <i>et al.</i> ⁶⁷ Katznelson <i>et al.</i> ⁸⁹
KP 2: knowledge and skills	10	Walsh <i>et al.</i> ⁶⁸ Bradley <i>et al.</i> ⁷¹ Thomas <i>et al.</i> ⁹⁰ Lemke <i>et al.</i> ⁹¹ Kalidindi <i>et al.</i> ⁹² Ben-Ari <i>et al.</i> ⁹³ Auerbach <i>et al.</i> ⁹⁴ Abualenain <i>et al.</i> ⁹⁵ Bischof <i>et al.</i> ⁸⁰ Sørensen <i>et al.</i> ⁸⁸
KP 3: behaviours	27	Abu-Sultaneh <i>et al.</i> ⁹⁶ Amiel <i>et al.</i> ²⁸ Armstrong <i>et al.</i> ⁹⁷ Auerbach <i>et al.</i> ³⁸ Barker <i>et al.</i> ⁹⁸ Bayouth <i>et al.</i> ⁹⁹ Bredmose <i>et al.</i> ³⁵ Campbell <i>et al.</i> ¹⁰⁰ Coggins <i>et al.</i> ¹⁰¹ Farah <i>et al.</i> ¹⁰² Generoso <i>et al.</i> ¹⁰³ Jørgens <i>et al.</i> ¹⁰⁴ Jung <i>et al.</i> ¹⁰⁵ Katznelson <i>et al.</i> ¹⁰⁶ Lakissian <i>et al.</i> ⁷³ Miller <i>et al.</i> ¹⁰⁷ O’Leary <i>et al.</i> ⁸⁷ Patterson <i>et al.</i> ⁶⁵ Petrosoniak <i>et al.</i> ³⁰ Pirie <i>et al.</i> ¹⁰⁸ Qian ¹⁰⁹ <i>et al.</i> Saqe-Rockoff <i>et al.</i> ¹¹⁰ Truta <i>et al.</i> ¹¹¹ Wong <i>et al.</i> ⁷⁴ Zern <i>et al.</i> ⁷⁷ Zimmermann <i>et al.</i> ⁷⁰ Walsh <i>et al.</i> ⁶⁸

Continued

**Table 3** Continued

Kirkpatrick level	n (%)	References
KP4: patient outcome	22	Abulebda <i>et al.</i> ²⁹ Aljahany <i>et al.</i> ⁷⁶ Barni <i>et al.</i> ¹¹² Chan <i>et al.</i> ³¹ Couto <i>et al.</i> ³⁶ Geis <i>et al.</i> ⁷² Gray <i>et al.</i> ⁴² Hamman <i>et al.</i> ⁵⁹ Hamman <i>et al.</i> ⁶⁰ Hargestam <i>et al.</i> ⁶¹ Kerner RL <i>et al.</i> ⁶² Kobayashi <i>et al.</i> ⁶³ Lavelle <i>et al.</i> ¹¹³ Paltved <i>et al.</i> ³² Patterson <i>et al.</i> ¹⁰ Shah <i>et al.</i> ³⁷ Shrestha <i>et al.</i> ³⁴ Steinemann <i>et al.</i> ⁴⁰ Wang <i>et al.</i> ¹¹⁴ Whitfill <i>et al.</i> ⁶⁹ Petrosoniak <i>et al.</i> ⁷⁵ Theilen <i>et al.</i> ⁴¹ Walsh <i>et al.</i> ⁶⁸
KP5: costs	2	Theilen <i>et al.</i> ⁴¹ Petrosoniak <i>et al.</i> ⁷⁵
Other, NA	10	Hunt <i>et al.</i> ⁴⁴ Kobayashi <i>et al.</i> ⁵² Mondrup <i>et al.</i> ³³ Kessler <i>et al.</i> ¹¹⁵ Siegel <i>et al.</i> ⁴³ Stevens <i>et al.</i> ¹¹⁶ Wieck <i>et al.</i> ¹¹⁷ Nadkarni <i>et al.</i> ¹¹⁸ Auerbach <i>et al.</i> ¹¹⁹ O'Leary <i>et al.</i> ⁸⁷
NA, not available.		

impact of ISS on ongoing patient care, clinical adverse events, wait times or patients leaving without being seen. Some editorials warn against these risks,^{16–18} but we found no studies exploring nor demonstrating the potential consequences or risks of ISS in a busy ED for the ongoing care of patients. Most studies seemingly overvalue the ‘future’ safety of patients while undervaluing the ‘ongoing’ safety of ED patients during ISS. This seems crucial because ISS is increasingly used to teach and train EM professionals. As for any method aiming at improving the quality of care, the risks for patients should be controlled. In that sense, any type of clinical research may jeopardise patient safety. For example, the time it takes to consent and enrol a patient in a therapeutic trial could impact ED safety measures. The time-sensitive nature of EM research has been reported in research primers.¹⁴ Our scoping review shows the absence of scientific evidence from studies with rigorous methodology to confirm the absence of risks for patients. The Foundation

for Healthcare Simulation Safety published a 10-item ‘pledge’ of ‘best practices’ for simulation programmes to reduce simulation-related hazards.^{17 46} However, even if those principles seem logical, they are not evidence based.

Educators and simulation teams should consider guidelines from existing literature on QI to ensure the sustainability of these interventions. In the case of ISS, the ongoing safety of patients could be referred to as a ‘balancing measure’. In the QI literature, balancing measures represent checkpoints that ensure there are no potential unintended consequences (risk to ongoing patient safety) resulting from an intervention (ISS).^{37 47–51}

Patient-reported measures, which appear to be underutilised based on our study results, represent a useful lens to understand the value of ISS more comprehensively. For example, at one investigator’s institution (AP), a Code Orange simulation during which dedicated volunteers checked in with patients waiting and patients had positive perceptions regarding this ongoing training (unpublished data).

Outcomes

One would think that being closer to the patient and to the patient care facility would make it easier and more intuitive to assess ISS-related patient outcomes, but only two studies assessed direct patient outcomes.^{41 42} This raises the question of the relevance of the Kirkpatrick pyramid. Indeed, we deemed appropriate to classify the identification of LST as a level 4 on the KP scale, because it is directly linked to improving the quality of care. This decision was reached on consensus among the authors and is debatable because direct patient outcomes are not measured. However, we found data exploring the improvement of care following the identification of LSTs to validate our choice of classification.^{41 42} We believe even though this was not directly measured in every study, the identification of LSTs leads to improved patient outcomes by reducing medical errors.⁵² We were unable to classify a great number of studies using this tool. As other authors have previously suggested, it may be wise to adopt a new lexicon and new endpoints to assess the impact of ISS.^{53 54} Some studies did, however, explore level KP4 and even KP5.⁴¹ Their inspiring results should encourage researchers to pursue their efforts and lead projects aiming at exploring higher levels of the KP pyramid.

Study methodology limitations

As highlighted in previous studies, studies exploring the impact of simulation-based interventions are still needed.^{2 55–57} We observed that even though the included studies were quite recent, as ISS seems to have emerged in the last 20 years, the vast majority used observational prospective designs. With ISS, the proximity between the intervention and the possible patient-centred outcomes is inspiring and educators should seek collaboration with methodologists to elaborate research protocols leaning

to prove the translational dimension of ISS. Simulation should be about improving patient outcomes, and the causative effect of isolating ISS should be a priority.

Strengths and limitations

Because there is no mesh word for 'ISS' we conducted a search on simulation and EM followed by more detailed reviews for 'in situ' specific articles, which made the screening process more complex. Furthermore, the terms used to describe ISS are highly heterogeneous. It is possible that this resulted in missed articles that otherwise should have been included. However, the corollary is this process required a close assessment of each article resulting in a more thorough review.

Also, several systematic reviews were excluded from our analysis. However, we reviewed and cross-referenced their reference list against our literature search. We found no discrepancies, supporting the accuracy of our approach.

Conclusion

Our scoping review identified major knowledge gaps related to the safe delivery of ISS in the ED, which require further investigation. Patient safety during ISS is often evaluated in the context of identifying or mitigating LSTs with little focus on the potential impact and risks to actual ED patients. For all these reasons, our research team sought to address this issue in a mix-method study, which will aim at demonstrating the safety and impact of ISS.⁵⁸

Author affiliations

¹Département de médecine familiale et de médecine d'urgence, Université Laval Faculté de médecine, Québec, Québec, Canada

²Emergency Department, CHU de Québec-Université Laval, Québec, Québec, Canada

³Emergency Department, CHU Cochin- Université de Paris, APHP, Paris, France

⁴Axe Santé des populations et pratiques optimales en santé, Centre de recherche du CHU de Québec-Université Laval, Québec, Québec, Canada

⁵Centre d'excellence sur le vieillissement de Québec, Québec, Québec, Canada

⁶Division of Emergency Medicine, Department of Medicine, University of Toronto, Toronto, Ontario, Canada

⁷Department of Emergency Medicine, St. Michael's Hospital, Toronto, Ontario, Canada

Twitter Marcel Emond @marcel_mond

Acknowledgements The authors would like to thank Chantal Beaugard for the valuable help on conducting a rigorous library search. The authors would like to thank the Société Française de Médecine d'Urgence (SFMU) for supporting this research project through a research award (JT). We would also like to thank the Fondation du CHU de Québec-Université Laval and the Emergency Physicians from the Hôpital de l'Enfant-Jésus, who also supported this work.

Contributors JT designed the study with the help of VB and under the supervision of MÉ. MÉ obtained research funding. JT, WL, GM and EJ assessed each study and performed data collection. JT drafted the manuscript, and VB, WL, GM, EJ, ER-D, AP and MÉ contributed substantially to its revision. MÉ accepts full responsibility for the work and/or the conduct of the study, had access to the data and controlled the decision to publish.

Funding This work was supported by the Fondation du CHU de Québec-Université Laval, grant number 3967. JT received a Fellowship award from the Société française de médecine d'urgence.

Disclaimer The sponsors were in no way involved in the design of this study protocol.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Ethics approval Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Marcel Emond <http://orcid.org/0000-0001-7158-8110>

REFERENCES

- Freund Y, Goulet H, Leblanc J, *et al.* Effect of systematic physician cross-checking on reducing adverse events in the emergency department: the CHARMED cluster randomized trial. *JAMA Intern Med* 2018;178:812–9.
- Cheng A, Bhanji F. A call to action: the future of simulation-based research in emergency medicine in Canada. *CJEM* 2020;22:8–10.
- Bond WF, Lammers RL, Spillane LL, *et al.* The use of simulation in emergency medicine: a research agenda. *Acad Emerg Med* 2007;14:353–63.
- Carenzo L, Ragozzino F, Colombo D. Virtual laboratory and imaging: an online simulation tool to enhance Hospital disaster preparedness training experience. *Eur J Emerg Med* 2018;25:128–33.
- Lee J, Song Y, Oh J, *et al.* Smartwatch feedback device for high-quality chest compressions by a single rescuer during infant cardiac arrest: a randomized, controlled simulation study. *Eur J Emerg Med* 2019;26:266–71.
- Chauvin A, Truchot J, Bafeta A. Randomized controlled trials of simulation-based interventions in emergency medicine: a methodological review. *Intern Emerg Med* 2018;13:433–44.
- Anderson R, Sebaldt A, Lin Y, *et al.* Optimal training frequency for acquisition and retention of high-quality CPR skills: a randomized trial. *Resuscitation* 2019;135:153–61.
- Boet S, Borges BCR, Naik VN, *et al.* Complex procedural skills are retained for a minimum of 1 yr after a single high-fidelity simulation training session †. *Br J Anaesth* 2011;107:533–9.
- Truchot J. Point of care simulation: towards new pragmatic simulation training. *Eur J Emerg Med* 2020;27:79–80.
- Patterson MD, Geis GL, Falcone RA, *et al.* In situ simulation: detection of safety threats and teamwork training in a high risk emergency department. *BMJ Qual Saf* 2013;22:468–77.
- Kobayashi L, Patterson MD, Overly FL, *et al.* Educational and research implications of portable human patient simulation in acute care medicine. *Acad Emerg Med* 2008;15:1166–74.
- Durning SJ, Artino AR. Situativity theory: a perspective on how participants and the environment can interact: AMEE guide No. 52. *Med Teach* 2011;33:188–99.
- Couto T, Kerrey BT, Taylor RG. Teamwork skills in actual, in situ, and in-center pediatric emergencies: performance levels across settings and perceptions of comparative educational impact. *Simul Healthc* 2015;10:76–84.
- American College of Emergency Physicians. Emergency Care Research - A Primer 2012 [2022-05-16]. Available: <https://www.emfoundation.org/globalassets/general/pdfs/acep-research-primer-book-pdf.pdf>
- School of Medicine and Public Health - University of Wisconsin-Madison. Conducting Research in the ED 2022 [2022-05-16]. Available: <https://www.emed.wisc.edu/conducting-research-ed>

- 16 Bajaj K, Minors A, Walker K, *et al.* "No-Go Considerations" for In Situ Simulation Safety. *Simul Healthc* 2018;13:221–4.
- 17 Raemer D, Hannenberg A, Mullen A. Simulation safety first: an imperative. *Adv Simul* 2018;3:25.
- 18 Torrie J, Cumin D, Sheridan J, *et al.* Fake and expired medications in simulation-based education: an underappreciated risk to patient safety. *BMJ Qual Saf* 2016;25:917–20.
- 19 Newstrom JW. Evaluating training programs: the four levels, by Donald L. Kirkpatrick. (1994). San Francisco: Berrett-Koehler. 229 PP. *Hum Resour Dev Q* 1995;6:317–20.
- 20 Phillips JJ, Stone R, Phillips PP. *The human resources Scorecard: measuring the return on Investment 2001*, 2001.
- 21 Peters MDJ, Godfrey CM, Khalil H, *et al.* Guidance for conducting systematic scoping reviews. *Int J Evid Based Healthc* 2015;13:141–6.
- 22 Liberati A, Altman DG, Tetzlaff J, *et al.* The PRISMA statement for reporting systematic reviews and meta-analyses of studies that evaluate healthcare interventions: explanation and elaboration. *BMJ* 2009;339:b2700.
- 23 Patterson MD, Blike GT, Nadkarni VM. Advances in Patient Safety In Situ Simulation: Challenges and Results. In: *Advances in patient safety: new directions and alternative approaches (vol 3: performance and tools)*. Rockville (MD): Agency for Healthcare Research and Quality (US), 2008.
- 24 Aggarwal R, Mytton OT, Derbrew M, *et al.* Training and simulation for patient safety. *Quality and Safety in Health Care* 2010;19:i34–43.
- 25 MacKinnon RJ, Pukk-Härenstam K, Kennedy C, *et al.* A novel approach to explore Safety-I and Safety-II perspectives in in situ simulations—the structured what if functional resonance analysis methodology. *Adv Simul* 2021;6:21.
- 26 Sørup CM, Jacobsen P, Forberg JL. Evaluation of emergency department performance – a systematic review on recommended performance and quality-in-care measures. *Scand J Trauma Resusc Emerg Med* 2013;21:62.
- 27 Hansen K, Boyle A, Holroyd B, *et al.* Updated framework on quality and safety in emergency medicine. *Emerg Med J* 2020;37:437–42.
- 28 Amiel I, Simon D, Merin O, *et al.* Mobile in situ simulation as a tool for evaluation and improvement of trauma treatment in the emergency department. *J Surg Educ* 2016;73:121–8.
- 29 Abulebda K, Lutfi R, Whitfill T, *et al.* A collaborative in situ simulation-based pediatric readiness improvement program for community emergency departments. *Acad Emerg Med* 2018;25:177–85.
- 30 Petrosoniak A, Ryzynski A, Lebovic G, *et al.* Cricothyroidotomy in situ simulation curriculum (CRIC study): training residents for rare procedures. *Simul Healthc* 2017;12:76–82.
- 31 Chan S, Babcock L, Geis G, *et al.* In situ simulation to mitigate threats to participation in a multicenter clinical trial in High-Acuity, low-frequency setting. *Sim Healthc* 2019;14:1–9.
- 32 Paltved C, Bjerregaard AT, Krogh K, *et al.* Designing in situ simulation in the emergency department: evaluating safety attitudes amongst physicians and nurses. *Adv Simul* 2017;2:4.
- 33 Mondrup F, Brabrand M, Folkestad L, *et al.* In-Hospital resuscitation evaluated by in situ simulation: a prospective simulation study. *Scand J Trauma Resusc Emerg Med* 2011;19:55.
- 34 Shrestha R, Shrestha AP, Shrestha SK, *et al.* Interdisciplinary in situ simulation-based medical education in the emergency department of a teaching hospital in Nepal. *Int J Emerg Med* 2019;12:19.
- 35 Bredmose PP, Hagemo J, Røislien J, *et al.* In situ simulation training in helicopter emergency medical services: feasible for on-call crews? *Adv Simul* 2020;5:7.
- 36 Couto TB, Barreto JKS, Marcon FC, *et al.* Detecting latent safety threats in an interprofessional training that combines in situ simulation with task training in an emergency department. *Adv Simul* 2018;3:23.
- 37 Shah SJ, Cusumano C, Ahmed S. In situ simulation to assess pediatric tracheostomy care safety: a novel multicenter quality improvement program. *Otolaryngol Head Neck Surg* 2020;163:250–8.
- 38 Auerbach M, Roney L, Aysseh A, *et al.* In situ pediatric trauma simulation: assessing the impact and feasibility of an interdisciplinary pediatric in situ trauma care quality improvement simulation program. *Pediatr Emerg Care* 2014;30:884–91.
- 39 Petrosoniak A, Fan M, Hicks CM, *et al.* Trauma resuscitation using in situ simulation team training (trust) study: latent safety threat evaluation using framework analysis and video review. *BMJ Qual Saf* 2021;30:739–746.
- 40 Steinemann S, Berg B, Skinner A, *et al.* In situ, multidisciplinary, simulation-based teamwork training improves early trauma care. *J Surg Educ* 2011;68:472–7.
- 41 Theilen U, Fraser L, Jones P, *et al.* Regular in-situ simulation training of paediatric medical emergency team leads to sustained improvements in hospital response to deteriorating patients, improved outcomes in intensive care and financial savings. *Resuscitation* 2017;115:61–7.
- 42 Gray A, Chartier LB, Pavenski K, *et al.* The clock is ticking: using in situ simulation to improve time to blood administration for bleeding trauma patients. *CJEM* 2021;23:54–62.
- 43 Siegel NA, Kobayashi L, Dunbar-Viveiros JA, *et al.* In situ medical simulation investigation of emergency department procedural sedation with randomized trial of experimental bedside clinical process guidance intervention. *Simul Healthc* 2015;10:146–53.
- 44 Hunt EA, Hohenhaus SM, Luo X, *et al.* Simulation of pediatric trauma stabilization in 35 North Carolina emergency departments: identification of targets for performance improvement. *Pediatrics* 2006;117:641–8.
- 45 Fan M, Petrosoniak A, Pinkney S, *et al.* Study protocol for a framework analysis using video review to identify latent safety threats: trauma resuscitation using in situ simulation team training (trust). *BMJ Open* 2016;6:e013683.
- 46 Foundation for Healthcare simulation safety. Foundation for Healthcare simulation safety 2021 [2021/09]. Available: <https://healthcaresimulationsafety.org>
- 47 Boulkedid R, Abdoul H, Loustau M, *et al.* Using and reporting the Delphi method for selecting healthcare quality indicators: a systematic review. *PLoS One* 2011;6:e20476.
- 48 Colman N, Newman JW, Nishisaki A, *et al.* Translational simulation improves compliance with the NEAR4KIDS airway safety bundle in a single-center PICU. *Pediatr Qual Saf* 2021;6:e409.
- 49 McKinley KW, Babineau J, Roskind CG, *et al.* Discrete event simulation modelling to evaluate the impact of a quality improvement initiative on patient flow in a paediatric emergency department. *Emerg Med J* 2020;37:193–9.
- 50 Tate K, Lee S, Rowe BH, *et al.* Quality indicators for older persons' transitions in care: a systematic review and Delphi process. *Can J Aging* 2022;41:1–15.
- 51 Chartier LB, Stang AS, Vaillancourt S, *et al.* Quality improvement primer Part 2: executing a quality improvement project in the emergency department. *CJEM* 2018;20:532–8.
- 52 Kobayashi L, Dunbar-Viveiros JA, Devine J, *et al.* Pilot-phase findings from high-fidelity in situ medical simulation investigation of emergency department procedural sedation. *Simul Healthc* 2012;7:81–94.
- 53 Brazil V. Translational simulation: not 'where?' but 'why?' A functional view of in situ simulation. *Adv Simul* 2017;2:20.
- 54 Posner GD, Clark ML, Grant VJ. Simulation in the clinical setting: towards a standard lexicon. *Adv Simul* 2017;2:15.
- 55 Cook DA. How much evidence does it take? A cumulative meta-analysis of outcomes of simulation-based education. *Med Educ* 2014;48:750–60.
- 56 Cook DA, Brydges R, Zendejas B, *et al.* Technology-enhanced simulation to assess health professionals: a systematic review of validity evidence, research methods, and reporting quality. *Acad Med* 2013;88:872–83.
- 57 Cheng A, Kessler D, Mackinnon R, *et al.* Reporting guidelines for health care simulation research: extensions to the CONSORT and STROBE statements. *Simul Healthc* 2016;11:238–48.
- 58 Truchot J, Boucher V, Raymond-Dufresne Eliane, *et al.* Evaluation of the feasibility and impacts of in situ simulation in emergency medicine—a mixed-method study protocol. *BMJ Open* 2021;11:e040360.
- 59 Hamman WR, Beaudin-Seiler BM, Beaubien JM, *et al.* Using simulation to identify and resolve threats to patient safety. *Am J Manag Care* 2010;16:e145–50.
- 60 Hamman WR, Beaudin-Seiler BM, Beaubien JM, *et al.* Using in situ simulation to identify and resolve latent environmental threats to patient safety: case study involving operational changes in a labor and delivery ward. *Qual Manag Health Care* 2010;19:226–30.
- 61 Hårgestam M, Lindkvist M, Jacobsson M, *et al.* Trauma teams and time to early management during in situ trauma team training. *BMJ Open* 2016;6:e009911.
- 62 Kerner RL, Gallo K, Cassara M, *et al.* Simulation for operational readiness in a new freestanding emergency department: strategy and tactics. *Simul Healthc* 2016;11:345–56.
- 63 Kobayashi L, Parchuri R, Gardiner FG, *et al.* Use of in situ simulation and human factors engineering to assess and improve emergency department clinical systems for timely telemetry-based detection of life-threatening arrhythmias. *BMJ Qual Saf* 2013;22:72–83.
- 64 O'Leary F, McGarvey K, Christoff A, *et al.* Identifying incidents of suboptimal care during paediatric emergencies—an observational

- study utilising in situ and simulation centre scenarios. *Resuscitation* 2014;85:431–6.
- 65 Patterson MD, Geis GL, LeMaster T, *et al.* Impact of multidisciplinary simulation-based training on patient safety in a paediatric emergency department. *BMJ Qual Saf* 2013;22:383–93.
- 66 Sørensen JL, Van der Vleuten C, Lindschou J, *et al.* 'In situ simulation' versus 'off site simulation' in obstetric emergencies and their effect on knowledge, safety attitudes, team performance, stress, and motivation: study protocol for a randomized controlled trial. *Trials* 2013;14:220.
- 67 Ullman E, Kennedy M, Di Delupis FD, *et al.* The Tuscan mobile simulation program: a description of a program for the delivery of in situ simulation training. *Intern Emerg Med* 2016;11:837–41.
- 68 Walsh BM, Gangadharan S, Whitfill T, *et al.* Safety threats during the care of infants with hypoglycemic seizures in the emergency department: a multicenter, simulation-based prospective cohort study. *J Emerg Med* 2017;53:467–74.
- 69 Whitfill T, Gawel M, Auerbach M. A simulation-based quality improvement initiative improves pediatric readiness in community hospitals. *Pediatr Emerg Care* 2018;34:431–5.
- 70 Zimmermann K, Holzinger IB, Ganassi L, *et al.* Inter-professional in-situ simulated team and resuscitation training for patient safety: description and impact of a programmatic approach. *BMC Med Educ* 2015;15:189.
- 71 Bradley NL, Innes K, Dakin C, *et al.* Multidisciplinary in-situ simulation to evaluate a rare but high-risk process at a level 1 trauma centre: the "Mega-Sim" approach. *Can J Surg* 2018;61:357–60.
- 72 Geis GL, Pio B, Pendergrass TL, *et al.* Simulation to assess the safety of new healthcare teams and new facilities. *Simul Healthc* 2011;6:125–33.
- 73 Lakissian Z, Sabouneh R, Zeineddine R, *et al.* In-Situ simulations for COVID-19: a safety II approach towards resilient performance. *Adv Simul* 2020;5:15.
- 74 Wong AH, Auerbach MA, Ruppel H, *et al.* Addressing dual patient and staff safety through a team-based standardized patient simulation for agitation management in the emergency department. *Simul Healthc* 2018;13:154–62.
- 75 Petrosoniak A, Fan M, Hicks CM, *et al.* Trauma resuscitation using in situ simulation team training (trust) study: latent safety threat evaluation using framework analysis and video review. *BMJ Qual Saf* 2021;30:739–46.
- 76 Aljahany M, Alassaf W, Alibrahim AA, *et al.* Use of in situ simulation to improve emergency department readiness for the COVID-19 pandemic. *Prehosp Disaster Med* 2021;36:6–13.
- 77 Zern SC, Marshall WJ, Shewokis PA, *et al.* Use of simulation as a needs assessment to develop a focused team leader training curriculum for resuscitation teams. *Adv Simul* 2020;5:6.
- 78 Shrestha AP, Shrestha A, Sonnenberg T, *et al.* COVID-19 Emergency Department Protocols: Experience of Protocol Implementation Through in-situ Simulation] & *Open Access Emerg Med* 2020;12:293–303.
- 79 Sørensen JL, Navne LE, Martin HM, *et al.* Clarifying the learning experiences of healthcare professionals with in situ and off-site simulation-based medical education: a qualitative study: Table 1. *BMJ Open* 2015;5:e008345.
- 80 Bischof JJ, Panchal AR, Finnegan GI, *et al.* Creation and validation of a novel mobile simulation laboratory for high fidelity, prehospital, difficult airway simulation. *Prehosp Disaster Med* 2016;31:465–70.
- 81 Burke RV, Demeter NE, Goodhue CJ, *et al.* Qualitative assessment of simulation-based training for pediatric trauma resuscitation. *Surgery* 2017;161:1357–66.
- 82 Davison M, Kinnear FB, Fullbrook P. Evaluation of a multiple-encounter in situ simulation for orientation of staff to a new paediatric emergency service: a single-group pretest/post-test study. *BMJ Simul Technol Enhanc Learn* 2017;3:149–53.
- 83 Gangadharan S, Tiyyagura G, Gawel M, *et al.* A Grounded theory qualitative analysis of interprofessional Providers' perceptions on caring for critically ill infants and children in pediatric and general emergency departments. *Pediatr Emerg Care* 2018;34:578–83.
- 84 Gundrosen S, Andenæs E, Aadahl P, *et al.* Team talk and team activity in simulated medical emergencies: a discourse analytical approach. *Scand J Trauma Resusc Emerg Med* 2016;24:135.
- 85 Mannenbach MS, Fahje CJ, Sunga KL, *et al.* An in situ simulation-based training approach to active Shooter response in the emergency department. *Disaster Med Public Health Prep* 2019;13:345–52.
- 86 Meurling L, Hedman L, Lidfeldt K-J, *et al.* Comparison of high- and low equipment fidelity during paediatric simulation team training: a case control study. *BMC Med Educ* 2014;14:221.
- 87 O'Leary FM, Hokin B, Enright K, *et al.* Treatment of a simulated child with anaphylaxis: an in situ two-arm study. *J Paediatr Child Health* 2013;49:541–7.
- 88 Sørensen JL, van der Vleuten C, Rosthøj S, *et al.* Simulation-Based multiprofessional obstetric anaesthesia training conducted in situ versus off-site leads to similar individual and team outcomes: a randomised educational trial. *BMJ Open* 2015;5:e008344.
- 89 Katznelson JH, Mills WA, Forsythe CS, *et al.* Project Cape: a high-fidelity, in situ simulation program to increase critical access hospital emergency department provider comfort with seriously ill pediatric patients. *Pediatr Emerg Care* 2014;30:397–402.
- 90 Thomas AA, Uspal NG, Oron AP, *et al.* Perceptions on the impact of a Just-in-Time room on trainees and Supervising physicians in a pediatric emergency department. *J Grad Med Educ* 2016;8:754–8.
- 91 Lemke D, Feux D, Doughty C. A novel mechanism for simulation of partial seizures in an infant. *Simul Healthc* 2012;7:359–61.
- 92 Kalidindi S, Kirk M, Griffith E. In-Situ simulation enhances emergency preparedness in pediatric care practices. *Cureus* 2018;10:e3389.
- 93 Ben-Ari M, Chayen G, Steiner IP, *et al.* The effect of in situ simulation training on the performance of tasks related to patient safety during sedation. *J Anesth* 2018;32:300–4.
- 94 Auerbach M, Kessler D, Foltin JC. Repetitive pediatric simulation resuscitation training. *Pediatr Emerg Care* 2011;27:29–31.
- 95 Abualenain JT, Al-Alawi MM. Simulation-Based training in Ebola personal protective equipment for healthcare workers: experience from King Abdulaziz university hospital in Saudi Arabia. *J Infect Public Health* 2018;11:796–800.
- 96 Abu-Sultaneh S, Whitfill T, Rowan CM, *et al.* Improving simulated pediatric airway management in community emergency departments using a collaborative program with a pediatric academic medical center. *Respir Care* 2019;64:1073–81.
- 97 Armstrong P, Peckler B, Pilkinton-Ching J, *et al.* Effect of simulation training on nurse leadership in a shared leadership model for cardiopulmonary resuscitation in the emergency department. *Emerg Med Australas* 2021;33:255–61.
- 98 Barker LT, Bond WF, Vincent AL, *et al.* A novel in situ simulation framework for introduction of a new technology: the 3-Act-3-Debrief model. *Adv Simul* 2020;5:25.
- 99 Bayouth L, Ashley S, Brady J, *et al.* An in-situ simulation-based educational outreach project for pediatric trauma care in a rural trauma system. *J Pediatr Surg* 2018;53:367–71.
- 100 Campbell DM, Poost-Foroosh L, Pavenski K, *et al.* Simulation as a toolkit—understanding the perils of blood transfusion in a complex health care environment. *Adv Simul* 2016;1:32.
- 101 Coggins AR, Nottingham C, Byth K, *et al.* Randomised controlled trial of simulation-based education for mechanical cardiopulmonary resuscitation training. *Emergency Medicine Journal* 2019;36:266–72.
- 102 Farah R, Stiner E, Zohar Z, *et al.* Cardiopulmonary resuscitation surprise drills for assessing, improving and maintaining cardiopulmonary resuscitation skills of hospital personnel. *Eur J Emerg Med* 2007;14:332–6.
- 103 Generoso JR, Latoures RE, Acar Y, *et al.* Simulation training in early emergency response (steer). *J Contin Educ Nurs* 2016;47:255–63.
- 104 Jörgens M, Königer J, Kanz K-G, *et al.* Testing mechanical chest compression devices of different design for their suitability for prehospital patient transport - a simulator-based study. *BMC Emerg Med* 2021;21:18.
- 105 Jung D, Carman M, Aga R, *et al.* Disaster preparedness in the emergency department using in situ simulation. *Adv Emerg Nurs J* 2016;38:56–68.
- 106 Katznelson JH, Wang J, Stevens MW, *et al.* Improving pediatric preparedness in critical access hospital emergency departments: impact of a longitudinal in situ simulation program. *Pediatr Emerg Care* 2018;34:17–20.
- 107 Miller D, Crandall C, Washington C, *et al.* Improving teamwork and communication in trauma care through in situ simulations. *Acad Emerg Med* 2012;19:608–12.
- 108 Pirie J, Cardenas S, Seleem W, *et al.* The use of statistical process control charts to evaluate interprofessional education sessions embedded into a pediatric emergency in situ resuscitation program. *Simul Healthc* 2019;14:121–8.
- 109 Qian J, Wang Y, Zhang Y, *et al.* A Survey of the First-Hour Basic Care Tasks of Severe Sepsis and Septic Shock in Pediatric Patients and an Evaluation of Medical Simulation on Improving the Compliance of the Tasks. *J Emerg Med* 2016;50:239–45.
- 110 Saqe-Rockoff A, Ciardiello AV, Schubert FD. Low-Fidelity, in-situ pediatric resuscitation simulation improves RN competence and self-efficacy. *J Emerg Nurs* 2019;45:538–44.



- 111 Truta TS, Boeriu CM, Copotoiu S-M, *et al.* Improving nontechnical skills of an interprofessional emergency medical team through a one day crisis resource management training. *Medicine* 2018;97:e11828.
- 112 Barni S, Mori F, Giovannini M, *et al.* In situ simulation in the management of anaphylaxis in a pediatric emergency department. *Intern Emerg Med* 2019;14:127–32.
- 113 Lavelle M, Attoe C, Tritschler C, *et al.* Managing medical emergencies in mental health settings using an interprofessional in-situ simulation training programme: a mixed methods evaluation study. *Nurse Educ Today* 2017;59:103–9.
- 114 Wang CJ, Lin SY, Tsai SH, *et al.* Implications of long-term low-fidelity in situ simulation in acute care and association with a reduction in unexpected cardiac arrests: a retrospective research study. *PLoS One* 2019;14:e0213789.
- 115 Kessler DO, Walsh B, Whitfill T, *et al.* Disparities in adherence to pediatric sepsis guidelines across a spectrum of emergency departments: a multicenter, cross-sectional observational in situ simulation study. *J Emerg Med* 2016;50:403–15.
- 116 Stevens AD, Hernandez C, Jones S, *et al.* Color-Coded prefilled medication syringes decrease time to delivery and dosing errors in simulated prehospital pediatric resuscitations: a randomized crossover trial. *Resuscitation* 2015;96:85–91.
- 117 Wieck MM, McLaughlin C, Chang TP, *et al.* Self-Assessment of team performance using T-NOTECHS in simulated pediatric trauma resuscitation is not consistent with expert assessment. *The American Journal of Surgery* 2018;216:630–5.
- 118 Nadkarni LD, Roskind CG, Auerbach MA, *et al.* The development and validation of a Concise instrument for formative assessment of team leader performance during simulated pediatric resuscitations. *Sim Healthcare* 2018;13:77–82.
- 119 Auerbach M, Brown L, Whitfill T, *et al.* Adherence to pediatric cardiac arrest guidelines across a spectrum of fifty emergency departments: a prospective, in situ, simulation-based study. *Acad Emerg Med* 2018;25:1396–408.