



# Dauricine Attenuates Vascular Endothelial Inflammation Through Inhibiting NF- $\kappa$ B Pathway

Ji Hu<sup>1,2†</sup>, Ru Chen<sup>1,2†</sup>, Jie An<sup>1,2,3</sup>, Yilong Wang<sup>1,2,4</sup>, Minglu Liang<sup>1,2</sup> and Kai Huang<sup>1,2\*</sup>

<sup>1</sup>Department of Cardiology, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China, <sup>2</sup>Clinic Center of Human Gene Research, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, China, <sup>3</sup>Department of Cardiology, Handan First Hospital, Handan, China, <sup>4</sup>Department of Cardiology, The First Affiliated Hospital of Sun Yat-Sen University, Guangzhou, China

## OPEN ACCESS

### Edited by:

Yun Zhang,  
Shandong University, China

### Reviewed by:

Shuangxi Wang,  
Shandong University, China  
Wared Nour-Eldine,  
Qatar Biomedical Research Institute,  
Qatar  
Mingxiang Zhang,  
Shandong University, China

### \*Correspondence:

Kai Huang  
huangkai\_hust@126.com

<sup>†</sup>These authors contributed equally to this work and shared first authorship

### Specialty section:

This article was submitted to  
Inflammation Pharmacology,  
a section of the journal  
Frontiers in Pharmacology

**Received:** 15 August 2021

**Accepted:** 29 October 2021

**Published:** 01 December 2021

### Citation:


Hu J, Chen R, An J, Wang Y, Liang M and Huang K (2021) Dauricine Attenuates Vascular Endothelial Inflammation Through Inhibiting NF- $\kappa$ B Pathway. *Front. Pharmacol.* 12:758962. doi: 10.3389/fphar.2021.758962

Endothelial cells are the fundamental components of blood vessels that regulate several physiological processes including immune responses, angiogenesis, and vascular tone. Endothelial dysfunction contributes to the development of various diseases such as acute lung injury, and endothelial inflammation is a vital part of endothelial dysfunction. Dauricine is an extract isolated from *Menispermum dauricum* DC, a traditional Chinese medical plant that can be used for pharyngitis. In this work, we found that IL-1 $\beta$ -induced overexpression of intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and E-selectin was inhibited by dauricine in primary human umbilical vein endothelial cells (HUVECs). Correspondingly, adhesion of human acute monocytic leukemia cell line (THP-1) to HUVECs was decreased by dauricine. Further studies showed that dauricine inhibited the activation of nuclear factor- $\kappa$ B (NF- $\kappa$ B) pathway in HUVECs stimulated with IL-1 $\beta$ . *In vivo*, dauricine protected mice from lipopolysaccharide (LPS)-induced acute lung injury. In lung tissues, the activation of NF- $\kappa$ B pathway and the expression of its downstream genes (ICAM-1, VCAM-1, and E-selectin) were decreased by dauricine, consistent with what was found *in vitro*. In summary, we concluded that dauricine could alleviate endothelial inflammation by suppressing NF- $\kappa$ B pathway, which might serve as an effective candidate for diseases related with endothelial inflammation.

**Keywords:** dauricine, endothelial dysfunction, NF- $\kappa$ B pathway, acute lung injury, inflammation

## INTRODUCTION

The vascular endothelium forms the inner surface of the cardiovascular system. It is not only a natural blood-organ barrier but also an endocrine tissue that plays pivotal roles in immune responses, angiogenesis, hemostasis, and the regulation of vascular tone (Boulanger, 2016; Sturtzel, 2017). Endothelial dysfunction has been noticed in various pathological states, including atherosclerosis, hypertension, kidney disease, and sepsis (Gimbrone and Garcia-Cardena, 2016; Konukoglu and Uzun, 2017; Jourde-Chiche et al., 2019; Joffe et al., 2020). Aggravated endothelial inflammation, which is characterized by overexpressed cytokines and adhesion molecules such as intercellular adhesion molecule-1 (ICAM-1), vascular cell adhesion molecule-1 (VCAM-1), and selectins (Konukoglu and Uzun, 2017), is an important pathological process in endothelial dysfunction. Several molecules and pathways that regulate endothelial inflammation have been elucidated, among which the classic nuclear factor- $\kappa$ B (NF- $\kappa$ B) proinflammatory pathway attracted wide attention (Brasier, 2010; Rao et al., 2019).



Dauricine is an alkaloid extracted from the roots of *Menispermum dauricum* DC, and this compound is a traditional Chinese medicine that has been used to treat rheumatism (Du et al., 1986). In early studies, dauricine was found to exert an inhibitory effect in several inflammatory mouse models (Du et al., 1986). Later, multiple functions of dauricine were reported. Dauricine inhibited viability and induced cell apoptosis by inhibiting the PI3K/Akt pathway in renal carcinoma cells (Zhang et al., 2018). Lipopolysaccharide (LPS)-induced bone loss was inhibited by dauricine as mediated via the ROS/PP2A/NF-κB axis (Park et al., 2020). In a cerebral ischemia/reperfusion rat model, dauricine attenuated the inflammatory process by downregulating the expression of ICAM-1, TNF-α, and IL-1β (Yang et al., 2007). Studies performed both *in vivo* and *in vitro* have proven that dauricine protected against *Streptococcus pneumoniae* coinfecting with influenza virus H5N1 through NF-κB pathway (Li et al., 2018). In our previous work, we found that dauricine negatively regulated LPS-induced acute lung injury and immune response of macrophages (Qiao et al., 2019). However, the effect of dauricine on endothelial inflammation remains unclear.

Since endothelial inflammation is involved in multiple inflammatory diseases, targeting the inflammatory process in endothelial cells may provide a potential therapeutic strategy. Dauricine, an anti-inflammatory drug that has been proven effective in treating the aforementioned inflammatory diseases, might be an effective inhibitor of endothelial inflammation. In this work, we intended to discover the function and potential mechanism of dauricine in endothelial inflammation.

## MATERIALS AND METHODS

### Cell Culture

Human umbilical vein endothelial cells (HUVECs) were cultured in specific endothelial medium (ScienCell, Carlsbad, CA,


United States) containing 1% endothelial cell growth supplement, 5% fetal bovine serum, and 1% penicillin-streptomycin solution in a cell incubator (37°C with 5% CO<sub>2</sub>). *In vitro* experiments were performed on HUVECs between passages 2 and 6. The human leukemia monocytic cell line (THP-1) and human embryonic kidney-293T (293T) cell line were obtained from American Type Culture Collection (ATCC, Manassas, VA, United States).

### Reagents

LPS was purchased from Sigma (St. Louis, MO, United States). Dauricine was obtained from Shanghai Yuanye Bio-Technology Co., Ltd (Shanghai, China) and diluted in dimethyl sulfoxide (DMSO). Four concentrations (5, 10, 20, and 40 µM) were finally reached in further experiment (Yang et al., 2010). The solvent DMSO was used as vehicle control, and its final concentration in the cell culture medium was controlled to be less than 0.1%. Ultrafiltered water was used to dissolve the interleukin-1β (final concentration, 10 ng/µl), and then it was filtered again by a 0.22-µm filter (Zhao and Liang, 2019). Acetoxymethyl ester (BCECF-AM) was purchased from the Beyotime Institute of Biotechnology (Jiangsu, China).

### Monocyte–Endothelial Cell Adhesion Assay

HUVECs were seeded into 6-well plates and grown until almost 80% confluence. Then, the cells were incubated with dauricine at different concentrations for 8 h before 4-h stimulation with IL-1β (10 ng/ml). Before use, THP-1 cells were harvested and resuspended in 1640 medium after being labeled with BCECF-AM for approximately 30 min. The collected THP-1 cells were then added to each well containing HUVECs. After being incubated for 30 min, the suspended THP-1 cells were removed with phosphate-buffered saline (PBS). The fluorescence signals of each well were photographed and then analyzed.



**FIGURE 2 |** Dauricine inhibited monocyte–endothelial cell interaction. **(A)** Representative images of THP-1 cells (green) that adhered to HUVECs (scale bar: 100  $\mu$ m). **(B)** Quantitative analysis of THP-1 cells that adhered to HUVECs. Data are expressed as mean  $\pm$  SD. ### $p$  < 0.001 versus the vehicle group; \*\*\* $p$  < 0.001 versus the vehicle + IL-1 $\beta$  group. DMSO, dimethyl sulfoxide; IL-1 $\beta$ , interleukin-1 $\beta$ ; HUVECs, human umbilical vein endothelial cells.


## Real-Time Polymerase Chain Reaction Assay

Primary HUVECs were pretreated with dauricine (5, 10, 20, and 40  $\mu$ M) for 8 h. IL-1 $\beta$  was then administered for 4 h to induce endothelial inflammation. After the treatment, HUVECs were harvested for RNA extraction using a TRIzol reagent (TaKaRa, Dalian, China; 9108). Total RAN was then transcribed into cDNA by a PCR kit (TaKaRa) according to the manufacturer's protocol. Real-time PCR (RT-PCR) was conducted using commercial kits (TaKaRa, RR820Q) as instructed by the manufacturer. Relative mRNA expression levels were normalized and evaluated by the

$2^{-\Delta\Delta CT}$  method. The primers used for target genes are all listed in **Supplementary Table S1**.

## Western Blotting

Western blotting assays were carried out as described previously (Du et al., 2017). Primary antibodies in the experiments were applied as follows: ICAM-1 (diluted at 1:1,000, ProteinTech, Chicago, IL, United States), VCAM-1 (diluted at 1:1,000, ProteinTech), E-selectin (diluted at 1:1,000, R&D Systems, Minneapolis, MN, United States),  $\beta$ -tubulin (diluted at 1:1,000, ProteinTech), NF- $\kappa$ B-p65 (diluted at 1:1,000, CST, Danvers, MA,



United States), NF- $\kappa$ B-pp65 (diluted at 1:1,000, CST), I $\kappa$ B $\alpha$  (diluted at 1:1,000, CST), and phospho-I $\kappa$ B $\alpha$  (diluted at 1:1,000, CST). Then the secondary antibodies with horseradish peroxidase coupling (1:10,000 dilution) obtained from ProteinTech were used for 2 h at room temperature. Electrochemiluminescence (ECL) detection reagents were purchased from Millipore (Billerica, MA, United States). Target proteins in the membrane were visualized with a Bio-Rad (Hercules, CA, United States) exposure imaging system.

### Nuclear and Cytoplasmic Protein Extraction

HUVECs were grown to 90% confluence in a 60-mm cell culture dish. Then vehicle (DMSO) or dauricine (40  $\mu$ M) was given to the cells for 8 h. After that, IL-1 $\beta$  (10 ng/ml) was administered for another 4 h. Subsequently, cells were harvested for nuclear and cytosolic proteins extraction by using an extraction kit (Beyotime, P0027) according to the manufacturer's instructions.

### Immunofluorescence

HUVECs were given a stimulation by IL-1 $\beta$  (10 ng/ml) for 30 min after the pretreatment with dauricine (5, 10, 20, and 40  $\mu$ M) for 8 h. Cells were then harvested for immunofluorescence assay as


previously described (Zhong et al., 2020). A specific antibody targeting p65 and a secondary antibody (Alexa Fluor<sup>®</sup>488, 1:800) were obtained from CST. Images were taken with a fluorescence microscope (Olympus, Tokyo, Japan). Immunofluorescence staining of paraffin-embedded lung sections was performed using a similar procedure after dewaxing and antigen recovery.

### Luciferase Reporter Assay

A p65-overexpressing vector and luciferase reporter gene vector containing the promoters for ICAM-1, VCAM-1, and E-selectin were constructed as described before (Zhao and Liang, 2019). Then, the vectors were introduced into 293T cells when the cell density reached 70%–80% confluence. After incubation for 24 h, the cells were administered with dauricine at a concentration of 40  $\mu$ M for 8 h, and then cells were harvested. The Dual-Glo luciferase Assay System (Promega, Madison, WI, United States) was used to detect the activity of luciferase.

### Chromatin Immunoprecipitation Assay

Chromatin immunoprecipitation (ChIP) assay was carried out by a ChIP assay kit (Millipore). Rabbit IgG was used as the negative control. The fractured DNA samples were pulled down by an



anti-NF- $\kappa$ B p65 antibody (CST). Subsequently, the enriched DNA fragments were detected by PCR using specific primers as shown in **Supplementary Table S2** (Rao et al., 2019).

## Animal Experiment

All animal experiments were approved by the Animal Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology. Male C57BL/6 mice (8–10 weeks) were obtained from Wuhan Beiente Biotechnology Company (Hubei, China). DMSO or dauricine was administered orally with a straight needle (12-gauge, 55 mm) once per day at the same time in seven successive days. LPS (15 mg/kg, Sigma) was then intraperitoneally injected to induce acute lung inflammation. The mice were euthanized to collect lung tissues 24 h after the administration of LPS.

## Statistical Analysis

All experiments were performed at least in triplicate. GraphPad Prism software was used for data analysis and output. ImageJ was used for the quantification of images from Western blotting assays. Quantitative data were presented as the mean  $\pm$  standard error of mean (SEM). Student's *t*-tests were

performed to assess the statistical significance between two groups.  $p < 0.05$  was considered to be statistically significant.

## RESULTS


### Dauricine Was Nontoxic to Human Umbilical Vein Endothelial Cells

The chemical structural formula of dauricine is shown in **Figure 1A**. To investigate the effect of dauricine on endothelial cells, we first conducted the MTT assay to explore its cytotoxicity on HUVECs. As shown in **Figure 1B**, dauricine was safe to use for HUVECs at doses ranging from 5 to 40  $\mu$ M.

### Dauricine Inhibited the Interactions Between Monocyte–Endothelial Cells

To explore the role of dauricine on IL-1 $\beta$ -induced endothelial inflammation, first, we assessed its function on monocyte–endothelial cell adhesion. An increased amount of THP-1 cells was attracted to HUVECs stimulated by IL-1 $\beta$  (10 ng/ml) in comparison with the nontreated HUVECs.





However, when HUVECs were pretreated with dauricine, they attracted markedly fewer THP-1 cells (Figures 2A,B).

## Dauricine Decreased the Expression of Cell Adhesion Molecules

Adhesion molecules are vital in the recruitment of inflammatory cells to endothelium. To further elucidate the role of dauricine on monocyte–endothelial cell interactions, we detected the expression levels of typical cell adhesion molecules in endothelial cells. The mRNA levels of ICAM-1, VCAM-1, and E-selectin were significantly upregulated in IL-1 $\beta$ -treated HUVECs (Figure 3A). However, treatment with dauricine suppressed the overexpression of these molecules (Figure 3A). Western blotting assays confirmed the decreased expression of ICAM-1, VCAM-1, and E-selectin in cells pretreated with dauricine (Figures 3B,C).


## Dauricine Inhibited IL-1 $\beta$ Induced Activation of NF- $\kappa$ B

The NF- $\kappa$ B pathway has been proven to regulate the expression of cell adhesion molecules in endothelial cells (Rao et al., 2019). To

determine the mechanism how dauricine regulated the expression of ICAM-1, VCAM-1, and E-selectin, we detected the activation of NF- $\kappa$ B pathway in HUVECs stimulated with IL-1 $\beta$ . The degradation of I $\kappa$ B $\alpha$  and the activation of p65 were significantly inhibited by dauricine (Figures 4A,B). To further clarify the role of dauricine on p65 translocation, we detected the distribution of p65 protein by immunostaining. Correspondingly, we found that dauricine effectively inhibited p65 translocation in IL-1 $\beta$ -treated HUVECs (Figure 4C). Next, nuclear and cytoplasmic proteins were separately extracted to detect the level of p65 in both fractions. As shown in Figure 4D, the level of cytoplasmic p65 was decreased, and endonuclear p65 was significantly increased in IL-1 $\beta$ -treated HUVECs, while incubation with 40  $\mu$ M of dauricine significantly inhibited the IL-1 $\beta$ -induced nuclear localization of p65.


## Dauricine Suppressed the Expression of ICAM-1, VCAM-1, and E-Selectin Through the NF- $\kappa$ B Pathway

Reporter gene constructs targeting ICAM-1, VCAM-1, and E-selectin were constructed for use in a dual-luciferase reporter assay. As shown in Figure 5A, the luciferase activities



of E-selectin, ICAM-1, and VCAM-1 were obviously enhanced in 293T cells cotransfected with a p65 overexpression vector, while the administration of dauricine remarkably reduced their

transcriptional activity. Furthermore, ChIP experiments were performed to determine the effect of dauricine on the DNA-binding activity of p65. Dauricine decreased the binding of p65 to



its target gene promoters, including the promoters of VCAM-1, ICAM-1, and E-selectin, in IL-1 $\beta$ -treated endothelial cells (Figure 5B).

## Dauricine Attenuated Lipopolysaccharide-Induced Acute Lung Injury

Based on previous studies that demonstrated the anti-inflammatory role of dauricine in IL-1 $\beta$ -treated endothelial cells, we established an acute lung injury model induced by LPS to further verify whether dauricine attenuated endothelial inflammation *in vivo*. Previously, we reported that dauricine alleviated LPS-induced macrophage infiltration in lung tissues and decreased serum levels of IL-1 $\beta$ , IL-6, and TNF- $\alpha$  (Qiao et al., 2019). In this work, we found that pretreatment with dauricine (20 mg/kg) effectively protected mice from LPS-induced acute inflammation (Figure 6A). Moreover, the expression of ICAM-1, VCAM-1, and E-selectin in lung tissues was decreased by dauricine pretreatment in LPS-treated mice (Figures 6B,C). Additionally, the phosphorylation of I $\kappa$ B $\alpha$  and p65 was also inhibited (Figures 6B,C). We performed immunofluorescence assay on lung tissues to determine the levels of ICAM-1, VCAM-1, and E-selectin expressed in endothelial cells. The expression of ICAM-1 was increased in the endothelial cells of lung tissues from LPS-treated mice, while it was attenuated in dauricine-treated mice (Figures 6D,F). Similar results were observed in the level of VCAM-1 and E-selectin (Supplementary figure S1A,B,C). Moreover, we found that p65 translocation in lung endothelial cells was significantly reduced by dauricine pretreatment

compared with the LPS-treated group (Figures 6E,G). We concluded that dauricine could alleviate the inflammatory response in lung endothelial cells, which corresponded with our *in vitro* findings (Figure 7).

## DISCUSSION

The relationship between endothelial inflammation and various diseases, including metabolic syndrome, chronic kidney disease, rheumatoid arthritis, and sepsis, has been debated for a long time (Yang et al., 2016; Grandl and Wolfrum, 2018; Diaz-Ricart et al., 2020; Lupu et al., 2020). During vascular inflammation, the expression of adhesion molecules, including ICAM-1, VCAM-1, and E-selectin, is elevated, promoting leukocyte recruitment to the endothelium (Zhao and Liang, 2019). Excessive leukocyte recruitment to the endothelium aggravates local inflammation and contributes to endothelial inflammation-related diseases such as atherosclerosis and dysfunctional microcirculation (Hernandez et al., 2013; Schmitt et al., 2014). Molecules and drugs targeting endothelial inflammation may be potential candidates for various similar diseases.

Traditional Chinese medicine has been used in inflammatory diseases for a long time (Ren et al., 2017; Chiang et al., 2020). And in recent years, many extracts from traditional herbs were proved to regulate several signaling pathways involved in inflammation (Spelman et al., 2006; Huang et al., 2016; Zhang et al., 2019). Dauricine, an alkaloid isolated from *M. dauricum*, has been found to protect against several inflammation-related diseases. Previous studies have reported that dauricine inhibited NF- $\kappa$ B activation



in macrophages and colon cancer cells (Yang et al., 2010; Qiao et al., 2019). In a cerebral ischemia/reperfusion rat model treated with dauricine, ICAM-1 was proven to be significantly reduced. In this work, we demonstrated that dauricine repressed IL-1 $\beta$ -induced NF- $\kappa$ B activation and reduced the expression of its downstream genes in endothelial cells. Therefore, dauricine could be considered for use in diseases related with endothelial inflammation.

The NF- $\kappa$ B pathway is vital to endothelial inflammation (Oates, 2015). Usually, the activation of cytoplasmic NF- $\kappa$ B is regulated by inhibitors of  $\kappa$ B (I $\kappa$ B). Extracellular proinflammatory stimulus promotes the phosphorylation and degradation of I $\kappa$ B; subsequently, the active unit, P65, is translocated to the nucleus, where it regulates the expression of downstream genes (Lee et al., 2009). However, the mechanism by which dauricine regulates the activation of the NF- $\kappa$ B pathway remains unknown. Further study is needed to determine whether dauricine regulates NF- $\kappa$ B directly by binding with its subunit or through other pathways.

Endothelial cells play pivotal roles in LPS-induced inflammation (Flavahan, 1992). We tested the anti-inflammatory function of dauricine in a mouse model of acute lung injury induced by LPS. We showed that mice pretreated with dauricine exhibited dramatically decreased expression of ICAM-1, VCAM-1, and E-selectin in lung vascular beds. Activation of the NF- $\kappa$ B pathway was also downregulated in lung tissue after pretreatment with dauricine. These findings suggested that dauricine reduced acute lung inflammation by inhibiting the NF- $\kappa$ B pathway. However, whether dauricine can prevent endothelial inflammation in other vascular-rich organs, such as the kidney, or in other inflammatory diseases, such as atherosclerosis, remains to be explored.

In addition, we found that MAPK pathways (the p38, ERK1/2, and JNK pathways) were activated in IL-1 $\beta$ -treated HUVECs and that dauricine could inhibit the phosphorylation of P38, JNK, and ERK (data not shown). MAPK pathways are ubiquitous and have been reported to regulate the expression of adhesion molecules, including ICAM-1, VCAM-1, and E-selectin in inflammatory diseases (Min and Pober, 1997; Pietersma et al., 1997; Yan et al., 2002). Therefore, dauricine might also exert its anti-inflammatory effect by inhibiting the MAPK pathway.

However, whether dauricine could alleviate the activation of the MAPK pathway *in vivo* and to what extent the MAPK and NF- $\kappa$ B pathway take part individually in the anti-inflammatory effect of dauricine remain unknown.

Collectively, the findings from our present study suggest that dauricine suppresses endothelial inflammation through or partially through inhibition of the NF- $\kappa$ B pathway. We propose that dauricine might serve as a potential therapeutic candidate for various diseases related to endothelial inflammation.

## DATA AVAILABILITY STATEMENT

The original contributions presented in the study are included in the article/**Supplementary Material**. Further inquiries can be directed to the corresponding author.

## ETHICS STATEMENT

The animal study was reviewed and approved by Ethics Committee of Tongji Medical College, Huazhong University of Science and Technology.

## AUTHOR CONTRIBUTIONS

KH contributed to the conception of the study. JH and JA performed the experiment. JH and RC contributed significantly to the analysis and manuscript preparation. JH and RC performed the data analyses and wrote the manuscript. YW and ML helped perform the analysis with constructive discussions. All authors gave final approval.

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fphar.2021.758962/full#supplementary-material>

## REFERENCES

- Boulanger, C. M. (2016). Endothelium. *Arterioscler Thromb. Vasc. Biol.* 36 (4), e26–31. doi:10.1161/atvbaha.116.306940
- Brasier, A. R. (2010). The Nuclear Factor- $\kappa$ B-Interleukin-6 Signaling Pathway Mediating Vascular Inflammation. *Cardiovasc. Res.* 86 (2), 211–218. doi:10.1093/cvr/cvq076
- Chiang, C. C., Cheng, W. J., Lin, C. Y., Lai, K. H., Ju, S. C., Lee, C., et al. (2020). Kan-Lu-Hsiao-Tu-Tan, a Traditional Chinese Medicine Formula, Inhibits Human Neutrophil Activation and Ameliorates Imiquimod-Induced Psoriasis-like Skin Inflammation. *J. Ethnopharmacol.* 246, 112246. doi:10.1016/j.jep.2019.112246
- Diaz-Ricart, M., Torramade-Moix, S., Pascual, G., Palomo, M., Moreno-Castaño, A. B., Martínez-Sánchez, J., et al. (2020). Endothelial Damage, Inflammation and Immunity in Chronic Kidney Disease. *Toxins (Basel)* 12 (6). doi:10.3390/toxins12060361
- Du, M., Yuan, L., Tan, X., Huang, D., Wang, X., Zheng, Z., et al. (2017). The LPS-Inducible lncRNA Mirt2 Is a Negative Regulator of Inflammation. *Nat. Commun.* 8 (1), 2049. doi:10.1038/s41467-017-02229-1
- Du, Z. H., Liu, H. G., Chai, C. Y., Luo, L. Y., and Hu, C. J. (1986). Anti-inflammatory Effect of Dauricine. *Zhongguo Yao Li Xue Bao* 7 (5), 419–422.
- Flavahan, N. A. (1992). Atherosclerosis or Lipoprotein-Induced Endothelial Dysfunction. Potential Mechanisms Underlying Reduction in EDRF/nitric Oxide Activity. *Circulation* 85 (5), 1927–1938. doi:10.1161/01.cir.85.5.1927
- Gimbrone, M. A., Jr., and García-Cardeña, G. (2016). Endothelial Cell Dysfunction and the Pathobiology of Atherosclerosis. *Circ. Res.* 118 (4), 620–636. doi:10.1161/circresaha.115.306301
- Grandl, G., and Wolfrum, C. (2018). Hemostasis, Endothelial Stress, Inflammation, and the Metabolic Syndrome. *Semin. Immunopathol.* 40 (2), 215–224. doi:10.1007/s00281-017-0666-5
- Hernandez, G., Bruhn, A., and Ince, C. (2013). Microcirculation in Sepsis: New Perspectives. *Curr. Vasc. Pharmacol.* 11 (2), 161–169. doi:10.2174/1570161111311020006

- Huang, Y., Cai, T., Xia, X., Cai, Y., and Wu, X. Y. (2016). Research Advances in the Intervention of Inflammation and Cancer by Active Ingredients of Traditional Chinese Medicine. *J. Pharm. Pharm. Sci.* 19 (1), 114–126. doi:10.18433/j3sg7k
- Joffre, J., Hellman, J., Ince, C., and Ait-Oufella, H. (2020). Endothelial Responses in Sepsis. *Am. J. Respir. Crit. Care Med.* 202 (3), 361–370. doi:10.1164/rccm.201910-1911TR
- Jourde-Chiche, N., Fakhouri, F., Dou, L., Bellien, J., Burtey, S., Frimat, M., et al. (2019). Endothelium Structure and Function in Kidney Health and Disease. *Nat. Rev. Nephrol.* 15 (2), 87–108. doi:10.1038/s41581-018-0098-z
- Konukoglu, D., and Uzun, H. (2017). Endothelial Dysfunction and Hypertension. *Adv. Exp. Med. Biol.* 956, 511–540. doi:10.1007/5584\_2016\_90
- Lee, Y. M., Seon, M. R., Cho, H. J., Kim, J. S., and Park, J. H. (2009). Benzyl Isothiocyanate Exhibits Anti-inflammatory Effects in Murine Macrophages and in Mouse Skin. *J. Mol. Med. (Berl)* 87 (12), 1251–1261. doi:10.1007/s00109-009-0532-6
- Li, H., Chen, X., and Zhou, S. J. (2018). Dauricine Combined with Clindamycin Inhibits Severe Pneumonia Co-infected by Influenza Virus H5N1 and Streptococcus Pneumoniae In vitro and In vivo through NF-Kb Signaling Pathway. *J. Pharmacol. Sci.* 137 (1), 12–19. doi:10.1016/j.jphs.2018.01.011
- Lupu, F., Kinasewitz, G., and Dormer, K. (2020). The Role of Endothelial Shear Stress on Haemodynamics, Inflammation, Coagulation and Glycocalyx during Sepsis. *J. Cel Mol Med* 24 (21), 12258–12271. doi:10.1111/jcmm.15895
- Min, W., and Pober, J. S. (1997). TNF Initiates E-Selectin Transcription in Human Endothelial Cells through Parallel TRAF-NF-Kappa B and TRAF-RAC/CDC42-JNK-c-Jun/ATF2 Pathways. *J. Immunol.* 159 (7), 3508–3518.
- Oates, J. C. (2015). Endothelial Dysfunction in Injury and Inflammation. *Am. J. Med. Sci.* 349 (1), 2. doi:10.1097/maj.0000000000000401
- Park, H. J., Gholam Zadeh, M., Suh, J. H., and Choi, H. S. (2020). Dauricine Protects from LPS-Induced Bone Loss via the ROS/PP2A/NF-κB Axis in Osteoclasts. *Antioxidants (Basel)* 9 (7), 588. doi:10.3390/antiox9070588
- Pietersma, A., Tilly, B. C., Gaestel, M., de Jong, N., Lee, J. C., Koster, J. F., et al. (1997). p38 Mitogen Activated Protein Kinase Regulates Endothelial VCAM-1 Expression at the post-transcriptional Level. *Biochem. Biophys. Res. Commun.* 230 (1), 44–48. doi:10.1006/bbrc.1996.5886
- Qiao, B., Wang, H., Wang, C., Liang, M., Huang, K., and Li, Y. (2019). Dauricine Negatively Regulates Lipopolysaccharide- or Cecal Ligation and Puncture-Induced Inflammatory Response via NF-Kb Inactivation. *Arch. Biochem. Biophys.* 666, 99–106. doi:10.1016/j.abb.2019.03.018
- Rao, C., Liu, B., Huang, D., Chen, R., Huang, K., Li, F., et al. (2021). Nucleophosmin Contributes to Vascular Inflammation and Endothelial Dysfunction in Atherosclerosis Progression. *J. Thorac. Cardiovasc. Surg.* 161, e377–e393. doi:10.1016/j.jtcvs.2019.10.152
- Ren, Y., Qiao, W., Fu, D., Han, Z., Liu, W., Ye, W., et al. (2017). Traditional Chinese Medicine Protects against Cytokine Production as the Potential Immunosuppressive Agents in Atherosclerosis. *J. Immunol. Res.* 2017, 7424307. doi:10.1155/2017/7424307
- Schmitt, M. M., Megens, R. T., Zerneck, A., Bidzhekov, K., van den Akker, N. M., Rademakers, T., et al. (2014). Endothelial Junctional Adhesion Molecule-A Guides Monocytes into Flow-dependent Predilection Sites of Atherosclerosis. *Circulation* 129 (1), 66–76. doi:10.1161/circulationaha.113.004149
- Spelman, K., Burns, J., Nichols, D., Winters, N., Ottersberg, S., and Tenborg, M. (2006). Modulation of Cytokine Expression by Traditional Medicines: a Review of Herbal Immunomodulators. *Altern. Med. Rev.* 11 (2), 128–150.
- Sturtzel, C. (2017). Endothelial Cells. *Adv. Exp. Med. Biol.* 1003, 71–91. doi:10.1007/978-3-319-57613-8\_4
- Yan, W., Zhao, K., Jiang, Y., Huang, Q., Wang, J., Kan, W., et al. (2002). Role of P38 MAPK in ICAM-1 Expression of Vascular Endothelial Cells Induced by Lipopolysaccharide. *Shock* 17 (5), 433–438. doi:10.1097/00024382-200205000-00016
- Yang, X., Chang, Y., and Wei, W. (2016). Endothelial Dysfunction and Inflammation: Immunity in Rheumatoid Arthritis. *Mediators Inflamm.* 2016, 6813016. doi:10.1155/2016/6813016
- Yang, X. Y., Jiang, S. Q., Zhang, L., Liu, Q. N., and Gong, P. L. (2007). Inhibitory Effect of Dauricine on Inflammatory Process Following Focal Cerebral Ischemia/reperfusion in Rats. *Am. J. Chin. Med.* 35 (3), 477–486. doi:10.1142/s0192415x07004990
- Yang, Z., Li, C., Wang, X., Zhai, C., Yi, Z., Wang, L., et al. (2010). Dauricine Induces Apoptosis, Inhibits Proliferation and Invasion through Inhibiting NF-kappaB Signaling Pathway in colon Cancer Cells. *J. Cel Physiol* 225 (1), 266–275. doi:10.1002/jcp.22261
- Zhang, J., Zhang, Q., Liu, G., and Zhang, N. (2019). Therapeutic Potentials and Mechanisms of the Chinese Traditional Medicine Danshensu. *Eur. J. Pharmacol.* 864, 172710. doi:10.1016/j.ejphar.2019.172710
- Zhang, S., Ren, Y., and Qiu, J. (2018). Dauricine Inhibits Viability and Induces Cell Cycle Arrest and Apoptosis via Inhibiting the PI3K/Akt Signaling Pathway in Renal Cell Carcinoma Cells. *Mol. Med. Rep.* 17 (5), 7403–7408. doi:10.3892/mmr.2018.8732
- Zhao, S., Liang, M., Wang, Y., Hu, J., Zhong, Y., Li, J., et al. (2019). Chrysin Suppresses Vascular Endothelial Inflammation via Inhibiting the NF-Kb Signaling Pathway. *J. Cardiovasc. Pharmacol. Ther.* 24 (3), 278–287. doi:10.1177/1074248418810809
- Zhong, Y., He, K., Huang, M., and Liang, M. (2020). Neferine Suppresses Vascular Endothelial Inflammation by Inhibiting the NF-Kb Signaling Pathway. *Arch. Biochem. Biophys.* 696, 108595. doi:10.1016/j.abb.2020.108595

**Conflict of Interest:** The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

**Publisher's Note:** All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

Copyright © 2021 Hu, Chen, An, Wang, Liang and Huang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.