



OPEN ACCESS

RESEARCH PAPER

# Systematic review and meta-analysis on outcome differences among patients with TICI2b versus TICI3 reperfusion: success revisited

Johannes Kaesmacher,<sup>1</sup> Tomas Dobrocky,<sup>1</sup> Mirjam R Heldner,<sup>2</sup> Sebastian Bellwald,<sup>2</sup> Pascal J Mosimann,<sup>1</sup> Pasquale Mordasini,<sup>1</sup> Sandra Bigi,<sup>3</sup> Marcel Arnold,<sup>2</sup> Jan Gralla,<sup>1</sup> Urs Fischer<sup>2</sup>

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jnnp-2017-317602>).

<sup>1</sup>University Institute of Diagnostic and Interventional Neuroradiology, University of Bern, Inselspital, Bern, Switzerland

<sup>2</sup>Department of Neurology, University of Bern, Inselspital, Bern, Switzerland

<sup>3</sup>Division of Child Neurology, Department of Pediatrics, University of Bern, Inselspital, Bern, Switzerland

## Correspondence to

Dr Urs Fischer, Department of Neurology, University of Bern, Inselspital, Bern CH-3010, Switzerland; [urs.fischer@insel.ch](mailto:urs.fischer@insel.ch)

JG and UF contributed equally.

Received 7 November 2017

Revised 1 February 2018

Accepted 7 February 2018

Published Online First

8 March 2018

## ABSTRACT

**Objective** A reperfusion quality of thrombolysis in cerebral infarction (TICI)  $\geq 2b$  has been set as the therapeutic angiography target for interventions in patients with acute ischaemic stroke. This study addresses whether the distinction between TICI2b and TICI3 reperfusion shows a clinically relevant difference on functional outcome.

**Methods** A systematic literature review and meta-analysis was carried out and presented in conformity with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses criteria to test the primary hypothesis that TICI2b and TICI3 reperfusion are associated with different rates of modified Rankin Scale (mRS)  $\leq 2$  at day 90. Secondary endpoints included rates of haemorrhagic transformations, mortality and excellent functional outcome (mRS  $\leq 1$ ). Summary estimates of ORs (sOR) with 95% CI were calculated using the inverse variance heterogeneity model accounting for multiple true effect sizes.

**Results** Fourteen studies on 2379 successfully reperfused patients were included (1131 TICI3, 1248 TICI2b). TICI3 reperfusion was associated with higher rates of functional independence (1.74, 95% CI 1.44 to 2.10) and excellent functional outcomes (2.01, 95% CI 1.60 to 2.53), also after including adjusted estimates. The safety profile of patients with TICI3 was superior, as demonstrated by lower rates of mortality (sOR 0.59, 95% CI 0.37 to 0.92) and symptomatic intracranial haemorrhages (sOR 0.42, 95% CI 0.25 to 0.71).

**Conclusion** TICI3 reperfusion is associated with superior outcome and better safety profiles than TICI2b reperfusion. This effect seems to be independent of time and collaterals. As reperfusion quality is the most important modifiable predictor of patients' outcome, a more conservative definition of successful therapy and further evaluation of treatment approaches geared towards achieving TICI3 reperfusion are desirable.

## INTRODUCTION

Quality of reperfusion is one of the most important and potentially modifiable determinants of clinical outcome in patients treated with thrombolysis following acute ischaemic stroke.<sup>1</sup> It is commonly evaluated by applying the five-step thrombolysis in cerebral infarction (TICI) grading scale.<sup>2</sup> Grades 2b and 3 are routinely termed 'successful reperfusion' as this was shown to be the most favourable

cut-off for predicting good outcome at 90 days with non-significant differences between grades 2b and 3.<sup>3,4</sup> Consequently, the target angiographic endpoint has been set to TICI  $\geq 2b$ .<sup>5</sup> However, differences in outcome between patients with TICI2b and TICI3 reperfusion have mostly been neglected, as they are often subtle or may have simply been overlooked, because both grades have been routinely subsumed under the term 'successful'. Recently, some studies have suggested superior outcomes for TICI3 versus TICI2b reperfusion, thus putting into question whether the definition of success should be revised.<sup>6-8</sup> Due to the relatively small number of patients included, it currently remains unclear whether the distinction between TICI2b and TICI3 is clinically relevant. The primary objective of this analysis was therefore to identify and quantify all available observational data on clinical outcomes between TICI2b and TICI3 reperfusion. Furthermore, we aimed to review and discuss recent refinements and modification of the TICI score.

## METHODS


The meta-analysis conducted adheres to the reporting guidelines laid down by the Preferred Reporting Items for Systematic Reviews and Meta-Analyses report (PRISMA statement)<sup>9</sup> and Meta-analysis of Observational Studies in Epidemiology checklist.<sup>10</sup> Several versions of the TICI scale exist and are referred to as original, modified or extended TICI scale (abbreviated as oTICI, mTICI and eTICI, respectively, see online supplementary table I). To avoid confusion, oTICI grade 2b is defined as antegrade reperfusion of at least two-thirds of the target territory.<sup>2</sup> mTICI grade 2b refers to antegrade reperfusion of at least half of the target territory.<sup>5</sup> Grade 2b in mTICI and eTICI is the same, but an additional TICI2c grade has been employed in eTICI referring to 'near complete perfusion except for slow flow in a few distal cortical vessels, or presence of small distal cortical emboli'.<sup>11,12</sup> Lastly, Liebeskind *et al* recently suggested the oTICI2c scale which subdivides the grade 2b into 2b with 50%–66% reperfusion and 2b with 67%–90% reperfusion. Here, reperfusion of 90%–99% is referred to as grade 2c<sup>13</sup> (see online supplementary table I).

## Literature search and data extraction

PubMed and Web of Science databases (from inception to 18 October 2017) were accessed using a

Check for updates

**To cite:** Kaesmacher J, Dobrocky T, Heldner MR, *et al*. *J Neurol Neurosurg Psychiatry* 2018;**89**:910–917.



**Figure 1** Flow chart according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) recommendations. mRS, modified Rankin Scale; TICI, thrombolysis in cerebral infarction.

predefined search strategy formulated according to the Population, Intervention, Comparison and Outcome (PICO) format (see online supplementary tables II and III).<sup>14</sup> Full-text articles and conference abstracts were reviewed. No language restriction was applied. Eligibility of the studies for the quantitative analysis was rated by two independent readers (JK and TD). Studies were included into the quantitative synopsis if the study reported on (1) the primary outcome stratified according to TICI2b versus TICI3 reperfusion or (2) the primary outcome stratified according to TICI2b versus TICI2c/3 reperfusion. Manual searching of reference lists of the included studies was coupled with a search of all articles citing the included articles using Google Scholar metadata (<https://scholar.google.ch/>). If available, TICI2c was pooled together with TICI3, as preliminary evidence has suggested that both are associated with a comparable clinical course.<sup>15–17</sup> In a sensitivity analysis, comparisons of TICI2c/3 and TICI2b were excluded leading to a comparison confined to mTICI2b versus mTICI3 in order to rule out the possibility that potential differences are only discernible when applying the more detailed eTICI scale. Studies that did not meet the eligibility criteria but appraised the topic of outcome differences between TICI2b and TICI3 reperfusion were included into further semi-quantitative or qualitative analyses.

Primary outcome was the rate of patients achieving functional independence at day 90, defined as modified Rankin Scale (mRS)  $\leq 2$ . Predefined secondary outcomes were excellent functional outcome (mRS  $\leq 1$ ), mRS shift, all-cause mortality (during hospitalisation or day 90, depending on the reporting standards), final infarct volume, rates of symptomatic intracranial haemorrhage (sICH) and rates of any intracranial haemorrhage (any ICH). If a study did not report on a clinical definition of sICH, rates of parenchymal haematomas (PH1/2 or PH2 only, depending on the way of reporting) were evaluated as

radiological surrogate, according to the European Acute Stroke Study definition.<sup>18</sup>

If available, the following parameters were extracted and calculated: type of study, unadjusted odds for the primary and secondary outcomes, results from adjusted analyses with a description of parameters adjusted for, type of TICI scale applied and presence of differences in baseline characteristics. Two independent raters extracted the data (JK and MRH). All extracted raw frequency counts can be found in the online supplementary dataset 1.

### Statistical analysis

The inverse variance heterogeneity model was used to calculate summary estimates of effect sizes (summarised ORs, sOR),<sup>19</sup> since included studies used different inclusion and exclusion criteria and require to account for multiple true effect sizes. To calculate unadjusted ORs, the prevalence of different endpoints was extracted from the published data for each arm. Summarised point estimates are displayed together with 95% CIs to express the odds for a comparison between TICI3 and TICI2b. Adjusted ORs were summarised separately, if available. Heterogeneity was explored using Cochrane's Q and I<sup>2</sup>.<sup>20</sup> Visual inspection of funnel plots and Doi plots and calculation of the Luis Furuya-Kanamori (LFK) index were used for the evaluation of publication bias regarding the primary endpoint.<sup>21</sup> Data analysis was performed using the software package MetaXL (EpiGear International, Sunrise Beach, Queensland, Australia) for Microsoft Excel.


### Risk of bias and quality assessment

The risk of bias was evaluated as per Cochrane Collaboration tool.<sup>22</sup> Additionally, the following quality criteria were specifically evaluated: (1) specification of inclusion criteria; (2)

**Table 1** Characteristics of studies included into quantitative analysis

Study	Study type	Multicentre	n				TICI grading provided by	Definition of mortality provided	Differences in baseline characteristics	Definition of sICH	Sites of occlusion included	Adjusted analysis available	
			TICI	TICI2b	TICI2c	TICI3							
Dargazani <i>et al</i> <sup>7</sup>	RO	Yes	mTICI	109	–	113	One neurologist; blinded to the interventionalists' gradings and clinical outcomes	3-month mortality	Better collaterals, less cardioembolic aetiology, shorter onset to reperfusion metrics, lower number of passes in TICI3 patients	Presence of ICH with NIHSS worsening of $\geq 4$ within 24 hours or prompting death	ICA, M1, M2	Yes	Centre, age, diabetes, admission NIHSS score, prior use of intravenous thrombolysis, site of occlusion, favourable collateral flow, aetiology and onset to reperfusion time
Almekhlafi <i>et al</i> <sup>17</sup>	RO	No	mTICI and eTICI	29	32	29	Two neurointerventionalists (blinding not specified, one graded eTICI, one graded mTICI)	3-month mortality	No P values available, however, faster imaging to reperfusion times and higher number treated with intravenous tPA in TICI2/3 patients	SITS-MOST	Anterior circulation stroke, not further specified	No	NA
Carvalho <i>et al</i> <sup>25</sup>	RO	Yes	mTICI	97	–	53	Consensus reading of neurointerventionalists and stroke neurologists with significant clinical and research experience in intra-arterial treatment	NA	NA	NA	M1	No	NA
Yoo <i>et al</i> <sup>3</sup>	RO	No	mTICI	68	–	110	Operator measured, non-blinded	3-month mortality	Not statistically significant shorter onset to reperfusion metrics in TICI3 patients	ECASS	ICA, M1, M2, tandem	Yes	Age, sex, diabetes, hypertension, left hemisphere stroke, symptom to reperfusion time
Kaesmacher <i>et al</i> <sup>16</sup>	RO	No	eTICI	117	84	45	Two neuroradiologists in consensus; blinded to interventionalists' grading and clinical outcomes	In-hospital mortality	Not statistically significant shorter onset to reperfusion metrics in TICI2/3 patients	PH1/2 (radiological)	M1/M2	Yes	Age, admission NIHSS, onset to reperfusion time, pretreatment alteplase
Chamorro <i>et al</i> <sup>23</sup>	RO	No	mTICI	51	–	74	Experienced radiologists and interventionalists (number not specified), blinded to all clinical data	3-month mortality	Shorter onset to reperfusion metrics, lower number of passes and better collaterals in TICI3 patients	Presence of ICH and NIHSS worsening of $\geq 4$	ICA, M1, M2, tandem	Yes	Age, sex, admission NIHSS, target occlusion location, ASPECTS, pretreatment alteplase and collaterals
Linfante <i>et al</i> <sup>16</sup>	RO	Yes	mTICI	107	–	127	Site-specific grading, not further specified	NA	NA	NA	M1, M2, ICA, basilar (10%)	No	NA
Rangaraju <i>et al</i> <sup>27</sup>	RO	No	mTICI	64	–	35	Operator measured, non-blinded	3-month mortality	None	NA	ICA, M1	Yes	Age, NIHSS and ASPECTS
Schmitz <i>et al</i> <sup>24</sup>	PH-RCT	Yes	mTICI	94	–	6	Core lab adjudicated (IMS II)	NA	NA	NA	ICA, M1, M2	No	NA
Humphries <i>et al</i> <sup>28*</sup>	RO	Yes	Not specified	37	–	52	Operator measured, non-blinded	NA	NA	NA	M1, M2, ICA, basilar (10%)	No	NA
Massari <i>et al</i> <sup>29</sup>	RO	No	mTICI	18	–	23	Operator measured, non-blinded	NA	NA	NA	M1, M2, ICA, basilar/vertebrobasilar junction (~10%)	No	NA
Marks <i>et al</i> <sup>30</sup>	PH-RCT	Yes	mTICI	28	–	18	Core lab adjudicated (DEFUSE 2)	NA	NA	NA	M1, M2, M3, ICA	No	NA
Goyal <i>et al</i> <sup>31</sup>	RO	Yes	Not specified	200	–	216	Not specified	3-month mortality	None	Not specified	NA	Yes	Not specified ('potential confounders')
Liebkeind <i>et al</i> <sup>13</sup>	PH-RCT	Yes	oTICI2c	44	258	44	Core lab adjudicated (HERMES)	3-month mortality	NA	NA	ICA, M1	No	NA

\*There are discrepancies regarding the true rate of TICI3 and TICI2b reperfusion. While table 3 of the respective publication suggests that there are 52 TICI3 cases and 37 TICI2b cases, the text states that there were 46 TICI3 cases. The numbers used for quantitative analysis are derived from table 3 of this publication. ASPECTS, Alberta Stroke Program Early CT Score; DEFUSE 2, Diffusion and Perfusion Imaging Evaluation for Understanding Stroke Evolution Study 2; ECASS, European Cooperative Acute Stroke Study; eTICI, extended TICI scale; HERMES, Highly Effective Reperfusion Evaluated in Multiple Endovascular Stroke Trials; ICA, internal carotid artery; ICH, intracranial haemorrhage; IMS II, Interventional Management of Stroke II; M1/M2/M3, segment one/two/three of the middle cerebral artery; mTICI, modified TICI scale; NA, not applicable; NIHSS, National Institutes of Health Stroke Scale; oTICI, original TICI scale; PH, parenchymal haematoma; PH-RCT, post hoc analysis of randomised controlled trial data; RO, retrospective observational; sICH, symptomatic intracranial haemorrhage; SITS-MOST, Safe Implementation of Thrombolysis in Stroke: Monitoring Study; TICI, thrombolysis in cerebral infarction; tPA, tissue plasminogen activator; total number of patients n=2461, n=2379 with available follow-up for the primary outcome (mRS at day 90).



**Figure 2** Summary OR TIC12c/3 versus TIC12b for d90 modified Rankin Scale (mRS)  $\leq 2$ . \*Used extended TICI (eTICI) scale with TIC12c; TIC12c and TIC13 were subsumed under TIC13; for grading used in Liebeskind *et al* see online supplementary table I. TICI, thrombolysis in cerebral infarction.

comprehensive reporting of baseline characteristics; (3) availability of adjusted analyses; and (4) core lab adjudicated reperfusion grading. Both ratings were performed independently by two readers. In cases of discrepancies a consensus was reached ( $n=6/126$  items).


## RESULTS

### Quantitative analyses

Fourteen studies with a total of 2379 successfully reperfused patients (1131 TIC13, 1248 TIC12b) with available follow-up were included in the quantitative analysis (see figure 1 for PRISMA flow chart).<sup>3 7 13 16 17 23–31</sup> During the eligibility rating process, five discrepancies arose, which could be resolved by a third rater (see online supplementary table IV). Three conference abstracts<sup>13 27 31</sup> and 11 research articles met the inclusion criteria. Eleven of the 14 studies were retrospective observational studies. One study examined different degrees of successful reperfusion in the Highly Effective Reperfusion Evaluated in Multiple Endovascular Stroke Trials collaboration of recent endovascular trials.<sup>13</sup> Eleven studies provided comparisons of mTIC12b


and mTIC13, one study reported a comparison of mTIC12b and mTIC13 together with mTIC12b and mTIC12c/3,<sup>17</sup> one compared mTIC12b with mTIC12c/3,<sup>16</sup> and one study provided comparison of TIC12b and TIC12c/3 using the TICI scale revised by Liebeskind *et al*.<sup>13</sup> The reporting frequencies and respective patient numbers for the primary and secondary endpoints can be found in online supplementary table V. An overview of characteristics of the included studies can be found in table 1 and online supplementary table VI). The most frequently observed differences between patients with TIC12c/3 and TIC12b reperfusion were shorter onset to reperfusion metrics and better collaterals in the TIC12c/3 group (see table 1). No differences regarding the rates of preinterventional intravenous tissue plasminogen activator (tPA) administration between both groups were reported (see online supplementary table VI).

TIC12c/3 reperfusion was more frequently associated with functional independence at day 90 than TIC12b (sOR 1.74, 95% CI 1.44 to 2.10, figure 2), without substantial heterogeneity ( $I^2$  13%,  $Q=15.00$ ,  $P=0.31$ ). This association remained statistically tangible if analysis was confined to studies that used



**Figure 3** Summary OR TIC12c/3 versus TIC12b for d90 modified Rankin Scale (mRS)  $\leq 1$ . \*Used extended TICI (eTICI) scale with TIC12c; TIC12c and TIC13 were subsumed under TIC13; for grading used in Liebeskind *et al* see online supplementary table I. TICI, thrombolysis in cerebral infarction.






**Figure 4** Summary OR TIC12c/3 versus TIC12b for any intracranial haemorrhage (ICH) and symptomatic intracranial haemorrhage (sICH). (A) Any type of ICH. (B) Symptomatic ICH. §Parenchymal haematomas (PH1/2) defined as sICH. \*Used extended TICI (eTICI) scale with TIC12c; TIC12c and TIC13 were subsumed under TIC13. TICI, thrombolysis in cerebral infarction.

the mTICI scale (mTIC13 vs mTIC12b, sOR 1.82, 95% CI 1.41 to 2.34, online supplementary figure I). Importantly, this association also reached statistical significance after summarising the adjusted ORs of the respective studies (adjusted sOR 2.36, 95% CI 1.69 to 3.30, online supplementary figure II). The overall event rate of mRS ≤2 calculated from the studies reviewed was ~55%. Assuming this as a reference level, achieving TIC13 instead of TIC12b in five cases, would result in one additional patient reaching functional independence according to the adjusted estimates. Furthermore, excellent functional outcomes were more commonly observed in patients with TIC12c/3 reperfusion (unadjusted sOR 2.01, 95% CI 1.60 to 2.53, adjusted sOR 2.70, 95% CI 1.71 to 4.25, figure 3 and online supplementary figure III, respectively). This was also the case when analysis was limited to studies using the mTICI scale (mTIC13 vs

mTIC12b, sOR 2.27, 95% CI 1.67 to 3.08, online supplementary figure IV).

TIC12c/3 reperfusion was associated with reduced haemorrhagic transformations (sOR 0.48, 95% CI 0.26 to 0.90 for any ICH, figure 4A), including symptomatic ICH (sOR 0.42, 95% CI 0.25 to 0.71, figure 4B, adjusted sOR 0.23, 95% CI 0.11 to 0.48, available in two studies). Correspondingly, there was reduced outcome fatality in patients in whom TIC12c/3 reperfusion was achieved (sOR 0.59, 95% CI 0.37 to 0.92, see figure 5). All of the above-mentioned associations were also present in an analysis confined to studies applying mTICI (mTIC13 vs mTIC12b, data not shown).

No asymmetry was noted for the analyses concerning the rates of functional independence at day 90, as revealed by funnel and Doi plot inspection (online supplementary figure V). The LFK index was indicative of no asymmetry (0.98).



**Figure 5** Summary OR TIC12c/3 versus TIC12b for mortality. \*Used extended TICI (eTICI) scale with TIC12c; TIC12c and TIC13 were subsumed under TIC13; for grading used in Liebeskind *et al* see online supplementary table I. TICI, thrombolysis in cerebral infarction.

Due to the nature of the topic under review, no study with random sequence allocation was available. A substantial risk of bias was observed in most studies, however, the most common being incomplete outcome data, selective reporting and blinding of participants and personnel (online supplementary table VII). The most common features reducing the quality of the respective studies were lack of core lab adjudicated reperfusion grading and lack of reporting on adjusted analyses (online supplementary table VII).

### Further semiquantitative and qualitative synopsis

One study provided a comparison of oTICI2b and oTICI3 patients but was not included because of a significant overlap of the study cohort with another analysis.<sup>6 16</sup> This article remains of interest, however, as it uses the more conservative oTICI scale in its definition of grade 2b. Nonetheless, a significant outcome difference was present, suggesting that the outcome discrepancies recognised in the above outlined meta-analysis are also present when applying the strictest scale.

A recent observational study failed to prove the existence of significant outcome differences among TICI3 and TICI2b patients when applying the oTICI or mTICI scale, although a clear trend was recognisable.<sup>15</sup> However, significant differences between TICI2c/3 and TICI2b were noted if the eTICI scale was applied. The study was excluded from quantitative analyses because no dichotomised mRS values were provided.<sup>15</sup>

One study included in the quantitative analysis group did not disclose an adjusted analysis for dichotomised analysis, although an ordinal regression analysis was performed. After correction for age, sex, pretreatment National Institutes of Health Stroke Scale score, target occlusion, infarct core and pretreatment alteplase, TICI3 was independently associated with a favourable mRS shift at day 90.<sup>23</sup>

Two of the included studies also provided analyses of tissue outcomes. Rangaraju *et al*<sup>27</sup> reported a significant reduction in final infarct volume in patients achieving TICI3 as compared with TICI2b reperfusion (6.2cc vs 22.5cc,  $P=0.007$ ). Corroborating this finding, Chamorro *et al* reported smaller final infarct volumes and reduced infarct growth in patients with TICI3 reperfusion. Importantly, this association remained statistically tangible after the correction for covariates, including infarct core on CT perfusion.<sup>23</sup>

## DISCUSSION

The study-level meta-analysis incorporating data from 2379 patients provides further evidence that the outcome of patients with TICI3 reperfusion is superior to that of patients in whom TICI2b reperfusion is achieved. This discrepancy was evident from multiple endpoints and even more pronounced when analyses were restricted to adjusted estimates. The observed effect remains significant irrespective which TICI score is applied. Logically, the better the TICI score, the more tissue is reperfused, and the smaller the chance for the penumbra to evolve into infarct.<sup>32 33</sup> However, the present analysis has substantiated that this specifically holds true also for different degrees of successful reperfusion (ie, TICI2b vs TICI3). This implies that TICI3 should be reported separately from TICI2b reperfusion in all future studies and calls into question whether the definition of successful reperfusion should be refined.<sup>6 25</sup>

Various versions of TICI scales already exist.<sup>34 35</sup> So far, the three most commonly used are the oTICI,<sup>2</sup> the mTICI<sup>5</sup> and the eTICI with the implementation of grade TICI2c.<sup>11 12</sup> All of these scales have an acceptable inter-rater reliability.<sup>15 36 37</sup> The TICI2c

score was first mentioned by Noser *et al*<sup>11</sup> and later revisited by Goyal *et al*.<sup>12</sup> Its primary intention was to better characterise and subcategorise successful reperfusion. So far, substantial evidence suggests that patients with a TICI2c reperfusion follow the same clinical course as TICI3 patients.<sup>15–17</sup> Some TICI2c reperfusion would be classified as TICI2b according to the mTICI and oTICI systems.<sup>15 17</sup> The eTICI systems therefore appear to be the best biomarker scale to predict patient outcome more accurately.<sup>15 17</sup> Furthermore, the clinical impact of TICI2b might be influenced by the eloquence of the non-reperfused area, a factor currently neglected. Distinguishing eloquent (TICI2b<sup>E</sup>) from non-eloquent (TICI2b<sup>NE</sup>) reperfusion might have added value but may also add unnecessary complexity to the scale.

Numerous reasons for successful but incomplete reperfusion are conceivable. The most common reason is probably iatrogenic distal embolisation during the thrombectomy manoeuvre, since preinterventional thrombus fragmentation with multiple emboli prior to thrombectomy is only rarely observed.<sup>38</sup> Another explanation could be microcirculatory failure due to vascular dysregulation or progressive oedema.<sup>39</sup> Another aspect to consider is that full parenchymal reperfusion may occur retrogradely via well-developed pial collaterals, despite some very distal emboli impeding antegrade flow, corresponding to the definition of eTICI2c, a functional equivalent of TICI3. Numerous studies reported that good collaterals favour excellent angiographic results.<sup>7 40–42</sup> However, the impact of TICI3 reperfusion on outcome seems to be independent of good collaterals and independent of time until reperfusion is achieved (cf adjusted estimates). Recent evidence from a large registry supports the notion that the advancement in technical equipment and operators' experience results in increasing rates of TICI3 reperfusion.<sup>43</sup> Increasing rates of TICI3 result from protection devices and techniques preventing distal embolisation, or due to the operators' dedication to treat remaining distal emboli.<sup>6</sup> A recent meta-analysis has shown that balloon guiding catheters (BGC) increase good angiographic outcomes.<sup>44</sup> Moreover, there are emerging techniques combining BGC and distal aspiration with stent retrievers or stent retriever-assisted vacuum-locked extraction of clots aimed at achieving maximum protection.<sup>45 46</sup> Given these results, it seems reasonable that technical efforts should be maximised to reduce the risks of periprocedural thrombus fragmentation. However, a well-balanced consideration of risks associated with these techniques should be made.

We have not found evidence that pretreatment with intravenous tPA favours achieving TICI3 instead of TICI2b reperfusion, as no differences in the rates of intravenous tPA administration between TICI2b and TICI2c/3 patients were observed. Results from an animal study have suggested a benefit of intravenous tPA in reducing downstream microvascular thrombosis during large vessel recanalisation.<sup>47</sup> However, equal rates of TICI3 reperfusion between patients treated with direct mechanical thrombectomy and bridging have been reported in recent observational studies.<sup>48–50</sup> Results from currently enrolling randomised controlled trials evaluating direct mechanical thrombectomy versus bridging (SWIFT-DIRECT, NCT03192332 and MR CLEAN-NO IV, ISRCTN80619088) will provide further high-quality evidence regarding the potential value of intravenous tPA regarding this issue.

Additionally, we cannot give a general recommendation to treat vascular occlusions causing TICI2b rather than TICI3 reperfusion only because TICI3 reperfusion is associated with better outcomes. Although a recent publication has addressed the technical feasibility and safety of manoeuvres aiming to improve TICI2b reperfusion to TICI3 reperfusion,<sup>16</sup>

this topic deserves further evaluation in a prospective design. In summary, the future direction of research should aim at evaluating strategies to increase the rate of TICI3 reperfusion, the ultimate angiographic benchmark of best clinical success and outcome.

### Strengths and limitations

So far, this is the largest pooled patient sample comparing the clinical outcome of patients with TICI2b and TICI3 reperfusion. However, this analysis has several limitations, mostly reflecting the limitations of the included studies. Most of these studies were retrospective observational analyses, giving them scope for selection, publication and detection bias. Furthermore, most reperfusion statuses were not core lab adjudicated, nor was the clinical endpoint assessment blinded. Additionally, outcome differences between TICI2b and TICI3 depend on the inclusion and exclusion criteria applied at each site. Although we tried to account for this heterogeneity using a more conservative statistical approach, we cannot exclude that this affected our analyses. Further evaluation derived from large registries may ultimately clarify whether core lab evaluated TICI3 vs TICI2b reperfusion is as clinically relevant as it appears on outcome and independent of potential covariates.

### CONCLUSION

Without considerable heterogeneity and across a wide range of clinical and biomarker endpoints analysed, TICI3 reperfusion is associated with superior outcome and safety compared with TICI2b. This effect seems to be independent of potential confounders (eg, time to reperfusion, collaterals). Data regarding the interaction and interdependence of these factors, however, are sparse. As reperfusion quality is the most important modifiable predictor of patient outcome, a more conservative definition of therapy success and further evaluation of treatment approaches geared towards achieving TICI3 reperfusion by preventing or treating distal emboli more efficiently are warranted.

**Acknowledgements** We thank Susan Kaplan for the writing assistance and language editing.

**Contributors** Study design: JK, JG, UF. Study supervision: UF, JG. Statistical analysis: JK. Initial draft: JK. Revision of the draft for important intellectual content: all authors. Interpretation of results: all authors. Eligibility rating and quality assessment: JK, TD, MRH. Extraction of raw frequency counts: JK, MRH.

**Funding** This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** Unrelated: JG is a global PI of STAR, CEC member of the PROMISE study (Penumbra), Consultancy; is the global PI for the SWIFT DIRECT study (Medtronic), Consultancy; and receives SNSF grants for magnetic resonance imaging in stroke. UF is the global PI for the SWIFT DIRECT study (Medtronic), Consultancy, and receives research grants from the Swiss National Science Foundation (SNSF). All other authors have nothing to disclose.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** All data are available within the respective analyses included into this meta-analysis. Extracted raw frequencies are available in the online supplementary dataset 1.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2018. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

### REFERENCES

- Manning NW, Chapot R, Meyers PM. Endovascular stroke management: key elements of success. *Cerebrovasc Dis* 2016;42:170–7.
- Higashida RT, Furlan AJ, Roberts H, et al. Trial design and reporting standards for intraarterial cerebral thrombolysis for acute ischemic stroke. *Journal of Vascular and Interventional Radiology* 2003;14:E1–E31.
- Yoo AJ, Simonsen CZ, Prabhakaran S, et al. Refining angiographic biomarkers of revascularization: improving outcome prediction after intra-arterial therapy. *Stroke* 2013;44:2509–12.
- Jayaraman MV, Grossberg JA, Meisel KM, et al. The clinical and radiographic importance of distinguishing partial from near-complete reperfusion following intra-arterial stroke therapy. *AJNR Am J Neuroradiol* 2013;34:135–9.
- Zaidat OO, Yoo AJ, Khatri P, et al. Recommendations on angiographic revascularization grading standards for acute ischemic stroke: a consensus statement. *Stroke* 2013;44:2650–63.
- Kleine JF, Wunderlich S, Zimmer C, et al. Time to redefine success? TICI 3 versus TICI 2b recanalization in middle cerebral artery occlusion treated with thrombectomy. *J Neurointerv Surg* 2017;9:117–21.
- Dargazanli C, Consoli A, Barral M, et al. Impact of modified TICI 3 versus Modified TICI 2b reperfusion score to predict good outcome following endovascular therapy. *AJNR Am J Neuroradiol* 2017;38:90–6.
- Kaesmacher J. Striving for the best: how far should we go? regarding "impact of modified tici 3 versus modified tici 2b reperfusion score to predict good outcome following endovascular therapy". *AJNR Am J Neuroradiol* 2017;38:E39.
- Moher D, Liberati A, Tetzlaff J, et al. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009;6:e1000097.
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *Jama* 2000;283:2008–12.
- Noser EA, Shaltoni HM, Hall CE, et al. Aggressive mechanical clot disruption: a safe adjunct to thrombolytic therapy in acute stroke? *Stroke* 2005;36:292–6.
- Goyal M, Fargen KM, Turk AS, et al. 2C or not 2C: defining an improved revascularization grading scale and the need for standardization of angiography outcomes in stroke trials. *J Neurointerv Surg* 2014;6:83–6.
- Liebelskind DS, Jovin TG, Majoie CB, et al. TICI Reperfusion in HERMES: Success in Endovascular Stroke Therapy. *Stroke* 2017;48(Suppl 1):A128.
- Schardt C, Adams MB, Owens T, et al. Utilization of the PICO framework to improve searching PubMed for clinical questions. *BMC Med Inform Decis Mak* 2007;7:16.
- Tung EL, McTaggart RA, Baird GL, et al. Rethinking Thrombolysis in Cerebral Infarction 2b. *Stroke* 2017;48:2488–93.
- Kaesmacher J, Maegerlein C, Zibold F, et al. Improving mTICI2b reperfusion to mTICI2c/3 reperfusion: A retrospective observational study assessing technical feasibility, safety and clinical efficacy. *Eur Radiol* 2018;28.
- Almekhlafi MA, Mishra S, Desai JA, et al. Not all "successful" angiographic reperfusion patients are an equal validation of a modified TICI scoring system. *Interv Neuroradiol* 2014;20:21–7.
- Fiorelli M, Bastianello S, Kummer V, et al. Hemorrhagic transformation within 36 hours of a cerebral infarct. *Stroke* 1999;30:2280–4.
- Doi SA, Barendregt JJ, Khan S, et al. Advances in the meta-analysis of heterogeneous clinical trials I: The inverse variance heterogeneity model. *Contemp Clin Trials* 2015;45:130–8.
- Higgins JPT, Thompson SG. Quantifying heterogeneity in a meta-analysis. *Stat Med* 2002;21:1539–58.
- SAR D, Barendregt JJ. Meta-analysis I. In: SAR D, Williams GM, eds. *Methods of Clinical Epidemiology*. Berlin, Heidelberg: Springer, 2013:229–52.
- Higgins JP, Altman DG, Gøtzsche PC, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *BMJ* 2011;343:d5928.
- Chamorro Á, Blasco J, López A, et al. Complete reperfusion is required for maximal benefits of mechanical thrombectomy in stroke patients. *Sci Rep* 2017;7:11636.
- Schmitz ML, Yeatts SD, Tomsick TA, et al. Recanalization and Angiographic Reperfusion Are Both Associated with a Favorable Clinical Outcome in the IMS III Trial. *Interv Neurol* 2016;5:118–22.
- Carvalho A, Santos T, Cunha A, et al. Need for refining successful revascularization in endovascular treatment of acute ischemic stroke: Data from real-world. *J Neurol Sci* 2018;384.
- Linfante I, Starosciak AK, Walker GR, et al. Predictors of poor outcome despite recanalization: a multiple regression analysis of the NASA registry. *J Neurointerv Surg* 2016;8:224–9.
- Rangaraju S, Aghaebrahim A, Streib C, et al. Abstract T MP7: TICI 2B vs. TICI 3: differences in infarct volumes and clinical outcomes in proximal intracranial large vessel occlusions treated with endovascular therapy. *Stroke* 2014;45:ATMP7 [http://stroke.ahajournals.org/content/45/Suppl\\_1/ATMP7.abstract](http://stroke.ahajournals.org/content/45/Suppl_1/ATMP7.abstract).
- Humphries W, Hoit D, Doss VT, et al. Distal aspiration with retrievable stent assisted thrombectomy for the treatment of acute ischemic stroke. *J Neurointerv Surg* 2015;7:90–4.
- Massari F, Henninger N, Lozano JD, et al. ARTS (Aspiration-Retriever Technique for Stroke): Initial clinical experience. *Interv Neuroradiol* 2016;22:325–32.

- 30 Marks MP, Lansberg MG, Mlynash M, *et al.* Correlation of AOL recanalization, TIMI reperfusion and TIC1 reperfusion with infarct growth and clinical outcome. *J Neurointerv Surg* 2014;6:724–8.
- 31 Goyal N, Tsivgoulis G, Frei D, *et al.* O-008 Comparison between tici 2b and tici 3. *J Neurointerv Surg* 2017;9(Suppl 1):A5–A6 [http://jn.is.bmj.com/content/9/Suppl\\_1/A5.2.abstract](http://jn.is.bmj.com/content/9/Suppl_1/A5.2.abstract).
- 32 Marks MP, Lansberg MG, Mlynash M, *et al.* Angiographic outcome of endovascular stroke therapy correlated with MR findings, infarct growth, and clinical outcome in the DEFUSE 2 trial. *Int J Stroke* 2014;9:860–5.
- 33 Manning NW, Campbell BC, Oxley TJ, *et al.* Acute ischemic stroke: time, penumbra, and reperfusion. *Stroke* 2014;45:640–4.
- 34 Fugate JE, Klunder AM, Kallmes DF. What Is Meant by “TICI(“)? *Am J Neuroradiol* 2013;34:1792–7.
- 35 Gerber JC, Miaux YJ, von Kummer R. Scoring flow restoration in cerebral angiograms after endovascular revascularization in acute ischemic stroke patients. *Neuroradiology* 2015;57:227–40.
- 36 Suh SH, Cloft HJ, Fugate JE, *et al.* Clarifying differences among thrombolysis in cerebral infarction scale variants: is the artery half open or half closed? *Stroke* 2013;44:1166–8.
- 37 Volny O, Cimflova P, Szeder V. Inter-Rater Reliability for Thrombolysis in Cerebral Infarction with TIC1 2c Category. *J Stroke Cerebrovasc Dis* 2017;26:2–4.
- 38 Gratz PP, Schroth G, Gralla J, *et al.* Whole-brain susceptibility-weighted thrombus imaging in stroke: fragmented thrombi predict worse outcome. *AJNR Am J Neuroradiol* 2015;36:1277–82.
- 39 Arsava EM, Arat A, Topcuoglu MA, *et al.* Angiographic microcirculatory obstructions distal to occlusion signify poor outcome after endovascular treatment for acute ischemic stroke. *Transl Stroke Res* 2018;9.
- 40 Liebeskind DS, Jahan R, Nogueira RG, *et al.* Impact of collaterals on successful revascularization in Solitaire FR with the intention for thrombectomy. *Stroke* 2014;45:2036–40.
- 41 Singer OC, Berkefeld J, Nolte CH, *et al.* Collateral vessels in proximal middle cerebral artery occlusion: the ENDOSTROKE study. *Radiology* 2015;274:851–8.
- 42 Leng X, Fang H, Leung TW, *et al.* Impact of collateral status on successful revascularization in endovascular treatment: a systematic review and meta-analysis. *Cerebrovasc Dis* 2016;41:27–34.
- 43 Zaidat OO, Castonguay AC, Nogueira RG, *et al.* TREVO stent-retriever mechanical thrombectomy for acute ischemic stroke secondary to large vessel occlusion registry. *J Neurointerv Surg* 2017. doi: 10.1136/neurintsurg-2017-013328. [Epub ahead of print].
- 44 Brinjikji W, Starke RM, Murad MH, *et al.* Impact of balloon guide catheter on technical and clinical outcomes: a systematic review and meta-analysis. *J Neurointerv Surg* 2017. doi: 10.1136/neurintsurg-2017-013179. [Epub ahead of print].
- 45 Stampfl S, Pfaff J, Herweh C, *et al.* Combined proximal balloon occlusion and distal aspiration: a new approach to prevent distal embolization during neurothrombectomy. *J Neurointerv Surg* 2017;9:346–51.
- 46 Maus V, Behme D, Kabbasch C, *et al.* Maximizing First-pass complete reperfusion with SAVE. *Clin Neuroradiol* 2017.
- 47 Desilles JP, Loyau S, Syvannarath V, *et al.* Alteplase reduces downstream microvascular thrombosis and improves the benefit of large artery recanalization in stroke. *Stroke* 2015;46:3241–8.
- 48 Coutinho JM, Liebeskind DS, Slater LA, *et al.* Combined intravenous thrombolysis and thrombectomy vs thrombectomy alone for acute ischemic stroke: a pooled analysis of the SWIFT and STAR Studies. *JAMA Neurol* 2017;74:268–74.
- 49 Dávalos A, Pereira VM, Chapot R, *et al.* Retrospective multicenter study of Solitaire FR for revascularization in the treatment of acute ischemic stroke. *Stroke* 2012;43:2699–705.
- 50 Bellwald S, Weber R, Dobrocky T, *et al.* Direct mechanical intervention versus bridging therapy in stroke patients eligible for intravenous thrombolysis: a pooled analysis of 2 registries. *Stroke* 2017;48:3282–8.