

## (Z)-3-(3,4-Dimethoxybenzyl)-1,5-benzo-thiazepin-4(5H)-one

R. Selvakumar,<sup>a</sup> M. Bakthadoss,<sup>a,b</sup>‡ S. Vijayakumar<sup>c</sup> and S. Murugavel<sup>d</sup>\*

<sup>a</sup>Department of Organic Chemistry, University of Madras, Marina Campus, Chennai 600 025, India, <sup>b</sup>Department of Chemistry, Pondicherry University, Puducherry 605 014, India, <sup>c</sup>Department of Physics, Sri Balaji Chokkalingam Engineering College, Arni, Thiruvannamalai 632 317, India, and <sup>d</sup>Department of Physics, Thanthai Periyar Government Institute of Technology, Vellore 632 002, India

Correspondence e-mail: smurugavel27@gmail.com


Received 1 April 2013; accepted 8 April 2013

Key indicators: single-crystal X-ray study;  $T = 293\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$ ;  $R$  factor = 0.060;  $wR$  factor = 0.175; data-to-parameter ratio = 62.7.

In the title compound,  $\text{C}_{18}\text{H}_{17}\text{NO}_3\text{S}$ , the thiazepine ring adopts a slightly distorted twist-boat conformation. The dihedral angle between the mean plane of the benzothiazepine ring system and the benzene ring is  $60.3(1)^\circ$ . In the crystal, molecules are linked by two pairs of inversion-related  $\text{N}-\text{H}\cdots\text{O}$  and  $\text{C}-\text{H}\cdots\text{O}$  hydrogen bonds, generating alternating  $R_2^2(8)$  and  $R_2^2(6)$  ring motifs, respectively, in a zigzag supramolecular chain that runs along the  $c$  axis. These chains stack along the  $a$  axis via  $\text{S}\cdots\text{C}$  [3.424 (2)  $\text{\AA}$ ] contacts. A three-dimensional supramolecular network is consolidated by  $\text{C}-\text{H}\cdots\pi$  and  $\pi-\pi$  interactions [inter-centroid distance between dimethoxybenzene rings = 3.815 (1)  $\text{\AA}$ ]. The crystal studied was a non-merohedral twin, with a refined value of the minor twin fraction of 0.2477 (6).

### Related literature

For background to the biology of thiazepine derivatives and for a related structure, see: Bakthadoss *et al.* (2013). For ring-puckering parameters, see: Cremer & Pople (1975).



### Experimental

#### Crystal data

$\text{C}_{18}\text{H}_{17}\text{NO}_3\text{S}$	$V = 3212.0(11)\text{ \AA}^3$
$M_r = 327.39$	$Z = 8$
Orthorhombic, $Pbcn$	Mo $K\alpha$ radiation
$a = 19.966(4)\text{ \AA}$	$\mu = 0.22\text{ mm}^{-1}$
$b = 10.355(2)\text{ \AA}$	$T = 293\text{ K}$
$c = 15.536(3)\text{ \AA}$	$0.35 \times 0.20 \times 0.15\text{ mm}$

#### Data collection

Bruker APEXII CCD diffractometer	13227 measured reflections
Absorption correction: multi-scan ( <i>TWINABS</i> ; Sheldrick, 1997)	13227 independent reflections
$T_{\min} = 0.927$ , $T_{\max} = 0.968$	8413 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.066$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.060$	211 parameters
$wR(F^2) = 0.175$	H-atom parameters constrained
$S = 1.02$	$\Delta\rho_{\text{max}} = 0.25\text{ e \AA}^{-3}$
13227 reflections	$\Delta\rho_{\text{min}} = -0.33\text{ e \AA}^{-3}$

**Table 1**  
Hydrogen-bond geometry ( $\text{\AA}$ ,  $^\circ$ ).

$Cg1$  and  $Cg2$  are the centroids of the C11–C16 and C2–C7 rings, respectively.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
N1–H1A…O1 <sup>i</sup>	0.86	2.02	2.863 (2)	167
C18–H18B…O3 <sup>ii</sup>	0.96	2.53	3.445 (3)	159
C4–H4…Cg1 <sup>iii</sup>	0.93	2.57	3.450 (2)	158
C10–H10B…Cg2 <sup>iv</sup>	0.97	2.82	3.725 (2)	155

Symmetry codes: (i)  $-x, -y, -z + 2$ ; (ii)  $-x, -y + 1, -z + 1$ ; (iii)  $-x + \frac{1}{2}, -y + \frac{1}{2}, z + \frac{1}{2}$ ; (iv)  $x, -y, z - \frac{3}{2}$ .

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97* and *PLATON* (Spek, 2009).

The authors thank Dr Babu Vargheese, SAIF, IIT, Madras, India, for his help with the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK5214).

### References

- Bakthadoss, M., Selvakumar, R., Manikandan, N. & Murugavel, S. (2013). *Acta Cryst. E69*, o562–o563.
- Bruker (2004). *APEX2, SAINT* and *XPREP*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cremer, D. & Pople, J. A. (1975). *J. Am. Chem. Soc.* **97**, 1354–1358.
- Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.
- Sheldrick, G. M. (1997). *TWINABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A64*, 112–122.
- Spek, A. L. (2009). *Acta Cryst. D65*, 148–155.

‡ Additional correspondence author, e-mail: bhakthadoss@yahoo.com.

# supplementary materials

*Acta Cryst.* (2013). E69, o708 [doi:10.1107/S1600536813009598]

## (Z)-3-(3,4-Dimethoxybenzyl)-1,5-benzothiazepin-4(5H)-one

R. Selvakumar, M. Bakthadoss, S. Vijayakumar and S. Murugavel

### Comment

The background to the biology and related structure of thiazepin derivatives, has been described recently (Bakthadoss *et al.*, 2013). In view of this biological importance, the crystal structure of the title compound has been carried out and the results are presented here.

Fig. 1. shows a displacement ellipsoid plot of (I), with the atom numbering scheme. The seven membered thiazepine ring (N1/S1/C1/C2/C7/C8/C9) adopts slightly distorted twist-boat conformation as indicated by puckering parameters (Cremer & Pople, 1975): QT = 0.9884 (11) Å,  $\varphi_2 = 359.3$  (1)° and  $\varphi_3 = 356.5$  (4)°. The dihedral angle between the benzothiazepin ring system and the benzene ring is 60.3 (1)°. The atom O1 deviates by -0.895 (1) Å from the least-squares plane of the thiazepin ring. The sum of angles at N1 atom of the thiazepin ring (359.9°) is in accordance with  $sp^2$  hybridization. The geometric parameters of the title molecule agree well with those reported for a similar structure (Bakthadoss *et al.*, 2013).

In the crystal, molecules are linked by two pairs of inversion-related N1—H1A…O1 and C18—H18B…O3 hydrogen bonds, generating alternate  $R_2^2(8)$  and  $R_2^2(6)$  ring motifs, respectively, resulting in a zigzag supramolecular chain running along the *c* axis. These chains stack along the *a* axis by S1…C4<sup>v</sup> = 3.424 (2) Å (Symmetry code: (v) = 1/2 - *x*, -1/2 + *y*, *z*) short contacts (Fig. 2 and Table 1). A three-dimensional supramolecular network is consolidated by C4—H4…Cg1<sup>iii</sup> (symmetry code: (iii) = 1/2 - *x*, 1/2 - *y*, 1/2 + *z*) and C10—H10B…Cg2<sup>iv</sup> (symmetry code: (iv) = *x*, -*y*, -1/2 + *z*) hydrogen bonds and Cg1—Cg1<sup>vi</sup> = 3.815 (1) Å (symmetry code: (vi) = -*x*, *y*, 3/2 - *z*) interactions (Fig. 3 and Table 1; Cg1 and Cg2 are the centroids of the C11–C16 and C2–C7 benzene rings, respectively).

### Experimental

A mixture of (*Z*)-methyl 2-(bromomethyl)-3-(3,4-dimethoxyphenyl)acrylate 2 mmol) and *o*-aminothiophenol (2 mmol) in the presence of potassium *tert*-butoxide (4.8 mmol) in dry THF (10 ml) was stirred at room temperature for 1 h. After the completion of the reaction as indicated by TLC, the reaction mixture was concentrated and the resulting crude mass was diluted with water (20 ml) and extracted with ethyl acetate (3 x 20 ml). The organic layer was washed with brine (2 x 20 ml) and dried over anhydrous sodium sulfate. The organic layer was concentrated, which successfully provide the crude final product ((*Z*)-3-(3,4-dimethoxybenzyl)benzo[*b*][1,4]thiazepin-4(5*H*)-one). This product was purified by column chromatography on silica gel to afford the title compound in good yield (44%). Single crystals suitable for X-ray diffraction were obtained by slow evaporation of its ethylacetate solution at room temperature.


### Refinement

The investigated crystal was found to be a two-component rotational twin. The data for both components were integrated using SAINT and scaled with TWINABS. Final refinement was performed using a HKLF5 file generated by TWINABS with a BASF parameter (0.2477 (6)). Owing to poor agreement, the reflections (-7 5 5), (-7 5 7), (-8 4 - 5), (8 5 5), (2 0 0), (-8 -

$(5\ 5)$ ,  $(8\ 5 - 5)$ ,  $(8 - 5\ 5)$ ,  $(-8 - 5 - 5)$ ,  $(-8 - 6\ 5)$ ,  $(-7\ 5\ 3)$ ,  $(-8 - 6 - 5)$ ,  $(7 - 5 - 3)$ ,  $(14\ 2\ 2)$  and  $(8 - 6\ 5)$  were omitted from the final cycles of refinement. All the H atoms were positioned geometrically and constrained to ride on their parent atom with C—H = 0.93–0.97 Å and N—H = 0.86 Å, and with  $U_{\text{iso}}(\text{H})=1.5U_{\text{eq}}$  for methyl H atoms and  $1.2U_{\text{eq}}(\text{C})$  for other H atoms.


### Computing details

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREP* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *SHELXL97* (Sheldrick, 2008) and *PLATON* (Spek, 2009).




**Figure 1**

Molecular structure of the title compound showing displacement ellipsoids at the 30% probability level. H atoms are presented as a small spheres of arbitrary radii.

**Figure 2**

A view of the supramolecular chain showing N—H···O (red dotted lines) and C—H···O (blue dotted lines) hydrogen bonds. The chains stack along the the *a* axis via S···C (green dotted lines) short contacts.

**Figure 3**

View of three-dimensional supramolecular network down the *c* axis. The N—H···O, C—H···O, S···C, C—H··· $\pi$  and  $\pi$ — $\pi$  interactions are shown as red, blue, green, orange and purple dotted lines, respectively. Cg1 and Cg2 are the centroids of the C11—C16 and C2—C7 benzene rings, respectively.

#### (Z)-3-(3,4-Dimethoxybenzyl)-1,5-benzothiazepin-4(5*H*)-one

##### Crystal data

C<sub>18</sub>H<sub>17</sub>NO<sub>3</sub>S

*M*<sub>r</sub> = 327.39

Orthorhombic, *Pbcn*

Hall symbol: -P 2n 2ab

*a* = 19.966 (4) Å

*b* = 10.355 (2) Å

*c* = 15.536 (3) Å

*V* = 3212.0 (11) Å<sup>3</sup>

*Z* = 8

*F*(000) = 1376

*D*<sub>x</sub> = 1.354 Mg m<sup>-3</sup>

Mo *K* $\alpha$  radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 2790 reflections

$\theta$  = 2.2–24.9°

$\mu$  = 0.22 mm<sup>-1</sup>

*T* = 293 K

Block, colourless

0.35 × 0.20 × 0.15 mm

*Data collection*

Bruker APEXII CCD  
diffractometer  
Radiation source: fine-focus sealed tube  
Graphite monochromator  
Detector resolution: 10.0 pixels mm<sup>-1</sup>  
 $\omega$  scans  
Absorption correction: multi-scan  
(TWINABS; Sheldrick, 1997)  
 $T_{\min} = 0.927$ ,  $T_{\max} = 0.968$

13227 measured reflections  
13227 independent reflections  
8413 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.066$   
 $\theta_{\max} = 24.9^\circ$ ,  $\theta_{\min} = 2.2^\circ$   
 $h = -23 \rightarrow 23$   
 $k = -12 \rightarrow 9$   
 $l = -18 \rightarrow 18$

*Refinement*

Refinement on  $F^2$   
Least-squares matrix: full  
 $R[F^2 > 2\sigma(F^2)] = 0.060$   
 $wR(F^2) = 0.175$   
 $S = 1.02$   
13227 reflections  
211 parameters  
0 restraints  
Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map  
Hydrogen site location: inferred from  
neighbouring sites  
H-atom parameters constrained  
 $w = 1/[\sigma^2(F_o^2) + (0.0582P)^2]$   
where  $P = (F_o^2 + 2F_c^2)/3$   
 $(\Delta/\sigma)_{\max} = 0.001$   
 $\Delta\rho_{\max} = 0.25 \text{ e } \text{\AA}^{-3}$   
 $\Delta\rho_{\min} = -0.32 \text{ e } \text{\AA}^{-3}$

*Special details*

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) etc. and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

*Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )*

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
C2	0.21158 (10)	0.01584 (18)	1.02679 (11)	0.0320 (5)
C3	0.26554 (10)	0.0619 (2)	1.07376 (13)	0.0401 (5)
H3	0.3072	0.0225	1.0683	0.048*
C4	0.25791 (10)	0.1650 (2)	1.12822 (13)	0.0434 (6)
H4	0.2939	0.1935	1.1611	0.052*
C5	0.19688 (11)	0.2264 (2)	1.13416 (12)	0.0405 (5)
H5	0.1921	0.2978	1.1700	0.049*
C6	0.14302 (10)	0.18270 (19)	1.08740 (11)	0.0339 (5)
H6	0.1020	0.2249	1.0913	0.041*
C7	0.14978 (9)	0.07580 (18)	1.03449 (10)	0.0265 (4)
C8	0.07914 (10)	-0.00895 (17)	0.91369 (11)	0.0300 (5)
C9	0.13353 (9)	-0.00907 (17)	0.84820 (11)	0.0277 (5)
C1	0.19328 (11)	-0.05969 (19)	0.86307 (11)	0.0359 (5)
H1	0.2230	-0.0622	0.8170	0.043*
C10	0.11450 (10)	0.04089 (18)	0.76116 (11)	0.0332 (5)
H10A	0.0735	-0.0016	0.7432	0.040*

H10B	0.1493	0.0170	0.7206	0.040*
C11	0.10408 (8)	0.18544 (17)	0.75624 (10)	0.0267 (4)
C12	0.12408 (10)	0.2698 (2)	0.81859 (11)	0.0369 (5)
H12	0.1440	0.2384	0.8685	0.044*
C13	0.11523 (10)	0.4015 (2)	0.80865 (12)	0.0406 (5)
H13	0.1286	0.4574	0.8522	0.049*
C14	0.08704 (10)	0.45013 (18)	0.73534 (12)	0.0345 (5)
C15	0.06728 (9)	0.36546 (19)	0.67025 (11)	0.0305 (5)
C16	0.07503 (9)	0.23488 (19)	0.68158 (10)	0.0298 (5)
H16	0.0607	0.1785	0.6388	0.036*
C18	0.02274 (13)	0.3361 (2)	0.52964 (13)	0.0661 (8)
H18A	0.0616	0.2902	0.5099	0.099*
H18B	0.0042	0.3852	0.4830	0.099*
H18C	-0.0100	0.2756	0.5502	0.099*
C17	0.08134 (13)	0.6638 (2)	0.78994 (14)	0.0715 (8)
H17A	0.0536	0.6325	0.8359	0.107*
H17B	0.0665	0.7483	0.7730	0.107*
H17C	0.1270	0.6684	0.8091	0.107*
N1	0.09093 (8)	0.02648 (15)	0.99504 (9)	0.0325 (4)
H1A	0.0573	0.0180	1.0291	0.039*
O1	0.02201 (7)	-0.04165 (14)	0.89186 (8)	0.0423 (4)
O3	0.04118 (7)	0.42072 (13)	0.59744 (8)	0.0462 (4)
O2	0.07680 (8)	0.57865 (13)	0.71876 (9)	0.0498 (4)
S1	0.22192 (3)	-0.12256 (6)	0.96130 (3)	0.04692 (18)

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C2	0.0340 (13)	0.0353 (12)	0.0267 (10)	0.0018 (10)	-0.0032 (9)	0.0069 (9)
C3	0.0285 (13)	0.0482 (15)	0.0437 (12)	0.0032 (10)	-0.0051 (9)	0.0068 (11)
C4	0.0369 (15)	0.0530 (15)	0.0404 (12)	-0.0140 (12)	-0.0120 (10)	0.0064 (11)
C5	0.0447 (14)	0.0402 (13)	0.0366 (11)	-0.0090 (11)	-0.0044 (10)	-0.0018 (10)
C6	0.0347 (13)	0.0359 (13)	0.0311 (10)	-0.0005 (10)	0.0000 (9)	0.0014 (9)
C7	0.0256 (12)	0.0326 (12)	0.0214 (9)	-0.0028 (9)	-0.0011 (8)	0.0067 (9)
C8	0.0302 (13)	0.0278 (12)	0.0320 (10)	0.0017 (9)	-0.0025 (9)	0.0044 (9)
C9	0.0310 (12)	0.0249 (11)	0.0271 (9)	0.0008 (9)	-0.0016 (8)	-0.0015 (8)
C1	0.0391 (14)	0.0386 (13)	0.0300 (10)	0.0034 (10)	0.0048 (9)	-0.0031 (9)
C10	0.0394 (13)	0.0351 (12)	0.0252 (9)	-0.0004 (9)	0.0006 (8)	-0.0022 (9)
C11	0.0263 (12)	0.0297 (11)	0.0242 (9)	-0.0046 (9)	0.0029 (8)	0.0008 (9)
C12	0.0463 (14)	0.0395 (13)	0.0248 (10)	-0.0052 (11)	-0.0030 (9)	-0.0008 (9)
C13	0.0524 (15)	0.0357 (14)	0.0336 (11)	-0.0129 (11)	0.0002 (10)	-0.0083 (10)
C14	0.0406 (14)	0.0246 (12)	0.0384 (11)	-0.0057 (9)	0.0126 (10)	-0.0006 (10)
C15	0.0324 (13)	0.0324 (12)	0.0265 (10)	0.0009 (9)	0.0052 (8)	0.0046 (9)
C16	0.0319 (13)	0.0310 (12)	0.0263 (10)	-0.0039 (9)	0.0006 (8)	-0.0034 (9)
C18	0.111 (2)	0.0510 (16)	0.0358 (13)	0.0186 (15)	-0.0196 (13)	-0.0014 (11)
C17	0.118 (2)	0.0342 (15)	0.0621 (16)	-0.0121 (15)	0.0208 (15)	-0.0165 (13)
N1	0.0229 (10)	0.0474 (11)	0.0273 (8)	-0.0062 (8)	0.0031 (7)	0.0024 (8)
O1	0.0294 (9)	0.0628 (10)	0.0346 (7)	-0.0108 (7)	-0.0031 (6)	-0.0053 (7)
O3	0.0675 (11)	0.0373 (9)	0.0338 (8)	0.0099 (7)	-0.0051 (7)	0.0047 (7)
O2	0.0797 (12)	0.0244 (8)	0.0452 (8)	-0.0014 (8)	0.0113 (8)	-0.0036 (7)

S1	0.0543 (4)	0.0454 (4)	0.0411 (3)	0.0223 (3)	-0.0086 (3)	0.0000 (3)
----	------------	------------	------------	------------	-------------	------------

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

C2—C3	1.386 (3)	C10—H10B	0.9700
C2—C7	1.386 (2)	C11—C12	1.364 (2)
C2—S1	1.770 (2)	C11—C16	1.394 (2)
C3—C4	1.371 (3)	C12—C13	1.383 (3)
C3—H3	0.9300	C12—H12	0.9300
C4—C5	1.377 (3)	C13—C14	1.367 (3)
C4—H4	0.9300	C13—H13	0.9300
C5—C6	1.374 (3)	C14—O2	1.371 (2)
C5—H5	0.9300	C14—C15	1.395 (3)
C6—C7	1.385 (3)	C15—O3	1.370 (2)
C6—H6	0.9300	C15—C16	1.372 (3)
C7—N1	1.420 (2)	C16—H16	0.9300
C8—O1	1.237 (2)	C18—O3	1.419 (2)
C8—N1	1.337 (2)	C18—H18A	0.9600
C8—C9	1.488 (3)	C18—H18B	0.9600
C9—C1	1.323 (3)	C18—H18C	0.9600
C9—C10	1.497 (2)	C17—O2	1.417 (2)
C1—S1	1.7549 (19)	C17—H17A	0.9600
C1—H1	0.9300	C17—H17B	0.9600
C10—C11	1.513 (2)	C17—H17C	0.9600
C10—H10A	0.9700	N1—H1A	0.8600
C3—C2—C7	119.50 (18)	C16—C11—C10	117.55 (16)
C3—C2—S1	119.38 (16)	C11—C12—C13	120.97 (18)
C7—C2—S1	121.07 (14)	C11—C12—H12	119.5
C4—C3—C2	120.44 (19)	C13—C12—H12	119.5
C4—C3—H3	119.8	C14—C13—C12	120.58 (18)
C2—C3—H3	119.8	C14—C13—H13	119.7
C3—C4—C5	119.93 (19)	C12—C13—H13	119.7
C3—C4—H4	120.0	C13—C14—O2	125.16 (17)
C5—C4—H4	120.0	C13—C14—C15	119.27 (18)
C6—C5—C4	120.3 (2)	O2—C14—C15	115.57 (17)
C6—C5—H5	119.8	O3—C15—C16	124.07 (17)
C4—C5—H5	119.8	O3—C15—C14	116.31 (17)
C5—C6—C7	120.03 (19)	C16—C15—C14	119.61 (17)
C5—C6—H6	120.0	C15—C16—C11	121.03 (17)
C7—C6—H6	120.0	C15—C16—H16	119.5
C6—C7—C2	119.72 (17)	C11—C16—H16	119.5
C6—C7—N1	117.57 (17)	O3—C18—H18A	109.5
C2—C7—N1	122.54 (17)	O3—C18—H18B	109.5
O1—C8—N1	119.77 (17)	H18A—C18—H18B	109.5
O1—C8—C9	119.02 (17)	O3—C18—H18C	109.5
N1—C8—C9	121.21 (18)	H18A—C18—H18C	109.5
C1—C9—C8	122.60 (17)	H18B—C18—H18C	109.5
C1—C9—C10	121.56 (17)	O2—C17—H17A	109.5
C8—C9—C10	115.60 (16)	O2—C17—H17B	109.5

C9—C1—S1	126.33 (15)	H17A—C17—H17B	109.5
C9—C1—H1	116.8	O2—C17—H17C	109.5
S1—C1—H1	116.8	H17A—C17—H17C	109.5
C9—C10—C11	114.99 (15)	H17B—C17—H17C	109.5
C9—C10—H10A	108.5	C8—N1—C7	130.72 (16)
C11—C10—H10A	108.5	C8—N1—H1A	114.6
C9—C10—H10B	108.5	C7—N1—H1A	114.6
C11—C10—H10B	108.5	C15—O3—C18	116.96 (16)
H10A—C10—H10B	107.5	C14—O2—C17	116.60 (16)
C12—C11—C16	118.50 (17)	C1—S1—C2	99.30 (9)
C12—C11—C10	123.88 (16)		
C7—C2—C3—C4	0.7 (3)	C11—C12—C13—C14	-1.0 (3)
S1—C2—C3—C4	-176.75 (15)	C12—C13—C14—O2	-179.20 (17)
C2—C3—C4—C5	-2.3 (3)	C12—C13—C14—C15	-0.1 (3)
C3—C4—C5—C6	1.7 (3)	C13—C14—C15—O3	-178.15 (17)
C4—C5—C6—C7	0.5 (3)	O2—C14—C15—O3	1.0 (2)
C5—C6—C7—C2	-2.1 (3)	C13—C14—C15—C16	1.5 (3)
C5—C6—C7—N1	173.30 (16)	O2—C14—C15—C16	-179.36 (16)
C3—C2—C7—C6	1.5 (3)	O3—C15—C16—C11	177.80 (17)
S1—C2—C7—C6	178.92 (13)	C14—C15—C16—C11	-1.8 (3)
C3—C2—C7—N1	-173.68 (16)	C12—C11—C16—C15	0.7 (3)
S1—C2—C7—N1	3.7 (2)	C10—C11—C16—C15	-176.41 (16)
O1—C8—C9—C1	-133.7 (2)	O1—C8—N1—C7	-175.13 (18)
N1—C8—C9—C1	46.7 (3)	C9—C8—N1—C7	4.5 (3)
O1—C8—C9—C10	40.8 (2)	C6—C7—N1—C8	133.3 (2)
N1—C8—C9—C10	-138.76 (18)	C2—C7—N1—C8	-51.4 (3)
C8—C9—C1—S1	-5.3 (3)	C16—C15—O3—C18	-1.4 (3)
C10—C9—C1—S1	-179.56 (14)	C14—C15—O3—C18	178.15 (18)
C1—C9—C10—C11	-113.1 (2)	C13—C14—O2—C17	-15.3 (3)
C8—C9—C10—C11	72.2 (2)	C15—C14—O2—C17	165.61 (18)
C9—C10—C11—C12	14.1 (3)	C9—C1—S1—C2	-58.0 (2)
C9—C10—C11—C16	-168.94 (16)	C3—C2—S1—C1	-125.91 (16)
C16—C11—C12—C13	0.7 (3)	C7—C2—S1—C1	56.68 (17)
C10—C11—C12—C13	177.59 (18)		

*Hydrogen-bond geometry (Å, °)*

Cg1 and Cg2 are the centroids of the C11—C16 and C2—C7 rings, respectively.

D—H···A	D—H	H···A	D···A	D—H···A
N1—H1A···O1 <sup>i</sup>	0.86	2.02	2.863 (2)	167
C18—H18B···O3 <sup>ii</sup>	0.96	2.53	3.445 (3)	159
C4—H4···Cg1 <sup>iii</sup>	0.93	2.57	3.450 (2)	158
C10—H10B···Cg2 <sup>iv</sup>	0.97	2.82	3.725 (2)	155

Symmetry codes: (i)  $-x, -y, -z+2$ ; (ii)  $-x, -y+1, -z+1$ ; (iii)  $-x+1/2, -y+1/2, z+1/2$ ; (iv)  $x, -y, z-3/2$ .