


Emergence of Almost Identical F36:A-:B32 Plasmids Carrying *bla*_{NDM-5} and *qepA* in *Escherichia coli* from Both Pakistan and Canada

This article was published in the following Dove Press journal:
Infection and Drug Resistance

Zulqarnain Baloch ^{1,*}
Luchao Lv^{1,2,*}
Lingxian Yi^{1,2}
Miao Wan^{1,2}
Bilal Aslam³
Jun Yang^{1,2}
Jian-Hua Liu^{1,2}

¹College of Veterinary Medicine, South China Agricultural University, Guangzhou 510642, People's Republic of China;

²Guangdong Provincial Key Laboratory of Veterinary Pharmaceuticals Development and Safety Evaluation, South China Agricultural University, Guangzhou, People's Republic of China; ³Government College University, Faisalabad 54000, Pakistan

*These authors contributed equally to this work

Abstract: The New Delhi Metallo-β-lactamase (NDM) producing *Enterobacteriaceae* is spreading worldwide. Although the *bla*_{NDM} gene has been identified in animal associated *Enterobacteriaceae* isolates in many countries, little is known about its occurrence in animal products in Pakistan. In this study, 13 *Escherichia coli* isolates were collected from chicken meat samples in Pakistan. Two isolates, 15978 and C4109, exhibited reduced susceptibility (MIC ≥1 μg/mL) to imipenem, and carried *bla*_{NDM-5} and *bla*_{NDM-7} gene, respectively. Whole-genome sequencing and Oxford Nanopore MinION sequencing revealed that 15978 and C4109 belonged to ST156 and ST167, respectively. *bla*_{NDM-7} was carried by an IncX3 plasmid that has disseminated worldwide, whereas *bla*_{NDM-5} was located on an F36: A-: B32 plasmid, which shared high identity with two plasmids carried by *E. coli* isolates from other countries (one from a patient in Canada). To the best of our knowledge, this is the first report characterizing *bla*_{NDM}-carrying plasmids from chicken meat samples in Pakistan. The dissemination of almost identical *bla*_{NDM-5}-bearing F36:A-B32 and *bla*_{NDM-7}-bearing IncX3 plasmids in different countries highlights the importance of international trade and travel in the spread of antimicrobial resistance strains and plasmids worldwide.

Keywords: plasmid, animal food, carbapenemase, *bla*_{NDM}

Introduction

Carbapenems are last-resort drugs for treating infections caused by multidrug-resistant (MDR) bacteria. However, resistance to carbapenems in gram-negative bacteria, especially *Enterobacteriaceae*, has increased rapidly over the last decade and poses an increasing threat to global public health.^{1,2} Carbapenem resistance in *Enterobacteriaceae* is primarily attributed to carbapenemase enzymes, especially *Klebsiella pneumoniae* carbapenemase (KPC) and the New Delhi metallo-β-lactamase (NDM).³ *bla*_{NDM} was firstly discovered from a Swedish patient in India during 2007.⁴ Since then, it has been increasingly identified throughout the world and was found epidemic in the Indian subcontinent including Pakistan, Afghanistan, and the Balkans regions etc.⁵ NDM is able to hydrolyze almost all β-lactams, and the hydrolytic activity of NDM enzymes cannot be weakened by β-lactamase inhibitors, such as clavulanate, tazobactam, sulbactam, and avibactam, leaving limited therapeutic options for infections caused by NDM-producing *Enterobacteriaceae*.⁶ *bla*_{NDM} genes are usually located on plasmids capable of efficient transfer between bacterial species and hosts in and out of hospitals.³

Correspondence: Jian-Hua Liu
College of Veterinary Medicine, South China Agricultural University, Guangzhou, Guangdong, People's Republic of China
Email jhliu@scau.edu.cn

Though carbapenems are not legally prescribed for use in livestock production, the occurrence of carbapenem-resistant *Enterobacteriaceae* (CRE), especially NDM-producing *Enterobacteriaceae* has been increasingly reported in livestock and meat products in the world.^{7–9} Yet, systematic study on the prevalence and characterization of NDM-producing *Enterobacteriaceae* from food animals and animal-derived foods remain to be sporadic in Pakistan.¹⁰ Here, for the first time, we characterized two NDM-producing *Escherichia coli* strains that were recovered from retail chicken meat samples in Pakistan.

Materials and Methods

Bacterial Isolation, Antimicrobial Susceptibility Testing, and Detection of the *bla*_{NDM} Gene

In March 2018, fourteen chicken meat samples were collected from local broiler meat outlets in Faisalabad, Pakistan. Antibiotic-free MacConkey agar plates were used to isolate *E. coli* strains. The isolates were identified by MALDI-TOF MS (Shimadzu-Biotech Corp., Kyoto, Japan).

The MICs of 14 antimicrobial agents, including ampicillin, cefotaxime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, ceftazidime, were assessed by either agar dilution or broth microdilution method (colistin and tigecycline) with the *E. coli* strain ATCC 25922 as the control according to CLSI guideline.¹¹ Isolates that showed reduced susceptibility to imipenem (MIC ≥ 1 $\mu\text{g/mL}$) were selected for PCR screening of carbapenemase genes.¹² PCR products were confirmed by sequencing.

Whole Genome Sequencing (WGS)

Whole genome DNA of NDM-producing *E. coli* isolates, 15978 and C4109, was extracted and sequenced using HiSeq (Illumina, San Diego, CA, USA) platforms. Afterwards, SOAPdenovo (version 2.04) was used to assemble sequence reads into contigs and to extract *bla*_{NDM}-bearing plasmid contigs. To obtain the complete sequence of the *bla*_{NDM-5}-carrying plasmid, we then sequenced *E. coli* 15978 on Oxford Nanopore MinION. The assemblies of long Nanopore reads and the short Illumina reads were combined via Unicycler version 0.4.3.¹³ The resistance genes, chromosomal mutations, virulence genes, plasmid type, and MLST of the two *bla*_{NDM} positive strains were analyzed by ResFinder 3.2, PointFinder, VirulenceFinder, PlasmidFinder, and MLST (<https://cge.cbs.dtu.dk/services/>), respectively. Comparative analysis of

*bla*_{NDM}-carrying plasmids was carried out using BLAST tools and BLAST Ring Image Generator (BRIG).¹⁴

Results and Discussion

A total of 13 *E. coli* isolates were recovered from 13 retail chicken meat samples. Two isolates, 15978 and C4109, showed reduced susceptibility to imipenem (MIC ≥ 1 $\mu\text{g/mL}$), and were identified to carry *bla*_{NDM-5} and *bla*_{NDM-7}, respectively (Table 1). The two isolates showed resistance to cefotaxime, ceftazidime, and ceftazidime, but remained susceptible to fosfomycin, colistin, amikacin, gentamycin, and tigecycline (Table 1). In addition, C4109 exhibited resistance to streptomycin and ciprofloxacin, while 15978 showed resistance to doxycycline and ciprofloxacin. *bla*_{NDM-5} and *bla*_{NDM-7} were successfully transferred to recipients *E. coli* C600 or *E. coli* DH5 α by conjugation and transformation, respectively.

WGS showed that *E. coli* 15978 and C4109 belonged to ST156 and ST167, respectively (Table 1). These two *E. coli* sequence types were also related to *bla*_{NDM} dissemination worldwide in humans,^{15,16} animals,^{17,18} and food.¹⁹ The resistance genes, chromosomal mutations, and virulence genes of the two isolates were displayed in Table 1. It showed that 15978 and C4109 carried eight and five other resistance genes, respectively.

In *E. coli* C4109, *bla*_{NDM-7} was carried by a 49,828-bp IncX3 plasmid pHN4109c (MK088485), which shared 92% coverage and 99% identity with pKW53T-NDM (KX214669) from clinical *E. coli* in Kuwait, pNDM5_IncX3 (KU761328) from clinical *K. pneumoniae* in China, and tig00000260 (CP021738) from *E. coli* in USA (Figure 1A). IncX3 has dominated the spread of *bla*_{NDM} in many countries, particularly in Asian countries, such as China,⁹ Korea,²⁰ Myanmar,²¹ and India.²² To the best of our knowledge, this study was the first to identify *bla*_{NDM}-positive IncX3 plasmid in Pakistan. Similar to other reports, *bla*_{NDM-7} was embedded in an IS26-*bla*_{NDM}- Δ Tn2 transposition unit which inserted into *umuD* gene in pHN4109c. However, a 3664-bp transposon Tn5403 was inserted in the IS3000 gene, which formed a unique genetic structure together with 5-bp direct repeats (TACAT) (Figure 1B).

The *bla*_{NDM-5}-carrying plasmid pHN15978 (MK291500) was a 128,762-bp F36:A-B32 plasmid containing 151 ORFs. It was comprised of a typical IncF-type backbone, encoding genes for replication, transfer, maintenance, stability functions, and a multidrug resistance region of 28145-bp. BLAST homology analysis demonstrated that the sequence of pHN15978 showed 99% identity and 100% query coverage with *E. coli* strain AR_452 plasmid unnamed1 (CP030329.1) and

Table 1 Characterization of NDM-Producing *Escherichia coli* Isolates and Transconjugant or Transformant

Strain	C4109	C4109 Transformant	<i>E. coli</i> DH5 α	15978	15978 Transconjugant	<i>E. coli</i> C600
ST	ST167			ST156		
Ampicillin	>128	>128	8	>128	>128	4
Cefotaxime	>128	64	0.03	>128	32	0.125
Ceftazidime	>128	>128	0.125	>128	>128	0.06
Cefoxitin	>128	>128	2	>128	128	4
Florfenicol	16	2	1	16	2	2
Fosfomycin	32	8	2	32	4	2
Streptomycin	256	>128	1	8	>128	>128
Doxycycline	4	1	0.125	128	128	0.5
Ciprofloxacin	>64	0.008	0.002	>64	0.125	0.008
Imipenem	4	2	0.125	1	1	0.125
Resistance genes	<i>bla</i> _{NDM-7} , <i>bla</i> _{CTX-M-15} , <i>aph</i> (3'')-Ib, <i>aph</i> (6)-Id, <i>qnrS1</i> , <i>sulI</i>	<i>bla</i> _{NDM-7}	-	<i>bla</i> _{NDM-5} , <i>bla</i> _{TEM-1b} , <i>aadA2</i> , <i>qepA</i> , <i>mph</i> (A), <i>mdf</i> (A), <i>sulI</i> , <i>tet</i> (B), <i>dfrA12</i>	<i>bla</i> _{NDM-5} , <i>bla</i> _{TEM-1b} , <i>aadA2</i> , <i>qepA</i> , <i>mph</i> (A), <i>sulI</i> , <i>tet</i> (B), <i>dfrA12</i>	-
Chromosomal point mutations	GyrA: S83L, D87N ParE: S458A ParC: S80I			GyrA: S83L, D87N ParC: S80I, E84G		
Virulence genes	<i>capU</i> , <i>iss</i>			<i>gad</i> , <i>iss</i> , <i>lpfA</i>		

Note: All isolates were susceptible to colistin, amikacin, gentamycin, and tigecycline.

FDAARGOS_448 plasmid unnamed1 (CP023959.1), with only 3-bp and 5-bp nucleotide differences, respectively (Figure 1B). *E. coli* AR_452, with an unknown geographic origin, was from human and retained in the CDC & FDA Antibiotic Resistance (AR) Isolate Bank (<https://www.cdc.gov/drugresistance/resistance-bank/>). FDAARGOS_448 was isolated from a patient in Canada in August 2014 and was stored in Database for Reference Grade Microbial Sequences (FDA-ARGOS) database (<https://www.fda.gov/MedicalDevices/ScienceandResearch/DatabaseforReferenceGradeMicrobialSequences/default.htm>). Of note, *E. coli* AR_452 was also assigned to ST156 and FDAARGOS_448 belonged to ST405. It seemed that *bla*_{NDM-5} gene might be circulating among human and food by *E. coli* ST156 clones or pHN15978-like plasmids in different regions. Unlike epidemic IncX3 plasmids, F36:A-:B32 plasmid is less related to the spread of *bla*_{NDM} and this is the first time to report *bla*_{NDM}-positive F36:A-:B32 plasmid. Thus, the identification of almost identical F36:A-:B32 plasmids carrying *bla*_{NDM-5} in

geographically far away countries, Pakistan and Canada, is surprising. Though there is no clear epidemiological link between *E. coli* 15978, *E. coli* strain AR_452, and FDAARGOS_448, poultry trade between Pakistan and Canada might partly explain these findings considering the fact that Pakistan poultry industry was built with the help of Canada based company shaver poultry breeding farms in 1962. In addition, international travel and migratory birds might also be responsible for the global dissemination of this F36:A-:B32 plasmid.^{23,24}

The multidrug resistance region of pHN15978 was mainly composed of three mobile modules (Figure 1C). The first part harbored β -lactam and macrolide resistance genes. More specifically, it consisted of a derivative of Δ Tn2 (*bla*_{TEM-1}) and an IS26-*mph*(A)-*mrx*-*mphR*(A)-IS6100 unit. The resistance region was also identified in *E. coli* plasmid pCARB35_02 (CP031655.1, dog, UK) and *K. pneumoniae* plasmid pCRKP-1215_2 (CP024840.1, human, Korea). In the second part, *bla*_{NDM-5} gene was found embedded in an ISCR1 complex




Figure 1 Comparisons of *bla*_{NDM}-positive plasmids and multidrug resistance region of pHN15978. **(A)** comparisons of *bla*_{NDM}-positive IncX3 plasmids. The innermost ring was the reference plasmid pHN4109c. The light blue, dark blue, yellow, green, grey and pink ring represents pKW53T-NDM (KX214669) from *E. coli* strain from human in Kuwait, pNDM_MGR194 (KF220657.1) from *K. pneumoniae* strain from human in India, pNDM5_IncX3 (KU761328) from *K. pneumoniae* strain from human in China, pOM26-1 (KP776609.1) from *E. coli* strain from human in Oman, p977-NDM (MG825382) from *E. coli* strain from pork in China, and tig00000260 (CP021738) from *E. coli* strain from USA, respectively. **(B)** comparisons of *bla*_{NDM-5}-positive F36:A-B32 plasmids. The innermost ring was the reference plasmid pHN15978 (128762bp, CP030329) and *E. coli* strain FDAARGOS_448 plasmid unnamed1 (128761bp, CP023959) from Canada (human urine, 2014.8), respectively. **(C)** Comparisons of multidrug resistance region of pHN15978. Regions of $\geq 99.0\%$ nucleotide sequence identity are shaded grayish blue.

class 1 integron, which was sequentially organized as IS26- Δ ISAb125-*bla*_{NDM-5}-*ble*_{MBL}-*trpF*-*tat*-ISCR1-*qacE* Δ 1-*sul1*-*aad2*-*hp*-*dfrA12*-*dintI1*, which was identical with the *E. coli* plasmid pM309-NDM5 (F36:A4:B- or F36:A20:B-, AP018833.1), pNDM-d2e9 (F2:A-:B-, CP026201.1), and pAMA1167-NDM-5 (F1:A1:B49, CP024805.1) from Myanmar, USA, and Denmark, respectively.^{21,25} The third part only contained one resistance gene, *qepA*. This genetic structure was sequentially organized as *groEL*/ Δ *intI1*-ISCR3-*qepA*- Δ *intI1*-IS26- Δ Tn2- Δ IS1, and was highly similar to that of pHN3A11 (JX997935.2, *E. coli*, cat, China),²⁶ pMH16-367M_1 DNA (AP018565.1, *Morganella morganii*, human, Vietnam), and pJJ1887-5 (CP014320.1, *E. coli*, human, USA).²⁷

Conclusion

In summary, we firstly characterized two *bla*_{NDM}-carrying plasmids from chicken meat samples in Pakistan. The

dissemination of almost identical *bla*_{NDM-5}-bearing F36:A-B32 plasmids and *bla*_{NDM-7}-bearing IncX3 plasmids in different countries highlights the importance of international trade and travel in the spread of antimicrobial resistance strains and plasmids worldwide.

Accession Number

The complete nucleotide sequence of plasmid pHN4109c and pHN15978 have been deposited in GenBank under accession no. MK088485 and MK291500, respectively.

Acknowledgments

The study was supported in part by grants from the National Natural Science Foundation of China (No. 31625026 and 81661138002).

Disclosure

The author reports no conflicts of interest in this work.

References

- Rodríguez-Baño J, Gutiérrez-Gutiérrez B, Machuca I, et al. Treatment of infections caused by extended-spectrum-beta-lactamase-, AmpC-, and carbapenemase-producing *Enterobacteriaceae*. *Clin Microbiol Rev*. 2018;31:e00079–e00017.
- Madec JY, Haenni M, Nordmann P, et al. Extended-spectrum beta-lactamase/AmpC- and carbapenemase-producing *Enterobacteriaceae* in animals: a threat for humans? *Clin Microbiol Infect*. 2017;23:826–833. doi:10.1016/j.cmi.2017.01.013
- Logan LK, Weinstein RA. The epidemiology of carbapenem-resistant *Enterobacteriaceae*: the impact and evolution of a global menace. *J Infect Dis*. 2017;215(Suppl 1):S28–S36. doi:10.1093/infdis/jiw282
- Yong D, Toleman MA, Giske CG, et al. Characterization of a new metallo-beta-lactamase gene, *bla_{NDM-1}*, and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother*. 2009;53(12):5046–5054. doi:10.1128/AAC.00774-09
- Johnson AP, Woodford N. Global spread of antibiotic resistance: the example of New Delhi metallo-beta-lactamase (NDM)-mediated carbapenem resistance. *J Med Microbiol*. 2013;62(4):499–513. doi:10.1099/jmm.0.052555-0
- Wu W, Feng Y, Tang G, et al. NDM metallo-beta-lactamases and their bacterial producers in health care settings. *Clin Microbiol Rev*. 2019;32(2):pii: e00115-18.
- Lv L, Zeng Z, Song Q, et al. Emergence of XDR *Escherichia coli* carrying both *bla_{NDM}* and *mcr-1* genes in chickens at slaughter and the characterization of two novel *bla_{NDM}*-bearing plasmids. *J Antimicrob Chemother*. 2018;73(8):2261–2263. doi:10.1093/jac/dky176
- Köck R, Daniels-Haardt I, Becker K, et al. Carbapenem-resistant *Enterobacteriaceae* in wildlife, food-producing, and companion animals: a systematic review. *Clin Microbiol Infect*. 2018;24(12):1241–1250. doi:10.1016/j.cmi.2018.04.004
- Zhang Q, Lv L, Huang X, et al. Rapid increase in carbapenemase-producing *Enterobacteriaceae* in retail meat driven by the spread of the *bla_{NDM-5}*-Carrying IncX3 plasmid in China from 2016 to 2018. *Antimicrob Agents Chemother*. 2019;63(8):e00573–e00619. doi:10.1128/AAC.00573-19
- Younas M, ur Rahman S, Shams S, et al. Multidrug resistant carbapenemase-producing *Escherichia coli* from chicken meat reveals diversity and co-existence of carbapenemase encoding genes. *Pak Vet J*. 2019;39(2):241–245.
- Clinical and Laboratory Standards Institute. *Performance Standards for Antimicrobial Susceptibility Testing. Twenty-Seventh Informational Supplement M100-S27*. Wayne, PA, USA: CLSI; 2017.
- Poirel L, Walsh TR, Cuvillier V, et al. Multiplex PCR assays for detection of acquired carbapenemase genes. *Diagn Microbiol Infect Dis*. 2011;70(1):119–123. doi:10.1016/j.diagmicrobio.2010.12.002
- Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome assemblies from short and long sequencing reads. *PLoS Comput Biol*. 2017;13(6):e1005595. doi:10.1371/journal.pcbi.1005595
- Alikhan NF, Petty NK, Zakour NLB, Beatson SA. BLAST Ring Image Generator (BRIG): simple prokaryote genome comparisons. *BMC Genomics*. 2011;12(1):402. doi:10.1186/1471-2164-12-402
- Ranjan A, Shaik S, Mondal A, et al. Molecular epidemiology and genome dynamics of New Delhi metallo-beta-lactamase-producing extraintestinal pathogenic *Escherichia coli* strains from India. *Antimicrob Agents Chemother*. 2016;60(11):6795–6805. doi:10.1128/AAC.01345-16
- Zhang R, Liu L, Zhou H, et al. Nationwide surveillance of clinical carbapenem-resistant *Enterobacteriaceae* (CRE) strains in China. *EBioMedicine*. 2017;19:98–106. doi:10.1016/j.ebiom.2017.04.032
- Wang Y, Zhang R, Li J, et al. Comprehensive resistome analysis reveals the prevalence of NDM and MCR-1 in Chinese poultry production. *Nat Microbiol*. 2017;2:16260. doi:10.1038/nmicrobiol.2016.260
- Tang B, Chang J, Cao L, et al. Characterization of an NDM-5 carbapenemase-producing *Escherichia coli* ST156 isolate from a poultry farm in Zhejiang, China. *BMC Microbiol*. 2019;19(1):82. doi:10.1186/s12866-019-1454-2
- Yao X, Doi Y, Zeng L, Lv L, Liu JH. Carbapenem-resistant and colistin-resistant *Escherichia coli* co-producing NDM-9 and MCR-1. *Lancet Infect Dis*. 2016;16(3):288–289. doi:10.1016/S1473-3099(16)00057-8
- Yoon EJ, Kang DY, Yang JW, et al. New Delhi metallo-beta-lactamase-producing *Enterobacteriaceae* in South Korea between 2010 and 2015. *Front Microbiol*. 2018;9:571. doi:10.3389/fmicb.2018.00571
- Sugawara Y, Akeda Y, Hagiya H, et al. Spreading patterns of NDM-producing *Enterobacteriaceae* in clinical and environmental settings in Yangon, Myanmar. *Antimicrob Agents Chemother*. 2019;63(3):e01924–18. doi:10.1128/AAC.01924-18
- Wang Y, Tong MK, Chow KH, et al. Occurrence of highly conjugative IncX3 epidemic plasmid carrying *bla_{NDM}* in *Enterobacteriaceae* isolates in geographically widespread areas. *Front Microbiol*. 2018;9:2272. doi:10.3389/fmicb.2018.02272
- Schwartz KL, Morris SK. Travel and the spread of drug-resistant bacteria. *Curr Infect Dis Rep*. 2018;20(9):29. doi:10.1007/s11908-018-0634-9
- Wang J, Ma ZB, Zeng ZL, Yang XW, Huang Y, Liu JH. The role of wildlife (wild birds) in the global transmission of antimicrobial resistance genes. *Dongwuxue Yanjiu*. 2017;38(2):55. doi:10.24272/j.issn.2095-8137.2017.024
- Overballe-Petersen S, Roer L, Ng K, et al. Complete nucleotide sequence of an *Escherichia coli* sequence type 410 strain carrying *bla_{NDM-5}* on an IncF multidrug resistance plasmid and *bla_{OXA-181}* on an IncX3 plasmid. *Genome Announc*. 2018;6(5):e01542–17. doi:10.1128/genomeA.01542-17
- Chen X, He L, Li Y, et al. Complete sequence of a F2: A-: B- plasmid pHN3A11 carrying *rmtB* and *qepA*, and its dissemination in China. *Vet Microbiol*. 2014;174(1–2):267–271. doi:10.1016/j.vetmic.2014.08.023
- Johnson TJ, Aziz M, Liu CM, et al. Complete genome sequence of a CTX-M-15-producing *Escherichia coli* strain from the H30Rx subclone of sequence type 131 from a patient with recurrent urinary tract infections, closely related to a lethal urosepsis isolate from the patient's sister. *Genome Announc*. 2016;4(3):e00334–16. doi:10.1128/genomeA.00334-16

Infection and Drug Resistance

Publish your work in this journal

Infection and Drug Resistance is an international, peer-reviewed open-access journal that focuses on the optimal treatment of infection (bacterial, fungal and viral) and the development and institution of preventive strategies to minimize the development and spread of resistance. The journal is specifically concerned with the epidemiology of

antibiotic resistance and the mechanisms of resistance development and diffusion in both hospitals and the community. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use. Visit <http://www.dovepress.com/testimonials.php> to read real quotes from published authors.

Submit your manuscript here: <https://www.dovepress.com/infection-and-drug-resistance-journal>

Dovepress