

Perceptual learning with hand – eye coordination as an effective tool for managing amblyopia: A prospective study

Shreya Shah, Mehul Shah, Anand Purohit, Vaibhav Khanguda

Purpose: Amblyopia is a serious condition resulting in monocular impairment of vision. Although traditional treatment improves vision, we attempted to explore the results of perceptual learning in this study. **Methods:** This prospective cohort study included all patients with amblyopia who were subjected to perceptual learning. The presenting data on vision, stereopsis and contrast sensitivity were documented in a pretested online format, and the pre- and post-treatment information was compared using descriptive, cross-tabulation and comparative methods on SPSS 2.2. The mean values were obtained, and $P < 0.05$ was considered statistically significant. **Results:** The cohort consisted of 47 patients (23 females and 24 males) with a mean age of 14.11 ± 7.13 years. A statistically significant improvement was detected in visual acuity after the perceptual learning session, and the median follow-up period was 17 days. Also, significant improvements were observed in stereopsis but not in the visual outcomes among the age groups. **Conclusion:** Perceptual learning with hand-eye coordination is an effective method for managing amblyopia. This approach can improve vision in all age groups. However, visual improvement is significantly influenced by ocular alignment.

Key words: Amblyopia, amblyopia management in elderly, perceptual learning, visual recovery

Amblyopia is a cortical development disorder secondary to abnormal visual inputs in each eye early in life (during the cortical plasticity stage), wherein dissimilar action potentials (in amplitude or time or both) generated in the retina reach the cortex. These cortical changes entice the visual cortex to prefer one eye over the other, leading to several functional deficiencies in the eye. These include altered visual functions, such as decreased vernier acuity, impaired contrast sensitivity, especially in detecting high-spatial frequency stimuli, and impaired motor signs, such as hand-eye coordination and spatial localization. These defects can be uni- or bilateral.^[1,2] Amblyopia is the most common cause of monocular visual impairment that affects 2%–5% of the general population.

The population-based studies revealed that the prevalence of the disease in 6–71-month-old children was 0.73%–1.9%,^[2] whereas school-based studies on older (> 5 years) children reported higher rates (1.0%–5.5%)^[2] depending on the population studied and the definition used.^[2–14] Bilateral amblyopia is less frequent than unilateral disease; however, the reported proportion varies considerably from 5% to 41% of all cases of amblyopia^[4,11–16] and is 2.7–18 times greater in the presence of strabismus.^[12,15,17–19]

The risk factors for amblyopia are premature birth, small for gestational age,^[20–24] developmental delay^[24] and having a first-degree relative with amblyopia.^[25,26] Some studies have reported that environmental factors, including maternal substance abuse during pregnancy, are associated with an increased risk of amblyopia or strabismus,^[27–32] while other studies have refuted this claim.^[12,15,33]

Drashti Netralaya, Dahod, Gujarat, India

Correspondence to: Dr. Mehul Shah, Drashti Netralaya, Chakalia Road, Dahod, Gujarat, India. E-mail: omtrustdahod@gmail.com

Received: 04-Apr-2022

Revision: 05-May-2022

Accepted: 05-Jul-2022

Published: 30-Sep-2022

Access this article online

Website:

www.ijo.in

DOI:

10.4103/ijo.IJO_852_22

Quick Response Code:



Treatment options for amblyopia in children are optical correction of significant refractive errors, patching, pharmacological treatment, refractive surgery and alternative therapies.

These findings strongly support the need to combine a pharmacological approach with personalized behavioral training aimed at targeting plasticity within specific brain regions or cortical circuits. However, additional studies are essential to further elucidate these approaches.

Polat *et al.*^[34] suggested that perceptual learning can augment the visual function in adult amblyopes. Also, pretest to posttest performance and gains in visual acuity (VA) improved in a series of 77 adult amblyopes in a learning trial of Gabor signals. The neural basis for this observation has been postulated to be lateral inhibition within the brain as a result of training.^[1] However, the gains in the test outcome in the amblyopic eye could not be transferred to novel situations, and improvement was only observed in the task practiced.

Saccades are short, with rapid eye movements lasting for 30–80 milliseconds working through an internal feedback loop based on the efferent motor commands sent to the ocular motor neurons. Sensory-motor integration occurs in the posterior parietal cortex (PPC), also referred to as an associative cortical

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

Cite this article as: Shah S, Shah M, Purohit A, Khanguda V. Perceptual learning with hand – eye coordination as an effective tool for managing amblyopia: A prospective study. Indian J Ophthalmol 2022;70:3630-3.

region, during the generation of the saccade. The lateral occipital vision-mediated cortex activation is present only during the targeted saccadic condition, while internal cognitive saccades are not associated with a lateral occipital cortical activity. The retina between the fovea and the stimulated peripheral retinal point is suppressed during a saccade. The fovea is alerted about the peripheral retinal stimulus a few milliseconds before the onset of the saccade. The top-down impulses originate in the prefrontal cortex and alter the attention mechanism, which in turn affects the striate cortex.

Overt attention is defined as the moment when the peripheral stimulus draws the attention of the eye and ensures foveal fixation. On the other hand, covert attention is when the patient is fixating on the central light and still can see the peripheral lights. Both covert and overt attention stimulate the lateral occipital cortex via the top-down impulses. The above motor and sensory aspects of vision are the principles used for developing the Orthoptek Magnocellular Stimulator (OMS), an advanced treatment modality for amblyopia. The stimulation of magnocellular cells also activates the fovea a few milliseconds before the saccadic movements begin and last after the completion of the saccade. The further away from the fovea the image is located on the retina, the greater the eye movement to bring the peripheral retinal image onto the fovea and the greater is the retinomotor value of the receptor. Hence, the fovea has zero retinomotor value,^[1] which is a crucial principle for instrument design [Fig. 1].

Perceptual learning is yet to gain widespread support. Most studies were conducted on only a small number of participants. The effects of perceptual learning have been demonstrated to last from hours to months without continued practice, albeit long-term follow-up is lacking. Additionally, binocular/dichoptic therapy and a comparison of various treatment modalities are essential. Currently, no studies have compared the conventional modalities of amblyopia management, such as patching and penalization, with the newer modalities, such as dichoptic therapy and liquid crystal glasses.



Figure 1: Electronic panel for hand eye coordination with laser pointer

Methods

This prospective cohort study was approved by the Ethics Committee of DxxxHospital (xx 21/22). All patients in the ocular motility clinic diagnosed with amblyopia of any cause (strabismic and anisometropic) were enrolled in this study. All eyes with organic lesions were excluded. All enrolled patients underwent complete cycloplegic refraction and anterior and posterior segment examination. Also, distance as well near stereopsis, contrast sensitivity and handwriting were examined, after which they were subjected to hand-eye coordination training for a designated period (20 sessions of 30 minutes each). All patients were followed up at the end of one month, and the information was documented in a pretested online format before and after the treatment.

Pre- and post-treatment VA, distance and near stereopsis, angle of deviation and contrast sensitivity were compared using the Statistical Package for the Social Sciences (SPSS) version 22, descriptive analysis, cross-tabulation and analysis of variance.

Results

This method was applied to 47 patients (23 [48.9%] women and 24 [51.1%] men) with a mean age of 14.11 ± 7.13 23 years [Table 1].

The mean follow-up period was 17 days.

Prior to the procedure, 28 (59.6%) patients were orthophoric, whereas after the procedure, 30 (63.8%) had orthophoria. A significant difference was detected for distance and near in the deviation by Prism bar cover test [Table 2].

Nonetheless, no significant difference was observed between the two age groups for distance as well as near ($P=0.449, 0.271$). Visual improvement was also observed in the older age group.

Table 1: Age and sex distribution

	Sex		Total
	F	M	
6-10	4	13	17
>10	19	11	30
Total	23	24	47

Table 2: Comparative study of pre- and post-study values

Variable	P
Pre post vision OD [#]	0.000
Pre post vision OS [§]	0.000
Pre post stereopsis for near	0.000
Pre post stereopsis distance	0.001
Strabismus	0.001
Type of strabismus	0.503
PBCT* OD [#] vs post procedure	0.007
PBCT* OS [§] vs post procedure	0.016
PBCT* near deviation OD [#] vs post procedure	0.045
PBCT* near deviation OS [§] vs post procedure	0.028
Contrast sensitivity	0.000

*PBCT, Prism bar cover test. [#]OD, Right eye. [§]OS, Left eye

Interestingly, strabismic amblyopia was detected in 48.9% of patients, and refractive amblyopia was noted in 46.8% of patients. However, no significant difference was observed in the visual outcome ($P = 0.178$) following training, but the pre- and post-treatment vision, near and distance stereopsis and contrast sensitivity differed significantly in the two subgroups [Tables 2 and 3].

Discussion

No significant difference was observed among the various age groups. Marianne *et al.* reported several changes in the perspective of the recovery period for amblyopia. Previous studies and clinical practice demonstrated that conventional treatment improved VA in the amblyopic eye in children aged >7 years. These media-described cases within the community eye care domain suggested that the literature on amblyopia treatment in older children (aged >7 years) should be screened to address the misconceptions and focus on the current issues faced by clinicians while treating newly diagnosed amblyopia in this age group.^[34,35]

Notably, VA improved significantly after the treatment [Table 2]. Grant *et al.*^[36] reported that a combined therapy of perceptual learning-based visual training and patching was effective in improving VA in children with amblyopia who did not regain their vision with patching alone or had poor patching compliance. This preliminary outcome should be confirmed in future clinical trials.

Similar findings have been reported by Bonaccorsi *et al.*^[37] on the impact of visual perceptual learning on amblyopia, with a focus on a new experimental model of perceptual learning in amblyopic rats.

Non-human animal models have demonstrated that selective serotonin re-uptake inhibitors (SSRIs) enhance the plasticity within the mature visual cortex and enable recovery from amblyopia.^[38,39]

The strategies employed by children with amblyopia and abnormal binocularity to attain precision grasping alter with age. The visual feedback during the “inflight” approach was pronounced during the age of 5–6 years, which increased the reliance on tactile/kinaesthetic feedback from object contact at an age of 7–9 years. However, regaining binocularity improved the hand–eye coordination speed and accuracy in an age-dependent manner, which is a better predictor of these fundamental performance measures than the degree of VA loss.^[36]

Antonio *et al.* Rodan A *et al.* reported that several treatments differing in design (for example, type of stimulus, context

used and duration of the training) and with a wide age group, including adults, have been applied. Most of the studies showed an improvement in some monocular and binocular visual functions, such as VA, contrast sensitivity and stereopsis. Currently, it is recommended to use these processes as an alternative or as a complement to the traditional passive therapy.^[40]

The strength of our study is its prospective nature, while the small sample size and the short follow-up period are some limitations. Thus, we may try a combination of therapies, such as patching and perceptual learning or randomized, controlled trial for alternative therapies.^[40]

Conclusion

Perceptual learning with hand–eye coordination is an effective method for managing amblyopia in older children. The technique is utilised, irrespective of age, and the visual outcome has improved in patients with orthophoria.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

References

- Gopal SKS, Kelkar J, Kelkar A, Pandit A. Simplified updates on the pathophysiology and Gopal recent developments in the treatment of amblyopia: A review. *Indian J Ophthalmol* 2019;67:1392-9.
- Martin PR, Lee BB, White AJ, Solomon SG, Rüttiger L. Chromatic Sensitivity of ganglion cells in the peripheral primate retina. *Nature* 2001;410:933-6.
- Snowdon S, Stewart-Brown S. Effectiveness of preschool vision Screening. *Nurs Stand* 1997;12:44-5.
- Pediatric Eye Disease Investigator Group. A randomized trial of near versus distance activities while patching for amblyopia in Children aged 3 to less than 7 years. *Ophthalmology* 2008;115:2071-8.
- Pediatric Eye Disease Investigator Group Writing Committee. A randomized trial comparing Bangerter filters and patching for the treatment of moderate amblyopia in children. *Ophthalmology* 2010;117:998-1004.e6.
- Bhola R, Keech RV, Kutschke P, Pfeifer W, Scott WE. Recurrence of amblyopia after occlusion therapy. *Ophthalmology* 2006; 113:2097-100.
- Tacagni DJ, Stewart CE, Moseley MJ, Fielder AR. Factors affecting the stability of visual function following cessation of occlusion therapy for amblyopia. *Graefes Arch Clin Exp Ophthalmol* 2007;245:811-6.
- Walsh LA, Hahn EK, LaRoche GR. The method of treatment cessation and recurrence rate of amblyopia. *Strabismus* 2009;17:107-16.
- Scheiman MM, Hertle RW, Beck RW, Edwards AR, Birch E, Cotter SA, *et al.* Randomized trial of treatment of amblyopia in children aged 7 to 17 years. *Arch Ophthalmol* 2005;123:437-47.
- Repka MX, Kraker RT, Beck RW, Birch E, Cotter SA, Holmes JM, *et al.* Treatment of severe amblyopia with weekend atropine: Results from 2 randomized clinical trials. *J AAPOS* 2009;13:258-63.
- Scheiman MM, Hertle RW, Kraker RT, Beck RW, Birch EE, Felius J, *et al.* Patching vs atropine to treat amblyopia in children aged 7 to 12 years: A randomized trial. *Arch Ophthalmol* 2008;126:1634-42.
- Pediatric Eye Disease Investigator Group. A randomized trial of

Table 3: Mean of variables

Variable	Mean	St deviation
Pre distance stereopsis	474.38	426.21
Pre near stereopsis	347.89	367.27
Post distance stereopsis	343.83	256.92
Post near stereopsis	299.56	256.21
Pre contrast	0.82	0.126
Post contrast	0.80	0.000

- Atropine versus patching for treatment of moderate amblyopia: Follow-up at 10 years of age. *Arch Ophthalmol* 2008;126:1039-44.
13. Medghalchi AR, Dalili S. A randomized trial of atropine vs patching for treatment of moderate amblyopia. *Iran Red Crescent Med J* 2011;13:578-81.
 14. Menon V, Shailesh G, Sharma P, Saxena R. Clinical trial of patching Versus atropine penalization for the treatment of anisometropic Amblyopia in older children. *J AAPOS* 2008;12:493-7.
 15. Feliuss J, Chandler DL, Holmes JM, Chu RH, Cole SR, Hill M, *et al.* Evaluating the burden of amblyopia treatment from the parent And child's perspective. *J AAPOS* 2010;14:389-95.
 16. Holmes JM, Pediatric Eye Disease Investigator Group. A randomized Pilot study of near activities versus non-near activities during patching therapy for amblyopia. *J AAPOS* 2005;9:129-36.
 17. Alotaibi AG, Fawazi SM, Alenazy BR, Abu-Amro KK. Outcomes of 3 hours part-time occlusion treatment combined with near activities among children with unilateral amblyopia. *Saudi Med J* 2012;33:395-8.
 18. Moseley MJ, Neufeld M, McCarry B, Charnock A, McNamara R, Rice T, *et al.* Remediation of refractive amblyopia by optical correction alone. *Ophthalmic Physiol Opt* 2002;22:296-9.
 19. Cleary M, Moody AD, Buchanan A, Stewart H, Dutton GN. Assessment of a computer-based treatment for older amblyopes: The Glasgow pilot study. *Eye (Lond)* 2009;23:124-31.
 20. Li RW, Ngo C, Nguyen J, Levi DM. Video-game play induces plasticity in the visual system of adults with amblyopia. *PLoS Biol* 2011;9:e1001135.
 21. Herbison N, Cobb S, Gregson R, Ash I, Eastgate R, Purdy J, *et al.* Interactive binocular treatment (I-BiT) for amblyopia: Results of a pilot study of 3D shutter glasses system. *Eye (Lond)* 2013; 27:1077-83.
 22. Li J, Thompson B, Deng D, Chan LY, Yu M, Hess RF, *et al.* Dichoptic training enables the adult amblyopic brain to learn. *Curr Biol* 2013; 23:308-9.
 23. Spiegel DP, Li J, Hess RF, Byblow WD, Deng D, Yu M, *et al.* Transcranial direct current stimulation enhances recovery of stereopsis in adults with amblyopia. *Neurotherapeutics* 2013;10:831-9.
 24. Leguire LE, Rogers GL, Walson PD, Bremer DL, McGregor ML. Occlusion and levodopa-carbidopa treatment for childhood amblyopia. *J AAPOS* 1998;2:257-64.
 25. Mohan K, Dhankar V, Sharma A. Visual acuities after levodopa administration in amblyopia. *J Pediatr Ophthalmol Strabismus* 2001;38:62-7.
 26. Repka MX, Kraker RT, Beck RW, Atkinson CS, Bacal DA, Bremer DL, *et al.* Pilot study of levodopa dose as treatment for residual amblyopia in children aged 8 years to younger than 18 years. *Arch Ophthalmol* 2010;128:1215-7.
 27. Dadeya S, Vats P, Malik KP. Levodopa/carbidopa in the treatment of amblyopia. *J AAPOS* 2009;46:87-90.
 28. Morishita H, Miwa JM, Heintz N, Hensch TK. Lynx1, a Cholinergic brake, limits plasticity in adult visual cortex. *Science* 2010;330:1238-40.
 29. Zhao J, Lam DS, Chen LJ, Wang Y, Zheng C, Lin Q, *et al.* Randomized controlled trial of patching vs acupuncture for anisometropic amblyopia in children aged 7 to 12 years. *Arch Ophthalmol* 2010;128:1510-7.
 30. Lam DS, Zhao J, Chen LJ, Wang Y, Zheng C, Lin Q, *et al.* Adjunctive Effect of acupuncture to refractive correction on anisometropic amblyopia: One-year results of a randomized crossover trial. *Ophthalmology* 2011;118:1501-11.
 31. Althomali TA. Posterior chamber toricphakic IOL implantation for the management of pediatric anisometropic amblyopia. *J Refract Surg* 2013;29:396-400.
 32. Paysse EA, Coats DK, Hussein MA, Hamill MB, Koch DD. Long-term outcomes of photorefractive keratectomy for anisometropic amblyopia in children. *Ophthalmology* 2006;113:169-76.
 33. Joslin CE, McMahon TT, Kaufman LM. The effectiveness of occluder contact lenses in improving occlusion compliance in Patients that have failed traditional occlusion therapy. *Optom Vis Sci* 2002;79:376-80.
 34. Pradeep A, Proudlock FA, Awan M, Bush G, Collier J, Gottlob I, *et al.* An educational intervention to improve adherence to high-dosage patching regimen for amblyopia: A randomised controlled trial. *Br J Ophthalmol* 2014;98:865-70.
 35. Piano MEF, Simmers AJ. 'It's too late'. Is it really? Considerations for amblyopia treatment in older children. *Ther Adv Ophthalmol* 2019;11:2515841419857379.
 36. Grant S, Suttle C, Melmoth DR, Conway ML, Sloper JJ. Age- and stereovision-dependent eye-hand coordination deficits in children with amblyopia and abnormal binocularity. *Invest Ophthalmol Vis Sci* 2014;55:5687-7015.
 37. Bonaccorsi J, Berardi N, Sale A. Treatment of amblyopia in the adult: Insights from a new rodent model of visual perceptual learning. *Front Neural Circuits* 2014;8:82.
 38. Hernández-Rodríguez CJ, Fukumitsu H, Ruiz-Fortes P, Soto-Negro R, Merino-Suárez M, Piñero DP. Efficacy of perceptual learning-based vision training as an adjuvant to occlusion therapy in the management of amblyopia: A pilot study. *Vision (Basel)* 2021;5:15.
 39. Lagas AK, Black JM, Russell BR, Kydd RR, Thompson B. The effect of combined patching and citalopram on visual acuity in adults with amblyopia: A randomized, crossover, placebo-controlled trial. *Neural Plast* 2019;2019:5857243.
 40. Rodán A, Candela Marroquín E, Jara García LC. An updated review about perceptual learning as a treatment for amblyopia. *J Optom* 2022;15:3-34.