

In conclusion, we found that *KIF20A* was an independent predictor for ccRCC. *KIF20A* expression was upregulated in database and clinical cases. Cell experiments found that suppression of *KIF20A* could inhibit the malignant characteristics of renal cancer cells. These results demonstrated that *KIF20A* could be a potential novel prognostic molecule and may become a treatment target for ccRCC. However, a large number of samples are still needed to verify its clinical value in the future.

Declaration of patient consent

The authors certify that they have obtained all appropriate patient consent forms. In the form, the patient(s)/or his/her guardian has/have given his/her/their consent for his/her/their images and other clinical information to be reported in the journal. The patients or his/her guardian understand that his/her/their name(s) and initials will not be published and due efforts will be made to conceal his/her/their identity, but anonymity cannot be guaranteed.

Funding

This project was supported by grants from the National Natural Science Foundation of China (No. 81902588) and the National Key Scientific Instrument Development Project (No. 81927807).

Conflicts of interest

None.

References

1. Xiao W, Xiong Z, Xiong W, Yuan C, Xiao H, Ruan H, *et al*. Melatonin/PGC1A/UCP1 promotes tumor slimming and represses tumor progression by initiating autophagy and lipid browning. *J Pineal Res* 2019;67:e12607. doi: 10.1111/jpi.12607.
2. Lai Y, Tang F, Huang Y, He C, Chen C, Zhao J, *et al*. The tumour microenvironment and metabolism in renal cell carcinoma targeted or immune therapy. *J Cell Physiol* 2020. doi: 10.1002/jcp.29969.[ahead of print].
3. Wu PF, Gao WW, Sun CL, Ma T, Hao JQ. Suberoylanilide hydroxamic acid overcomes erlotinib-acquired resistance via phosphatase and tensin homolog deleted on chromosome 10-mediated apoptosis in non-small cell lung cancer. *Chin Med J* 2020;133:1304–1311. doi: 10.1097/CM9.0000000000000823.
4. Yuan L, Chen L, Qian K, Qian G, Wu CL, Wang X, *et al*. Co-expression network analysis identified six hub genes in association with progression and prognosis in human clear cell renal cell carcinoma (ccRCC). *Genomics data* 2017;14:132–140. doi: 10.1016/j.gdata.2017.10.006.
5. Wei S, Zheng Y, Jiang Y, Li X, Geng J, Shen Y, *et al*. The circRNA circPTPRA suppresses epithelial-mesenchymal transitioning and metastasis of NSCLC cells by sponging miR-96-5p. *EBioMedicine* 2019;44:182–193. doi: 10.1016/j.ebiom.2019.05.032.

How to cite this article: Xiao W, Chen K, Liang HG, Zhang XP. Identification of *KIF20A* as a tumor biomarker and forwarder of clear cell renal cell carcinoma. *Chin Med J* 2021;134:2137–2139. doi: 10.1097/CM9.0000000000001331

Corrigendum

Corrigendum: Amyloid and tau positive mild cognitive impairment: clinical and biomarker characteristics of dementia progression

DOI: 10.1097/CM9.0000000000001765

In the article “Amyloid and tau positive mild cognitive impairment: clinical and biomarker characteristics of dementia progression”, which appeared in vol.134, issue14, page 1709 of *Chinese Medical Journal*, the following words “for Alzheimer’s disease Neuroimaging Initiative” should be added to the author section. The full authors should be corrected as “Hong-Chun Wei¹, Bing Li¹, Kok Pin Ng², Qing-Xi Fu³, Sheng-Jie Dong⁴, Mao-Wen Ba¹, Min Kong⁵; for Alzheimer’s disease Neuroimaging Initiative”. And in page 1717, the Funding Part should be corrected as “The study was supported by grants from the Shandong Provincial key research and development project (No.2018GSF118235), the Shandong Provincial Natural Science Foundation (No.ZR2016HL16) and Youth Research Start-up Fund of Yantai Yuhuangding Hospital Affiliated to Qingdao University (No.2020-25).”

Reference

1. Wei HC, Li B, Ng KP, Fu QX, Dong SJ, Ba MW, Kong M. Amyloid and tau positive mild cognitive impairment: clinical and biomarker characteristics of dementia progression. *Chin Med J* 2021; 134:1709–1719. doi: 10.1097/CM9.0000000000001496.