

Research

Distinct gene expression profiles of human type 1 and type 2 T helper cells

Heli Hamalainen^{*†}, Hua Zhou^{†‡}, William Chou[‡], Hideki Hashizume[‡], Renu Heller[‡] and Riitta Lahesmaa^{*}

Addresses: ^{*}Turku Centre for Biotechnology, University of Turku and Abo Akademi University, FIN-20521 Turku, Finland. [†]Roche Bioscience, Inflammatory Diseases Unit, 3401 Hillview Avenue, Palo Alto, CA 94304, USA. [‡]These authors contributed equally to this work.

Correspondence: Riitta Lahesmaa. E-mail: riitta.lahesmaa@btk.utu.fi

Published: 21 June 2001

Genome **Biology** 2001, **2**(7):research0022.1-0022.11

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2001/2/7/research/0022>

© 2001 Hamalainen et al., licensee BioMed Central Ltd
(Print ISSN 1465-6906; Online ISSN 1465-6914)

Received: 14 March 2001

Revised: 26 April 2001

Accepted: 15 May 2001

Abstract

Background: The development and activation of CD4⁺ helper T cell (Th) subsets with distinct patterns of unbalanced production of cytokines play an important part in infectious, allergic and autoimmune diseases. Human neonatal cord blood CD4⁺ Th cells can be polarized into type 1 or type 2-like effector cells *in vitro* by culturing them in the presence of interleukin (IL)-12 or IL-4, respectively. We have exploited this experimental system to identify marker genes that are differentially expressed by polarized Th1 and Th2 cells. An oligonucleotide microarray specifically designed to screen for inflammation-related candidate genes was used and the differential expression was further validated with a quantitative real-time RT-PCR method.

Results: In addition to the previously described marker genes of Th cells, we report subtle changes in the expression of several other genes that represent growth factors, receptors and other signaling molecules in polarized Th1 and Th2 cell subsets. Additionally, we describe a novel set of genes as Th1/Th2 differentiation markers for cells activated by anti-CD3 and anti-CD28 antibodies.

Conclusions: This study demonstrates the power of the targeted use of microarrays in combination with quantitative real-time RT-PCR in identifying and validating new marker genes for gene expression studies.

Background

The functionally distinct repertoire of secreted cytokines of type 1 and type 2 CD4⁺ T helper cells (Th1 and Th2 cells) has been shown to play an important role in the pathogenesis of human allergy and inflammatory diseases [1]. Effector Th1 cells secrete predominantly interferon- γ (IFN- γ) and interleukin-2 (IL-2) and regulate cell-mediated immunity against intracellular pathogens, whereas differentiated Th2 cells

produce IL-4 and IL-5 and promote antibody-mediated humoral immune responses [2,3]. In several immunological disorders the balance between the type 1 and type 2 cells is disturbed and favors either a predominant Th1 response (autoimmune diseases) or enhanced Th2 response (allergic inflammation and atopic disorders) [1]. Cytokines IL-12 and IL-4 play a major role in selectively regulating the development and differentiation of CD4⁺ T cell subsets to IFN- γ and

IL-2 producing Th1 or IL-4 and IL-5 secreting Th2, respectively. This differentiation process can be mimicked *in vitro*.

DNA microarray techniques can be used to study gene expression on a large scale [4,5]. The expression of a large number of genes can be monitored simultaneously and the expression profiles in different samples compared. Recent approaches in immunology exploiting this technology include the expression monitoring of changes caused by cytokines [6,7], viruses [8,9], inflammation [10,11] or cancers of hematopoietic origin [12-14]. The profiles of mRNA expression in human type 1 and type 2 Th cells are still largely unknown. An earlier report describing differentiating Th cells using an oligonucleotide array analysis led to the production of a large volume of data [15], but because of the experimental design the gene expression profile of Th2 cells remained obscure. We chose to study the expression of polarized T helper cells and describe the comparative analysis of effector Th1/Th2 cells. As a preliminary screen, a restricted number (250) of inflammation-related genes was included for validation. In a parallel analysis a quantitative reverse transcription polymerase chain reaction (RT-PCR) method (TaqMan) was used to evaluate the results. We show that an oligonucleotide microarray complemented by quantitative RT-PCR is an excellent method for gene expression profiling.

Results

Probe selection and oligonucleotide array design

The Roche PA-1 oligonucleotide array contained 66,176 features that correspond to 517 inflammation-related genes and was fabricated by Affymetrix. Among these, 250 genes were of human origin whereas the others represented mouse and rat genes. Each gene is monitored by 64 pairs of features that consist of 25-mer oligonucleotides, one perfectly complementary to the gene sequence and the other identical except for a single mismatch located in the center of the oligonucleotide.

Specificity, sensitivity and linearity of the detection methods used

The specificity and sensitivity of the current type of oligonucleotide array used are based on several criteria. First, the design of the antisense oligonucleotides was targeted toward a well annotated full-length cDNA sequence (without introns) for each gene. Second, the number of both perfectly and mismatching oligonucleotides (features) for each sequence was 64. When directly converted, the 32 specific oligonucleotides (25-mers) overlap 800 bp of sequence data. Third, for this preliminary screen the amount of mRNA used for *in vitro* transcription reaction was rather high, 1 μ g. In principle this amount should represent even the low abundant messages needed for difference detection [16-18]. Lastly, the hybridization mixture was used only once and discarded. However, from our earlier studies [18] it was known that for abundant messages (such as IFN- γ) or very rare transcripts (IL-4) the linearity of detection in this

type of hybridization-based method is largely compromised (10^2 to 10^3). We therefore have chosen a real-time RT-PCR method that facilitates quantitation of the transcripts over at least five orders of magnitude [19]. To confirm the linearity of the detection for each gene dilution curves were created (see Materials and methods). In addition, on the basis of our earlier work [20], we knew that the differences in the expression for several of the key genes in effector type 1 and type 2 cells would be very small. To minimize individual variation the oligonucleotide array was first hybridized twice with a type 1 or type 2 sample that had been previously validated with known marker genes. A second aliquot of the same mRNA was then independently prepared and hybridized to a new oligonucleotide array manufactured within the same batch. After this, the analyzed data was confirmed in samples from other individuals with a real-time RT-PCR method.

Expression profiles of Th1 and Th2 cells

The expression of a total of 250 human inflammation-related genes was screened with an oligonucleotide array by performing two independent hybridizations with a Th1 or Th2 cell sample. The phenotype of these cells had been determined using ELISA as type 1 or type 2 (secretion of high IFN- γ , low IL-4, or high IL-4, low IFN- γ , respectively, data not shown). The differentially expressed human RNA transcripts of Th1 and Th2 cells are shown in Figures 1 and 2. The level of expression is shown as the average fluorescence intensity value for each gene in each sample. Collectively, 34 transcripts were recorded as differentially expressed in these cells (14 in type 1 and 20 in type 2 cells). The transcript of a proinflammatory cytokine INF- γ is abundant in type 1 T cells and very low in type 2 cells (Figure 1a). IFN- γ mRNA had routinely been measured with real-time RT-PCR from our cell cultures and the actual difference between the type 1 and type 2 cells was in the order of 10^3 - to 10^5 -fold. An oligonucleotide array, however, indicated less than 10^2 -fold difference in type 1 and type 2 cells (78-fold, calculated from average fluorescence intensity values for IFN- γ : Th1 1567.8, SD 129.1 and Th2 < 20, SD 3.3). In clear contrast to IFN- γ are the transcripts of the IL-4/IL-13 cluster known to promote humoral type 2 responses. Despite their all (IL-4, IL-5 and IL-13) being detected as differentially expressed in T-cell samples by an oligonucleotide array (Figure 2b), reliable detection of both IL-4 and IL-5 failed in a customized RT-PCR setup, probably because of their low level of expression.

The other transcripts expressed at a higher level by Th1 than Th2 cells are presented in Figure 1a,b. In addition to IFN- γ , these include IL-8, tumor necrosis factor- α (TNF- α) and granzyme B (fluorescence intensity > 400) as well as granulocyte-macrophage colony-stimulating factor (GM-CSF), macrophage inhibitory protein-1 α (MIP-1 α), MIP-1 β , RANTES, IL-7 receptor (IL-7R), IL-12R β 2, SLAM, CCR1, CCR2, CCR5, I κ B, TIMP-1, c-Jun, IRF-1, caspase-1 and





Figure 1

Genes preferentially expressed in human Th1 cells as measured by oligonucleotide microarrays. Genes expressed at higher level in Th1 cells (black bars) as compared to Th2 cells (white bars) at day 14 are shown. Fluorescence intensity values for (a) genes with increased expression levels of > 400, and (b) genes with expression levels increased by < 400. Error bars indicate variation in two hybridizations from duplicate preparations of the same sample.

clusterin (fluorescence intensity < 400). Transcripts of MIF, IL-4R and STAT4 gave the highest relative fluorescence intensity values (> 300) in Th2 samples and their relative expression is shown as a separate graph (Figure 2a). The fluorescence values for all other genes that were expressed at

higher level in Th2 cells compared to Th1 cells were lower (< 300; Figure 2b). In Th2 cells, preferentially expressed genes include in addition to IL-4, IL-5 and IL-13, the transcripts for IFN- α β R, IFN- γ R β , fibroblast growth factor receptor (FGFR), CCR4, Smad2, BAX, CAS and caspase-6.

**Figure 2**

Genes preferentially expressed in human Th2 cells as measured by microarrays. Genes expressed at a higher level in Th2 cells (black bars) as compared to Th1 cells (white bars) at day 14 are presented. Fluorescence intensity values for **(a)** genes with increased expression levels of > 300, and **(b)** genes with expression levels of < 300. Error bars indicate variation in two hybridizations from duplicate preparations of the same sample.

Gene expression according to real-time RT-PCR

Hybridizations of oligonucleotide arrays were used to screen a panel of genes in T cells that are differentially expressed by polarized Th1 or Th2 cells (Figures 1,2). To validate the observations, the same Th1 and Th2 samples were quantitated together with samples of cultured Th1/Th2 cells originating from other individuals by using a real-time RT-PCR (TaqMan) method. We designed probe and primer sets for the quantitative detection of mRNAs for 14 genes (Table 1; caspase-1, CCR1, CCR2, CCR4, CCR5, clusterin, FGFR, GM-CSF, I κ B, IL-4R, RANTES, STAT4, TIMP-1 and TNF- α). In addition, we had earlier confirmed a differential expression of SLAM, IFN- γ R β and IL-12R β 2 transcripts in these cells [20]. All TaqMan RT-PCR measurements were carried out

for each gene in duplicate. After three individual measurements the fold change in mRNA expression in Th1 cells compared to Th2 cells was calculated (Table 2). The overall transcriptional preferences recorded with a real-time RT-PCR method correspond well with the differences recorded by the oligonucleotide microarrays. The measurements from Th1 or Th2 cultures of other individuals further confirm this. When the calculated fold differences by the different methods in Th1 and Th2 samples are compared they are in most cases reminiscent but not equal. Because of a broader dynamic range of the real-time RT-PCR detection, we consider TaqMan results to be more reliable in this respect. In the case of the chemokine receptors CCR1, CCR2 and CCR3, where the expression level in Th2 cells is low, the

Table 1

TaqMan primers and probes used in this study		
Gene (human)	Accession number	1) 5' - (FAM)-probe-(TAMRA) - 3' 2) 5' - primer 1 - 3' 3) 5' - primer 2 - 3'
Caspase-1	M87507	1) AA GACGTGTGCGGCTTGACTTGTC 2) AATACTGTCAAATTCCTCATTGCAGATAAT 3) AAGTCGGCAGAGATTTATCCAATAA
CCR1	L10918	1) TGGCCATCGTCCACGCCG 2) TCCTGCTGACGATTGACAGGTA 3) GTGCCCGCAAGGCAAAC
CCR2	U03882	1) AGTGCTTCGCAGATGTCCTTGATGCTC 2) GCGTTTAATCACATTCGAGTGTTC 3) CCACTGGCAAATTAGGGAACAA
CCR4	X85740	1) TTA CTCTGCTGACACCCCGCTCATC 2) CAATACTGTGGGCTCCTCCAA 3) ATCCATGGTGGACTGCGTGTA
CCR5	U54994	1) TGCAAAAGGCTGAAGAGCATGACTGACAT 2) GCTGGTCATCCTCATCCTGATAA 3) ATGGCCAGGTTGAGCAGGTA
Clusterin	X14728	1) TGCCAGCCGGGACACCCAG 2) CTTGCGCTTTCGTGAGGT 3) GAGCAGCTGAACGAGCAGTTT
FGFR	X51803	1) CCGTAGCTCCATATTGGACATCCCGAG 2) AGATAACACCAAACCAAACCGTATG 3) GCATGCAATTTCTTTCCATCTT
GM-CSF	M11222	1) CCCCTTTGACTGCTGGGAGCCAG 2) CCTGAAGGACTTTCTGCTTGCTCA 3) CTCATCTGGCCGGTCTCACT
IκB	M69043	1) TTCTAGTGTGACAGTGGCCAGCTGC 2) TCTCTGGCAGCATCTGAAGGT 3) CCCAAGCACCCGGATACAG
IL-4R	X52425	1) AATGCCACTGCCAGGCTGTCC 2) GTGGCAGGTAAGGGCTGAGTAGA 3) CACCTCGCAGTCCGCAGA
RANTES	M21121	1) TTGGCACACACTTGGCGGTTCTTC 2) TCCCGAACCCATTTCTTCTCT 3) CCCAGCAGTCGTCTTTGTCA
STAT4	L78440	1) AGTCTCGCAGGATGTCAGCGAATGG 2) GCTGAGAGCTGTAGTGTTTACCGA 3) AATAAAGGCCGTTGTCTGCT
TIMP-1	X03124	1) CAATTCCGACCTCGTCATCAGGGC 2) CACCCACAGACGGCCTTCT 3) TCTGGTGTCCCCACGAACTT
TNF-α	M10988	1) CATCTTCTCGAACCCGAGTGACAAGC 2) TGGCCCAGGCAGTCAGA 3) GGTTTGCTACAACATGGGCTACA

Abbreviations: FAM, 6-carboxyfluorescein; TAMRA, 6-carboxytetramethylrhodamine.

sensitivity of the method becomes limiting and leads to higher values of standard deviation. The reason for the significantly higher fold difference values recorded by the oligonucleotide arrays for TIMP-1 and IL-4 is currently unknown. Most notably, even when the fold difference for some of the target genes (CCR5, IκB, STAT4) is low

(< twofold) according to an oligonucleotide array analysis, the transcriptional preference in all Th1/Th2 samples is similar and the statistical significance high ($p < 0.01$).

Activation-induced changes in polarized T helper cells

Because cell activation is known to affect the transcription of genes in T cells [21], we next asked whether the cell activation also changes the level of transcription of genes in polarized T helper cells. Real-time RT-PCR measurements of gene expression were carried out with a set of 14 genes (Table 1) on three separate T-cell cultures. Polarized (day 14 cells) CD4⁺ Th1 or Th2 cells were triggered with antibodies for CD3 and CD28 for 2, 6 or 24 hours and gene expression was quantitated (Figure 3). During the time course of activation a general trend of downregulation of transcription was observed in the genes studied. The expression of a house-keeping gene, however, was unaffected (data not shown). The downregulation occurred within 24 hours in both type 1 and type 2 cells. Even the transcripts for GM-CSF, TNF-α and IκB, which were rapidly upregulated in the early activation period were downregulated soon after. Downregulation of expression was also seen for CCR1, CCR2, CCR5, caspase-1 and clusterin. Both T-cell subsets seem to exhibit very similar patterns of changes in the mRNA levels during activation of these genes, with the following exceptions. First, the early induction of GM-CSF and TNF-α expression is more pronounced in Th1 cells than in Th2 cells. Second, the downregulation of transcripts at 24 hours of activation for CCR1, CCR2 and CCR5 in Th2 cells is more effective compared to Th1 cells. The mRNA levels for RANTES, CCR4, FGFR and STAT4 in Th1 or Th2 cells did not respond to an anti-CD3/CD28 mediated activation signal or the observed changes were low (< two- to threefold). In conclusion, genes that responded promptly to the activation signal were GM-CSF, TNF-α, IκB, CCR1, CCR2, CCR5, caspase-1 and clusterin. Most importantly, activation-induced changes in expression of these genes did not affect the preferential expression pattern in Th1 and Th2 cells.

Discussion

We have used high-density oligonucleotide arrays to profile gene expression of cultured polarized human Th1 and Th2 cells. Many research groups, including ours, have intensively searched for genes that would be differentially regulated in human type 1 or type 2 cells and would potentially explain their distinct functions in immunological disorders [22]. Until recently, only a few genes, such as those for cytokines, have been defined as differentially expressed in human CD4⁺ T helper subsets [1,2]. Classically, the definition of a type 1 or type 2 Th cell was based on measurements of a limited number of cytokines secreted by these cells after activation (by phorbol myristoyl acetate and calcium ionophore). Current developments in RNA technologies enable the basal level of gene expression to be studied in these cells. The research has been inspired by the discoveries of differentially




Figure 3 Activation-induced changes of expression of genes in human Th1 and Th2 cells as measured with real-time RT-PCR. **(a-h)** Fold changes of expression of (a) GM-CSF, (b) TNF- α , (c) I κ B, (d) CCR1, (e) CCR2, (f) CCR5, (g) caspase-1 and (h) clusterin in human Th1 cells (black bars) and Th2 cells (white bars). Activation was induced by coligating CD3 and CD28 in human Th1 and Th2 cells for 2, 6 or 24 h. Data are from samples from Th1 and Th2 cell cultures originating from three individuals. Each measurement was carried out in duplicate and repeated.

Table 2**Fold differences in gene expression in human Th1/Th2 cells**

Preference	Gene	Oligonucleotide array	Real-time RT-PCR		SD	deltaC _T (Th2) (average of individuals 1,2,3)	SD	Fold difference [†] (average of individuals 1,2,3)	Statistical significance [§]
		Fold difference by microarray [†] (individual 1) [‡]	Fold difference by real-time RT-PCR [†] (individual 1) [‡]	deltaC _T (Th1) (average of individuals 1,2,3)					
Th1 > Th2	GM-CSF	4.1	3.6	6.73	0.80	8.86	0.83	4.4	**
	TNF- α	12.3	6.2	2.91	0.28	4.40	0.92	2.8	**
	RANTES	3.1	4.0	1.23	0.72	3.88	0.71	6.3	**
	CCR1	5.1	122.0	7.18	0.85	12.73	1.95	46.7	**
	CCR2	8.9	30.7	7.57	0.55	10.95	2.10	10.4	**
	CCR5	1.9	109.0	10.64	0.46	15.08	1.92	21.7	**
	I κ B	1.7	1.4	1.76	0.65	2.92	0.55	2.2	**
	TIMP-1	6.6	1.2	3.39	0.48	3.48	0.80	1.1	n.s.
	Caspase-1	2.9	1.9	8.08	0.72	8.87	0.75	1.7	**
	Clusterin	2.4	1.4	3.50	0.60	4.02	0.12	1.4	**
Th2 > Th1	IL-4R	7.0	2.3	7.90	0.48	6.31	1.01	3.0	n.s.
	FGFR	3.8	1.9	10.10	0.53	9.35	0.18	1.7	**
	CCR4	3.1	3.2	7.37	0.47	5.23	0.18	4.3	**
	STAT4	1.8	1.7	7.07	0.43	6.06	0.55	2.0	**

[†]For calculations, see Materials and methods. [‡]Same sample measured with both methods. [§]Data from Th1/Th2 culture samples representing three individuals were used. According to analysis of variance: n.s., not significant ($p > 0.05$); *significant ($0.01 < p < 0.05$); **highly significant ($p < 0.01$).

expressed genes, first in mouse cells and then often followed by confirmatory discoveries in human T helper cells. These discoveries in human cell systems have included the differential expression of the receptor chains IL-12R β 2 and IFN- γ R β [23,24], chemokine receptors CCR2, CCR4, CCR5 [25,26] and SLAM [20]. Several attempts have been made to delineate a specific Th1/Th2 transcription factor, which would determine the development of different T-cell phenotypes [27]. The role of such transcription factors as c-Maf [28] and GATA-3 [29] in Th2-cell development and T-bet [30] in Th1 development is currently being extensively studied mainly in mouse models. In order to broaden the view on gene expression in human type 1 and type 2 cells we designed a microarray to screen for 250 inflammation-related human genes. With careful experimental considerations we are able to report subtle differences in the expression of 34 genes in polarized human CD4⁺ T helper cells that characterize their phenotypes and participate in their function.

The immune response is mediated by the coordinated expression of growth factors and chemokines. It is characterized by the presence of several secreted mediator molecules (ligands) in variable doses surrounding the cell simultaneously. Therefore, both the type and the quantity of the ligands surrounding the cell, as well as the receptors on the cell surface, dictate the early events in a signaling cascade. Microarray profiles of human type 1 and type 2 cells reveal a large number of molecules known to potentiate inflammation

responses. In addition to the proinflammatory molecules IFN- γ and TNF- α , the transcripts for GM-CSF, IL-8, MIP-1 α , MIP-1 β and RANTES were also preferentially expressed in Th1 cells as compared to Th2 cells. GM-CSF is often produced together with the proinflammatory molecules TNF- α and IFN- γ and its production by activated human T cells has been reported [1,31]. The expression profile of the type 2 cells has somewhat fewer inflammatory mediators than Th1 cells. In addition to interleukines IL-4, IL-5 and IL-13, only MIF mRNA was found to be expressed at a higher level in Th2 cells than in Th1 cells. The simultaneous expression of the classical Th2 cytokines IL-4, IL-5 and IL-13 is not surprising however, as it was recently discovered that they are all transcribed from a single gene cluster as a result of coordinated regulation [32,33]. Increased expression of MIF has been observed in activated mouse Th2 cell clones [34].

Chemokines IL-8, MIP-1 α , MIP-1 β and RANTES are known to recruit neutrophils, eosinophils, macrophages and other lymphocytes to the site of inflammation [35]. MIP-1 α , MIP-1 β and RANTES have previously been shown to be direct chemoattractants of Th1 cells [36,37]. These chemokines are promiscuously used by several receptors, but all three are ligands for CCR5, which has been shown to be preferably expressed in type 1 cells [25,38,39]. Therefore, it is of interest to note that the CC-chemokines and their receptors are coexpressed in Th1 cells. The coexpression is not only Th1 cell specific, but the similar expression pattern

applies to Th2 cells that express CCR4 (this study, and [15,25,26]) and its ligand TARC [15] to which Th2 cells are also responsive [39]. Because the gene expression profiles in this study reflect pure cell cultures of effector cell populations, it seems plausible that these chemotactic signals are actively used within the type 1 or type 2 cells.

The comparison of the expression profiles of polarized T helper cells also reveals several differences in the expression of signaling receptor chains. Differences in expression of IL-12R β 2 [23], IFN- γ R β [24] and SLAM [20] in human type 1 and type 2 cells have been previously described. T cells differed also in their expression for IL-7R, IFN- α β R and FGFR. Signaling through all three receptors is poorly characterized. Both IL-7 and type I interferons increase the survival of activated T cells [40,41]. Interestingly, it was recently shown that a specific effector dendritic cell (DC2) that induces primarily type 2 differentiation in T cells is the natural source of type 1 interferon in human blood [42]. Our results indicate that at least the receptor chain IFN- α β R of the IFN type I interferons is preferentially expressed in Th2 cells and in lesser amounts in Th1 cells.

At present it is not known why the basal level of transcription of STAT4 appears to be higher in Th2 cells compared to Th1 cells. This is surprising because under the culture conditions used for Th2 cells, the IL-12 is neutralized in the medium with an antibody, whereas in the Th1 cells the STAT4 is strongly stimulated. Also, there is no literature available as to whether the STAT4 expression levels are affected by signals from the type I interferon receptors. Functional receptors for FGF in some human T cells have been reported by Zhao *et al.* [43]. However, our finding that FGFR is preferentially expressed in Th2 cells conflicts with a report by Rogge *et al.* [15], in which a 6.9-fold higher expression in Th1 cells was found. It is possible that the expression of FGFR does vary in the course of T helper cell development, depending on the state of differentiation and cell activation, because the observations in these papers are from different phases of T-cell differentiation.

The oligonucleotide array used here measured the transcripts for relatively few transcription factors and, for example, c-Maf, GATA-3 and T-bet were not included. However, the differential expression of these transcription factors during human T helper cell differentiation has been confirmed in these cells (E. Ylikoski and R. Lund, unpublished observations). In the Th2 effector cell profile the preferential expression of Smad2 is still worth mentioning. The details of transforming growth factor- β (TGF- β) receptor signaling in T cell development, for which the Smads form the signaling cascade to the nucleus, are still poorly characterized. It has been suggested that TGF- β modulates growth and development of T-cell precursors during the early differentiation phase [44]. In addition, more recent papers show that the inhibition of type 2 T-cell differentiation by

TGF- β actually occurs by inhibiting the expression of GATA-3 in mouse [45,46] and that TGF- β regulates the Th2-related airway hyperreactivity in an animal model of asthma [47].

The differential expression of several cell death- and apoptosis-related genes in Th1 and Th2 cells is also evident from the profiles created. Previous studies have indicated that T cells exploit different pathways for cell death [37,38]. Mouse Th1 cells were recently shown to be susceptible to activation-induced cell death (AICD) by Fas ligand, whereas Th2 cells were shown to be resistant to it [48,49]. This resistance was suggested to be characteristic of asthma [48,50]. A distinct panel of cell death-related genes preferentially expressed in Th1 cells includes those for granzyme B, caspase-1 and clusterin, whereas Th2 cells preferably expressed Bax, CAS and caspase-6. The differential expression of clusterin that was also sustained during T-cell activation is also interesting in the light of recent data suggesting that clusterin can act as a secreted chaperone in mammalian cells [51] and protects cells from the cytotoxicity of TNF- α [52]. The preferential expression of TNF- α and clusterin in Th1 cells, together with the differentially expressed I κ B and c-Jun, suggests that in addition to Fas/Fas ligand-mediated death, signaling through the TNF receptor family could also serve as an additional pathway to control cell survival and death in these cells. Systematic studies on cell death in Th1 and Th2 cells are needed to clarify this.

To identify differentially expressed genes for the human T helper subsets that would not be affected by cell activation status, the effect of activation through CD3 and CD28 in human Th1 or Th2 cells was studied. During activation, most of the genes studied responded by downregulating the transcription to a basal or even lower level of transcription. This was also true for the transcripts of GM-CSF, TNF- α and I κ B, which had been rapidly induced and subsequently downregulated back to the basal level. Most importantly, the transitional preference favoring Th1 cells, which have a higher level of expression of GM-CSF, TNF- α , I κ B, CCR1, CCR2, CCR5, caspase-1 and clusterin compared to Th2 cells, did not change during activation, a feature potentially important with clinical samples of mixed cell phenotypes.

In conclusion, microarray technology in combination with quantitative RT-PCR proved to be powerful in profiling the expression of inflammation-related genes in human CD4⁺ Th1 and Th2 cell populations. The expression patterns of type 1 and type 2 cells described here potentially reflect also the transcriptional patterns common in other type 1 and type 2 cells of the immune response [53]. What we have shown here is that differences in the basal levels of gene transcription can be extremely small and yet can indicate essential functional outcomes of that particular cell type. We believe that the results presented here will provide valuable new insights into gene array discovery. In addition, they also

provide a challenge to bioinformaticians to develop accurate algorithms for expression profiling.

Materials and methods

Human CD4⁺ T cells

Human CD4⁺ T helper cells were isolated from neonatal cord blood and cultured in polarizing conditions for 14 days as previously described. Briefly, the stimulation of the cells was carried out by plating 1.0×10^6 T cells per ml and 0.5×10^6 per ml of irradiated (6400 rad) CD32-B7 transfected mouse L fibroblasts [54] and 100 ng/ml PHA (Murex Diagnostics, France) on 24-well flat-bottomed plates (Lindbro, ICN Biomedicals). Cells were grown in Yssel's medium supplemented with 1% of human AB serum (Gemini Bioproducts) and 100 U/ml human recombinant IL-2 (Cellular Products). Cell differentiation was primed with either 2.5 ng/ml human recombinant IL-12 (R & D Systems) or with 10 ng/ml human recombinant IL-4 (R & D Systems,) for Th1 or Th2 cells, respectively. Culture media for Th2 cells contained 10 μ g/ml human recombinant α -IL-12 (R & D Systems). The cells were fed every other day and split at day 3-4 after stimulation. At day 7, cells were restimulated and cultured as described above for another 7 days. When cells were harvested for RNA isolation, an additional immunomagnetic purification step of CD4⁺ cells was performed (DYNABEADS M-450 CD4, DYNAL A.S). The protocol for activating the cells through CD3/CD28 for 2, 6 and 24 h has been previously described [55]. Monoclonal antibodies for CD3 and CD28 were from Immunotech and goat F(ab')₂ antibody from BioSource International. The phenotype of the cells was defined by measuring secreted IFN- γ and IL-4, corresponding to type 1 and type 2 cells, respectively, by ELISA (Genzyme).

RNA sample preparation and hybridization

Poly(A)⁺ mRNA (1 μ g) was isolated by two rounds of Oligotex purification (Qiagen). cDNA synthesis of mRNA, *in vitro* transcription and the production of a biotin-labeled RNA probe for oligonucleotide arrays were performed as previously described [8]. The hybridization mixture contained 12.5 μ g of the fragmented sample cRNA together with defined amounts of control *Escherichia coli* and bacteriophage P1 gene cRNAs, produced for spiking purposes in an identical manner as the sample, and a biotin labeled control oligo (5'-biotin-GTCAAGATCGTACCGTTTCAG-3') from Genset. Subsequent hybridization and washing steps were done as described by Wodicka *et al.* [17]. Chips were washed with GeneChip WashB program on a fluidics station and scanned with the GeneChip scan program (Affymetrix).

Oligonucleotide microarray data analysis

The principles of quantitative analysis of hybridization intensities have previously been described [16,17,56]. Data analysis was carried out under low or high stringency using the GeneChip software program. Low-stringency parameter

values were: difference threshold (DT) = 20; ratio threshold (RT) = 1.5; change threshold (CT) = 15, and percent change threshold (PCT) = 30. High-stringency parameter values were: DT = 50; RT = 1.5; CT = 50 and PCT = 30. Comparative data were obtained by analyzing results from IL-4-treated RNA samples (Th2 cells) compared to those from IL-12-treated RNA samples (Th1 cells) as baseline. All intensity values below 20 were assigned a threshold value 20. The data reported here is from the more rigorous high stringency analysis (cutoff range ≥ 1.5 -fold). The GenBank accession numbers of the sequences used for the initial microarray design and reported here are as presented in Table 1 and as follows: BAX (U19599); c-Jun (JO4111); granzyme B (M28879); IFN- $\alpha\beta$ R (X77722); IFN- γ (X13274); IFN- γ R β (U05875); IL-4 (M13982); IL-5 (X04688); IL-8 (M17017); IL-13 (L06801); IL-7R (M29696); IL-12R β 2 (U64198); IRF-1 (X14454); MIF (M25639); MIP-1 α (M23452); MIP-1 β (JO4130); SLAM (U33017) and Smad2 (U59911).

Real-time quantitative RT-PCR

The principles of the real-time RT-PCR detection with hydrolysis probes (TaqMan) have been previously described [19,57]. TaqMan probes and primers for the quantitative detection of target mRNAs were designed (Table 1) by using Primer Express computer software (Applied Biosystems). The real-time measurements were carried out with the ABI PRISM 7700 SDS instrument (Applied Biosystems) as described in Hamalainen *et al.* [20]. Samples were analyzed in duplicate in three independent runs. Statistical significance was determined by ANOVA model. The C_T value is defined as the cycle number in which the detected fluorescence exceeds the threshold value [19,57]. In all experiments the threshold value used to determine C_T during analysis was kept constant. For each probe and primer pair the linearity of detection was confirmed to have a correlation coefficient of at least 0.98 over the detection area by measuring a fourfold dilution curve with cDNA isolated from T helper cells. Fold difference in expression in Th1 and Th2 cells was therefore calculated assuming 100% efficient PCR where each C_T was normalized to GAPDH:

$$(1) \text{ Fold difference} = \frac{2^{|C_{T1}(\text{target}) - C_{T1}(\text{GAPDH})|}}{2^{|C_{T2}(\text{target}) - C_{T2}(\text{GAPDH})|}}$$

$$(2) \text{ Fold difference} = \frac{2^{|\Delta C_{T1}|}}{2^{|\Delta C_{T2}|}}$$

Where C_{T1}(target) and C_{T2}(target) represent the C_T values for the target gene of Th1 and Th2 samples, respectively. C_{T1}(GAPDH) and C_{T2}(GAPDH) represent the C_T values for the GAPDH gene of the Th1 and Th2 samples, respectively.

Acknowledgements

We thank Tuija Kyrola and Paula Suominen for technical assistance and Nina Johansson for critical review of the manuscript. This study was sup-

ported by grants from the Academy of Finland, National Technology Agency (TEKES) and Turku University Hospital Fund.

References

- Romagnani S: **Lymphokine production by human T cells in disease states.** *Annu Rev Immunol* 1994, **12**:227-257.
- Seder RA, Paul WE: **Acquisition of lymphokine-producing phenotype by CD4⁺ T cells.** *Annu Rev Immunol* 1994, **12**:635-673.
- Abbas AK, Murphy KM, Sher A: **Functional diversity of helper T lymphocytes.** *Nature* 1996, **383**:787-793.
- Marshall A, Hodgson J: **DNA chips: an array of possibilities.** *Nat Biotechnol* 1998, **16**:27-31.
- Duggan DJ, Bittner M, Chen Y, Meltzer P, Trent JM: **Expression profiling using cDNA microarrays.** *Nat Genet* 1999, **21**:10-14.
- Der SD, Zhou A, Williams BR, Silverman RH: **Identification of genes differentially regulated by interferon alpha, beta, or gamma using oligonucleotide arrays.** *Proc Natl Acad Sci USA* 1998, **95**:15623-15628.
- Iwata M, Vieira J, Byrne M, Horton H, Torok-Storb B: **Interleukin-1 (IL-1) inhibits growth of cytomegalovirus in human marrow stromal cells: inhibition is reversed upon removal of IL-1.** *Blood* 1999, **94**:572-578.
- Zhu H, Cong JP, Mamtora G, Gingeras T, Shenk T: **Cellular gene expression altered by human cytomegalovirus: global monitoring with oligonucleotide arrays.** *Proc Natl Acad Sci USA* 1998, **95**:14470-14475.
- Geiss GK, Bumgarner RE, An MC, Agy MB, van't Wout AB, Hamersmark E, Carter VS, Upchurch D, Mullins JI, Katze MG: **Large-scale monitoring of host cell gene expression during HIV-1 infection using cDNA microarrays.** *Virology* 2000, **266**:8-16.
- Heller RA, Schena M, Chai A, Shalon D, Bedilion T, Gilmore J, Woolley DE, Davis RW: **Discovery and analysis of inflammatory disease-related genes using cDNA microarrays.** *Proc Natl Acad Sci USA* 1997, **94**:2150-2155.
- Kaminski N, Allard JD, Pittet JF, Zuo F, Griffiths MJ, Morris D, Huang X, Sheppar D, Heller RA: **Global analysis of gene expression in pulmonary fibrosis reveals distinct programs regulating lung inflammation and fibrosis.** *Proc Natl Acad Sci USA* 2000, **97**:1778-1783.
- Amundson SA, Bittner M, Chen Y, Trent J, Meltzer P, Fornace AJ, Jr: **Fluorescent cDNA microarray hybridization reveals complexity and heterogeneity of cellular genotoxic stress responses.** *Oncogene* 1999, **18**:3666-3672.
- Golub TR, Slonim DK, Tamayo P, Huard C, Gaasenbeek M, Mesirov JP, Coller H, Loh ML, Downing JR, Caligiuri MA, et al.: **Molecular classification of cancer: class discovery and class prediction by gene expression monitoring.** *Science* 1999, **286**:531-537.
- Alizadeh AA, Eisen MB, Davis RE, Ma C, Lossos IS, Rosenwald A, Boldrick JC, Sabet H, Tran T, Yu X, et al.: **Distinct types of diffuse large B-cell lymphoma identified by gene expression profiling.** *Nature* 2000, **403**:503-511.
- Rogge L, Bianchi E, Biffi M, Bono E, Chang SY, Alexander H, Santini C, Ferrari G, Sinigaglia L, Seiler M, et al.: **Transcript imaging of the development of human T helper cells using oligonucleotide arrays.** *Nat Genet* 2000, **25**:96-101.
- Lockhart DJ, Dong H, Byrne MC, Follettie MT, Gallo MV, Chee MS, Mittmann M, Wang C, Kobayashi M, Horton H, Brown, EL: **Expression monitoring by hybridization to high-density oligonucleotide arrays.** *Nat Biotechnol* 1996, **14**:1675-1680.
- Wodicka L, Dong H, Mittmann M, Ho MH, Lockhart DJ: **Genome-wide expression monitoring in *Saccharomyces cerevisiae*.** *Nat Biotechnol* 1997, **15**:1359-1367.
- Heller RA, Allard J, Zuo F, Lock C, Wilson S, Klonowski P, Gmuender H, van Hart H, Booth R: **Gene chips and microarrays: applications in disease profiles, drug target discovery, drug action and toxicity.** In *Secondary Gene Chips and Microarrays: Applications in Disease Profiles, Drug Target Discovery, Drug Action and Toxicity*. Edited by Schena M. New York: Oxford University Press, 1999, 187-202.
- Heid CA, Stevens J, Livak KJ, Williams PM: **Real time quantitative PCR.** *Genome Res* 1996, **6**:986-994.
- Hamalainen H, Meissner S, Laheesmaa R: **Signaling lymphocytic activation molecule (SLAM) is differentially expressed in human Th1 and Th2 cells.** *J Immunol Methods* 2000, **242**:9-19.
- Teague TK, Hildeman D, Kedl RM, Mitchell T, Rees W, Schaefer BC, Bender J, Kappler J, Marrack P: **Activation changes the spectrum but not the diversity of genes expressed by T cells.** *Proc Natl Acad Sci USA* 1999, **96**:12691-12696.
- Shanafelt MC, Yssel H, Soderberg C, Steinman L, Adelman DC, Peltz G, Laheesmaa R: **CD45 isoforms on human CD4⁺ T-cell subsets.** *J Allergy Clin Immunol* 1996, **98**:433-440.
- Rogge L, Barberis-Maino L, Biffi M, Passini N, Presky DH, Gubler U, Sinigaglia F: **Selective expression of an interleukin-12 receptor component by human T helper 1 cells.** *J Exp Med* 1997, **185**:825-831.
- Groux H, Sornasse T, Cottrez F, de Vries JE, Coffman RL, Roncarolo MG, Yssel H: **Induction of human T helper cell type 1 differentiation results in loss of IFN-gamma receptor beta-chain expression.** *J Immunol* 1997, **158**:5627-5631.
- Bonecchi R, Bianchi G, Bordignon PP, D'Ambrosio D, Lang R, Borsatti A, Sozzani S, Allavena P, Gray PA, Mantovani A, Sinigaglia F: **Differential expression of chemokine receptors and chemotactic responsiveness of type 1 T helper cells (Th1s) and Th2s.** *J Exp Med* 1998, **187**:129-134.
- Imai T, Nagira M, Takagi S, Kakizaki M, Nishimura M, Wang J, Gray PW, Matsushima K, Yoshie O: **Selective recruitment of CCR4-bearing Th2 cells toward antigen-presenting cells by the CC chemokines thymus and activation-regulated chemokine and macrophage-derived chemokine.** *Int Immunol* 1999, **11**:81-88.
- Glimcher LH, Murphy KM: **Lineage commitment in the immune system: the T helper lymphocyte grows up.** *Genes Dev* 2000, **14**:1693-1711.
- Ho IC, Hodge MR, Rooney JW, Glimcher LH: **The proto-oncogene c-maf is responsible for tissue-specific expression of interleukin-4.** *Cell* 1996, **85**:973-983.
- Zheng W, Flavell RA: **The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells.** *Cell* 1997, **89**:587-596.
- Szabo SJ, Kim ST, Costa GL, Zhang X, Fathman CG, Glimcher LH: **A novel transcription factor, T-bet, directs Th1 lineage commitment.** *Cell* 2000, **100**:655-669.
- Romagnani S: **Biology of human TH1 and TH2 cells.** *J Clin Immunol* 1995, **15**:121-129.
- Agarwal S, Rao A: **Modulation of chromatin structure regulates cytokine gene expression during T cell differentiation.** *Immunity* 1998, **9**:765-775.
- Kelly BL, Locksley RM: **Coordinate regulation of the IL-4, IL-13, and IL-5 cytokine cluster in Th2 clones revealed by allelic expression patterns.** *J Immunol* 2000, **165**:2982-2986.
- Bacher M, Metz CN, Calandra T, Mayer K, Chesney J, Lohoff M, Gemsa D, Donnelly T, Bucala R: **An essential regulatory role for macrophage migration inhibitory factor in T-cell activation.** *Proc Natl Acad Sci USA* 1996, **93**:7849-7854.
- Baggiolini M, Dewald B, Moser B: **Human chemokines: an update.** *Annu Rev Immunol* 1997, **15**:675-705.
- Schrum S, Probst P, Fleischer B, Zipfel PF: **Synthesis of the CC-chemokines MIP-1alpha, MIP-1beta, and RANTES is associated with a type 1 immune response.** *J Immunol* 1996, **157**:3598-3604.
- Siveke JT, Hamann A: **T helper 1 and T helper 2 cells respond differentially to chemokines.** *J Immunol* 1998, **160**:550-554.
- Loetscher P, Uguccioni M, Bordoli L, Baggiolini M, Moser B, Chizzolini C, Dayer JM: **CCR5 is characteristic of Th1 lymphocytes.** *Nature* 1998, **391**:344-345.
- Sallusto F, Lenig D, Mackay CR, Lanzavecchia A: **Flexible programs of chemokine receptor expression on human polarized T helper 1 and 2 lymphocytes.** *J Exp Med* 1998, **187**:875-883.
- Soares MV, Borthwick NJ, Maini MK, Janossy G, Salmon M, Akbar AN: **IL-7-dependent extrathymic expansion of CD45RA⁺ T cells enables preservation of a naive repertoire.** *J Immunol* 1998, **161**:5909-5917.
- Marrack P, Kappler J, Mitchell T: **Type 1 interferons keep activated T cells alive.** *J Exp Med* 1999, **189**:521-530.
- Siegal FP, Kadowaki N, Shodell M, Fitzgerald-Bocarsly PA, Shah K, Ho S, Antonenko S, Liu YJ: **The nature of the principal type 1 interferon-producing cells in human blood.** *Science* 1999, **284**:1835-1837.
- Zhao XM, Byrd VM, McKeehan WL, Reich MB, Miller GG, Thomas JW: **Costimulation of human CD4⁺ T cells by fibroblast growth factor-1 (acidic fibroblast growth factor).** *J Immunol* 1995, **155**:3904-3911.

44. Swain SL, Huston G, Tonkonogy S, Weinberg A: **Transforming growth factor-beta and IL-4 cause helper T cell precursors to develop into distinct effector helper cells that differ in lymphokine secretion pattern and cell surface phenotype.** *J Immunol* 1991, **147**:2991-3000.
45. Heath VL, Murphy EE, Crain C, Tomlinson MG, O'Garra A: **TGF-beta1 down-regulates Th2 development and results in decreased IL-4- induced STAT6 activation and GATA-3 expression.** *Eur J Immunol* 2000, **30**:2639-2649.
46. Gorelik L, Fields PE, Flavell RA: **Cutting edge: TGF-beta inhibits Th type 2 development through inhibition of GATA-3 expression.** *J Immunol* 2000, **165**:4773-4777.
47. Hansen G, McIntire JJ, Yeung VP, Berry G, Thorbecke GJ, Chen L, DeKruyff RH, Umetsu DT: **CD4(+) T helper cells engineered to produce latent TGF-beta1 reverse allergen-induced airway hyperreactivity and inflammation.** *J Clin Invest* 2000, **105**:61-70.
48. Zhang X, Brunner T, Carter L, Dutton RW, Rogers P, Bradley L, Sato T, Reed JC, Green D, Swain SL: **Unequal death in T helper cell (Th)1 and Th2 effectors: Th1, but not Th2, effectors undergo rapid Fas/FasL-mediated apoptosis.** *J Exp Med* 1997, **185**:1837-1849.
49. Varadhachary AS, Perdow SN, Hu C, Ramanarayanan M, Salgame P: **Differential ability of T cell subsets to undergo activation-induced cell death.** *Proc Natl Acad Sci USA* 1997, **94**:5778-5783.
50. Jayaraman S, Castro M, O'Sullivan M, Bragdon MJ, Holtzman MJ: **Resistance to Fas-mediated T cell apoptosis in asthma.** *J Immunol* 1999, **162**:1717-1722.
51. Humphreys DT, Carver JA, Easterbrook-Smith SB, Wilson MR: **Clusterin has chaperone-like activity similar to that of small heat shock proteins.** *J Biol Chem* 1999, **274**:6875-6881.
52. Humphreys D, Hochgrebe TT, Easterbrook-Smith SB, Tenniswood MP, Wilson MR: **Effects of clusterin overexpression on TNF-alpha- and TGFbeta-mediated death of L929 cells.** *Biochemistry* 1997, **36**:15233-15243.
53. Harris DP, Haynes L, Sayles PC, Duso DK, Eaton SM, Lepak NM, Johnson LL, Swain SL, Lund FE: **Reciprocal regulation of polarized cytokine production by effector B and T cells.** *Nat Immunol* 2000, **1**:475-482.
54. Yang LP, Byun DG, Demeure CE, Vezzio N, Delespesse G: **Default development of cloned human naive CD4 T cells into interleukin-4- and interleukin-5- producing effector cells.** *Eur J Immunol* 1995, **25**:3517-3520.
55. Laheesmaa R, Allsup A, Soderberg C, Jackman J, Findell P, Peltz G: **Modulation of the Grb2-associated protein complex in human CD4+ T cells by receptor activation.** *J Immunol* 1995, **155**:3815-3822.
56. Lipshutz RJ, Fodor SP, Gingeras TR, Lockhart DJ: **High density synthetic oligonucleotide arrays.** *Nat Genet* 1999, **21**:20-24.
57. Gibson UE, Heid CA, Williams PM: **A novel method for real time quantitative RT-PCR.** *Genome Res* 1996, **6**:995-1001.