

CORRECTION

Correction: Non-Invasive Delivery of dsRNA into De-Waxed Tick Eggs by Electroporation

Newton Ruiz, Leonardo Araujo de Abreu, Luís Fernando Parizi, Tae Kwon Kim, Albert Mulenga, Gloria Regina Cardoso Braz, Itabajara da Silva Vaz, Jr., Carlos Logullo

Fig 5, "Silencing of AKT and GSK changes glycogen content R. microplus eggs," is incorrect. Please see the correct Fig 5 and its caption here.



G OPEN ACCESS

Citation: Ruiz N, de Abreu LA, Parizi LF, Kim TK, Mulenga A, Cardoso Braz GR, et al. (2015) Correction: Non-Invasive Delivery of dsRNA into De-Waxed Tick Eggs by Electroporation. PLoS ONE 10(7): e0133949. doi:10.1371/journal.pone.0133949

Published: July 24, 2015

Copyright: © 2015 Ruiz et al. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.






Fig 5. Silencing of AKT and GSK changes glycogen content R. microplus eggs. Glycogen content was determined in egg homogenates obtained 7 days after electroporation with AKT (A) or GSK (B) dsRNA and compared with eggs treated with β Gal dsRNA. Statistical analysis was carried out using the Student t test (p<0.05), (Triplicate; n = 3).

doi:10.1371/journal.pone.0133949.g001

Reference

 Ruiz N, Abreu LAd, Parizi LF, Kim TK, Mulenga A, Braz GRC, et al. (2015) Non-Invasive Delivery of dsRNA into De-Waxed Tick Eggs by Electroporation. PLoS ONE 10(6): e0130008. doi: 10.1371/journal.pone.0130008 PMID: 26091260