

Review

## Extracellular matrix: Forum introduction

Daniel D Carson\*

Address: Forum coordinator, Department of Biological Sciences, University of Delaware, Newark, DE 19716, USA

Email: Daniel D Carson\* - dcarson@udel.edu

\* Corresponding author

Published: 07 January 2004

*Reproductive Biology and Endocrinology* 2004, 2:1This article is available from: <http://www.rbej.com/content/2/1/1>

Received: 29 July 2003

Accepted: 07 January 2004

© 2004 Carson; licensee BioMed Central Ltd. This is an Open Access article: verbatim copying and redistribution of this article are permitted in all media for any purpose, provided this notice is preserved along with the article's original URL.

The extracellular matrix (ECM) is now appreciated as a key regulator of cell and tissue behavior [1,2]. Support for this notion has been derived from correlative studies demonstrating large changes in the composition and distribution of extracellular matrix components during tissue development, differentiation and in response to hormone/growth factor/cytokine influences. In addition, a variety of functional studies demonstrate ECM-dependent cellular responses *in vitro* while genetic approaches clearly demonstrate critical requirements for genes encoding ECM components or modifiers of ECM, e.g., metalloproteases, in a wide variety of processes *in vivo* [3,4].

ECM produces these effects in multiple ways. One of these is via direct interactions with cell surface ECM receptors of various types, including integrins and syndecans. As a result of these interactions, cytoskeletal elements are reorganized which, alone, can strongly influence responses such as apoptosis [5]. In addition, interactions with these receptors activate a complex array of intracellular signal transduction cascades that modulate a variety of cellular functions including secretory activities and gene transcription. In this regard, the ECM must be considered to act in a continuum with the cell surface, cytoskeleton and, ultimately, the nucleus. ECM also serves as a reservoir of growth factors and cytokines. These key molecules may bind to either polysaccharide or protein constituents of ECM. In some cases, biochemical neutralizers of growth factor activities, e.g., noggin [6], bind strongly to ECM. These interactions further help limit the range of activities of growth factors and cytokines.

In most cases, even individual ECM components are, in fact, complexes of non-identical proteins, e.g., laminin, collagens. These complexes normally function in the con-

text of much larger macromolecular complexes including many other ECM components. Therefore, it is not surprising that the composition of ECM is usually quite dynamic. This not only reflects differential expression of genes encoding ECM components, but also various post-transcriptional processes. In the cases of proteoglycans, structural nuances of the glycosaminoglycan chains are often critical for protein binding activities. Thus, differences in the expression or activities of glycosyltransferases and sulfotransferases alter the function of proteoglycans. Moreover, secreted polysaccharide-degrading enzymes may modify these same polysaccharides after assembly into the ECM and can release bound growth factors/cytokines. In other examples, different isoforms of ECM polypeptide constituents are generated either by alternative mRNA splicing or differential expression of discrete genes encoding similar proteins. Additional variety or control of function is provided by the action of secreted or cell surface-associated metalloproteases that hydrolyze specific ECM components at specific sites [3]. These activities can have several consequences. New sites may be opened for the addition of newly synthesized ECM components, the ECM may be loosened allowing for new cell growth/proliferation or migration, or growth factors/cytokines may be released. Similar proteolytic enzymes also can act at the cell surface to trigger release of ectodomains of ECM, receptors, growth factor receptors or even growth factors themselves [4].

Finally, given that most cellular responses to ECM require engagement of cell surface receptors, factors that interfere with cell surface receptor interactions, e.g., mucins, also play key roles in regulating these events. Mucins also play complex roles since they not only limit certain cell-cell, cell-ECM interactions, but also can bind to other cell sur-

face receptors, e.g., selectins, or, in the case of transmembrane mucins, interact directly with intracellular signal transducers. As is the case for cell surface receptors, transmembrane mucins may be released by the action of cell surface proteases allowing for rapid modulation of mucin-dependent events.

In this miniseries, authors will consider the role of ECM in reproductive processes and cancer. A select group of topics has been chosen that, in some cases, will broadly consider the role of ECM in a specific biological context, i.e., prostate and prostate cancer. Other contributions will focus on the functions of specific ECM components or ECM receptors in particular contexts. These include the roles of syndecans in mammary epithelia, the functions of integrins in embryo implantation and the role of mucins in early embryonic development and certain cancers. Collectively, the reader should gain a perspective on the complex regulation of ECM expression and function in important, representative reproductive and endocrine systems.

## References

1. Schmeichel KL, Weaver VM, Bissell M: **Structural cues from the tissue microenvironment are essential determinants of the human mammary epithelial cell phenotype.** *J Mammary Gland Biol Neoplasia* 1998, **3**:201-213.
2. Hood JD, Cheresh DA: **Role of integrins in cell invasion and migration.** *Nat Rev Cancer* 2002, **2**:91-100.
3. Lynch CC, Matrisian LM: **Matrix metalloproteinases in tumor-host cell communication.** *Differentiation* 2002, **70**:561-573.
4. Blobel CP: **Functional and biochemical characterization of ADAMs and their predicted role in protein ectodomain shedding.** *Inflamm Res* 2002, **51**:83-84.
5. Huang S, Ingber DE: **The structural and mechanical complexity of cell-growth control.** *Nat Cell Biol* 1999, **1**:E131-E138.
6. Paine-Saunders S, Viviano BL, Economides AN, Saunders S: **Heparan sulfate proteoglycans retain Noggin at the cell surface: a potential mechanism for shaping bone morphogenetic protein gradients.** *J Biol Chem* 2002, **277**:2089-2096.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

